Published in final edited form as:

Org Biomol Chem. 2018 February 14; 16(7): 1073–1078. doi:10.1039/c7ob02931d.

Ribosomal incorporation of backbone modified amino acids via an editing-deficient aminoacyl-tRNA synthetase

Emil S. Iqbal^{1,2}, Kara K. Dods^{1,2}, and Matthew C.T. Hartman^{1,2,*}

¹Department of Chemistry, Virginia Commonwealth University (VCU), 1001 West Main Street, P.O. Box 842006, Richmond, Virginia 23284, United States

²Massey Cancer Center, Virginia Commonwealth University, 401 College Street, Richmond, Virginia 23298, United States

Abstract

The ability to incorporate non-canonical amino acids (ncAA) using translation offers researchers the ability to extend the functionality of proteins and peptides for many applications including synthetic biology, biophysical and structural studies, and discovery of novel ligands. Here we describe the high promiscuity of an editing-deficient valine-tRNA synthetase (ValRS T222P). Using this enzyme, we demonstrate ribosomal translation of 11 ncAAs including those with novel side chains, α,α -disubstitutions, and cyclic β -amino acids.

Graphical abstract

A promiscuous aminoacyl-tRNA synthetase charges diverse amino acids onto tRNAs, and enables their incorporation into peptides using in vitro transation.

Introduction

Life has evolved to utilize the chemistry of 20 proteinogenic amino acids (AAs) encoded by 61 codons. This genetic code, while highly conserved and safeguarded at multiple checkpoints, constrains biology to a fraction of the available chemical space. By manipulating individual translation components *in vitro* and *in vivo*, researchers have successfully created ribosomally translated peptides and proteins containing a few hundred ncAAs.¹⁻¹¹

Sense codon reassignment^{1,2,4,12} is a particularly powerful strategy to introduce ncAAs when it is coupled with a reconstituted translation system, such as the Protein Synthesis

* To whom correspondence should be addressed: mchartman@vcu.edu.

Using Recombinant Elements (PURE) translation system.¹³ In this system, a wild type AA and aminoacyl-tRNA synthetase (AARS) pair can be withheld and substituted with a tRNA charged with an ncAA to allow for incorporation of the ncAA in response to vacated codons. To prepare this ncAA-tRNA, researchers have turned to a variety of methods. The most general strategy for the synthesis of ncAA-tRNAs is the Flexizyme system developed by Suga and coworkers.^{14,15} This method utilizes activated amino acid esters and a ribozyme (Flexizyme) to catalyze the charging of an ncAA onto an acceptor tRNA. The Flexizyme system has allowed for the incorporation of a wide range of ncAAs^{15–17}, yet this technique does have some drawbacks. Flexizyme recognizes the invariable 3' end of a tRNA, and therefore cannot selectively charge a specific tRNA within a mixture. Since purified tRNAs are not readily available, *in vitro* transcribed tRNAs, which may not be as efficient as their fully modified counterparts are required.^{18–20} The Flexizyme system also requires the synthesis of activated amino acids.

To address these limitations, one could potentially use a promiscuous AARS^{5,21,22} in order to efficiently charge tRNAs with ncAAs. This strategy would have the advantage of using standard, non-activated amino acids and would be compatible with the readily available, wild-type *E. coli* tRNA mixture. Herein, we have focused our efforts on applying this strategy to *E. coli* ValRS. The wild-type ValRS has been shown to charge a moderate number of ncAAs so we conjectured that deleting the editing activity would broaden the substrate scope.^{23–25} For example, Döring et al. showed that an editing deficient ValRS T222P in *E. coli* resulted in global misincorporations of threonine, cysteine, and 2-aminobutyric acid.²⁶ We reasoned that ValRS T222P could potentially charge additional structurally similar amino acids to further expand the genetic code.

Results/Discussion

Based on this notion, we screened 29 ncAAs (Figure S1) for their ability to be charged by ValRS T222P using a MALDI charging assay.²⁷ This assay involves appending a triphenylphosphonium derivative onto the primary amino group of an AA-tRNA to enhance MALDI-TOF visualization (Figure S2). Eleven successful analogs (Figure 1 and Table S1, and Figure S3) were charged by ValRS T222P. The first group of ncAAs included L-amino acids containing side chain modifications: *trans* alkene (crotyl glycine, **CrG**), azide (Azidohomoalanine, **Aha**), nitrile (β -cyano Alanine, **BCA**), as well as L-*allo*-Threonine (**A-Thr**), Norleucine (**Nle**), and aminobutyric acid (**Abu**). Side chain analogs like these are a rich source of functional diversity and are often employed in medicinal chemistry efforts to optimize peptide activity or for use as chemical biology tools²⁸. The azide functional group, in particular is a useful handle for post-translational modifications using bioorthogonal chemistry.²⁹

ValRS T222P was also able to charge two α,α -disubstituted amino acids containing an α -methyl group: α -methyl Alanine (2-aminoisobutyric acid, **Aib**), and α -methyl Serine (**α Ser**). While some of the above amino acids are near structural analogs to known ValRS T222P substrates²⁶, we were surprised to find this enzyme is also able to charge constrained β -amino acids (*1R,2S*) 2-aminocyclohexane carboxylic acid (**β -1**), and (*1S,2S*) 2-aminocyclohexane carboxylic acid (**β -2**) (Figure 1). (*1S,2S*) 2-aminocyclopentane

carboxylic acid (**β -3**) was also charged, but less efficiently as evidenced by the presence of a significant additional peak corresponding to charging of background valine (Figure S4A). We determined that the source of this contaminating valine was the ValRS T222P enzyme itself, and this background charging could be minimized by diluting the enzyme (Figure S5).

Because the ValRS T222P mutation only blocks the distal editing site, each of the aforementioned charged ncAAs must be recognized by the wild-type catalytic site in ValRS. By comparing those ncAAs that were charged to those that were not substrates (Figure S1), some trends can be gathered. ValRS rejected *N*-alkylated amino acids, amino acids that contain longer unbranched side chains, or those that contain polar groups other than on the β -carbon. Considering that **Aib**, **α Ser**, **β -1**, **β -2**, and **β -3** were charged, it was surprising that the α -methyl analog of Valine as well as the β^3 homolog of Valine (Figure S1) were not.

We then sought to compare the catalytic efficiency of our ValRS T222P with flexizyme using the AARS charging assay developed by Wolfson and Uhlenbeck³⁰. We followed the progress of each charging assay over 2 h. Valine and **β -1** reached maximal charging after 5 min and 20 min, respectively (Figure S6). Charging of the corresponding activated AAs using Flexizyme proceeded much more slowly, and only showed a small degree of charging after 2 h, even at 128-fold higher catalyst (Figure S6).

We then sought to determine if the ribosome, Ef-Tu, and translation factors could tolerate these amino acids in mRNA-directed peptide synthesis using the PURE system. Because ValRS T222P is promiscuous towards canonical amino acids, we cannot add it directly to a PURE translation mixture, or scrambled peptides would result. To solve this problem, we supplemented our translation reactions with *E. coli* total tRNA previously charged with the amino acid of interest using ValRS T222P. This “precharged” total tRNA mixture was then added to a PURE translation reaction that lacked ValRS or ValRS T222P but included the other AAs and AARS enzymes (His/HisRS and Met/MetRS) required to read the codons in our mRNA template MYMH₆ (Figure 2A). The fidelity of translation was verified by MALDI-TOF analysis (Figure 2B, S7 and Table S1), and relative translation yields were measured by ³⁵S-Methionine incorporation (Figure 3 and Table S2). Importantly, translations supplemented with total tRNA precharged under conditions that lacked the ncAA gave very low yields (Figure 3, “-ncAA”), demonstrating that background readthrough of the valine codon was very low under these conditions.

We first tested the ncAAs without backbone modifications. Each of these is incorporated with moderate efficiency and high fidelity (Figure 2, 3A, S7, and Table S2). These ncAAs vary widely in side chain functional group diversity. Four have previously been shown to be decent substrates for the translation apparatus, although on different tRNA bodies: **Aha**¹, **CrG**³¹, **Nle**³¹, and **Abu**.²⁶ Two are novel ribosomal substrates: the nitrile containing **BCA**, and **A-Thr** which is an isomer of threonine rarely found in nature.

α -methyl amino acids facilitate helical secondary structures in peptides.³² This property makes them useful residues in peptide stapling techniques aiming to stabilize helical protein-protein interactions with synthetic peptides.³³ Our system incorporated **α Ser** and **Aib** with high fidelity (Figure 2 and S7) and moderate to high efficiency (Figure 3B). Of the two, only

Aib has been shown to be ribosomally incorporated³⁴, and its efficient incorporation required engineering of the translation apparatus.³⁵

Initially described by Gellman and coworkers, **β-1** and **β-2**, like other constrained β-amino acids have been shown to force peptides into constrained helical secondary structures.³⁶ In addition, β amino acids confer proteolytic resistance to peptides.³⁷ Combining these two traits, Checco et al. recently inserted **β-3** into a stapled peptide known to bind to several antiapoptotic Bcl-2 family proteins.³⁸ The result was an α/β peptide hybrid that maintained nanomolar binding affinity towards its target with a serum stability 150 fold greater over the L-α parent peptide. The incorporation of these β amino acids into peptides via *in vitro* translation would allow for the synthesis of conformationally defined, protease stable peptide libraries. While the ribosome has recently been shown to be compatible with β-AAAs, the applicable amino acids to this point have been the simple β³ homologs of the canonical AAs that do not offer the same conformational benefits.^{7,39} Our ability to charge **β-1**, **β-2**, and **β-3** onto tRNA^{Val} using ValRS T222P now offered us the possibility to test these as substrates for the translation apparatus. We were delighted to find that all three were substrates (Figure 2, 3B, S4B and S7), although in two cases (**β-2** and **β-3**) small peaks due to background valine incorporation were also observed.

Encouraged by these results we focused on monitoring incorporation of certain side chain analogs and backbone analogs **αSer**, **Aib**, **β-1**, and **β-2**, in response to other mRNA templates. The side chain analogs were incorporated efficiently into a longer template that required multiple incorporations (Figure S8). The backbone analogs proved more challenging. None of the backbone ncAAs could be incorporated into a template where we added a single Ala codon in between the initiator Met and Val codons, MAVMH₆. This template contains a C-terminal His-tag which prevents incompletely translated peptides from being captured and detected. To visualize if truncation was a problem for these backbone ncAAs, we used a template with an N-terminal tag, MH₆MVEP. The common product in these experiments was the peptide MH₆M (Figure S9), indicating the translation had terminated prior to the ncAA addition. These results with different templates show that the incorporation of the backbone ncAAs is challenging and mRNA-template dependent.

We then wondered if supplementing the translation system with additional tRNA^{Val} could solve this problem. We *in vitro* transcribed tRNA₁^{Val} (Table S3) which is known to recognize GUA, GUG, and GUU Val codons.⁴⁰ As a rigorous test, we created a longer template that required multiple incorporations of a particular ncAA as well as the ability to monitor for truncations, MHHHHHHMVAAAVEP. Using our standard pre-charging strategy and this template, we did not observe full length peptides with dual incorporation of **αSer** and **Aib** (not shown). However, when we supplemented the translation reaction with the *in vitro* transcribed tRNA₁^{Val}, we achieved efficient dual incorporation of **αSer** and **Aib** (Figure 4 and S2). These analogs could also be double incorporated into a more complicated peptide similar to those used in mRNA displayed libraries MYTNPDCFGNPYCGGGHHHHHH (Figure S10).⁴¹ For **β-1** and **β-2**, however, the results were different. Supplementation of the translation with tRNA₁^{Val} precharged with **β-1** or **β-2** did not lead to dual incorporations. In addition, we used a ribosomal mutant reported by Schepartz⁸ that has been shown to improve the incorporation efficiency of β-amino acids.

We could not observe incorporation with this ribosome mutant either. It is therefore likely that further engineering of the translation apparatus^{8,35,42,43} will be required for efficient incorporation of these β -amino acids in all contexts.

Most research with editing deficient AARS enzymes has focused on their fidelity towards canonical amino acids. In this report, we exploit the inherent promiscuity of one of these enzymes to incorporate 11 different ncAAs into peptides, 8 of which have not been reported before as substrates for the translation apparatus. By precharging these ncAAs onto tRNA^{Val} in the context of total tRNA and omitting ValRS T222P from the translation reaction, we prevented loss of fidelity due to co-translational charging of competing proteinogenic AAs. This strategy, unlike Flexizyme, allows for the attachment of an unactivated ncAA onto specific tRNAs within a commercial mix of total tRNA. Moreover, the ability to incorporate novel side and backbone analogs using translation will open access to diverse, yet structurally defined peptide libraries using mRNA-display technologies.^{44–49}

Methods

MALDI-AARS Screening Assay Procedure

Each charging reaction contained 30 mM HEPES-KOH (pH 7.4), 15 mM MgCl₂, 25 mM KCl, 2 mM 2-mercaptoethanol, 6 mM ATP, 0.09 mg/mL BSA (previously dialyzed into ddH₂O), 0.02 units/ μ L PP_iase, 350 μ M deacylated tRNA, 5 μ M ValRS T222P, 10 mM non-canonical amino acid, and ddH₂O to a final volume of 50 μ L. Each reaction was initiated by addition of ValRS T222P to the mixture and incubated for 30 minutes at 37°C. Each reaction was quenched by addition of 0.1 volume of NaOAc (3.0 M, pH 5.2) and tRNA was extracted with unbuffered phenol:chloroform:isoamyl alcohol solution (25:24:1). The aqueous layer was subjected to a second extraction with chloroform, followed by ethanol precipitation (3 vol.) with 0.1 volume NaOAc (3.0 M, pH 5.2). After centrifugation, the pellet was washed twice with 70% cold ethanol and resuspended in 200 mM NaOAc (pH 5.0) for use in AARS screening assay. tRNA for use in screening assay (6.75 μ L) was added to water (3.75 μ L), 30 mM (4-formylphenoxypropyl)triphenylphosphonium bromide in MeOH, 20 mM NaCNBH₃ in 50 mM NaOAc (pH 5.0) and tumbled at 37°C for 2 h. The reaction was quenched with 0.1 volume 4.4 M NH₄OAc pH 5.0 and 3 volumes of ice cold 100% EtOH, pelleted and washed twice with 70% EtOH and twice with 100% EtOH. The resulting pellet was resuspended in 2.25 μ L 200 mM NH₄OAc and digested with 0.25 μ L of 1 U/ μ L of Nuclease P1 (in 200 mM NH₄OAc, pH 5.0). After 20 minutes at room temperature, the reaction was quenched by placing on ice. One microliter was mixed with saturated CHCA (α -cyano-4-hydroxycinnamic acid) matrix in 1:1 MeCN:1% TFA (9 μ L). One microliter of the resulting suspension was spotted on a MALDI plate and analyzed.

tRNA Pre-charging

The MALDI-AARS screening assay procedure was followed up until the first ethanol precipitation step. The precipitated tRNA was then dissolved in 20 μ L 5 mM KOAc (pH 5.5) and stored at –80 °C until use.

In Vitro Translations

Each reaction (50 μ L) contained HEPES-KOH pH=7.6 (50 mM), spermidine (2 mM), potassium acetate (10 mM), magnesium acetate (6 mM), DTT (1 mM), inorganic pyrophosphatase (1 μ M), creatine kinase (40 μ g/mL), nucleotide diphosphate kinase (1.08 μ g/mL), (6R,S)-5,10-formyl-5,6,7,8-tetrahydrofolic acid (100 μ M), myokinase (3 μ g/mL), creatine phosphate (30 mM), ATP (1.5 mM), GTP (1.5 mM), *E. coli* total tRNA (2.4 mg/mL), IF-1 (2.7 μ M), IF-2 (0.4 μ M), IF-3 (1.5 μ M), EF-G (0.52 μ M), EF-TS (8 μ M), EF-Tu (10 μ M), RF-1 (0.3 μ M), RR-F (0.5 μ M), RF-3 (0.17 μ M), ribosomes (1.2 μ M), MTF (0.6 μ M), Methionine (5 μ M for radiolabeled samples, 100 μ M for non-radiolabeled samples), 0.2 μ M 35 S-Methionine (10 μ Ci, radiolabeled samples only) necessary amino acids and necessary AARS enzymes (0.1 μ M – 1.0 μ M)⁵⁰ and mRNA template (1 μ M). With the exception of dual incorporation of backbone modified amino acid assays, total tRNA precharged with a given amino acid (1.5 A₂₆₀ units) was added at the start of translation and again after 30 min. For dual incorporation of backbone modified amino acids, 25 μ M of in vitro transcribed precharged tRNA was added to the reaction. The translations were initiated by addition of the mRNA. After incubation for 1 h at 37°C, the reactions were quenched with 150 μ L TBS (50 mM Tris, 300 mM NaCl pH 8.0) with 5 mM BME. Ni-NTA resin (50 μ L) was added to a 500 μ L centrifugal filter (VWR) along with the quenched translation reaction, and the mixture was tumbled at room temperature. After 1 h, the resin was washed three times with 500 μ L TBS and eluted with 1% TFA (50 μ L). For reactions labeled with 35 S-Met, the yield was determined by scintillation counting of 47 μ L of the elution. The non-radiolabeled reactions were purified and concentrated by Zip-Tip® C₁₈ chromatography. The tips were first wetted with MeCN, followed by 1:1 MeCN:0.1% TFA, then with 0.1% TFA. Then the peptide was loaded onto the tip by pipetting up and down 15 times in the peptide solution. The tip was washed three times with 0.1% TFA, and then eluted with 5 μ L CHCA matrix (α -cyano-4-hydroxycinnamic acid in 1:1 MeCN:0.1% TFA). An aliquot (1 μ L) of the resulting suspension was spotted on a MALDI plate and analyzed.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. Sam Gellman (University of Wisconsin-Madison) and Dr. Justin Murray (Amgen) for the β -amino acids used in this study, and Dr. Alanna Schepartz (Yale University) for providing a pLK35 plasmid encoding the β -ribosomes. We thank Dr. David E. Hacker, Dr. Stacie L. Richardson, Nicolas Abrigo, and Patrick Dupart for critical review of the manuscript. We acknowledge the support of the NIH (R01CA166264) in this work. The MALDI-TOF instrument is part of the Massey Cancer Center Proteomics Resource supported by CCSG grant NCI 5P30CA16059-35.

References

1. Hartman MCT, Josephson K, Lin CW, Szostak JW. PLoS One. 2007; 2:e972. [PubMed: 17912351]
2. Johnson JA, Lu YY, Van Deventer JA, Tirrell DA. Curr. Opin. Chem. Biol. 2010; 14:774–780. [PubMed: 21071259]
3. Young TS, Schultz PG. J. Biol. Chem. 2010; 285:11039–11044. [PubMed: 20147747]
4. Forster AC, Tan Z, Nalam MNL, Lin H, Qu H, Cornish VW, Blacklow SC. Proc. Natl. Acad. Sci. U. S. A. 2003; 100:6353–6357. [PubMed: 12754376]

5. Guo L-T, Wang Y-S, Nakamura A, Eiler D, Kavran JM, Wong M, Kiessling LL, Steitz TA, O'Donoghue P, Söll D. *Proc. Natl. Acad. Sci.* 2014; 111:16724–16729. [PubMed: 25385624]
6. Ring CM, Iqbal ES, Hacker DE, Hartman MCT, Cropp TA. *Org. Biomol. Chem.* 2017; 15:4536–4539. [PubMed: 28517015]
7. Fujino T, Goto Y, Suga H, Murakami H. *J. Am. Chem. Soc.* 2016; 138:1962–1969. [PubMed: 26807980]
8. Melo Czekster C, Robertson WE, Walker AS, Söll D, Schepartz A. *J. Am. Chem. Soc.* 2016; 138:5194–5197. [PubMed: 27086674]
9. Ellman J, Mendel D, Schultz P. *Science* (80-). 1992; 255:197–200.
10. Frankel A, Millward SW, Roberts RW. *Chem. Biol.* 2003; 10:1043–1050. [PubMed: 14652071]
11. Achenbach J, Jahnz M, Bethge L, Paal K, Jung M, Schuster M, Albrecht R, Jarosch F, Nierhaus KH, Klussmann S. *Nucleic Acids Res.* 2015; 43:5687–5698. [PubMed: 26026160]
12. Josephson K, Hartman MCT, Szostak JW. *J. Am. Chem. Soc.* 2005; 127:11727–11735. [PubMed: 16104750]
13. Shimizu Y, Inoue A, Tomari Y, Suzuki T, Yokogawa T, Nishikawa K, Ueda T. *Nat. Biotechnol.* 2001; 19:751–755. [PubMed: 11479568]
14. Goto Y, Katoh T, Suga H. *Nat. Protoc.* 2011; 6:779–790. [PubMed: 21637198]
15. Ohuchi M, Murakami H, Suga H. *Curr. Opin. Chem. Biol.* 2007; 11:537–542. [PubMed: 17884697]
16. Terasaka N, Iwane Y, Geiermann AS, Goto Y, Suga H. *Int. J. Mol. Sci.* 2015; 16:6513–6531. [PubMed: 25803109]
17. Rogers JM, Suga H. *Org. Biomol. Chem.* 2015; 13:9353–9363. [PubMed: 26280393]
18. Saint-Léger A, Ribas De Pouplana L. *Biochimie.* 2015; 114:72–79. [PubMed: 25921436]
19. Agris PF. *Nucleic Acids Res.* 2004; 32:223–238. [PubMed: 14715921]
20. El Yacoubi B, Bailly M, de Crécy-Lagard V. *Annu. Rev. Genet.* 2012; 46:69–95. [PubMed: 22905870]
21. Young DD, Young TS, Jahnz M, Ahmad I, Spraggon G, Schultz PG. *Biochemistry.* 2011; 50:1894–1900. [PubMed: 21280675]
22. Brustad E, Bushey ML, Brock A, Chittuluru J, Schultz PG. *Bioorganic Med. Chem. Lett.* 2008; 18:6004–6006.
23. Jakubowski H, fersht AR. *Nucleic Acids Res.* 1981; 9:3105–3117. [PubMed: 7024910]
24. Mursinna RS, Lincecum TL, Martinis SA. *Biochemistry.* 2001; 40:5376–5381. [PubMed: 11331000]
25. Beebe K, De Pouplana LR, Schimmel P. *EMBO J.* 2003; 22:668–675. [PubMed: 12554667]
26. Döring V, Mootz HD, Nangle LA, Hendrickson TL, de Crécy-Lagard V, Schimmel P, Marlière P. *Science.* 2001; 292:501–504. [PubMed: 11313495]
27. Hartman MCT, Josephson K, Szostak JW. *P Natl Acad Sci Usa.* 2006; 103:4356–4361.
28. Stevenazzi A, Marchini M, Sandrone G, Vergani B, Lattanzio M. *Bioorg. Med. Chem. Lett.* 2014; 24:5349–5356. [PubMed: 25455481]
29. Kiick KL, Saxon E, Tirrell DA, Bertozzi CR. *Proc. Natl. Acad. Sci. U. S. A.* 2002; 99:19–24. [PubMed: 11752401]
30. Wolfson AD, Uhlenbeck OC. *Proc. Natl. Acad. Sci.* 2002; 99:5965–5970. [PubMed: 11983895]
31. Van Hest JCM, Kiick KL, Tirrell DA. *J. Am. Chem. Soc.* 2000; 122:1282–1288.
32. Toniolo C, Bonora GM, Barone V, Bavoso A, Benedetti E, Di Blasio B, Grimaldi P, Lelj F, Pavone V, Pedone C. *Macromolecules.* 1985; 18:895–902.
33. Schafmeister CE, Po J, Verdine GL. *J. Am. Chem. Soc.* 2000; 122:5891–6288.
34. Tan Z, Forster AC, Blacklow SC, Cornish VW. *J. Am. Chem. Soc.* 2004; 126:12752–12753. [PubMed: 15469251]
35. Katoh T, Iwane Y, Suga H. *Nucleic Acids Res.*
36. Appella DH, Christianson LA, Klein DA, Powell DR, Huang X, Barchi JJ, Gellman SH. *Nature.* 1997; 387:381–384. [PubMed: 9163422]

37. Hook DF, Bindschädler P, Mahajan YR, Sebesta R, Kast P, Seebach D. *Chem. Biodivers.* 2005; 2:591–632. [PubMed: 17192006]
38. Checco JW, Lee EF, Evangelista M, Sleebs NJ, Rogers K, Pettikiriachchi A, Kershaw NJ, Eddinger GA, Belair DG, Wilson JL, Eller CH, Raines RT, Murphy WL, Smith BJ, Gellman SH, Fairlie WD. *J. Am. Chem. Soc.* 2015; 137:11365–11375. [PubMed: 26317395]
39. Melo Czekster C, Robertson WE, Walker AS, Söll D, Schepartz A. *J. Am. Chem. Soc.* 2016; 138:5194–5197. [PubMed: 27086674]
40. Dong H, Nilsson L, Kurland CG. *J. Mol. Biol.* 1996; 260:649–663. [PubMed: 8709146]
41. Hacker DE, Hoinka J, Iqbal ES, Przytycka TM, Hartman MCT. *ACS Chem. Biol.* 2017; 12:795–804. [PubMed: 28146347]
42. Dedkova LM, Fahmi NE, Paul R, Del Rosario M, Zhang L, Chen S, Feder G, Hecht SM. *Biochemistry.* 2012; 51:401–415. [PubMed: 22145951]
43. Maini R, Nguyen DT, Chen S, Dedkova LM, Chowdhury SR, Alcalá-Torano R, Hecht SM. *Bioorganic Med. Chem.* 2013; 21:1088–1096.
44. Xu L, Aha P, Gu K, Kuimelis RG, Kurz M, Lam T, Lim AC, Liu H, Lohse PA, Sun L, Weng S, Wagner RW, Lipovsek D. *Chem. Biol.* 2002; 9:933–942. [PubMed: 12204693]
45. Schlippe YVG, Hartman MCT, Josephson K, Szostak JW. *J. Am. Chem. Soc.* 2012; 134:10469. [PubMed: 22428867]
46. Seelig B, Szostak JW. *Nature.* 2007; 448:828–831. [PubMed: 17700701]
47. Hacker DE, Hoinka J, Iqbal ES, Przytycka TM, Hartman MCT. *ACS Chem. Biol.* 2017; 12:795–804. [PubMed: 28146347]
48. Passioura T, Suga H. *Chem. Commun.* 2017; 53:1931–1940.
49. Millward SW, Fiacco S, Austin RJ, Roberts RW. *ACS Chem. Biol.* 2007; 2:625–634. [PubMed: 17894440]
50. White ER, Reed TM, Ma Z, Hartman MCT. *Methods.* 2013; 60:70–74. [PubMed: 23718982]

Figure 1. AARS Charging Assay Screen

MALDI-TOF spectra of derivatized ncAA-tRNA species after incubation with ValRS T222P.

A magnified view of the region around each major peak is shown in Figure S3.

Figure 2. MALDI-TOF spectra of *in vitro* translated peptides

A) Sequence of the coding region of the mRNA coding region used in translation experiments. The Val codon GUA is reassigned to the ncAA shown. B) MALDI-TOF spectra of M-ncAA-MH₆ peptides with the designated amino acids. Circles (°) indicate peptides of mass consistent with incorporation of the ncAA. Minor peaks consistent with background incorporation of Valine are highlighted with an arrow. A magnified view of the region around each major peak is shown in Figure S7.

Figure 3.

Translation Efficiency. Translation yields (pmols/50 μ L reaction) as measured by 35 S-Methionine incorporation into Ni-NTA capturable peptides. Yields of side chain modified (A) and backbone modified (B) incorporation. Colors indicate ncAAs of the same class: orange, side chain modified; red, β -ncAA; purple, α -methyl-ncAA; gray: precharged tRNA without ncAA; black, absence of precharged tRNA. Error bars represent the standard deviation of at least 4 experiments.

Figure 4. Dual incorporation of ncAAs on MH₆MVA₃VEP

(A) MALDI-TOF MS demonstrating dual incorporation of α -methyl Serine with 25 μ M

synthetic tRNA^{Val} Exp [M+H]⁺ 1772.75 m/z Obs 1772.45 m/z (B) MALDI-TOF

demonstrating dual incorporation of α -methyl Alanine (Aib) with 25 μ M synthetic tRNA^{Val}

Exp [M+H]⁺ 1740.76 m/z Obs 1740.36 m/z . (C) Incorporation Yields for dual incorporation

of α -methyl Serine and α -methyl Alanine compared to Valine.