Published in final edited form as:

Neurobiol Aging. 2018 August ; 68: 34–47. doi:10.1016/j.neurobiolaging.2018.03.030.

Preferential Consolidation of Emotionally Salient Information During a Nap is Preserved in Middle Age

Sara E. Alger^{a,*}, Elizabeth A. Kensinger^b, and Jessica D. Payne^a

^aUniversity of Notre Dame, Department of Psychology, 118 Haggard Hall, Notre Dame, IN, USA 46556

^bBoston College, Department of Psychology, McGuinn Hall Room 300, 140 Commonwealth Ave., Chestnut Hill, MA 02467

Abstract

Sleep preferentially preserves aspects of memory that are most salient and valuable to remember at the expense of memory for less relevant details. Daytime naps and nocturnal sleep enhance this emotional memory trade-off effect, with memory for emotional components correlated with slow wave sleep (SWS) during the day and rapid eye movement (REM) sleep overnight. However, these studies have primarily sampled from young adult populations. Sleep and memory are altered by middle age, and the aim of the present study was to examine how age affects sleep-based mechanisms of emotional memory prioritization, using a daytime nap protocol to compare young to middle-aged adults. In both age groups, a nap soon after encoding scenes that contained a negative or neutral object on a neutral background led to superior retention of emotional object memory at the expense of memory for the related backgrounds. Sleep spindle activity during SWS was related to memory for this emotionally salient information across the age range.

Keywords

Aging; Emotion; Middle age; Napping; Selective memory; Slow wave sleep

1. Introduction

Our memories are not exact replicas of our experiences. Rather, both intrinsic and extrinsic properties of events determine what gets prioritized and promoted for a place in our memory. For example, salient information, especially if it is relevant for future use, is selectively consolidated, often at the expense of memory for less important information (Stickgold & Walker, 2013). This type of memory trade-off observed when examining memory for complex emotional episodes, where memory for the emotionally salient focus of the episode is preferentially preserved while memory for neutral, contextual detail is forgotten or

*Corresponding Author: Sara E. Alger, 118 Haggard Hall, Notre Dame, IN 46556, sealger78@gmail.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

suppressed (Kensinger et al., 2007). Importantly, the magnitude of the emotional memory trade-off increases over a period of sleep (Payne et al., 2008a; Payne & Kensinger, 2011), demonstrating that this phenomenon is not simply the product of attentional factors during encoding, but to active processes unfolding during sleep (Bennion et al., 2015).

While several studies have substantiated this role for sleep in the selective consolidation of salient information, this research has primarily been conducted in young adults (Payne et al., 2008a; Payne et al., 2012). This limits our understanding of both the scope of the effect and the underlying sleep-based mechanisms. Both sleep quality and episodic memory naturally begin to decline as we approach middle age (e.g. Ohayon, Carskadon, Guilleminault, & Vitiello, 2004; Verhaeghen & Salthouse, 1997). Moreover, the habits we form in middle age, including cognitive and sleep habits, may set the stage for healthy or unhealthy mental, cognitive, and physical health as we age. As such, a major goal of the current study was to investigate emotional memory and sleep in this understudied age group. In order to more fully understand the gap in knowledge this present study aims to address, we first need to review what is known regarding how memory and sleep are altered with age.

The ability to encode, consolidate, and retrieve new episodic experiences declines across the lifespan (Rajah et al., 2015), although most evidence comes from much older adults and very little is known specifically about middle age. Among other brain areas, structural atrophy and functional changes have been revealed in regions heavily implicated in processing episodic memories (Raz et al., 1997; Buckner, 2004; Aizawa et al., 2009), notably the medial temporal lobe (MTL), specifically the hippocampus, and the prefrontal cortex (PFC). These changes begin to happen during the middle years of life (Raz et al., 2010). The profound reduction of cell density, synapses, and overall decreased plasticity in these areas impairs the ability to store new episodic information (Burke & Barnes, 2006). Beyond these structures individually, connectivity between the hippocampus and neocortex, vital for episodic memory consolidation and long-term retention, diminishes with age (Grady et al., 2003; Morcom et al., 2003; Duverne et al., 2009).

However, there is conflicting evidence regarding memory impairments with age as they relate to emotional memory. Emotionally salient information consists of intrinsic characteristics that evoke an emotional reaction and lead to a perception of the valence of the stimuli (e.g., negative, neutral, or positive) and a level of associated physiological arousal (e.g., calming, exciting, agitating). Many studies have demonstrated that emotion processing is maintained with age (Mather, 2006; Kensinger, 2008) and that emotional information continues to be preferentially preserved over neutral information (Kensinger et al., 2002, Waring & Kensinger, 2009). Much of the influence behind why emotional information is remembered better than neutral information involves differences in attentional focus during encoding, with more attentional resources dedicated toward the emotional input (Hamann, 2001). Although such factors trigger an emotional trade-off in memory (Kensinger et al., 2004; 2007), factors including post-stimulus elaboration (Steinmetz & Kensinger, 2013) and consolidation processes (Payne & Kensinger, 2010) modulate this initial effect. Evidence points to the idea that as we age, the emotional memory trade-off becomes more pervasive, perhaps due to the reduced ability to shift attention (Hasher, 2007), or a greater focus on the emotional center of the stimuli (Mather & Carstensen, 2003). While it appears that this type

of nuanced selective memory for emotionally salient information persists with age, what is unknown is the role of sleep in the emotional memory trade-off effect as we age.

Along with changes in neural plasticity and circuitry described above come alterations to sleep quality and quantity, which begin to occur as early as middle age (Ohayon et al., 2004). On a macro level, sleep latency, or the time to fall asleep, increases with age, as does the amount of time and duration of awakenings after sleep onset, leading to more fragmented and fragile sleep (Pace-Schott & Spencer, 2013). Sleep, overall, is shorter in duration, composed of increased lighter sleep (stages 1 and 2 sleep) as well as shorter non-rapid eye movement (NREM)-REM cycles. While there is a more subtle reduction in REM sleep (Van Cauter et al., 2000; Pace-Schott & Spencer, 2011), most notably reduced is the deeper NREM sleep, called slow wave sleep (SWS).

At the micro level, the most notable alterations to the properties of sleep beginning in middle age involve cortical slow wave activity (SWA, comprising slow oscillations 0.5 – 1 Hz and delta activity 1 – 4 Hz) and sleep spindles, which are thalamically generated bursts of activity (11 – 15 Hz) present throughout NREM sleep. Spectral power of cortical SWA is reduced with age, most prominently over the prefrontal cortex during the first cycle of nocturnal sleep. This is the time when homeostatic sleep pressure is highest, typically leading to maximal expression of SWA in younger adults (Mander et al., 2013). This reduction in prefrontal SWA indicates a possible impairment in homeostatic sleep regulation as age increases (Dijk et al., 2010). Properties of the slow waves are also reduced, with shorter amplitude and less density and frequency of the slow waves (Carrier et al., 2011; Massimini et al., 2004). Similarly, spectral power and frequency of sleep spindles are reduced beginning in middle age compared to young adults (De Gennaro & Ferrara, 2003; Mander et al., 2014). Additionally, spindle density and amplitude decrease, with the greatest reductions over the frontal cortex (Martin et al., 2012; Nicolas et al., 2001). While we are still discovering the reasons underlying these oscillatory alterations, recent research points to structural changes in brain regions with age. Of particular interest, but not exclusively, reduced grey matter density and atrophy in the prefrontal cortex is related to slow oscillation activity (Dubé et al., 2015; Varga et al., 2016). Additionally, as sleep spindles have been found to be temporally correlated with hippocampally generated sharp-wave ripples, bursts of activity of around 200 Hz (Siapas & Wilson 1998; Sirota et al. 2003), atrophy of hippocampal cells as age increases leads to sleep spindle impairment (Mander et al., 2017).

In young adults, the general theory of systems consolidation holds that reactivation of the neural networks associated with a learning experience leads to the strengthening of synaptic connections and a gradual redistribution of the memory into long-term cortical storage (see Diekelmann & Born, 2010, Payne et al., 2008b for review). It is thought that this reactivation optimally occurs during sleep, specifically SWS for declarative memories. Hippocampal sharp wave ripples, thought to induce long term potentiation in the neural circuitry, and thalamically-generated sleep spindles are temporally grouped by the upstates of the slow oscillations (Pavlidis & Winson, 1989; Buzsaki, 1989). This is hypothesized to be the complex mechanism leading to the reinstatement of the neural network and consolidation of the memory during sleep. Slow wave activity is also theorized to downscale the synaptic potentials that increase with activity and deplete neural resources, functionally resetting the

membrane potentials and increasing the plasticity of the cells (Tononi & Cirelli, 2003). At the same time, downscaling increases the signal-to-noise ratio, with highly potentiated networks representing learned experiences remaining above threshold and weaker traces falling below threshold, resulting in a memory trace that stands out after sleep.

The question, then, is whether the functional relationship between sleep and memory consolidation remains the same as we age. As the macro and micro properties of these sleep mechanisms begin to be reduced with age, does this, in turn, lead to reduced memory in parallel? Do the same relationships that we see in young adults exist between sleep and memory as we age? Recent research supports this idea, with shorter, shallower, sleep related to impaired cognitive performance even with healthy aging (Backhaus et al., 2007; Van Der Werf et al., 2009; Mander et al., 2014; Lo et al. 2016; Cavauto et al., 2016). Looking specifically at young and middle aged adults, Backhaus and colleagues (2007) found evidence to support maintenance of the functional relationship between SWS and paired-associates memory, in that they saw both sleep and memory decline in middle age, leading to reduced sleep-dependent consolidation. Moreover, although memory was equal between age groups prior to sleep (specifically early night SWS-rich sleep), memory declined more for the middle-age adults when retested after sleep. It may also be that the components of the complex sleep-based mechanism becomes uncoupled with age, such that slow oscillations and sleep spindles become uncoordinated, leading to memory impairments (Ladenbauer et al., 2017; Helfrich et al., 2017). However, it is possible that sleep and memory become functionally disassociated with age, with memory retention no longer predicted by sleep-based mechanisms, or not in the same manner, as in the young (Scullin, 2013; Baran et al., 2016). For example, Wilson and colleagues (2012) found no difference in sleep-dependent changes in memory for paired-associates between young, middle age, and older adults, even though the sleep physiology likely differed between these age groups. Although they found preserved performance for all, they did not record sleep, so could not conclude anything of the relationship between sleep features and memory. It is possible that these studies are limited by the lens of their analyses, such as focusing more on the macro alterations in sleep with age (e.g., overall percentage of SWS) rather than examining the sleep and memory relationship at the micro level (e.g., spectral power of oscillations, topographical differences) (Mander et al., 2017).

The current study was undertaken with several carefully considered study design decisions in mind. First, while the emotional memory trade-off effect has been well examined in both young and elderly adults, it is not well understood in middle age. We wanted first, to investigate emotional memory consolidation in this population, compared to well-studied young adults, and to then expand on the memory effect by exploring the impact of sleep on the selective consolidation of emotionally salient information. Second, as mentioned earlier, the work exploring the role of sleep in this selective consolidation has primarily sampled from young adults, typically undergraduates at universities and colleges. Using such “captive audience” samples is certainly convenient, and the reduced variability in such young, relatively healthy populations has provided a clear picture of the effect in this demographic. However, there is a critical need to expand research to the greater population across a wider age range to examine the impact on aging in a generalizable way. Therefore, we chose to

sample from the regional community rather than the student population to increase generalizability.

Third, while it has been well-established that the magnitude of the emotional memory trade-off increases over a period of nocturnal sleep, primarily through the preferential preservation of the memory for emotionally foci of complex scenes (Payne et al., 2008a; Payne & Kensinger, 2011), only one study has examined this effect across a daytime nap (Payne et al., 2015). Similar increases in the emotional memory trade-off were observed during a nap compared to an overnight design. However, different properties of sleep were involved in selective memory for emotional components. REM sleep across a night, both total amount in minutes and percentage of total sleep time spent in REM sleep, was positively related to memory for negative objects (Payne et al., 2012). Emotional memories have been found to benefit from REM sleep, with theta activity (4–7 Hz) during REM sleep and high functional connectivity between the amygdala and the medial temporal lobe resulting in stronger memory traces for emotional information (Wagner et al., 2001; Hu et al., 2006; Nishida et al., 2009). While there has been a tendency to associate SWS and REM sleep with neutral declarative and emotional memory consolidation, respectively, this strict dichotomy seems increasingly unlikely as literature emerges implicating REM sleep in neutral (Groch et al., 2011), and SWS in emotional memory processing. Importantly, as related to the emotional trade-off effect, we recently found that properties of SWS (percentage and delta frequency power) during a daytime nap were positively related to negative object memory (Payne et al., 2015). This suggests a possible functional difference between nocturnal and daytime sleep in memory processing that needs to be further explored. The current study was designed to extend previous work (Payne et al., 2015) and provide further evidence and clarity of the contribution of daytime sleep physiology in the selective consolidation of emotional components of memory.

Finally, the current study design employed a novel between-subjects daytime nap design, with a nap either occurring immediately after encoding in the late morning or in the afternoon with a delay after encoding. The purpose of this design was two-fold. We wanted to create two conditions that encoded and were tested at the same time of day that also had equal amounts of sleep and waking activity with the placement of the nap in time being the primary difference to address interference concerns that simply being “offline” during a nap passively protected the memory from external stimulation (see Figure 1). We also wanted to manipulate the sleep stage composition of the naps, with the later nap consisting of proportionally more SWS based on the idea that homeostatic increases in sleep drive increases SWS during the subsequent sleep period. This design was previously implemented and concluded that a daytime nap with more SWS (i.e., the later nap compared with naps earlier in the day) promotes declarative memory consolidation (Alger et al., 2010). If properties of SWS (e.g., amount, power, spindle activity) during a nap underlie selective consolidation of emotionally salient information as previously reported (Payne et al., 2015), we anticipated we would see this more clearly using this dual-nap design.

In the current study, we aimed to compare the emotional memory trade-off effect in young (18–39 years) and middle age (40–64 years) adults using a dual-nap design. We hypothesized that a daytime nap would better preserve memory for the emotionally salient

objects in encoded scenes at the expense of memory for the related neutral backgrounds, compared to remaining awake, thus increasing the magnitude of the negative memory tradeoff. We predicted that, similar to our previous nap study in young adults (Payne et al., 2015), memory for the emotionally salient objects would be related exclusively to features of SWS (e.g., amount of SWS, spectral frequency markers, sleep spindles). Stemming from this prediction, we anticipated that if the later nap resulted in a higher proportion of SWS within the composition of the nap, this could result in superior retention of the salient information. Finally, while we expected to replicate the emotional memory trade-off effect in both young and middle age adults, whether we would see a similar relationship between properties of sleep and preferential memory preservation was more of an exploratory question, especially as so little is known about sleep and memory in middle age. Perhaps we would see the relationship weakening with age or perhaps different aspects of sleep would benefit memory as age increased. Notably, as we anticipated more Stage 2 sleep in substitution for SWS with greater age, and spindle activity in NREM sleep is altered with age, perhaps this lighter stage of sleep is more involved in memory processing as we age.

2. Methods

2.1. Participants

Eighty participants (52 female) were recruited from the greater Northern Indiana community, through advertisements in local online publications (e.g., Craigslist), flyers around the University of Notre Dame campus targeting non-students, as well as through a preexisting pool of individuals who were concurrently participating in the longitudinal developmental Notre Dame Study of Health and Well-being. In order to increase external validity, this study aimed to minimize recruiting individuals currently enrolled at a four-year university. Subjects recruited were 18–64 years of age (mean $\pm$ SEM, 36.35 ± 1.39 years) and in good health as assessed by an in-depth screening self-report, with no history of sleep disturbance, learning disorders, or mental/emotional problems. They were free from all medications known to impair or facilitate sleep, mood, and attention. While we recruited evenly across this age range, we intended to compare a young cohort to middle age adults. Consistent with the categorical boundaries set in the Notre Dame Study of Health and Well-being, we considered young adults as 18–39 years old ($n = 45$, 32 female, mean age 26.91 ± 0.80) and middle age adults as 40–64 years old ($n = 35$, 20 female, mean age 48.49 ± 1.23). All subjects received monetary compensation at the completion of their participation. This study was conducted according to the principles expressed in the University of Notre Dame Human Subjects Institutional Review Board, with all subjects providing written informed consent.

Prior to arrival to the lab, participants were contacted via email to ensure eligibility by administering the screening form. Participants were instructed to get no less than 6 hours of sleep the night before the experimental day and refrain from caffeine, alcohol, nicotine, and unnecessary medication 24 hours before and for the duration of the study. In accordance with these restrictions, they kept a one-week sleep log to track their sleep/wake times, napping habits, and alcohol and caffeine intake prior to the experimental day. Upon arrival to the lab, participants were asked to sign a consent form explaining the nature of the research.

Throughout the experimental day, they completed a demographics questionnaire as well as several other questionnaires to assess sleep habits and rule out psychopathology. These included the Pittsburg Sleep Quality Index (PSQI; Buysse, Reynolds, Monk, Berman, & Kupfer, 1989) to assess general sleep quality, the Morningness–Eveningness Questionnaire (MEQ; Horne & Ostberg, 1976) to assess morningness–eveningness tendencies, the Beck Anxiety Inventory (BAI; Beck, Epstein, Brown, & Steer, 1988) and State-Trait Anxiety Inventory (STAI, Spielberger, 2010) to assess level of state (STAI-X1) and trait (STAI-X2) anxiety, the Beck Depression Inventory–II (BDI; Beck, Ward, & Mendelson, 1961) to assess depressive symptoms, the Positive and Negative Affect Schedule (PANAS; Watson, Clark, & Tellegen, 1988) to assess positive and negative affect, and the Stanford Sleepiness Scale (SSS; Hoddes, Zarcone, Smythe, Phillips, & Dement, 1973) to assess state sleepiness. These measures were used primarily to compare nap and wake groups to ensure they were similar to one another as well as to determine if any participant should be excluded because of scoring in elevated (i.e., clinical) ranges for depression or anxiety, which could impact both sleep and emotional memory formation.

2.2. Materials

2.2.1. Encoding materials—During the encoding task, participants viewed a set of 100 scenes that displayed either a neutral or negatively salient foreground object (50 of each valence) placed on a plausible neutral background (e.g., a deer or an angered grizzly bear in the forest, a taxi cab or a taxi accident on a street). For each of the scenes, we created four different sets by placing two similar neutral objects (e.g., two images of a deer) and two similar negative objects (e.g., two images of angry bears) on two neutral backgrounds (e.g., two forests) in order to create four different but related scenes (Payne et al., 2008a, 2012, 2015; Cunningham et al., 2014). We used similar images to avoid the possibility that specific images may be more memorable, thus skewing the results. By varying the type of object (neutral or negative) and the background version (one of the two paired backgrounds), four different lists of 68 scenes were created. Each participant saw only one list at encoding, which was randomly determined using a mixed Latin Square design.

2.2.2. Recognition materials—Memory for the images was tested during two sessions, baseline, which occurred immediately after encoding, and retest, which occurred later in the day. During both recognition tests, objects and backgrounds that had been viewed in the encoded complex scenes were presented separately (i.e., only the object or the background at one time) in random order. Each of these items had been previously studied was considered ‘old’. Intermixed with these ‘old’ images were an equal number of objects and backgrounds, equated on measured ratings of valence and arousal, that had not previously been presented and were considered ‘new’. Half of the encoded information was tested at baseline and the other half at retest. Therefore, each recognition test consisted of 25 ‘old’ negative objects, 25 ‘old’ neutral objects, 25 ‘old’ backgrounds previously paired with the negative objects, 25 ‘old’ backgrounds previously paired with the neutral objects, 25 ‘new’ images each of negative and neutral objects, and 50 ‘new’ backgrounds, for a total of 200 images presented during each test.

Most of the complex images used in this study have also been used in previous studies and have yielded normed ratings categorizing the scenes as either negative or neutral (Payne et al., 2008a, 2012, 2015; Cunningham et al., 2014). Because we altered a few of the images, we first conducted a pilot study ($n = 24$) in part to evaluate the valence and arousal of the images to guide their placement into a negative or neutral category. In this pilot study, sampling from the undergraduate community of the university (ages 18 – 23 years), images were assessed using 7-point scales. All negative images had received arousal ratings of 5 to 7 (with higher scores associated with more arousing ratings) and valence ratings higher than 5 (with higher scores representing a more negative image). All neutral items (objects and backgrounds) had been rated as un-arousing (arousal values lower than 3) and neutral (valence ratings between 3 and 5). Subjective ratings were also taken from participants in the current study for the full scenes at encoding, both to ensure the participants were attending to the stimuli, thus facilitating encoding, as well as to verify that the image ratings by this community population encompassing a wider age range were consistent with the undergraduate population ratings in the pilot study.

2.3. Procedure

At approximately 9:30am, participants arrived in the Sleep, Stress, and Memory Lab, signed informed consent, and completed a set of questionnaires (PSQI, MEQ, STAI-X2, BAI, STAI-XI, and PANAS), which also included the Stanford Sleepiness Scale (SSS). They then incidentally encoded all 100 scenes with instructions to attend to the scene and make subsequent ratings of valence and arousal. Each scene was displayed for 3000ms. After each scene was presented, participants were asked to make their self-paced ratings, first for valence and then level of arousal, on separate screens. Ratings were followed by a crosshair fixation point for 500ms in order to direct attention and prepare for the next scene presentation. A self-paced baseline recognition test followed immediately after encoding, during which an image of an individual scene component (i.e., object or background) was presented centrally. Participants were required to press '0' if the image was considered 'new', or '1' if they considered the image 'old' (see Figure 2). Participants were told at the end of the first session that they would be tested in this manner once more during the second session.

The current study was designed to have a wake control condition compared to two nap conditions in which participants obtained an equal amount of sleep and wake as one another, the primary difference being the positioning of the nap in time. Due to scheduling constraints, participants were randomly assigned into wake control, immediate nap, or delayed nap groups in advance of their participation date. Wake and delayed nap participants were allowed to leave the lab following the baseline test with the instruction not to nap or consume caffeine, alcohol, nicotine, or unnecessary medication during the retention period. Those in the immediate nap group were prepared for polysomnography (PSG). At around 11:00am, immediate nap subjects were given a 90-minute PSG-recorded sleep opportunity lasting until about 12:30pm. If participants obtained SWS or REM sleep during this time, they were allowed to sleep until they transitioned out of these stages in order to reduce sleep inertia and disorientation. After awakening, electrodes were removed and these nap participants were allowed to leave the lab until the second session, approximately 4 hours

later. Delayed nap participants returned to the lab at 2:30pm and were prepared in the same manner for a 90-minute PSG-monitored nap opportunity, beginning at 3pm. They were awakened at approximately 4:30pm and allowed 30 minutes to recover for sleep inertia before beginning the retest session (see Figure 1 for timeline).

All participants returned to the lab at 5pm for the retest session and began by completing a second set of questionnaires similar to the previous session (BDI, STAI-XI, PANAS, and SSS). Following these questionnaires, participants completed a psychomotor vigilance task (PVT, Dinges & Powell, 1985) to obtain objective measures of alertness and address the possible confound of an overall vigilance difference impacting task performance, rather than specifically sleep or lack of sleep accounting for performance differences. This cognitive test is a sustained reaction-time task that measures the average speed of a participants' response to a visual stimulus and demonstrates high test-retest reliability (median intraclass correlation coefficients/ICC > 0.8) and sensitivity (Dorrian et al. 2005, Basner & Dinges, 2012) to the effects of sleep deprivation (sensitivity = 93.7%), suggesting high convergent validity as well. After completing the PVT, participants were again tested via recognition on their memory for the remaining half of encoded material using the same method described above (see Figure 1 for protocol).

2.4. Polysomnography Recordings

Nap participants were monitored online using digital EEG acquisition software (Comet System-Grass/Twin PSG Clinical Software) using a standard polysomnographic montage, which included electroencephalography (EEG; F3, F4, C3, C4, Cz), electrooculography (EOG), and chin electromyography (EMG) channels, each referenced to contralateral mastoids. Each 30-s epoch of recorded sleep was scored blind to participants' behavioral task performance according to the standards of Rechtschaffen and Kales (1968). The PSG recording was scored visually as NREM Stages 1, 2, SWS, and REM sleep, or wakefulness. EEG data were then filtered at 0.3–35 Hz (with a 60 Hz notch filter), and artifacts were visually identified and removed using the EEGLAB 13.4 (Delorme & Makeig, 2004) toolbox for MATLAB 9.1 (The MathWorks Inc., Natick, MA, 2012). Spectral analysis was then applied to all artifact-free epochs of NREM and REM sleep during the nap. Fast Fourier transforms were carried out in MATLAB 9.1 to calculate mean absolute power spectral density ($\mu\text{V}^2/\text{Hz}$) in the slow oscillation (0.5–1 Hz), delta (1–4 Hz), theta (4–7 Hz), and sigma (11–15 Hz) frequency bands in order to examine the frequencies associated with NREM SWS, sleep spindles, and REM sleep. Sleep spindle counts were obtained by filtering all artifact-free epochs of Stage 2 sleep and SWS at 11–15 Hz and then applying the established automatic detection algorithm of Ferrarelli et al. (2007) using MATLAB 9.1. Total number of spindles, spindle density (spindles/total analyzed time in Stage 2 and SWS), duration, amplitude, and power of the spindle events were calculated for frontal and central scalp electrodes (F3, F4, C3, C4) as well as the average across electrodes. The spindle range of 11–15 Hz was chosen based on prior research using this definition (Schabus et al., 2007; Tamminen et al., 2010; Lewis & Durrant, 2011).

2.5. Data Analyses

We compared behavioral performance primarily within each age cohort to examine the impact of napping on memory in young (wake condition $n = 15$, immediate nap $n = 15$, delayed nap $n = 15$) and middle age (wake $n = 11$, immediate nap $n = 12$, delayed nap $n = 12$) adults. For both baseline and retest sessions, memory for the objects and backgrounds was calculated separately for each valence (negative and neutral) as the number of items accurately remembered (i.e. hits) divided by the number of items originally viewed. To correct for response bias, we calculated corrected memory by subtracting the proportion of false alarms ('old' judgments to new pictures) of each object and background type from the proportion of hits (Snodgrass and Corwin, 1988) to obtain the measures we will refer to below as memory for each component (e.g., negative object memory, neutral object memory, etc.). As a measure of the trade-off in memory, we further calculated the trade-off scores for each valence at each test by subtracting the memory for the associated backgrounds from the memory of the objects (e.g., corrected memory for negative objects minus corrected memory for backgrounds on which negative items were presented). These trade-off scores give us the magnitude of the difference between object and background memory. Performance changes over the retention period were the primary analyses conducted using repeated measured mixed ANOVA. Correlations were conducted between sleep variables and behavior to investigate the nature of the role of sleep in performance benefits.

3. Results

3.1. Questionnaires and Sleepiness Measures

All analyses to compare questionnaire data were conducted between the six groups encompassing both experimental conditions (i.e., wake, immediate nap, delayed nap) and age groups (i.e., young, middle age). A One-Way ANOVA was conducted to verify that there were no differences between conditions or age groups on any questionnaire responses or sleep log data, such as amount of sleep achieved the night before the experimental day. There were no significant differences between groups on any measure of anxiety, depression, positive or negative affect, sleep quality, or morningness/eveningness tendencies. On average, all groups scored similarly as neither morning nor evening type on the MEQ, with only 3 individuals in separate groups scoring as an extreme morning ($n = 2$) or extreme evening ($n = 1$) chronotype. Excluding these extreme types from the analyses below did not change any of our findings. As for sleep and wake activity prior to arriving for participation, all subjects were compliant in getting more than 6 hours of sleep the prior night, with group averages ranging from 7.42 ± 0.19 to 7.78 ± 0.30 hours ($p = 0.63$). All groups of subjects awoke, on average, between 6:45 – 7:45 AM the morning of the experiment, with no significant differences between groups. We found no difference in average number of daytime naps taken over the week prior to participation, either between age cohorts or between experimental conditions. With regard to sleep quality over the previous 1-month time interval, PSQI scores were not significantly different between groups (all $p > 0.78$).

In order to verify that there were no group differences in sleepiness that could explain performance variations, we compared scores on the Stanford Sleepiness Scale and found no difference between groups either at Session 1 ($F_{5,74} = 1.51$, $p = 0.20$) or Session 2 ($F_{5,74} =$

1.64, $p = 0.16$). In addition, PVT comparisons revealed no significant differences between groups at retest in basic vigilance and alertness ($F_{5,74} = 0.83$, $p = 0.53$).

As mentioned above, we previously conducted a pilot study to assess these images in part to determine the average valence and arousal ratings in order to categorize them into negative and neutral categories for the current study. We also had subjects in the current study rate the complete scenes at encoding, both to promote deeper encoding as well as verify our categorization of images. In examining the average image ratings per image in both the young and middle age cohorts, we determined that the image ratings were, overall, shifted in a more positive direction. Within each of the 4 different lists, we found that the average valence and arousal ratings resulted in 16 of the 50 neutral images being categorized as positive (valence ratings 1 to 3, mean 2.60). We also found that 9 scenes in each list that were originally categorized as negative were consistently rated as neutral and 1 neutral scene per list was rated as negative, thus leading us to recategorize these images for subsequent analyses. This yielded 42 scenes rated as neutral (valence 3.89, arousal 2.24) and 42 scenes rated as negative (valence 5.74, arousal 5.23) per list. In light of the literature regarding the possibility of a positivity bias with age, in which memory become biased to better remember more positive information compared to young adults (Mather & Carstensen, 2005), we examined whether there were differences in these ratings between young and middle age cohorts. As this task has been previously used to compare young to elderly adults with no sign of a positivity bias (Waring & Kensinger, 2009), we were not anticipating a difference. Correspondingly, between the cohorts, we found that it was the same images that fell outside of the original categorization, resulting in the same recategorization within each age cohort. In general, there was a slight shift toward positive in the middle age group for average valence ratings for both neutral (3.87 compared to 3.91 in the young group, $t_{78} = 0.28$, $p = 0.78$) and negative (5.68 compared to 5.79 in the young group, $t_{78} = 1.05$, $p = 0.30$) scenes. This positivity shift was not great enough to result in the image categorizations differing between groups.

3.2. The Impact of Napping on Memory

We first analyzed the impact of napping on the selective consolidation of emotional memory within each age cohort to determine if the previously reported effect of sleep increasing the magnitude of the emotional memory trade-off was present. For both the young and middle age cohorts, we conducted a 3 (condition: wake, immediate nap, delayed nap) x 2 (session: baseline, retest) x 2 (valence: negative, neutral) x 2 (component: object, baseline) mixed ANOVA with condition as a between-subject variable and session, valence, and component as within-subjects variables. In the young cohort, a main effect of session ($F_{1,42} = 341.41$, $p < 0.001$) indicated that memory, in general, declined between baseline and retest sessions (see Table 1). We also found a main effect of component ($F_{1,42} = 218.15$, $p < 0.001$) demonstrating superior memory for objects over backgrounds. A two-way interaction was found between valence x component ($F_{1,42} = 243.58$, $p < 0.001$), demonstrating the emotional memory trade-off effect, with better memory for negative objects (0.87 ± 0.01) compared to neutral (0.69 ± 0.02), but at the expense of memory for backgrounds paired with negative objects (0.41 ± 0.02) compared to those paired with neutral objects (0.57 ± 0.02). A second two-way interaction between session x component ($F_{1,42} = 27.48$, $p <$

0.001), indicated that memory for objects was better retained across sessions, and this was further qualified by a three-way interaction between session x component x group ($F_{2,42} = 4.12$, $p = 0.02$, $\eta_p^2 = 0.16$). Independent t-tests revealed that the immediate nap group retained memory over time for the objects significantly better than the wake control group ($t_{28} = -2.65$, $p = 0.01$), but not the delayed nap group ($t_{28} = 1.65$, $p = 0.11$). A three-way interaction was also found between session x valence x component ($F_{1,42} = 4.76$, $p = 0.04$, $\eta_p^2 = 0.10$), demonstrating that negative object memory is better retained over time than neutral object memory while memory for backgrounds paired with negative objects decays more than neutral backgrounds, thus increasing the magnitude of the emotional memory trade-off over time. However, this effect was further qualified by a significant four-way interaction ($F_{2,42} = 3.75$, $p = 0.03$, $\eta_p^2 = 0.15$). Post hoc comparisons revealed that the immediate nap group had significantly higher retention of memory for negative objects compared to both the wake ($t_{28} = -3.22$, $p = 0.003$) and delayed nap ($t_{28} = 2.22$, $p = 0.03$) conditions, and significantly more decay in memory for backgrounds associated with negative objects compared to the wake condition ($t_{28} = 2.14$, $p = 0.04$). In other words, immediate napping resulted in the greatest increase in the magnitude of the emotional memory trade-off in the young cohort.

In the middle age cohort, we found a similar pattern of results. Main effects of session ($F_{1,32} = 189.16$, $p < 0.001$) and component ($F_{1,32} = 282.36$, $p < 0.001$) demonstrated a general decline in memory over time and superior memory for objects. A two-way interaction was found between valence x component ($F_{1,32} = 142.95$, $p < 0.001$), again demonstrated the emotional memory trade-off effect. A second two-way interaction between session x component ($F_{1,32} = 10.55$, $p = 0.003$), indicated that memory for objects was better retained across sessions. Similar to the young cohort, middle aged adults showed a significant four-way interaction ($F_{2,32} = 4.53$, $p = 0.02$, $\eta_p^2 = 0.22$), again indicating a greater increase in the magnitude of the emotional memory trade-off in the immediate nap group. Post hoc comparisons revealed that the immediate nap group had significantly higher retention of memory for negative objects compared to the wake condition ($t_{21} = -2.04$, $p = 0.05$) and more decay in memory for backgrounds associated with negative objects - significantly so compared to the delayed nap condition ($t_{21} = -2.40$, $p = 0.03$), and at a trend level compared to the wake condition ($t_{21} = 1.92$, $p = 0.07$).

Figure 3 depicts the above findings for young and middle age using the calculated change in the memory trade-off scores (i.e., the difference in memory for objects and memory for backgrounds) across the retention period. The similarity of the pattern of the effect between the age groups for the negative memory trade-off can be seen with more clarity using this measure rather than examining object and background memory separately, as reported above. When comparing the increase in the magnitude of the trade-off scores over time using t-tests, no differences were found between conditions for neutral information for either young (all $p > 0.37$) or middle age (all $p > 0.32$). However, focusing on the increase in the negative trade-off, we see that for both age groups, the immediate nap condition had a significantly greater increase in the difference between memory for negative objects and background than both the wake (young $t_{28} = -3.14$, $p = 0.004$; middle age $t_{21} = -2.54$, $p = 0.02$) and delayed nap (young $t_{28} = 2.52$, $p = 0.02$; middle age $t_{21} = 2.17$, $p = 0.04$) conditions (Figure 3).

In order to get a clearer picture of the relationship between age and memory, we ran a correlation analysis between age (as a continuous variable across the entire age range of 18–64 years) and baseline performance for the negative information to determine how age affected the ability to encode new salient information. We found significant negative correlations between age and negative object and associated background memory ($r = -0.40$, $p < 0.001$; $r = -0.27$, $p = 0.02$, respectively) and background memory associated with neutral objects ($r = -0.30$, $p = 0.01$), but not neutral object memory ($r = -0.14$, $p = 0.20$), although also negatively correlated. However, when looking at consolidation of this information, measured by the change in performance across the two sessions, we did not see a significant correlation with any dependent variable of interest and age for any condition (wake group, all $r < 0.22$, all $p > 0.28$; immediate nap, all $r < 0.29$, all $p > 0.14$; delayed nap, all $r < 0.28$, all $p > 0.10$). These findings indicate that while encoding may be impaired with age in this task, the ability to consolidate the information does not appear to be.

3.3. Sleep Stage Parameters, EEG Spectral Power, and Sleep Spindles

As one of our predictions in using this dual-nap design was to examine the sleep stage composition between nap times, we first examined nap architecture in the immediate and delayed naps within each age cohort to determine if there was a shift in the relative amounts of Stage 2, SWS, and REM sleep, as a function of increased sleep pressure throughout the waking day (Alger et al., 2010). Table 2 summarizes sleep parameters for both age cohorts for the two naps. The PSG recording for one subject in the young delayed nap condition contained a gap that prevented calculations of the total sleep time or percentages of the specific stages relative to this time. Those data were thus excluded from subsequent sleep architecture comparisons and correlational analyses. Within the young cohort, we found no significant differences between immediate and delayed nap architecture (Table 2). Of particular interest, total sleep time (TST, $t_{27} = -0.03$, $p = 0.98$) and the percentages of Stage 2 ($p = 0.54$), SWS ($p = 0.66$), and REM sleep ($p = .18$) were all similar between naps. In the middle age cohort, we likewise found no significant differences in naps (TST $t_{22} = 0.82$, $p = 0.42$; Stage 2 $p = 0.44$; SWS $p = 0.78$; REM sleep $p = 0.65$). Therefore, we collapsed across the two nap times in order to compare young to middle age sleep quality based on prior research demonstrating altered sleep architecture with age. As detailed in Table 3, middle aged adults had significantly shorter total sleep time (TST, $t_{51} = 2.61$, $p = 0.01$), more wake after sleep onset (WASO, $t_{52} = -3.55$, $p = 0.001$), poorer sleep efficiency ($t_{51} = 2.94$, $p = 0.007$), more Stage 1 ($t_{51} = -3.93$, $p < 0.001$) and Stage 2 sleep (trending, $t_{51} = -1.87$, $p = .07$), and less SWS ($t_{51} = 4.35$, $p < 0.001$) compared to young adults.

A limited number of comparisons were planned between sleep and both negative and neutral memory measures. Based on our a priori hypotheses and the previous literature regarding differing correlations with REM sleep and SWS and memory for salient information, we focused our correlational sleep stage analyses on percentages of SWS and REM sleep. Because we were also interested in the impact of alterations to properties of Stage 2 sleep (sleep spindle activity), we included this stage in separate, secondary analyses. Looking across all nappers, we found a significant positive correlation between the percentage of SWS and memory for the negative objects at retest ($r_{53} = 0.35$, $p = 0.01$, Figure 4), which is similar to what was previously found using this emotional memory trade-off task and a

daytime nap protocol (Payne et al., 2015). This supports our hypothesis that SWS during a nap facilitates the selective consolidation of emotionally salient information. Of the total number of subjects in the napping conditions, 25 young and 10 middle age adults obtained SWS. When we removed those who did not reach SWS from the analyses, the correlation remained significant ($r_{35} = 0.34$, $p = 0.05$). No other significant correlations were found. When examining correlations between sleep stages and memory within each age cohort, we again saw a relationship between SWS and negative object memory in the young cohort both as a trend at retest ($r_{29} = 0.36$, $p = 0.06$) and significant assessing the change in performance across the retention period ($r_{29} = 0.38$, $p = 0.04$), indicating better preservation of negative object memory with more SWS. No other correlations in either cohort reached significance but showed similar direction of the relationships as when looking at sleep across the age continuum.

To further clarify the role of sleep physiology in selective memory processing, we examined mean absolute spectral power ($\mu V^2/Hz$), specifically looking at slow oscillations (.5–1 Hz), delta (1–4 Hz), theta (4–7 Hz), and sigma activity (11–15 Hz). Comparing the age cohorts on spectral power, we found that middle age adults had significantly less slow oscillation and delta power ($t_{32} = 2.85$, $p = 0.01$; $t_{32} = 2.85$, $p = 0.01$) but similar power in the sigma and theta ranges (both $p > 0.47$) as the young adults. Our correlational analyses revealed that the only significant relationships that emerged were from activity during SWS. We found that memory for negative objects at retest was significantly positively correlated with both slow oscillation power ($r_{24} = 0.45$, $p = 0.03$) and delta power ($r_{24} = 0.44$, $p = 0.03$) in the young adults. This complements the finding showing significantly higher slow oscillation and delta spectral power in the young vs. middle age adults. To rule out the possibility that these correlations were driven by a single electrode site, we confirmed that slow oscillation activity was significantly, or nearly significantly, related to memory at all four electrode sites (C3 $p = 0.07$, C4 $p = 0.04$, F3 $p = 0.09$, F4 $p = 0.01$). Similarly, delta power was related to negative object memory at all four electrodes as well (C3 $p = 0.04$, C4 $p = 0.02$, F3 $p = 0.09$, F4 $p = 0.02$). These relationships between negative object memory and slow oscillations/delta power were not significant in the middle age group.

Analyses of sleep spindle activity during NREM sleep yielded significant associations that were complementary to our overall findings in that the only significant relationships to emerge were from SWS. Across all nappers, we found that memory for negative objects was positively related to all measured aspects of sleep spindle activity, but only during SWS, including the total number of sleep spindles ($r_{33} = 0.46$, $p = 0.007$), spindle density ($r_{33} = 0.45$, $p = 0.009$), average duration of spindles ($r_{33} = 0.50$, $p = 0.003$), spindle amplitude ($r_{33} = 0.50$, $p = 0.003$), and power of the spindle events ($r_{33} = 0.52$, $p = 0.002$). As we examined the relationship between negative object memory and several different measures of spindle activity in one analysis, we corrected for multiple comparisons using the Bonferroni correction. All p-values surpassed the more conservative criteria of 0.01 to remain significant. To ensure that these associations were not the result of activity from one electrode, we verified that all electrode sites maintained the same significant positive relationship between negative object memory and all spindle variables, with the exception of sleep spindle density and the C3 and F4 sites, where only a trending relationship was seen ($r_{31} = 0.30$, $p = 0.13$; $r_{31} = 0.31$, $p = 0.09$, respectively). When examining the associates

within age cohorts, young adults showed significant positive relationships with the same variables (total spindle number, $r_{24} = 0.47$, $p = 0.02$; density, $r_{24} = 0.38$, $p = 0.07$; duration, $r_{24} = 0.53$, $p = 0.008$; amplitude, $r_{24} = 0.54$, $p = 0.006$; power, $r_{24} = 0.52$, $p = 0.01$). Middle age adults also demonstrated similar relationships between spindle activity and emotional object memory, but did not reach significance (all r values between 0.61 to 0.49, p values between 0.08 to 0.19). Using the Fisher r -to- z transformation, we compared the difference between the young and middle-age correlation coefficients for memory and spindle feature associations and found no significant difference between any association (total spindles: $z = -0.21$, $p = 0.83$; density: $z = -0.67$, $p = 0.50$; duration: $z = -0.21$, $p = 0.84$; amplitude: $z = 0.16$, $p = 0.87$; power: $z = -0.10$, $p = 0.92$). Comparing the young to middle age adults during SWS, the older group showed significantly fewer spindles ($t_{31} = 2.17$, $p = 0.04$) that were smaller amplitude ($t_{31} = 2.11$, $p = 0.04$) and nearly significantly briefer in duration ($t_{31} = 1.96$, $p = 0.06$) spindles. However, spindle density and power were similar between the age cohorts (both $p > 0.28$). Activity during SWS was our primary focus, but because we do observe a decrease in SWS while seeing an increase in Stage 2 sleep with age, we also ran separate, secondary spindle analyses on Stage 2 sleep. No significant correlations emerged from these analyses.

4. Discussion

The current study aimed to investigate the impact of daytime napping on the selective preservation of salient information across an age range from young to middle age. Specifically, we examined both how the behavioral emotional memory trade-off effect manifested between young and middle aged individuals and whether sleep soon after encoding played an active role in preferentially preserving negatively salient memories while simultaneously suppressing or forgetting neutral contextual details. We observed that memory in the middle-aged adults was numerically poorer than young adults, with age negatively correlated with ability to encode. However, memory for emotional objects was still promoted above neutral information. Beyond this, we assessed the quality and quantity of sleep for all ages and found that sleep was significantly worse in many ways in the middle-aged adults. Our observations fit the literature describing the alterations that develop in sleep architecture with age, with shorter total sleep time and more frequent arousals during a period of sleep. Furthermore, we confirmed that a middle-aged population obtains significantly less SWS and more of Stages 1 and 2 sleep, resulting in lighter and less consolidated sleep. However, despite these general declines in sleep and memory, we still revealed that napping immediately after learning selectively preserved memory, particularly emotional memory, and led to a greater increase in the magnitude of the emotional memory trade-off over time.

We note that our experimental groups had different experiences and possibly different expectations throughout the experiment, as the wake condition did not experience electrode hookup and did not undergo an in-laboratory nap with PSG recording. Aside from the activities specific to the recording of the nap, however, laboratory experiences were similar between groups. As we found no differences between conditions for any questionnaire measure, PVT score, or baseline memory performance, we do not think that the different experiences between conditions played a role in the results we obtained.

There is a dearth of research that focuses on the middle age in general, but specifically when examining the relationship between sleep and memory consolidation. The few studies that have looked at this population have produced equivocal findings, with some studies reporting preserved (Wilson et al., 2012) or reduced (Backhaus et al., 2007) sleep-dependent performance changes in declarative memory. To our knowledge, the current study is the first to focus on the middle age population examining the kind of nuanced, selective memory seen using the trade-off task, particularly in this highly novel napping paradigm. While studies tend to compare the extremes of young versus elderly adults to draw out the differences brought about by alterations in memory and sleep with aging, few studies investigate those of the age where these alterations begin to occur, leaving a gap in knowledge regarding this population.

One unexpected finding that this study revealed was that when sampling from a wider age range from the greater community, not focusing on a narrow population of young undergraduate university students, we did not observe the expected increase in the amount of SWS participants obtained on average during the late nap. Process S, or the homeostatic sleep drive posits that there is an increase in sleep pressure and need for homeostatic regulation, indexed by the amount of SWS, with SWS increasing exponentially as a function of how much time is spent awake prior to sleep (Knowles et al., 1986). Previous research has supported that this increase in SWS occurs (Carskadon & Dement, 2005, Tononi & Cirelli, 2006) and that it may be related to differential memory processing (Alger et al., 2010). However, research regarding the relationship between the time-based shift in sleep architecture as a modulator of memory processing has sampled heavily from the young student population. Perhaps the unique sleeping habits and intensive learning experienced by a typical undergraduate results in a more pronounced increase in SWS over time, particularly as evidence supports increases in properties of slow wave activity after intensive learning (Tononi & Cirelli, 2003; Huber et al., 2004; Eschenko et al., 2008). Future studies will need to replicate the lack of this shift in sleep architecture we observed in the current study to delve into the possible reasons behind it.

However, being that our study found that the sleep composition was relatively equal between the immediate and delayed naps in our age cohorts, the story here is about the benefits of immediately napping compared to the consequences of delaying sleep. Between the two nap conditions, the total amount of time asleep, as well as the time spent awake and engaged in everyday activity was equal, with the placement of the nap in time being the primary difference. This design helps to address potential criticisms that any benefits in the nap group are due to passive interference, such that the nap group performs better simply because they were protected from external stimuli during the whole of the retention period. If this were the only factor, then both naps in the current study would equally passively protect the memory and would perform similarly to one another. However, other factors may play a role, such as the natural deterioration of memory due to interference prior to a delayed nap that may reduce what memory is available to consolidate. Moreover, there is the idea that a limited window of time during which sleep can facilitate consolidation of labile memory traces, after which sleep will not be as effective (Mednick et al., 2011). The design we used is reminiscent of another common sleep study design in which learning, sleeping, and testing occurs across a 24- (or 48-) hour period. In these studies, sleep either occurs

immediately after learning in the evening (commonly called a sleep-first condition) or after a full day of wake filled with everyday waking interference following learning in the morning (called a wake-first condition), with testing occurring 24 hours after learning. Notably, these studies confirm that sleeping soon after learning leads to stabilization and protection of the memory from subsequent interference (Talamini et al., 2008; Van der Werf et al., 2009), including preferential preservation of emotional information (Payne et al., 2012). The current study demonstrates that these immediate benefits can even be observed over shorter retention periods, suggesting that this type of immediate sleep can be used as a technique for optimal memory.

The differences between young and middle-age adults in regard to the spectral composition of sleep falls in line with what has previously been found, particularly regarding the reduction of slow oscillatory and delta power with age. In addition to these reductions, we did not see the same correlations between memory for the emotional components of the images and power in these frequency ranges during SWS in the middle-aged subjects, as we observed in the young subjects. These findings may indicate that delta and slow oscillatory activity are less important or less actively involved in the preferential consolidation of salient information as we age. This could lend support to the idea that there is a weakening of the functional relationship between sleep and memory with age (Scullin, 2013). Indeed, while we found that the percentage of SWS during the nap was positively correlated with negative object memory across all participants, this was clearly driven by the young adults, with far weaker relationships in the middle-aged adults. However, looking to the sleep physiology within SWS, we see similar relationships between memory for negative information and sleep spindle features. This suggests, perhaps, that looking beyond the stage of SWS as a whole and focusing on specific features of sleep within this stage can reveal ongoing active beneficial processes across age.

There are a few possible explanations for our results. It is clear by looking at the behavioral data that sleep, overall, plays a significant role in the preferential preservation of salient information at the expense of memory for background detail, as demonstrated by the same pattern of the effect between age cohorts, with immediate sleep yielding superior memory. It may be that the underlying sleep-based mechanisms (i.e., collaboration between slow oscillations and sleep spindles actively leading to systems consolidation) are altered with age and that another, as of yet unknown, mechanism becomes more involved (e.g., sleep spindles alone). We may also be limited by how we are measuring sleep, relying on low-density EEG recordings, which may prevent us from uncovering the specifics of the change in mechanism. Using high-density EEG recordings in the future may allow us to tease out topographical differences in activity and uncover an altered mechanism. Extending this design over a wider age range that also encompasses older adults would also allow us to infer if the relationship between the known sleep-based mechanisms (i.e., SWS and spindles) and memory further changes with age. This is particularly important in light of new evidence that demonstrates the importance of the phase-coupling between slow oscillations and sleep spindles and how they may become uncoupled with age and/or cognitive impairment/Alzheimer's disease (Helfrich et al., 2017; Ladenbauer et al., 2017; Mander et al., 2017). There is great potential of developing an intervention strategy to boost sleep physiology and improve memory if the relationship persists with age. In young adults,

interventions such as increasing SWA (Westerberg et al., 2015) or sleep spindles (Mednick et al., 2013; Ladenbauer et al., 2016) through stimulation or pharmacological manipulations have enhanced memory performance, although whether this would be as successful in healthy older individuals has yet to be examined.

We caution, however, that the correlations we detected between sleep and performance, while intriguing, should be taken with a grain of salt, as our nap design resulted in many participants on the older end of the age range not obtaining any SWS. Only 10 of the 24 middle age adults obtained SWS in their naps, and only 9 of those subjects had detectable spindle activity. Thus, our observation of a dissociated relationship between memory and SWS (i.e., weaker with percentage and delta/slow oscillation power yet similar spindle power) may stem from these correlational analyses being underpowered, in contrast to our behavioral analyses, which were appropriately powered. In other words, our ability to fully perceive the functional relationship between sleep physiology and memory may have been impeded by the very nature of the aging brain in sleep, with reduced SWS, delta, slow oscillatory, and spindle activity reducing the number of participants included in the analyses.

Nevertheless, the current study adds support to the idea that daytime and nocturnal sleep contribute to the consolidation of salient information in functionally different ways, with SWS actively facilitating memory during daytime naps, most clearly in the younger participants. Similar to what we revealed in a previous study, memory for the emotionally salient focus on experiences was related to the relative amount of SWS and delta frequency power during this stage (Payne et al., 2015), with further evidence from the current study demonstrating a role for slow oscillatory power and sleep spindle activity during this stage as well. It is important to conclude from these sleep analyses that the *only* aspect of memory that was found to be related to sleep physiology was emotional object memory, not any form of neutral memory, reflecting preferential consolidation. These findings open the door to future studies designed to answer the question of why this differential contribution of SWS during the day and REM sleep overnight may occur. One thing to consider is that daytime naps have considerably less REM sleep than nocturnal sleep, given the way in which we cycle through sleep stages and the shorter duration of a nap. In the current study, 19 out of 30 young adults and 11 out of 24 middle-aged adults in the napping conditions obtained REM sleep. This is similar to, or more than in the case of the young adults, the number of subjects obtaining REM sleep in another nap study examining emotional memory consolidation. Nishida et al. (2008) found an association between REM sleep, and theta within REM sleep, and emotional memory consolidation. However, they did not examine selective consolidation, but rather consolidation of scenes as a whole. Our study not only brings further into question previously held ideas about REM sleep predominantly benefitting emotional memory, but also replicates our previous SWS and selective consolidation of emotional components findings (Payne et al., 2015). In our previous study, 16 out of 22 young adults obtained REM sleep. We speculate that perhaps circadian phase differences and factors such as neurohormonal cycles, such as the natural rise in cortisol that coincides with REM-rich nocturnal sleep, may be important in explaining the differential contribution of REM overnight and SWS during the day for selective emotional memory consolidation. It is important, though, in any future study, to dig down into the different levels of sleep, from gross sleep staging, to spectral composition, to phenomena occurring

during sleep such as sleep spindles, in order to gain a clear picture of the mechanisms of selective consolidation and how they may change with age.

In summary, we demonstrated the importance of obtaining sleep soon after learning for superior selective consolidation of emotional memories, an effect that was robustly seen across young and, for the first time, middle-aged adults. While we found that properties of SWS, such as the relative amount, frequency markers (i.e., delta, slow oscillations) were involved in this preferential preservation of salient information, these relationships were primarily driven by the young adults. Although sleep spindle activity during SWS appeared to be associated with memory for negative information in both young and middle-age adults, this relationship did not reach significance for the middle-age adults, likely due to far fewer middle-age adults producing spindles. The current study serves as a first step to discover whether the functional relationship between memory, particularly memory for important, adaptive experiences, and known mechanisms of sleep physiology-based consolidation persists into the middle age and beyond. Uncovering this relationship is crucial, as it is possible for the future development of interventions to boost properties of SWS, possibly leading to the improvement of memory that normally declines as we age.

Acknowledgments

Research reported in this publication was supported by the National Institute on Aging of the National Institutes of Health under Award Number F32AG047807. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

- Aizawa K, Ageyama N, Yokoyama C, Hisatsune T. Age-dependent alteration in hippocampal neurogenesis correlates with learning performance of macaque monkeys. *Experimental Animals*. 2009; 58(4):403–407. [PubMed: 19654438]
- Alger SE, Lau H, Fishbein W. Delayed onset of a daytime nap facilitates retention of declarative memory. *PloS One*. 2010; 5(8):e12131. [PubMed: 20808821]
- Backhaus J, Born J, Hoeckesfeld R, Fokuhl S, Hohagen F, Junghanns K. Midlife decline in declarative memory consolidation is correlated with a decline in slow wave sleep. *Learning & memory*. 2007; 14(5):336–341. [PubMed: 17522024]
- Baran B, Mantua J, Spencer RM. Age-related changes in the sleep-dependent reorganization of declarative memories. *Journal of Cognitive Neuroscience*. 2016; 28(6):792–802. [PubMed: 26918588]
- Basner M, Dinges DF. An adaptive-duration version of the PVT accurately tracks changes in psychomotor vigilance induced by sleep restriction. *Sleep*. 2012; 35(2):193–202. [PubMed: 22294809]
- Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting and Clinical Psychology*. 1988; 56(6):893–897. doi:<http://dx.doi.org.proxy.library.nd.edu/10.1037/0022-006X.56.6.893>. [PubMed: 3204199]
- Beck AT, Steer RA, Ball R, Ranieri WF. Comparison of Beck depression Inventories–IA and –II in psychiatric outpatients. *Journal of Personality Assessment*. 1996; 67(3):588–597. doi:http://dx.doi.org.proxy.library.nd.edu/10.1207/s15327752jpa6703_13. [PubMed: 8991972]
- Bennion KA, Payne JD, Kensinger EA. Selective effects of sleep on emotional memory: what mechanisms are responsible? *Translational Issues in Psychological Science*. 2015; 1(1):79.
- Buckner RL. Memory and executive function in aging and AD: multiple factors that cause decline and reserve factors that compensate. *Neuron*. 2004; 44(1):195–208. [PubMed: 15450170]

- Burke SN, Barnes CA. Neural plasticity in the ageing brain. *Nature Reviews Neuroscience*. 2006; 7(1): 30–40. [PubMed: 16371948]
- Buyssse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research*. 1989; 28(2):193–213. [PubMed: 2748771]
- Buzsáki G. The hippocampo-neocortical dialogue. *Cerebral Cortex*. 1996; 6(2):81–92. [PubMed: 8670641]
- Carrier J, Viens I, Poirier G, Robillard R, Lafortune M, Vandewalle G, ... Filipini D. Sleep slow wave changes during the middle years of life. *European Journal of Neuroscience*. 2011; 33(4):758–766. [PubMed: 21226772]
- Carskadon MA, Dement WC. Normal human sleep: an overview. *Principles and Practice of Sleep Medicine*. 2005; 4:13–23.
- Cavuto MG, Ong B, Pike KE, Nicholas CL, Bei B, Kinsella GJ. Objective but not subjective sleep predicts memory in community - dwelling older adults. *Journal of Sleep Research*. 2016; 25(4): 475–485. [PubMed: 26868539]
- Cunningham TJ, Crowell CR, Alger SE, Kensinger EA, Villano MA, Mattingly SM, Payne JD. Psychophysiological arousal at encoding leads to reduced reactivity but enhanced emotional memory following sleep. *Neurobiology of Learning and Memory*. 2014; 114:155–164. [PubMed: 24952130]
- De Gennaro L, Ferrara M. Sleep spindles: an overview. *Sleep Med Rev*. 2003; 7:423–440. [PubMed: 14573378]
- Delorme A, Makeig S. EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*. 2004; 134(1):9–21. [PubMed: 15102499]
- Diekelmann S, Born J. The memory function of sleep. *Nature Reviews Neuroscience*. 2010; 11(2): 114–126. [PubMed: 20046194]
- Dijk DJ, Groeger JA, Stanley N, Deacon S. Age-related reduction in daytime sleep propensity and nocturnal slow wave sleep. *Sleep*. 2010; 33:211–223. [PubMed: 20175405]
- Dinges DF, Powell JW. Microcomputer analyses of performance on a portable, simple visual RT task during sustained operations. *Behavior Research Methods*. 1985; 17(6):652–655.
- Dorrian, J., Rogers, NL., Dinges, DF. Psychomotor vigilance performance: a neurocognitive assay sensitive to sleep loss. In: Kushida, C., editor. *Sleep Deprivation: Clinical Issues, Pharmacology and Sleep Loss Effects*. New York: Marcel Dekker, Inc; 2005. p. 39-70.
- Dubé J, Lafortune M, Bedetti C, Bouchard M, Gagnon JF, Doyon J, ... Carrier J. Cortical thinning explains changes in sleep slow waves during adulthood. *Journal of Neuroscience*. 2015; 35(20): 7795–7807. [PubMed: 25995467]
- Duverne S, Motamedinia S, Rugg MD. Effects of age on the neural correlates of retrieval cue processing are modulated by task demands. *Journal of Cognitive Neuroscience*. 2009; 21(1):1–17. [PubMed: 18476757]
- Eschenko O, Ramadan W, Mölle M, Born J, Sara SJ. Sustained increase in hippocampal sharp-wave ripple activity during slow-wave sleep after learning. *Learning & Memory*. 2008; 15(4):222–228. [PubMed: 18385477]
- Ferrarelli F, Huber R, Peterson MJ, Massimini M, Murphy M, Riedner BA, Watson A, Bria P, Tononi G. Reduced sleep spindle activity in schizophrenia patients. *American Journal of Psychiatry*. 2007; 164(3):483–492. [PubMed: 17329474]
- Grady CL, McIntosh AR, Craik FI. Age - related differences in the functional connectivity of the hippocampus during memory encoding. *Hippocampus*. 2003; 13(5):572–586. [PubMed: 12921348]
- Groch S, Wilhelm I, Diekelmann S, Sayk F, Gais S, Born J. Contribution of norepinephrine to emotional memory consolidation during sleep. *Psychoneuroendocrinology*. 2011; 36(9):1342–1350. [PubMed: 21493010]
- Hamann S. Cognitive and neural mechanisms of emotional memory. *Trends in Cognitive Sciences*. 2001; 5(9):394–400. [PubMed: 11520704]

- Hasher, L. Inhibition: An attentional control mechanism. In: Roediger, HL., III, Dudai, Y., Fitzpatrick, SM., editors. *Science of Memory: Concepts*. New York: Oxford University Press; 2007. p. 291–294.
- Helfrich RF, Mander BA, Jagust WJ, Knight RT, Walker MP. Old Brains Come Uncoupled in Sleep: Slow Wave-Spindle Synchrony, Brain Atrophy, and Forgetting. *Neuron*. 2017; 97:1–10.
- Hoddes E, Zarcone V, Smythe H, Phillips R, Dement WC. Quantification of sleepiness: a new approach. *Psychophysiology*. 1973; 10(4):431–436. [PubMed: 4719486]
- Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *International Journal of Chronobiology*. 1976; 4(2):97–110. [PubMed: 1027738]
- Hu P, Stylos-Allan M, Walker MP. Sleep facilitates consolidation of emotional declarative memory. *Psychological Science*. 2006; 17(10):891–898. [PubMed: 17100790]
- Huber R, Ghilardi MF, Massimini M, Tononi G. Local sleep and learning. *Nature*. 2004; 430(6995):78–81. [PubMed: 15184907]
- Kensinger EA, Brierley B, Medford N, Growdon JH, Corkin S. Effects of normal aging and Alzheimer's disease on emotional memory. *Emotion*. 2002; 2(2):118. [PubMed: 12899186]
- Kensinger EA, Anderson A, Growdon JH, Corkin S. Effects of Alzheimer disease on memory for verbal emotional information. *Neuropsychologia*. 2004; 42(6):791–800. [PubMed: 15037057]
- Kensinger EA, Garoff-Eaton RJ, Schacter DL. Effects of emotion on memory specificity: Memory trade-offs elicited by negative visually arousing stimuli. *Journal of Memory and Language*. 2007; 56(4):575–591.
- Kensinger EA. Age differences in memory for arousing and nonarousing emotional words. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2008; 63(1):13–18.
- Knowles JB, MacLean AW, Salem L, Vetere C, Coulter M. Slow-wave sleep in daytime and nocturnal sleep: An estimate of the time course of "Process S". *Journal of Biological Rhythms*. 1986; 1(4):303–308. [PubMed: 2979592]
- Ladenbauer J, Külzow N, Passmann S, Antonenko D, Grittner U, Tamm S, Flöel A. Brain stimulation during an afternoon nap boosts slow oscillatory activity and memory consolidation in older adults. *Neuroimage*. 2016; 142:311–323. [PubMed: 27381076]
- Ladenbauer J, Ladenbauer J, Külzow N, de Boer R, Avramova E, Grittner U, Flöel A. Promoting sleep oscillations and their functional coupling by transcranial stimulation enhances memory consolidation in mild cognitive impairment. *Journal of Neuroscience*. 2017; 37(30):7111–7124. [PubMed: 28637840]
- Lewis PA, Durrant SJ. Overlapping memory replay during sleep builds cognitive schemata. *Trends in cognitive sciences*. 2011; 15(8):343–351. [PubMed: 21764357]
- Lo JC, Groeger JA, Cheng GH, Dijk DJ, Chee MW. Self-reported sleep duration and cognitive performance in older adults: a systematic review and meta-analysis. *Sleep medicine*. 2016; 17:87–98. [PubMed: 26847980]
- Mander BA, Rao V, Lu B, Saletin JM, Lindquist JR, Ancoli-Israel S, Jagust W, Walker MP. Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. *Nature Neuroscience*. 2013; 16:357–364. [PubMed: 23354332]
- Mander BA, Rao V, Lu B, Saletin JM, Ancoli-Israel S, Jagust WJ, Walker MP. Impaired prefrontal sleep spindle regulation of hippocampal-dependent learning in older adults. *Cerebral Cortex*. 2014; 24:3301–3309. [PubMed: 23901074]
- Mander BA, Winer JR, Walker MP. Sleep and Human Aging. *Neuron*. 2017; 94(1):19–36. [PubMed: 28384471]
- Martin N, Lafortune M, Godbout J, Barakat M, Robillard R, Poirier G, ... Carrier J. Topography of age-related changes in sleep spindles. *Neurobiology of Aging*. 2012; 34(2):468–76. [PubMed: 22809452]
- Massimini M, Huber R, Ferrarelli F, Hill S, Tononi G. The sleep slow oscillation as a traveling wave. *Journal of Neuroscience*. 2004; 24(31):6862–6870. [PubMed: 15295020]
- Mather, M. Why memories may become more positive as people age. In: Uttl, B, Ohta, N., Siegenthaler, AL., editors. *Memory and Emotion: Interdisciplinary Perspectives*. Malden, MA: Blackwell Press; 2006. p. 135–159.

- Mather M, Carstensen LL. Aging and attentional biases for emotional faces. *Psychological Science*. 2003; 14(5):409–415. [PubMed: 12930469]
- Mather M, Carstensen LL. Aging and motivated cognition: The positivity effect in attention and memory. *Trends in Cognitive Sciences*. 2005; 9(10):496–502. [PubMed: 16154382]
- Mednick SC, McDevitt EA, Walsh JK, Wamsley E, Paulus M, Kanady JC, Drummond SP. The critical role of sleep spindles in hippocampal-dependent memory: a pharmacology study. *Journal of Neuroscience*. 2013; 33(10):4494–4504. [PubMed: 23467365]
- Morcom AM, Good CD, Frackowiak RS, Rugg MD. Age effects on the neural correlates of successful memory encoding. *Brain*. 2003; 126(1):213–229. [PubMed: 12477708]
- Nicolas A, Petit D, Rompre S, Montplaisir J. Sleep spindle characteristics in healthy subjects of different age groups. *Clinical Neurophysiology*. 2001; 112(3):521–527. [PubMed: 11222974]
- Nishida M, Pearsall J, Buckner RL, Walker MP. REM sleep, prefrontal theta, and the consolidation of human emotional memory. *Cerebral Cortex*. 2008; 19(5):1158–1166. [PubMed: 18832332]
- Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV. Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep*. 2004; 27(7):1255–1273. [PubMed: 15586779]
- Pace-Schott EF, Shepherd E, Spencer RM, Marcello M, Tucker M, Propper RE, Stickgold R. Napping promotes inter-session habituation to emotional stimuli. *Neurobiology of Learning and Memory*. 2011; 95(1):24–36. [PubMed: 20969968]
- Pace-Schott EF, Spencer RM. Age-related changes in consolidation of perceptual and muscle-based learning of motor skills. *Frontiers in Aging Neuroscience*. 2013; 5
- Pavlidis C, Winson J. Influences of hippocampal place cell firing in the awake state on the activity of these cells during subsequent sleep episodes. *Journal of Neuroscience*. 1989; 9(8):2907–2918. [PubMed: 2769370]
- Payne JD, Kensinger EA. Sleep's role in the consolidation of emotional episodic memories. *Current Directions in Psychological Science*. 2010; 19(5):290–295.
- Payne JD, Kensinger EA. Sleep leads to changes in the emotional memory trace: evidence from fMRI. *Journal of Cognitive Neuroscience*. 2011; 23(6):1285–1297. [PubMed: 20521852]
- Payne JD, Stickgold R, Swanberg K, Kensinger EA. Sleep preferentially enhances memory for emotional components of scenes. *Psychological Science*. 2008a; 19(8):781–788. [PubMed: 18816285]
- Payne, JD., Ellenbogen, JM., Walker, MP., Stickgold, R. *Learning and Memory: A comprehensive reference*. New York: Elsevier; 2008b. The role of sleep in memory consolidation.
- Payne JD, Chambers AM, Kensinger EA. Sleep promotes lasting changes in selective memory for emotional scenes. *Frontiers in Integrative Neuroscience*. 2012; 6
- Payne JD, Kensinger EA, Wamsley EJ, Spreng RN, Alger SE, Gibler K, ... Stickgold R. Napping and the selective consolidation of negative aspects of scenes. *Emotion*. 2015; 15(2):176. [PubMed: 25706830]
- Rajah MN, Maillet D, Grady CL. Episodic memory in healthy older adults. *The Wiley Handbook on the Cognitive Neuroscience of Memory*. 2015:347–370.
- Raz N, Gunning F, Head D, Dupuis JH, McQuain J, Briggs SD, ... Acker JD. Selective aging of the human cerebral cortex observed in vivo: differential vulnerability of the prefrontal gray matter. *Cerebral Cortex*. 1997; 7(3):268–282. [PubMed: 9143446]
- Raz N, Ghisletta P, Rodrigue KM, Kennedy KM, Lindenberger U. Trajectories of brain aging in middle-aged and older adults: regional and individual differences. *Neuroimage*. 2010; 51(2):501–511. [PubMed: 20298790]
- Rechtschaffen A, Kales A. A manual of standardized terminology, techniques, and scoring systems for sleep stages of human subjects. 1968
- Schabus M, Dang-Vu TT, Albouy G, Balet E, Boly M, Carrier J, ... Phillips C. Hemodynamic cerebral correlates of sleep spindles during human non-rapid eye movement sleep. *Proceedings of the National Academy of Sciences*. 2007; 104(32):13164–13169.
- Scullin MK. Sleep, memory, and aging: the link between slow-wave sleep and episodic memory changes from younger to older adults. *Psychology and Aging*. 2013; 28(1):105. [PubMed: 22708533]

- Siapas AG, Wilson MA. Coordinated interactions between hippocampal ripples and cortical spindles during slow-wave sleep. *Neuron*. 1998; 21(5):1123–1128. [PubMed: 9856467]
- Sirota A, Csicsvari J, Buhl D, Buzsáki G. Communication between neocortex and hippocampus during sleep in rodents. *Proceedings of the National Academy of Sciences*. 2003; 100(4):2065–2069.
- Snodgrass JG, Corwin J. Pragmatics of measuring recognition memory: applications to dementia and amnesia. *Journal of Experimental Psychology: General*. 1988; 117(1):34. [PubMed: 2966230]
- Spielberger, CD. State - Trait anxiety inventory. John Wiley & Sons, Inc; 2010.
- Steinmetz KRM, Kensinger EA. The emotion-induced memory trade-off: More than an effect of overt attention? *Memory & Cognition*. 2013; 41(1):69–81. [PubMed: 22948959]
- Stickgold R, Walker MP. Sleep-dependent memory triage: evolving generalization through selective processing. *Nature Neuroscience*. 2013; 16(2):139–145. [PubMed: 23354387]
- Talamini LM, Nieuwenhuis IL, Takashima A, Jensen O. Sleep directly following learning benefits consolidation of spatial associative memory. *Learning & Memory*. 2008; 15(4):233–237. [PubMed: 18391183]
- Tamminen J, Payne JD, Stickgold R, Wamsley EJ, Gaskell MG. Sleep spindle activity is associated with the integration of new memories and existing knowledge. *Journal of Neuroscience*. 2010; 30(43):14356–14360. [PubMed: 20980591]
- Tononi G, Cirelli C. Sleep and synaptic homeostasis: a hypothesis. *Brain Research Bulletin*. 2003; 62(2):143–150. [PubMed: 14638388]
- Tononi G, Cirelli C. Sleep function and synaptic homeostasis. *Sleep Medicine Reviews*. 2006; 10(1):49–62. [PubMed: 16376591]
- Van Cauter E, Leproult R, Plat L. Age-related changes in slow wave sleep and REM sleep and relationship with growth hormone and cortisol levels in healthy men. *JAMA*. 2000; 284:861–868. [PubMed: 10938176]
- Van Der Werf YD, Van Der Helm E, Schoonheim MM, Ridderikhoff A, Van Someren EJ. Learning by observation requires an early sleep window. *Proceedings of the National Academy of Sciences*. 2009; 106(45):18926–18930.
- Varga AW, Ducca EL, Kishi A, Fischer E, Parekh A, Koushyk V, ... Burschtin OE. Effects of aging on slow-wave sleep dynamics and human spatial navigational memory consolidation. *Neurobiology of Aging*. 2016; 42:142–149. [PubMed: 27143431]
- Verhaeghen P, Salthouse TA. Meta-analyses of age–cognition relations in adulthood: Estimates of linear and nonlinear age effects and structural models. *Psychological bulletin*. 1997; 122(3):231. [PubMed: 9354147]
- Wagner U, Gais S, Born J. Emotional memory formation is enhanced across sleep intervals with high amounts of rapid eye movement sleep. *Learning & Memory*. 2001; 8(2):112–119. [PubMed: 11274257]
- Waring JD, Kensinger EA. Effects of emotional valence and arousal upon memory trade-offs with aging. *Psychology and Aging*. 2009; 24(2):412. [PubMed: 19485658]
- Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of Personality and Social Psychology*. 1988; 54(6):1063. [PubMed: 3397865]
- Westerberg CE, Florczak SM, Weintraub S, Mesulam MM, Marshall L, Zee PC, Paller KA. Memory improvement via slow-oscillatory stimulation during sleep in older adults. *Neurobiology of Aging*. 2015; 36(9):2577–2586. [PubMed: 26116933]
- Wilson JK, Baran B, Pace-Schott EF, Ivry RB, Spencer RM. Sleep modulates word-pair learning but not motor sequence learning in healthy older adults. *Neurobiology of Aging*. 2012; 33(5):991–1000. [PubMed: 22244090]

Highlights

- Young and middle age cohorts prioritize emotional memory at the expense of less important details.
- Emotional memory is preferentially preserved similarly for young and middle-age adults during a nap.
- Properties of slow wave sleep were related specifically to memory for salient information.
- Some relationships between aspects of sleep and selective memory appears to be altered with age.
- Sleep spindles were associated with selective consolidation of salient memories across the age range.

Figure 1. Protocol
All participants were trained and tested at the same time of day as one another, with encoding of the images at 10am, followed by baseline testing, and retest occurring at 5pm. Nap groups had a 90-minute nap opportunity either at 11am (immediate nap condition) or 3pm (delayed nap condition).

Encoding Session

Testing Sessions

Figure 2. Memory Trade-off Task

Participants viewed complex scenes comprising either negative (i.e., taxi cab accident) or neutral foreground images placed on plausible neutral backgrounds. The images were presented for 3000ms, followed by self-paced assessments of valence and arousal. Baseline testing occurred immediate after encoding, with half of the encoded objects and backgrounds, now presented individually and one at a time, intermixed with an equal amount of foils. Participants were required to decide if the images were ones they hadn't seen before (i.e., "new") or ones they had seen before (i.e., "old"), using the keys 0 and 1, respectively, to indicate their choice. Retest occurred approximately 7 hours after encoding in the same manner.

Figure 3. Emotional Memory Trade-off Effect Across Time by Age Cohort

The emotional memory trade-off score is the difference in negative object memory and related background memory, with a larger trade-off score representing a bigger difference in these memories. The figure above represents the increase in the magnitude of this trade-off score across the retention period for each condition for both the young adults (left) and middle-aged adults (right). The increase in emotional memory trade-off in the immediate nap condition was significantly greater than both the wake (young: $t_{28} = -3.14$, $p = 0.004$; middle age: $t_{21} = -2.54$, $p = 0.02$) and delayed nap (young: $t_{28} = 2.52$, $p = 0.02$; middle age: $t_{21} = 2.17$, $p = 0.04$) conditions.

Figure 4. Correlation Between SWS and Negative Object Memory

The percentage of SWS was positively correlated with memory for the negative objects at retest ($r_{53} = 0.35$, $p = 0.01$ across subjects (Young = squares, Middle Age = triangles). This replicates the relationship we previously found in young adults (Payne et al., 2015). When we removed those who did not reach SWS from the analyses, the correlation remained significant ($r_{35} = 0.34$, $p = 0.05$).

Figure 5. Correlations Between Negative Object Memory and Features of Sleep Spindles

Negative object memory was positively correlated with features of sleep spindles for young and middle-age adults. Spindle measurements were transformed into z-scores within each age cohort in order to represent the associations on the same scale across cohorts. Strength of associations was similar between age cohorts. The associations were significant within the young adults. A) Total number of spindles: Young, $r = 0.47, p = 0.02$; Middle Age, $r = 0.54, p = 0.13$, B) Spindle duration: Young, $r = 0.53, p = 0.008$; Middle Age, $r = 0.60, p = 0.12$, C) Spindle amplitude: Young $r = 0.54, p = 0.006$; Middle Age, $r = 0.49, p = 0.19$, D) Spindle power: Young, $r = 0.52, p = 0.01$; Middle Age, $r = 0.55, p = 0.12$.

Table 1

Behavioral Performance

Condition	Baseline (B)		Retest (R)		Change (R-B)		Trade-off Change (R-B)
	OB	BG	OB	BG	OB	BG	
Young Adults							
<i>Neutral Performance</i>							
<i>Wake</i>	0.75 ± 0.03	0.66 ± 0.04	0.56 ± 0.04	0.43 ± 0.04	-0.19 ± 0.03	-0.24 ± 0.02	0.05 ± 0.04
<i>Immediate</i>	0.79 ± 0.03	0.72 ± 0.03	0.65 ± 0.04	0.47 ± 0.04	-0.15 ± 0.03	-0.25 ± 0.04	0.10 ± 0.05
<i>Delayed</i>	0.78 ± 0.02	0.69 ± 0.04	0.61 ± 0.04	0.42 ± 0.05	-0.17 ± 0.03	-0.28 ± 0.04	0.10 ± 0.05
<i>Negative Performance</i>							
<i>Wake</i>	0.95 ± 0.01	0.47 ± 0.05	0.77 ± 0.04*	0.27 ± 0.03	-0.18 ± 0.03**	-0.21 ± 0.05*	0.03 ± 0.07**
<i>Immediate</i>	0.91 ± 0.02	0.59 ± 0.03	0.87 ± 0.02	0.25 ± 0.04	-0.04 ± 0.03	-0.34 ± 0.04	0.31 ± 0.05
<i>Delayed</i>	0.92 ± 0.02	0.58 ± 0.04	0.80 ± 0.03	0.31 ± 0.04	-0.13 ± 0.03*	-0.27 ± 0.04	0.14 ± 0.04*
Middle Age Adults							
<i>Neutral Performance</i>							
<i>Wake</i>	0.72 ± 0.03	0.63 ± 0.04	0.51 ± 0.05	0.38 ± 0.05	-0.21 ± 0.03	-0.26 ± 0.05	0.05 ± 0.05
<i>Immediate</i>	0.77 ± 0.03	0.60 ± 0.06	0.58 ± 0.06	0.39 ± 0.06	-0.19 ± 0.05	-0.21 ± 0.05	0.02 ± 0.06
<i>Delayed</i>	0.73 ± 0.04	0.56 ± 0.05	0.62 ± 0.04	0.36 ± 0.06	-0.11 ± 0.02	-0.21 ± 0.05	0.10 ± 0.05
<i>Negative Performance</i>							
<i>Wake</i>	0.87 ± 0.08	0.46 ± 0.06	0.67 ± 0.05#	0.24 ± 0.03	-0.20 ± 0.04*	-0.21 ± 0.05#	0.01 ± 0.08*
<i>Immediate</i>	0.88 ± 0.03	0.53 ± 0.07	0.79 ± 0.04	0.16 ± 0.05	-0.09 ± 0.04	-0.37 ± 0.06	0.28 ± 0.07
<i>Delayed</i>	0.84 ± 0.01	0.40 ± 0.05	0.75 ± 0.03	0.22 ± 0.05	-0.10 ± 0.02	-0.18 ± 0.04*	0.09 ± 0.06*
Mean ± SEM							
OB = Object memory							
BG = Background memory							
# p < 0.10							
* p < 0.05							

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

5005
d > 0.005
**

All noted significant comparisons are between the Immediate Nap condition and either the Wake or Delayed Nap condition, with notation adjacent to one of the latter conditions.
Trade-off Change was calculated by subtracting the difference between object and background memory at baseline from the difference between object and background memory at retest

Table 2

Sleep parameters for all conditions

	Young Adults			Middle Age Adults		
	<i>Immediate Nap</i>	<i>Delayed Nap</i>	<i>p-value</i>	<i>Immediate Nap</i>	<i>Delayed Nap</i>	<i>p-value</i>
<i>TST</i>	67.80 ± 3.56	68.00 ± 5.97	0.98	56.50 ± 6.34	47.88 ± 8.41	0.34
<i>Onset Latency</i>	10.27 ± 1.78	8.89 ± 2.55	0.73	15.38 ± 4.59	9.42 ± 2.34	0.17
<i>REM latency</i>	50.05 ± 5.16	57.17 ± 7.86	0.41	59.33 ± 7.76	61.90 ± 4.37	0.82
<i>WASO</i>	15.10 ± 2.98	13.68 ± 3.80	0.80	24.42 ± 3.92	33.96 ± 5.86	0.13
<i>Efficiency</i>	72.89 ± 3.84	75.06 ± 6.47	0.80	58.33 ± 6.39	52.38 ± 8.87	0.53
<i>Stage 1 %</i>	17.13 ± 1.37	16.77 ± 3.38	0.95	28.48 ± 4.83	35.73 ± 5.58	0.21
<i>Stage 2 %</i>	41.39 ± 3.86	44.16 ± 4.94	0.65	53.58 ± 3.96	48.52 ± 5.07	0.45
<i>SWS %</i>	29.19 ± 4.33	32.13 ± 5.84	0.66	10.35 ± 3.94	8.49 ± 5.24	0.80
<i>REM %</i>	12.28 ± 3.28	6.94 ± 1.87	0.16	7.58 ± 3.14	5.76 ± 2.37	0.66
Mean ± SEM						

Table 3

Comparison of Young and Middle Age Sleep

	Young Adults	Middle Age Adults	p-value
<i>TST</i>	67.90 ± 3.36	52.19 ± 5.23	0.02
<i>Onset Latency</i>	9.60 ± 1.51	12.40 ± 2.60	0.34
<i>REM latency</i>	53.42 ± 4.55	60.50 ± 4.48	0.31
<i>WASO</i>	14.41 ± 2.36	29.19 ± 3.59	0.001
<i>Efficiency</i>	73.94 ± 3.64	55.36 ± 5.38	0.007
<i>Stage 1 %</i>	16.96 ± 1.75	32.11 ± 3.69	0.001
<i>Stage 2 %</i>	42.73 ± 3.06	51.05 ± 3.19	0.07
<i>SWS %</i>	30.61 ± 3.54	9.42 ± 3.21	< 0.001
<i>REM %</i>	9.70 ± 1.96	6.67 ± 1.93	0.28

Mean ± SEM