Published in final edited form as:

J Med Chem. 2008 February 14; 51(3): 603–611. doi:10.1021/jm070910s.

Binding of serotonin and N₁-benzenesulfonyltryptamine-related analogs at human 5-HT₆ serotonin receptors: Receptor modeling studies

Małgorzata Dukat¹, Philip D. Mosier¹, Renata Kolanos¹, Bryan L. Roth², and Richard A. Glennon^{1,*}

¹Department of Medicinal Chemistry, School of Pharmacy, Medical College of Virginia Campus, Virginia Commonwealth University, Richmond, VA 23298-0540

²Departments of Pharmacology, Psychiatry, Division of Medicinal Chemistry and Natural Products, Neurosciences Center and Comprehensive Cancer Center Schools of Medicine and Pharmacy; University of North Carolina, Chapel Hill, North Carolina 27599 USA

Abstract

A population of 100 graphics models of the human 5-HT₆ serotonin receptor was constructed based on the structure of bovine rhodopsin. The endogenous tryptamine-based agonist serotonin (5-HT; **1**) and the benzenesulfonyl-containing tryptamine-derived 5-HT₆ receptor antagonist MS-245 (**4a**) were automatically docked with each of the 100 receptor models using a genetic algorithm approach. Similar studies were conducted with the more selective 5-HT₆ receptor agonist EMDT (**5**) and optical isomers of EMDT-related analog **8**, as well as with optical isomers of MS-245 (**4a**)-related and benzenesulfonyl-containing pyrrolidine **6** and aminotetralin **7**. Although associated with the same general aromatic/hydrophobic binding cluster, 5-HT (**1**) and MS-245 (**4a**) were found to preferentially bind with distinct receptor conformations, and did so with different binding orientations (*i.e.*, poses). A 5-HT pose/model was found to be common to EMDT (**5**) and its analogs, whereas that identified for MS-245 (**4a**) was found common to benzenesulfonyl-containing compounds. Specific amino acid residues were identified that can participate in binding, and evaluation of a sulfenamide analog of MS-245 indicates for the first time that the presence of the sulfonyl oxygen atoms enhances receptor affinity. The results indicate that the presence or absence of an N₁-benzenesulfonyl group is a major determinant of the manner in which tryptamine-related agents bind at 5-HT₆ serotonin receptors.

TABLE OF CONTENTS GRAPHIC

Keywords

Serotonin; 5-Hydroxytryptamine (5-HT); 5-HT₆ serotonin receptors; EMDT; MS-245; Benzenesulfonyltryptamines

Many actions of the neurotransmitter substance serotonin (5-hydroxytryptamine; 5-HT, **1**) are mediated by its interaction with various populations (5-HT₁ – 5-HT₇) of 5-HT receptors.¹ Apart from ionotropic 5-HT₃ receptors, other members of this receptor family are Class A rhodopsin-like G protein-coupled receptors (GPCRs).^{1,2} Human 5-HT₆ receptors, first cloned in 1996,³ are positively coupled to an adenylate cyclase second messenger system, and there is evidence that this receptor population is involved in cognition, obesity, and certain neurological and neuropsychiatric disorders including schizophrenia and depression.⁴ This receptor population is also of interest because clozapine and a number of other typical and atypical antipsychotic agents bind at 5-HT₆ receptors with high affinity.⁵ Only within the last six or seven years have 5-HT₆-selective agents been identified, and among some of the first antagonists described were Ro 04-6790 (**2**),⁶ SB-258510 (**3a**) and its *N*-desmethyl analog SB-271046 (**3b**),⁷ MS-245 (**4a**),^{8,9} and the agonist EMDT (**5**)⁸ (Chart 1) (reviewed^{4,10}).

Prior to the availability of selective agents, site directed mutagenesis studies with rat 5-HT₆ receptors identified conserved aspartate (Asp106 [3.32])^{*} and threonine (Thr196 [5.46]) moieties as being directly or indirectly involved in the actions of serotonergic agonists, and that their individual mutation substantially reduces both the affinity (radioligand binding) and efficacy (stimulation of adenylate cyclase action) of the nonselective agonists 5-HT (**1**) and (+)lysergic acid diethylamide (LSD).^{12,13}

^{*}There is a high degree of homology between rat and human 5-HT₆ receptors and their amino acid sequence numbering is identical for the amino acids described herein. For convenience, amino acids are numbered by their human sequence number followed, in brackets, by their numbering using the Ballesteros-Weinstein convention.¹¹

To date, there have been four attempts to provide insight into how 5-HT₆ receptor ligands interact with 5-HT₆ receptors using graphical models.^{14–17} Bromidge¹⁴ was first to describe a possible docking mode for a 5-HT₆ antagonist (i.e., SB-258510; **3**) to a 5-HT₆ receptor model and his epigrammatic description implicated Phe277 [6.44] and Trp281 [6.48] as participating in possible π - π stacking interactions (Table 1). Hirst et al.¹⁶ examined the docking of several antagonists including Ro 04-6790 (**2**) and SB-258510 (**3**) at mouse, rat, and human 5-HT₆ receptor models to explain the reduced affinity of certain antagonists (in particular Ro 04-6790) for mouse receptors. Implicated in possible π - π stacking interactions (rat model) were Phe188 [5.38], Trp281 [6.48], Phe284 [6.51], and Phe302 [7.35]. Pullagurla et al.,¹⁷ using a human 5-HT₆ receptor model, examined several MS-245 (**4a**) analogs and identified Phe284 [6.51], Phe285 [6.52], and Phe302 [7.35] as being involved in π - π interactions. More recently, Lopez-Rodriguez et al.¹⁵ studied the binding of a series of 5-HT₆ antagonists (including **2**, **3b** and **4**) using a human 5-HT₆ receptor model. An aromatic/ hydrophobic binding region was defined by Val107 [3.33], Phe188 [5.38], Ala192 [5.42], Phe198 [5.48], and Phe285 [6.52]. Despite using different approaches, each of the four studies identified certain common amino acids as possibly participating in the interaction of antagonists with 5-HT₆ receptors. That is, the studies all began with recognition that a role for Asp106 [3.32] is key to binding, and identified a common binding pocket or “cluster” consisting of aromatic/hydrophobic residues in TM5 and TM6 (e.g. Trp281, Phe284 and/or Phe285). However, orientation of the ligands *within* this region differed considerably. For example, the SO₂ oxygen atoms of the sulfonyl-containing antagonists have been variously suggested to be within hydrogen bonding distance to, for example, Asn288 [6.55], Ser193 [5.43], Leu182 (in the e2 loop), and Gln291 (at the TM6-e3 junction) (see Table 1).^{14–15}

Two investigations examined agonists,^{16,17} and only one of these examined the possible binding modes of 5-HT at human 5-HT₆ receptors.¹⁷ The two models implicated a role for Asp106 [3.32] and Thr196 [5.46] for interaction with the agonists 5-HT and/or LSD.^{16,17}

Using graphics models of the receptor, attempts have been made to identify how 5-HT₆ receptor antagonist ligands interact with various receptor amino acid features. For the most part, however, these studies involved antagonists with multiple basic amine groups (with consequent and inherent uncertainty, and assumptions, as to which amine specifically interacts with the TM3 aspartate moiety Asp106), employed agents with substantial conformational flexibility, and have assumed that the sulfonyl moiety (common to nearly all 5-HT₆ antagonists) participates in binding. From what we can discern, only one of the prior studies explicitly states that a non-manual docking method was utilized,¹⁷ and only a single study examined a possible binding mode for 5-HT.¹⁷

The present investigation was designed to specifically examine the binding of tryptamine-related analogs; the binding of other 5-HT₆ ligands will be examined in due course. In this investigation we examined possible docking modes for the nonselective agonist 5-HT (**1**) and of the more 5-HT₆-selective agonist EMDT (**5**) at human 5-HT₆ receptors. We also examined the docking of the tryptamine-derived antagonist MS-245 (**4a**) and structurally-related, and stereochemically- and/or conformationally-defined benzenesulfonyl analog pyrrolidine **6** and aminotetralin **7**. In light of empirical evidence that N₁-unsubstituted and

N_1 -benzenesulfonyltryptamines likely bind in a dissimilar manner at 5-HT₆ receptors,^{18,19} and with realization that agonists and antagonists might bind optimally with different receptor conformations (and that there might even be differences in the binding of different agonists at a given receptor),²⁰ we utilized a docking technique that should emphasize and account for such differences. Finally, we examined the binding of stereochemically-defined optical isomers of EMDT analog **8**, and also determined whether or not the sulfonyl oxygen atom(s) contribute to binding of benzenesulfonyltryptamines by examining sulfenamide **9** in comparison with its sulfonyl counterpart **4b**. As such, this investigation employed *a*) 5-HT₆ receptor ligands bearing a single basic amine, *b*) stereochemically- or (somewhat) conformationally-defined analogs of the 5-HT₆ ligands MS-245 (**4a**) and EMDT (**5**), and *c*) an automated and unbiased docking procedure that might account for differences in the binding of agonist *versus* antagonist analogs at human 5-HT₆ receptors.

Chemistry

The synthesis of certain compounds required for this investigation has been previously reported by us, including MS-245 (**4a**),⁸ **4b**,²¹ EMDT (**5**),⁸ *R*(+)**6**, and *S*(-)**6**.²² The synthesis of compounds **7R** and **7S** employed the isomers of **11** (Scheme 1). Isomers **11** were prepared in five steps from 5-amino-2-naphthol (**10**) following a literature procedure²³ that involved protection of the amine, and Birch reduction, and stereoselective synthesis of **11R** and **11S** from the resultant ketones. Dimethylation of the individual optical isomers of **11** afforded isomers **12** which were deprotected to isomers **13** and then sulfonylated to isomers **7** by reaction with benzenesulfonyl chloride (Scheme 1). Compounds **12** and **13** have been previously reported in the patent literature as their racemates;²³ although optical isomers of **13** also were mentioned, synthesis was not described nor were physicochemical properties recorded.²³

Isomers **8** were prepared from 2-ethyl-5-methoxyindole²⁴ in a manner that paralleled the synthesis of **6**,²² and related compounds.²⁵ Compound **9** was obtained by interaction of *N,N*-dimethyltryptamine (DMT) with *N*-(phenylthio)succinimide.²⁶

Results and Discussion

Human 5-HT₆ receptor affinities (K_i values) of certain compounds employed in this investigation already have been reported from our laboratories: MS-245 and its des-methoxy counterpart (**4a** and **4b**; K_i = 2.1 and 4.1 nM, respectively),²¹ EMDT (**5**; K_i = 16 nM),⁸ *R*(+)**6** (K_i = 0.3 nM), and *S*(-)**6** (K_i = 1.7 nM).²² Aminotetralin isomers **7R** and **7S** displayed little enantioselectivity (K_i = 49 ± 12 and 90 ± 20 nM, respectively), whereas isomers of the pyrrolidine counterpart of EMDT showed greater stereoselectivity of binding (**8R** K_i = 1.8 ± 0.2 nM; **8S** K_i = 220 ± 25 nM). For several structurally related chiral pyrrolidines, we have previously found greater enantioselectivity in the absence of the benzenesulfonyl group (up to 70-fold) than in its presence (<6-fold).²² Sulfenamide **9** (K_i = 90 ± 10 nM) displayed >20-fold reduced affinity relative to its sulfonamide counterpart **4b** (K_i = 4.1 nM).

Model construction and docking studies

Human 5-HT₆ receptor models were constructed and various docking solutions were explored for 5-HT (**1**), MS-245 (**4a**), EMDT (**5**), MS-245 analogs **6**, aminotetralins **7**, and EMDT-related optical isomers **8** such that both the receptor models and the ligands were allowed to interact in a fairly flexible manner. Receptor flexibility was explicitly addressed *via* the generation of a population of 100 h5-HT₆ receptor models whose conformational state varied among the members of the population. Automated docking routines were then used to flexibly place each ligand into each receptor model. The resulting receptor-ligand complexes were scored and subsequently ranked using the ChemScore^{27,28} fitness function. Thus, each ligand was allowed to select an optimal receptor conformation from the population, reducing the bias that might occur when using a single receptor model to dock different ligands. Two common receptor conformations were identified: one for agents lacking a benzenesulfonyl group (*i.e.*, 5-HT, EMDT and analogs), and another for agents possessing such a group (*i.e.*, MS-245 and related analogs). The top-scoring receptor conformation for each nonbenzenesulfonyl analog was identical. The top-scoring receptor conformation for benzenesulfonyl-containing analogs also was the same, except that for MS-245 (**4a**) the second highest top-scoring receptor conformation was selected. The latter (ChemScore = 39.32 relative to the top score of 41.20) was selected because of its commonality with the receptor conformation identified for the other benzenesulfonyl-containing analogs. In the top-scoring MS-245 receptor conformation, the hydrogen bonding component suffers at the expense of a greater lipophilic component, reflecting a somewhat different pose for the benzenesulfonyl moiety.

The primary differences in the two receptor models are depicted in Figure 1. The largest difference in side chain orientation occurs for Trp281 [6.48]. The phenyl portion of the indole nucleus of Trp281 is oriented toward transmembrane helix 5 (TM5) for the non-benzenesulfonyl-containing compounds, and toward transmembrane helix 3 (TM3) for benzenesulfonyl-containing compounds. In addition, residues comprising the second extracellular loop (e2) also exhibited a significantly different position and orientation between the two receptor models. This is not surprising since the overall movement of the e2 loop is expected to be greater than that for the transmembrane helical segments. A complete list of RMSD differences for individual residues of the binding cavity may be found in the Supporting Information section. The two top receptor solutions are described in more detail in the following sections as they relate to docked ligands.

5-HT and EMDT analogs

With the amine function docked within 2.7 Å of Asp106 [3.32], the indolic nucleus of 5-HT is situated in a cluster of amino acids consisting of Leu182 [e2], Trp281 [6.48], and Phe284 [6.51] (Figure 2). Previous models for docked 5-HT¹⁷ and LSD¹⁶ suggest that the indolic NH might hydrogen bond with Thr196 [5.46]. In the present model, the indolic NH is directed towards Arg181 [e2] which is <4 Å distant. This virtually turns the indolic nucleus “up-side down” relative to a prior binding motif described by us.¹⁷ Both Ser111 [3.37] and Thr196 [5.46] (Figure 2) are within hydrogen bond distance of the 5-HT hydroxyl group (O-O distance 3.0 – 3.1 Å) indicating that the hydroxyl group can form a hydrogen bond with

either one or, simultaneously, with both amino acids. This is consistent with the observation that removal of the 5-hydroxyl group of 5-HT results in several-fold decreased affinity.³⁰ These hydrogen bond interactions were stable during a 100-ps dynamics run (see Experimental section). Similar results were obtained with EMDT (**5**) except that the methoxy oxygen atom is a little closer (2.6 Å) to Ser111 [3.37] and Thr196 [5.46].

Both optical isomers of EMDT analog **8** were also docked in a similar manner. Of the one hundred docking solutions, the top solution (receptor conformation) for both isomers was the same docking model identified for 5-HT (**1**) and EMDT (**5**). The primary difference in docking modes for **8R** and **8S** is that the *N*-methyl group of **8R** was directed toward the extracellular region (opening) of the binding cavity, whereas for **8S** the *N*-methyl group faced the intracellular region (bottom) of the binding cavity. The pose distinctions for the isomers of **8** are reflected in their ChemScore values of 36.96 for **8R** and 35.13 for **8S**. The ChemScore is larger for the higher-affinity **8R**, which is consistent with the observed binding affinities for the two isomers. While the scores are similar in magnitude, there are some important but subtle differences in the components that contribute to the total ChemScore values. Analysis of the individual ChemScore components indicated that the hydrogen bond component was higher for the **R** isomer (2.62) than for the **S** (2.40). This suggests that the ammonium group of **8R** is able to form a stronger ionic/hydrogen bond with Asp106 than can the ammonium group of **8S**. Other differences in the ChemScore components occurred for the clash (**8R** = 3.84; **8S** = 3.30) and internal torsion (**8R** = 4.12; **8S** = 5.77) penalty terms. The clash penalty term, which is a measure of undesirable van der Waals overlap between atoms of the ligand and those of the receptor, was higher for **8R**, whereas the internal torsion penalty, which is a measure of the deviation from ideality of the rotatable bonds (*i.e.* gauche vs. eclipsed substituents), was higher for **8S**. Analyses of molecular dynamics simulation data in order to explore more fully the significance of these terms as they relate to the binding modes of the two isomers are ongoing.

MS-245 and analogs

A common binding model was identified for MS-245 (**4a**), pyrrolidine **6R**, and aminotetralin **7R** (as illustrated in Figure 3 for MS-245). With the basic amine of each ligand within 2.7 Å of Asp106, their aryl moieties are situated in essentially the same hydrophobic pocket described above for 5-HT (**1**); however, they are positioned differently than 5-HT. For example, the benzenesulfonyl aryl groups (see Aryl' in Table 1) are within 3.5 to 4.0 Å of Ala157, Ala192, and Phe285. Furthermore, the sulfonyl oxygen atoms are within hydrogen bond distance (O-O distance: 2.8 – 3.3 Å) to both Ser111 [3.37] and Thr196 [5.46], except for **6R** which is located within 3.0 Å of Ser111 and 3.8 Å away from Thr196. The model indicates that the sulfonyl oxygen atoms should be able to participate in hydrogen bond formation with at least one, if not both, of these amino acids. That the model is common to MS-245 (**4a**) and stereochemically- and/or conformationally-defined analogs **6R** and **7R** is telling, and suggests that the conformation of MS-245 shown in Figure 3 reflects a conformation optimal for binding. Furthermore, as shown in Figure 3, the benzenesulfonyl group is nearly 90° out of the plane of the indole nucleus; this is a conformation we had previously predicted would be optimal for binding on the basis of studies with conformationally-constrained benzenesulfonyl analogs.¹⁸

The current model, like prior models, shows that the sulfonyl oxygen atoms of sulfonyl-containing antagonists are positioned for a productive interaction with certain receptor-associated features. This raises the obvious question: Are the sulfonyl oxygen atoms actually required for binding? To address this issue, we prepared and examined sulfenamide **9** ($K_i = 90$ nM) which was found to bind with 20-fold lower affinity than its sulfonyl counterpart **4b** ($K_i = 4.1$ nM). This provides the first evidence that at least one of the sulfonyl oxygen atoms of MS-245-type analogs is required for enhanced affinity. Moreover, that the sulfonyl oxygen atom(s), and not the methoxy group, of MS-245 (**4a**; $K_i = 2.1$ nM) interacts with Thr196 is consistent with the finding that removal of the methoxy group has little impact on affinity (i.e., **4b** $K_i = 4.1$ nM). Also, as shown in Figure 3, there are no amino acid residues within about 7 Å that could potentially hydrogen bond with this methoxy group. Indeed, even an intact indole nucleus is not required for tryptamines to bind at 5-HT₆ receptors. This is evident from the binding of phenylethylamine **15** ($K_i = 52$ nM)³¹ and aminotetralins **7**. Compound **16** ($K_i = 15$ nM),³¹ which lacks the benzenoid ring and methoxy substituent of MS-245 (**4a**), binds with only 4-fold lower affinity than **4b**. This further supports the notion that the methoxy substituent of MS-245 is unlikely to interact with a specific receptor-associated feature. The somewhat reduced affinity of **16** relative to **4** can be explained by the fewer aromatic/hydrophobic contacts possible with the abbreviated heterocyclic nucleus.

Consistency with site-directed mutagenesis results

As already mentioned, Asp106 and Thr196 have been shown important for ligand binding at 5-HT₆ receptors. Two prior modeling studies with N₁-unsubstituted tryptamine analogs (i.e., 5-HT and LSD) have suggested that the indolic N₁-H moiety might be within hydrogen bonding distance to Thr196^{16,17} – a finding not inconsistent with earlier site-directed mutagenesis studies that showed the importance of this residue for agonist binding and agonist efficacy. That is, on the basis of site directed mutagenesis studies showing that the affinity of N₁-unsubstituted LSD-related ergolines (i.e., LSD, ergotamine, lisuride) is decreased, and that the affinity of N₁-methyl ergolines (i.e., methysergide and mesulergine, but not metergoline) is enhanced when Thr196 is mutated to alanine, it was speculated that the indolic NH of 5-HT (**1**) might participate in hydrogen bond formation with Thr196.¹³ This is certainly a possibility; however, it is not known how ergolines bind relative to 5-HT. In fact, comparing the affinities of various ergolines at wild type and several mutant 5-HT_{2A} receptors, it has been suggested that there are differences in binding modes even amongst the ergolines.³² Hence, there is no compelling or *a priori* reason to suspect, or assume, that 5-HT (**1**) and LSD-related ergolines must bind in a comparable manner to 5-HT receptors. Furthermore, we have found that N₁-substituted analogs of 5-HT-related tryptamines either display reduced or enhanced affinity for 5-HT₆ receptors (depending upon the nature of the N₁-substituent), and have previously attributed these affinity differences to different modes of binding.^{9,18,19}

The present model shows that the hydroxyl group, not the N₁-H, of 5-HT (**1**) is situated within hydrogen bonding distance to Thr196 [5.46] (and Ser111 [3.37]). Interesting to note is that there are two conserved hydroxyl-containing amino acids (Ser204 [5.43] and Ser207 [5.46]) common to rhodopsin-like G-protein coupled receptors that bind catecholamine agonists (i.e., dopamine receptors, adrenergic receptors),³³ and that it has been shown on the

basis of site-directed mutagenesis studies that these two residues are important for binding and agonist action and probably participate in hydrogen bond interactions with the hydroxyl groups of catecholaminergic agonists.^{33–35}

Other than for Asp106 and Thr196, few other single amino acid mutations have been reported. Mutation of the rat Trp102 [3.28] (i.e., W102F mutant) had only a small effect on the affinity of 5-HT (<2-fold) or on several nonselective antagonists (<6-fold).¹² Although Trp102 is not within 5 Å of docked 5-HT, it is within 4.5 to 5.0 Å of EMDT (**5**) and the other agents examined (see Supporting Information). Given its distance from the docked molecules, Trp102 might not be expected to exert a major influence on binding. In the rat, mutation of Ala154 [4.53] to serine had no effect on the receptor affinity or agonist action of 5-HT;¹² Ala154 was not found within 5.0 Å of docked 5-HT in the current model. Teitler and co-workers³⁶ found that mutation of Ser267 [6.34] to lysine enhances the affinity of 5-HT at human 5-HT₆ receptors. However, it is unlikely that Ser267 is directly involved in the binding of 5-HT because it is near the intracellular margin of the receptor. Furthermore, it has been demonstrated that the Ser267 mutation results in a constitutively active receptor,^{36,37} and it is known that constitutively active forms of 5-HT receptors characteristically display enhanced affinity for 5-HT.³⁸ Human 5-HT₆ receptor mutation constructs also have been examined in the intracellular loop regions.³⁹ On this basis, a particular residue (Lys265) has been shown instrumental in the coupling of 5-HT₆ receptors to Gas; however, there is no reason to suspect any direct involvement in ligand binding.

Taken together, new models are presented for the interaction of 5-HT and several tryptamine-related ligands at 5-HT₆ receptors. Docking solutions were identified for the interaction of agonist 5-HT (**1**), EMDT (**5**) and EMDT analog **8**, each at one hundred different receptor conformations. In each case, their preferred (based on ChemScore) docking solution was the same. With this model, the basic amines interact with Asp-106, the 5-hydroxy (or methoxy) group interacts with Thr196 and/or Ser111, and the indolic nucleus is located in an aromatic/hydrophobic pocket. Antagonist MS-245 (**4a**) binds differently; MS-245 utilizes the same aromatic/hydrophobic pocket, but otherwise binds in an altogether different fashion than 5-HT. Specifically, the sulfonyl oxygen atoms are situated within hydrogen bond distance of Thr196 and Ser111, and the benzenesulfonyl aryl group (Aryl'; Table 1) interacts with Ala157, Ala192, and Phe285. (Note: other nearby amino acid residues are either shown in Figure 3 or are listed in Supporting Information.)

Benzenesulfonyl compounds **6R** and **7R** were found to dock in a similar manner to this common model.

In summary, two relatively similar, but distinct, 5-HT₆ receptor binding models are described: one for the agonist 5-HT (**1**), and another for the antagonist MS-245 (**4a**). Both ligands utilize a common binding domain but are oriented quite differently within this region. There is consistency with the binding of other tryptamine-related analogs at one or the other of these two models depending upon the presence or absence of an N₁-benzenesulfonyl group, and the models are also consistent with structure-affinity findings and the results of site-directed mutagenesis. We already have provided empirical evidence that tryptamine analogs likely bind differently at 5-HT₆ receptors depending upon whether or not they possess a N₁-benzenesulfonyl group; thus, it is particularly gratifying that two

distinct modes of binding were identified by the automated docking procedure employed herein. The models will now be employed to examine the binding of other 5-HT₆ receptor agonist and antagonist ligands to define pharmacophore models for 5-HT₆ receptor agonists and antagonists.

Experimental

Chemistry

Melting points were taken in glass capillary tubes on a Thomas Hoover melting point apparatus and are uncorrected. ¹H NMR spectra were recorded with a Varian EM-390 spectrometer, and peak position are given in parts per million (δ) downfield from tetramethylsilane as internal standard. Microanalyses were performed by Atlantic Microlab (GA) for the indicated elements, and the results are within 0.4% of calculated values. Reactions and product mixtures were routinely monitored by thin-layer chromatography (TLC) on silica gel precoated F₂₅₄ Merck plates, and chromatographic separations were performed on silica gel columns.

***R*(+)-5-Benzenesulfonamido-2-(dimethylamino)tetralin Hydrochloride (7*R*)**

A solution of benzenesulfonyl chloride (0.08 g, 0.5 mmol) in dry CH₂Cl₂ (2 mL) was added in a dropwise manner at 0 °C to a solution of **13*R*** (0.09 g, 0.5 mmol) in dry CH₂Cl₂ (2 mL). The reaction mixture was allowed to stir at 0 °C for 3 h, solvent was evaporated under reduced pressure, and the residue was recrystallized from MeOH/Et₂O. The residue was converted to a salt by treatment of a dry methanolic solution with HCl-saturated anhydrous Et₂O. The precipitate was collected by filtration and recrystallized from MeOH/Et₂O to give **7*R*** (0.05 g, 28 %) as light-pink crystals: mp 194–196 °C; [α]_D²⁰ = +69.0° (*c* = 0.5, MeOH); ¹H NMR (DMSO-*d*₆) δ 1.50 (m, 1H, CH₂), 2.19 (m, 1H, CH₂), 2.47 (m, 1H, CH₂), 2.74–3.00 (m, 8H, CH₃, CH₂), 3.14 (d, 1H, CH₂), 3.36 (m, 1H, CH₂), 6.73 (d, 1H, ArH), 6.99–7.09 (m, 2H, ArH), 7.55–7.60 (m, 2H, ArH), 7.64–7.69 (m, 3H, ArH), 9.64 (bs, 1H, NH), 10.66 (bs, 1H, NH⁺). Anal. calcd for (C₁₈H₂₂N₂O₂S × HCl × 0.5 H₂O) C, H, N.

***S*(-)-5-Benzenesulfonamido-2-(dimethylamino)tetralin Hydrochloride (7*S*)**

Compound **7*S*** was prepared from **13*S*** in 32% yield in the same manner as **7*R***. The salt was obtained as light-pink crystals after recrystallization from MeOH/Et₂O: mp 195–197 °C; [α]_D²⁰ = -75.7° (*c* = 0.5, MeOH); ¹H NMR (DMSO-*d*₆) δ 1.51 (m, 1H, CH₂), 2.19 (m, 1H, CH₂), 2.47 (m, 1H, CH₂), 2.75–3.01 (m, 8H, CH₃, CH₂), 3.13 (d, 1H, CH₂), 3.35 (m, 1H, CH₂), 6.73 (d, 1H, ArH), 6.99–7.09 (m, 2H, ArH), 7.54–7.60 (m, 2H, ArH), 7.64–7.69 (m, 3H, ArH), 9.64 (bs, 1H, NH), 10.65 (bs, 1H, NH⁺). Anal. calcd for (C₁₈H₂₂N₂O₂S × HCl × 0.5 H₂O) C, H, N.

***R*(+)-2-Ethyl-5-methoxy-3-[(1-methylpyrrolidin-2-yl)methyl]-1*H*-indole Hydrochloride (8*R*)**

A solution of **14*R*** (0.54 g, 1.38 mmol) in dry THF (7 mL) was added in a dropwise manner at 0 °C to a stirred suspension of LiAlH₄ (0.24 g, 6.21 mmol) in dry THF (7 mL). The resulting mixture was heated under a N₂ atmosphere for 4 h. The reaction mixture was cooled, and sodium sulfate decahydrate (7 g) was added very carefully, portionwise, followed by H₂O (0.24 mL) and EtOAc (7 mL). The resulting mixture was allowed to stir at

room temperature under a N₂ atmosphere for 24 h. The reaction mixture was then filtered through Celite, and solvent was evaporated under reduced pressure. The residual oil was then chromatographed on a silica gel column (Aldrich silica gel 60) using first EtOAc (to remove benzyl alcohol), and then EtOAc/MeOH/NH₄OH (9:1:2 drops) as eluent to give a crude product (0.33 g, 88%) as an oil. The product was converted to its hydrochloride salt in anhydrous MeOH by addition of HCl/Et₂O to give an oil that solidified under high vacuum. The solid was recrystallized from CH₂Cl₂/Et₂O to give the product (0.27 g, 64%) as off-white crystals: mp 157–159 °C; ¹H NMR (DMSO-d₆) δ 1.25 (t, 3H, CH₃), 1.68–1.92 (m, 4H, CH₂), 2.72 (m, 2H, CH₂), 2.87 (s, 3H, CH₃), 2.92–3.06 (m, 2H, CH₂), 3.27 (dd, 1H, CH₂), 3.48–3.58 (m, 2H, CH₂), 3.77 (s, 3H, CH₃), 6.67 (d, 1H, ArH), 7.00 (s, 1H, ArH), 7.16 (d, 1H, ArH), 10.75 (s, 1H, NH); [α]_D²⁰ = +19.6 (c = 0.5, MeOH). Anal. calcd for (C₁₇H₂₄N₂O × HCl) C, H, N.

S(–)-2-Ethyl-5-methoxy-3-[(1-methylpyrrolidin-2-yl)methyl]-1*H*-indole Hydrochloride (**8S**)

Beginning with the acid chloride of *N*-(benzyloxycarbonyl)-L-proline (i.e., **14S**), the target compound was prepared in 61% yield in the same manner as its *R*(+)-enantiomer **8R**, and isolated as off-white crystals after recrystallization from CH₂Cl₂/Et₂O: mp 156.5–158.5 °C; ¹H NMR (DMSO-d₆) δ 1.24 (t, 3H, CH₃), 1.68–1.94 (m, 4H, CH₂), 2.73 (m, 2H, CH₂), 2.86 (s, 3H, CH₃), 2.94–3.08 (m, 2H, CH₂), 3.27 (dd, 1H, CH₂), 3.48–3.58 (m, 2H, CH₂), 3.77 (s, 3H, CH₃), 6.67 (d, 1H, ArH), 7.00 (s, 1H, ArH), 7.16 (d, 1H, ArH), 10.75 (s, 1H, NH); [α]_D = –24.6 (c = 0.5, MeOH). Anal. calcd for (C₁₇H₂₄N₂O × HCl) C, H, N.

1-Phenylthio-*N,N*-dimethyltryptamine Oxalate (**9**)

A solution of *N*-(phenylthio)succinimide²⁶ (0.21 g, 1.0 mmol) in CH₂Cl₂ was added to a vigorously stirred mixture of *N,N*-dimethyltryptamine (0.09 g, 0.5 mmol), tetrabutylammonium hydrogen sulfate (0.02 g, 0.05 mmol) and 50% aqueous KOH (0.5 mL) in CH₂Cl₂ (2.5 mL) over a 10- to 15-min period at room temperature. After half of the succinimide was added, additional tetrabutylammonium hydrogen sulfate (0.02 g, 0.05 mmol) was added to the reaction mixture. The reaction mixture was washed with H₂O (3 × 10 mL), the organic portion was separated and dried (Na₂SO₄), and solvent was evaporated under reduced pressure to give an oily yellow residue. Purification by chromatography on a silica gel column (Aldrich silica gel 60) using CH₂Cl₂/MeOH (9:1) as eluent gave **9** (free base; 0.10 g, 68%) as a yellow oil: ¹H NMR (CDCl₃) δ 2.37 (s, 6H, CH₃), 2.67 (t, 2H, CH₂), 2.96 (t, 2H, CH₂), 6.92 (d, 2H, ArH), 7.06 (s, 1H, ArH), 7.13–7.31 (m, 5H, ArH), 7.58–7.65 (m, 2H, ArH). A small sample of the free base was converted to the oxalate salt and recrystallized from MeOH to give the product as white crystals: mp 191–192 °C. Anal. calcd for (C₁₈H₂₀N₂S × C₂H₂O₄) C, H, N.

R-5-Acetamido-2-dimethylaminotetralin (**12R**)

A solution of aqueous H₂CO (37% w/w, 0.79 mL) was added in a dropwise manner to a cooled (ice bath) and stirred mixture of *R*-acetamido-2-aminotetralin²³ (**11R**) (0.51 g, 2.5 mmol), sodium cyanoborohydride (0.35 g, 5.6 mmol) and glacial HOAc (0.14 mL, 2.5 mmol) in a mixture of dry MeOH (30 mL) and MeCN (10 mL). The mixture was allowed to warm to room temperature and stirring was continued for 3 h before addition of saturated

aqueous K₂CO₃ (25 mL). Organic solvent was removed under reduced pressure, and the residue was diluted with H₂O (25 mL) and extracted with EtOAc (3 × 30 mL). The combined organic portion was washed with brine and dried (Na₂SO₄). Solvent was removed under reduced pressure to give the product (0.53 g, 91 %) as an oily residue that was used in the preparation of **13R**: ¹H NMR (CDCl₃) δ 1.72 (m, 1H, CH₂), 2.19–2.29 (m, 4H, CH₂, CH₃), 2.48–2.56 (m, 7H, CH₂, CH₃), 2.66 (m, 1H, CH₂), 2.82–2.94 (m, 2H, CH₂), 3.06 (d, 1H, CH₂), 6.97 (d, 1H, ArH), 7.14–7.19 (m, 2H, ArH, NH), 7.54 (d, 1H, ArH).

S-5-Acetamido-2-*N,N*-dimethylaminotetralin (12S)

The compound was prepared in 87% yield from *S*-5-acetamido-2-aminotetralin (**11S**)²³ in the same manner as **12R**, and isolated as an oily residue: ¹H NMR (CDCl₃) δ 1.72 (m, 1H, CH₂), 2.19–2.29 (m, 4H, CH₂, CH₃), 2.48–2.56 (m, 7H, CH₂, CH₃), 2.66 (m, 1H, CH₂), 2.82–2.94 (m, 2H, CH₂), 3.06 (d, 1H, CH₂), 6.97 (d, 1H, ArH), 7.14–7.19 (m, 2H, ArH, NH), 7.54 (d, 1H, ArH).

R-5-Amino-2-dimethylaminotetralin (13R)

Concentrated HCl (20 mL) was added to a solution of *R*-5-acetamido-2-dimethylaminotetralin (**12R**) (0.50 g, 2.2 mmol) in absolute EtOH (60 mL) and the reaction mixture was heated at reflux for 2 h. The mixture was cooled to 0 °C and basified with 40% NaOH to pH 12. The mixture was extracted with CH₂Cl₂ (3 × 50 mL), and the combined organic portion was dried (Na₂SO₄), and solvent was evaporated under reduced pressure to give the product (0.35 g, 86 %) as a semisolid material: ¹H NMR (CDCl₃) δ 1.72 (m, 1H, CH₂), 2.30 (m, 1H, CH₂), 2.41–2.56 (m, 7H, CH₃, CH₂), 2.68–2.86 (m, 3H, CH₂), 2.97 (d, 1H, CH₂), 3.60 (bs, 2H, NH₂), 6.56 (d, 1H, ArH), 6.59 (d, 1H, ArH), 6.99 (t, 1H, ArH). The compound was used without further purification in the synthesis of **7R**.

S-5-Amino-2-dimethylaminotetralin (13S)

Beginning with **12S**, compound **13S** was prepared in 78% yield in the same manner as **13R** and isolated as a semisolid material: ¹H NMR (CDCl₃) δ 1.71 (m, CH₂, 1H), 2.27 (m, 1H, CH₂), 2.44–2.55 (m, 7H, CH₃, CH₂), 2.63–2.84 (m, 3H, CH₂), 2.96 (d, 1H, CH₂), 3.60 (bs, 2H, NH₂), 6.55 (d, 1H, ArH), 6.59 (d, 1H, ArH), 6.98 (t, 1H, ArH).

R(+)-3-[[*N*-(Benzyloxycarbonyl)pyrrolidin-2-yl]carbonyl]-2-ethyl-5-methoxy-1*H*-indole (14R)

Oxalyl chloride (0.19 mL, 2.21 mmol) was added to a stirred solution of *N*-(benzyloxycarbonyl)-D-proline (0.35 g, 1.47 mmol) in dry CH₂Cl₂ (5 mL) with a trace of DMF (1 drop). The resulting effervescent solution was allowed to stir at room temperature under a N₂ atmosphere for 3 h. Solvent was evaporated under reduced pressure, dry hexane (3 mL) was added, and the resulting solution was again evaporated under reduced pressure to afford *N*-(benzyloxycarbonyl)-D-proline acid chloride; the crude product was dissolved in dry benzene (3 mL).

Concomitantly, a solution of ethyl magnesium bromide (3.0 M in Et₂O, 1.07 mL, 3.09 mmol) was added in a dropwise manner to a stirred solution of 2-ethyl-5-methoxyindole²⁴ (0.52 g, 2.94 mmol) in benzene (10 mL) at 0 °C under a N₂ atmosphere. The resulting mixture was allowed to stir at the same temperature for 15 min, then the solution of *N*-

(benzyloxycarbonyl)-D-proline acid chloride in benzene was added in a dropwise manner with vigorous stirring. The reaction mixture was allowed to stir at 0 °C under a N₂ atmosphere for 1 h, a saturated aqueous solution of NaHCO₃ (7 mL) and EtOAc (9 mL) were successively added, and the mixture was allowed to stir at room temperature for an additional 1 h. The organic layer was removed, and the aqueous portion was extracted with EtOAc (3 × 10 mL). The organic extracts were combined, dried (MgSO₄), and solvent was evaporated under reduced pressure. The residue was chromatographed on a silica gel column (Aldrich silica gel 60) using CH₂Cl₂/MeOH (195:5) as eluent to give a crude product (0.27 g, 47%) as a semi-solid material. The product was used without further characterization in the preparation of **8R**.

S(+)-3-[[N-(Benzyloxycarbonyl)pyrrolidin-2-yl]carbonyl]-2-ethyl-5-methoxy-1H-indole (14S)

Compound **14S** was prepared in 51% yield from *N*-(benzyloxycarbonyl)-L-proline in the same manner as **14R**, and isolated as a yellow semi-solid material. The product was used without further characterization in the preparation of **8S**.

Radioligand binding assay

The h5-HT₆ radioligand binding assays were performed as previously described.³ In brief, h5-HT₆ cDNA was transiently expressed in HEK-293 cells using Fugene6 according to the manufacturer's recommendations. Twenty-four h after transfection the medium was replaced; 24 h later, medium containing dialyzed serum (to remove 5-HT) was added. At 75 h after transfection, cells were harvested by scraping and centrifugation. Cells were then washed by centrifugation and re-suspension once in phosphate buffered saline (pH=7.40; PBS) and then frozen as tight pellets at -80 °C until use. Binding assays were performed at room temperature for 90 min in binding buffer (50 mM Tris-Cl, 10 mM MgCl₂, 0.1 mM EDTA, pH = 7.40) with [³H]LSD (1 nM final concentration) using 10 μM clozapine for non-specific binding. Various concentrations of unlabeled test agent were used for *K_i* determinations, with *K_i* values calculated using the program GraphPad Prism. Specific binding represented 80–90% of total binding. *K_i* values represent a minimum of triplicate determinations.

Computational methods

Molecular modeling studies were conducted using AMD 64 Dual-Core Opteron-based HP xw9400 and Intel Dual-Core Xeon-based SGI VSS40 Linux workstations utilizing SYBYL (version 7.3; Tripos Associates, Inc., St. Louis, MO).

The amino acid sequence of the human 5-HT₆ receptor (accession code P50406) was retrieved from the SwissProt-ExPASy databank and aligned to the sequence of bovine rhodopsin (P02699) based on the Bissantz *et al.*⁴⁰ multiple alignment profile of several diverse type A human GPCRs: dopamine D₃ (P35462), muscarinic acetylcholine m₁ (P11229), vasopressin V_{1a} (P37288), β₂-adrenoceptor (P07550), and δ-opioid (P41143). The ClustalX^{41,42} multiple alignment program was used; in short: *i*) the amino acid sequences of the GPCR profile were aligned to the sequence of bovine rhodopsin in two steps starting from the first transmembrane helix (TM1) to the first five residues of the third intracellular loop (i3) and then from the last five residues of the i3 loop to the end of the seventh

transmembrane helix (TM7) using the BLOSUM weight matrices and with a gap opening penalty of 15.0; *ii*) the alignment of extracellular loop e2 was manually adjusted to align the conserved disulfide bridge in the structure of bovine rhodopsin with the corresponding cysteines in all five aminergic receptors. This alignment provided an unambiguous mapping of residues in the TM regions containing no additions or deletions.

Using the alignment described above, 3-D models of the h5-HT₆ receptor were generated based on the 2.2-Å resolution X-ray crystal structure⁴³ of bovine rhodopsin ('A' chain of PDB code 1U19). Using MODELLER⁴⁴ (version 9.1), a population of one hundred models was generated whose members possessed varied side chain and, to a lesser extent, backbone conformations. The N- and C-termini were truncated prior to the first, and after the seventh TM helix, respectively. The e3 loop (which varies considerably both in terms of length and sequence identity between bovine rhodopsin and 5-HT₆) was modeled as a simple poly-Gly sequence whose initial backbone coordinates were taken from rhodopsin. Following the procedure of Xhaard, *et al.*,⁴⁷ the variation in the side chain conformations of the MODELLER-generated h5-HT₆ receptors was maximized by mutating to alanine the residues falling within 12.0 Å of the retinal ligand in the 1U19 MODELLER rhodopsin template. After adding hydrogen atoms to the one hundred MODELLER-generated h5-HT₆ models, the receptors were energy-minimized (Tripos Force Field, Powell minimization, termination criterion: energy gradient = 0.05 kcal/(mol·Å), Gasteiger-Hückel charges with a distance-dependent dielectric constant = 4.0, and a nonbonded cutoff of 8 Å) using a SYBYL Programming Language script written in-house).

The 3-D structures of the ligands were generated using the SKETCH MOLECULE command within SYBYL. For chiral compounds, the individual isomers were sketched as separate structures. Basic amines were protonated. All ligand structures were then energy-minimized using the same parameters as described above for the receptor models.

The automated docking program GOLD^{45,46} (Genetic Optimization for Ligand Docking, version 3.1, CCDC, Cambridge, UK) was employed to dock ligands into each of the one hundred h5-HT₆ receptor models. A 20-Å radius from the Asp106 [3.32] side chain C γ atom oxygen atoms was used to define the receptor binding cavity. Docking simulations were not restricted to early termination and the ligands were allowed to rotate around single bonds, and non-aromatic rings were free to flip. Ten genetic algorithm docking runs were performed for each ligand structure. The ChemScore fitness function (an estimation of the total free energy change that occurs on ligand binding) was used to rank the docked ligand-receptor complexes. Based on the resulting ChemScores, one of the one hundred MODELLER-generated receptor models was selected for each ligand. Two such receptors were consistently identified: one being common to ligands lacking the sulfonamide group and another common to ligands containing a sulfonamide moiety.

The PROTABLE facility within SYBYL and the PROCHECK program were employed to identify potential, unusual, and sterically unfavorable side chain geometries; these were iteratively corrected as necessary.

All receptor/ligand complexes were subjected to short molecular dynamics simulations using the DYNAMICS routine within SYBYL to provide additional evidence that the GOLD-docked poses were not artifactual (*i.e.*, to assess the stability of the bound pose of the ligands in the h5-HT₆ receptor models). Molecular dynamics simulations were carried out for 100 ps and snapshots were taken every 25 fs. Otherwise, the default settings for the DYNAMICS were employed. The energy setup was analogous to that described for energy minimization. To maintain the integrity of the ligand-receptor complexes, all residues except the ligand and amino acids within an 8-Å radius of the ligand were maintained as an aggregate; these atoms did not move during the course of the simulation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported, in part, by NIMH grant MH-60599 (RAG), the A. D. Williams Fund (RAG), the School of Pharmacy, Medical College of Virginia, Virginia Commonwealth University, MH-57635 (BLR), and the NIMH Psychoactive Drug Screening Program (BLR). RK was supported by T32 grant DA 007027.

Abbreviations

MS-245	N ₁ -Benzenesulfonyl-5-methoxy- <i>N,N</i> -dimethyltryptamine
EMDT	2-ethyl-5-methoxy- <i>N,N</i> -dimethyltryptamine
LSD	lysergic acid diethylamide
SB-258510	5-chloro- <i>N</i> -[4-methylpiperazin-1-yl]phenyl]-3-methyl-2-benzothiophenesulfonamide
Ro 04-6790	4-amino- <i>N</i> -(2,6- <i>bis</i> -methylamino-pyrimidin-4-yl)benzenesulfonamide

References

1. Glennon, RA., Dukat, M. Serotonin receptors and drugs affecting serotonergic neurotransmission. In: Williams, DA., Lemke, TL., editors. Foye's Principles of Medicinal Chemistry. Lippincott Williams & Wilkins; Philadelphia: 2002. p. 315-337.
2. Kroeze WK, Kristiansen K, Roth BL. Molecular biology of serotonin receptors structure and function at the molecular level. *Curr. Top. Med. Chem.* 2002; 2:507–528. [PubMed: 12052191]
3. Kohen R, Metcalf MA, Khan N, Druck T, Huebner K, Lachowicz JE, Meltzer HY, Sibley DR, Roth BL, Hamblin MW. Cloning, characterization, and chromosomal localization of a human 5-HT₆ serotonin receptor. *J. Neurochem.* 1996; 66:47–56. [PubMed: 8522988]
4. Glennon RA. Higher-end serotonin receptors: 5-HT₅, 5-HT₆, and 5-HT₇. *J. Med. Chem.* 2003; 46:2795–2812. [PubMed: 12825922]
5. Roth BL, Craig SC, Choudhary MS, Uluer A, Monsma FJ Jr, Shen Y, Meltzer HY, Sibley DR. Binding of typical and atypical antipsychotic agents to 5-hydroxytryptamine-6 and 5-hydroxytryptamine-7 receptors. *J. Pharmacol. Exp. Ther.* 1994; 268:1403–1410. [PubMed: 7908055]

6. Sleight AJ, Boess FG, Bös M, Levet-Trafit B, Riemer C, Bourson A. Characterization of Ro 04-6790 and Ro 63-0563: Potent and selective antagonists at human and rat 5-HT₆ receptors. *Br. J. Pharmacol.* 1998; 124:556–562. [PubMed: 9647481]
7. Bromidge SM, Brown AM, Clarke SE, Dodgson K, Gager T, Grassam HL, Jeffrey PM, Joiner GF, King FD, Middlemiss DN, Moss SF, Newman H, Riley G, Routledge C, Wyman P. 5-Chloro-N-(4-methoxy-3-piperazin-1-yl-phenyl)-3-methyl-2-benzothiophenesulfonamide (SB-271046): A potent, selective, and orally bioavailable 5-HT₆ receptor antagonist. *J. Med. Chem.* 1999; 42:202–205. [PubMed: 9925723]
8. Glennon RA, Lee M, Rangisetty JB, Dukat M, Roth BL, Savage JE, McBride A, Rauser L, Hufesien L, Lee DKH. 2-Substituted tryptamines: Agents with selectivity for 5-HT₆ serotonin receptors. *J. Med. Chem.* 2000; 43:1011–1018. [PubMed: 10715164]
9. Tsai Y, Dukat M, Slassi A, MacLean N, Demchyshyn L, Savage JE, Roth BL, Hufesein S, Lee M, Glennon RA. N₁-(Benzenesulfonyl)tryptamines as novel 5-HT₆ antagonists. *Bioorg. Med. Chem. Lett.* 2000; 10:2295–2299. [PubMed: 11055342]
10. a) Woolley ML, Marsden CA, Fone KCF. 5-HT₆ Receptors. *Curr. Drug Targets CNS Neurol. Disord.* 2004; 3:59–79. [PubMed: 14965245] b) Holenz J, Pauwels PJ, Diaz JL, Merce R, Codony X, Buschmann H. Medicinal chemistry strategies to 5-HT₆ ligands as potential cognitive enhancers and antiobesity agents. *Drug Discovery Today.* 2006; 11:283–299. [PubMed: 16580970]
11. Ballesteros JA, Weinstaein H. Integrated methods for the construction of three-dimensional models and computational probing of structure-function relations in G protein coupled receptors. *Methods Neurosci.* 1995; 25:366–428.
12. Boess FG, Monsma FJ, Sleight AJ. Identification of residues in transmembrane regions III and IV that contribute to the ligand binding site of the serotonin 5-HT₆ receptor. *J. Neurochem.* 1998; 71:2169–2177. [PubMed: 9798944]
13. Boess FG, Monsma FJ, Meyer V, Zwingelstein C, Sleight AJ. Interaction of tryptamine and ergoline compounds with threonine 196 in the ligand binding site of the 5-hydroxytryptamine₆ receptor. *Mol. Pharmacol.* 1997; 52:515–523. [PubMed: 9284367]
14. Bromidge SM. Potent and selective 5-HT₆ receptor antagonists. *Roy. Soc. Chem. Special Publication.* 2001; 264:101–119.
15. Lopez-Rodriguez ML, Benhamu B, de la Fuente T, Sanz A, Pardo L, Campillo M. A three-dimensional pharmacophore model for 5-hydroxytryptamine₆ (5-HT₆) receptor antagonists. *J. Med. Chem.* 2005; 48:4216–4219. [PubMed: 15974573]
16. Hirst WD, Minton JA, Bromidge SM, Moss SF, Latter AJ, Riley G, Routledge C, Middlemiss DN, Price GW. Characterization of [¹²⁵I]-SB-258585 binding to human recombinant and native 5-HT₆ receptors in rat, pig and human brain tissue. *Br. J. Pharmacol.* 2000; 130:1597–1605. [PubMed: 10928963]
17. Pullagurla MR, Westkaemper RB, Glennon RA. Possible differences in modes of agonist and antagonist binding at human 5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* 2004; 14:4569–4573. [PubMed: 15357994]
18. Kolanos R, Dukat M, Roth BL, Glennon RA. Interaction of N₁-unsubstituted and N₁-benzenesulfonyltryptamines at h5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* 2006; 16:5832–5835. [PubMed: 16945532]
19. Nyandege A, Kolanos R, Roth BL, Glennon RA. Further studies on the binding of N₁-substituted tryptamines at h5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* 2007; 17:1691–1694. [PubMed: 17239595]
20. Gether U. Uncovering molecular mechanisms involved in activation of G protein-coupled receptors. *Endocrine Rev.* 2000; 21:90–113. [PubMed: 10696571]
21. Pullagurla MR, Dukat M, Setola V, Roth B, Glennon RA. N₁-Benzenesulfonylgramine and N₁-benzenesulfonylskatole: Novel 5-HT₆ receptor ligand templates. *Bioorg. Med. Chem. Lett.* 2003; 13:3355–3359. [PubMed: 12951125]
22. Abate C, Kolanos R, Dukat M, Setola V, Roth BL, Glennon RA. Interaction of chiral MS-245 analogs at 5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* 2005; 15:3510–3513. [PubMed: 15990303]
23. a) Grauert, M., Merz, H., Mierau, J., Schingnitz, G., Schneider, C. EP. 402923 A2. Jun 13, 1990 b) Grauert, M., Merz, H., Mierau, J., Schingnitz, G., Schneider, C. US. 005196454 A. Mar 23, 1993

24. Dillard RD, Bach NJ, Draheim SE, Berry DR, Carlson DG, Chirgadze NY, Clawson DK, Hartley LW, Johnson LM, Jones ND, McKinney ER, Mihelich ED, Olkowski JL, Schevitz RW, Smith AC, Snyder DW, Sommers CD, Wert J-P. Indole inhibitors of human nonpancreatic secretory phospholipase A₂. 1. Indole-3-acetamides. *J. Med. Chem.* 1996; 39:5119–5136. [PubMed: 9005255]
25. Macor JE, Chenard BL, Post RJ. Use of 2,5-dimethylpyrrole as an amino-protecting group in an efficient synthesis of 5-amino-3-[(N-methylpyrrolidin-2(R)-yl)methyl]indole. *J. Org. Chem.* 1994; 59:7496–7498.
26. Reese, CB. Process for the preparation of oligodeoxyribonucleotide phosphorothioate triesters via coupling reaction. PCT. Int. Appl. 2001064702. Sep 07, 2001
27. Verdonk ML, Cole JC, Hartshorn MJ, Murray CW, Taylor RD. Improved Protein-Ligand Docking Using GOLD. *Proteins.* 2003; 52:609–623. [PubMed: 12910460]
28. Eldridge MD, Murray CW, Auton TR, Paolini GV, Mee RP. Empirical Scoring Functions: I. The Development of a Fast Empirical Scoring Function to Estimate the Binding Affinity of Ligands in Receptor Complexes. *J. Comp.-Aided Mol. Des.* 1997; 11:425–445.
29. Kabsch W, Sander C. Dictionary of Protein Secondary Structure: Pattern Recognition of Hydrogen-Bonded and Geometrical Features. *Biopolymers.* 1983; 22:2577–2637. [PubMed: 6667333]
30. Glennon RA, Bondarev M, Roth B. 5-HT₆ serotonin receptor binding of indolealkylamines: A preliminary structure-affinity investigation. *Med. Chem. Res.* 1999; 9:108–117.
31. Sikazwe D, Bondarev ML, Dukat M, Rangisetty JB, Roth BL, Glennon RA. Binding of sulfonyl-containing arylalkylamines at human 5-HT₆ serotonin receptors. *J. Med. Chem.* 2006; 49:5217–5225. [PubMed: 16913710]
32. Choudhary MS, Sachs N, Uluer A, Glennon RA, Westkaemper RB, Roth BL. Differential ergoline and ergopeptide binding to 5-hydroxytryptamine_{2A} receptors: Ergolines require an aromatic residue at position 340 for high affinity binding. *Mol. Pharmacol.* 1995; 47:450–457. [PubMed: 7700242]
33. Strader CC, Candelore MR, Hill WS, Sigal IS, Dixon RAF. Identification of two serine residues involved in agonist activation of the β -adrenergic receptor. *J. Biol. Chem.* 1989; 264:13572–13578. [PubMed: 2547766]
34. Cox BA, Henningsen RA, Spanoyannis A, Neve RL, Neve KA. Contributions of conserved serine residues to the interactions of ligands with dopamine D2 receptors. *J. Neurochem.* 1992; 59:627–635. [PubMed: 1321233]
35. Pollock NJ, Manelli AM, Hutchins CW, Steffey ME, MacKenzie RG, Frail DE. Serine mutations in transmembrane V of the dopamine D1 receptor affect ligand interactions and receptor activation. *J. Biol. Chem.* 1992; 267:17780–17786. [PubMed: 1355478]
36. Purohit A, Herrick-Davis K, Teitler M. Creation, expression, and characterization of a constitutively active mutant of the human 5-HT₆ receptor. *Synapse.* 2003; 47:218–224. [PubMed: 12494404]
37. Romero G, Pujol M, Perez P, Buschmann H, Pauwels PJ. Whole spectrum analysis of ligand efficacy at constitutively active human wild-type and S267K 5-HT₆ receptors in HEK-293F cells. *J. Pharmacol. Toxicol. Methods.* 2007; 55:144–150. [PubMed: 16769227]
38. Teitler M, Herrick-Davis K, Purohit A. Constitutive activity of G-protein coupled receptors: Emphasis on serotonin receptors. *Curr. Top. Med. Chem.* 2002; 2:529–538. [PubMed: 12052192]
39. Kang H, Lee WK, Choi YH, Vukoti KM, Bang WG, Yu YG. Molecular analysis of the interaction between intracellular loops of the human serotonin receptor type 6 (5-HT₆) and the α subunit of G_s protein. *Biochem. Biophys. Res. Commun.* 2005; 329:684–692. [PubMed: 15737640]
40. Bissantz C, Bernard P, Hibert M, Rognan D. Protein-based virtual screening of chemical databases. II. Are homology models of G-protein coupled receptors suitable targets? *Proteins: Struct., Funct., Genet.* 2003; 50:5–25. [PubMed: 12471595]
41. Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, Thompson JD. Multiple Sequence Alignment with the Clustal Series of Programs. *Nuc. Acids Res.* 2003; 31:3497–3500.
42. Higgins DG, Thompson JD, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 1994; 22:4673–4680. [PubMed: 7984417]

43. Okada T, Sugihara M, Bondar A-N, Elstner M, Entel P, Buss V. The retinal conformation and its environment in rhodopsin in light of a new 2.2 Å crystal structure. *J. Mol. Biol.* 2004; 342:571–583. [PubMed: 15327956]
44. Sali A, Blundell TL. Comparative protein modeling by satisfaction of spatial restraints. *J. Mol. Biol.* 1993; 234:779–815. [PubMed: 8254673]
45. Jones G, Willett P, Glen RC. Molecular recognition of receptor sites using a genetic algorithm with a description of desolvation. *J. Mol. Biol.* 1995; 245:43–53. [PubMed: 7823319]
46. Jones G, Willett P, Glen RC, Leach AR, Taylor R. Development and validation of a genetic algorithm for flexible docking. *J. Mol. Biol.* 1997; 267:727–748. [PubMed: 9126849]
47. Xhaard H, Nyrönen T, Rantanen V-V, Ruuskanen JO, Laurila J, Salminen T, Scheinin M, Johnson MS. Model structures of α -2 adrenoreceptors in complex with automatically docked antagonist ligands raise the possibility of interactions dissimilar from agonist ligands. *J. Struct. Biol.* 2005; 150:126–143. [PubMed: 15866736]

Figure 1.

Overlap of the two selected receptor conformations highlighting those side chains in the binding site that differ most substantially between the two sites. The position of the receptors relative to one another was determined by calculating the minimum RMSD of the alpha carbon atoms of residues in the binding pocket. Side chains that are colored by atom type belong to the receptor preferred by ligands that do not contain a benzenesulfonyl group (green ribbon). Side chains that are colored either red (RMSD ≥ 2.0 Å) or yellow (1.0 Å $\leq$ RMSD < 2.0 Å) belong to the receptor selected for the ligands that do contain a benzenesulfonyl group (purple ribbon). The conformations of other side chains in the binding site differed between the two receptors, but to a smaller degree. These have been omitted for clarity.

Figure 2.

Proposed binding mode of serotonin (5-HT; **1**) in the h5-HT₆ receptor, looking through the extracellular side of the helical bundle. TM5, TM6 and TM7 are nearest to the viewer. The backbone ribbon trace is color-coded based on the assignment of secondary structure using the Kabsch-Sander²⁹ algorithm (red = helix, blue = sheet, violet = turn, and yellow = coil). 5-HT (**1**) is rendered as a CPK-style space-filling model, and the side chains of residues whose heavy atoms fall within 4.0 Å of the ligand's heavy atoms are rendered as capped sticks.

Figure 3.

Proposed binding mode of MS-245 (**4a**) in the h5-HT₆ receptor, looking through the extracellular side of the helical bundle. TM5, TM6 and TM7 are nearest to the viewer. The MS-245 ligand is rendered as a ball-and-stick model with the carbon backbone in green, and the side chains of residues whose heavy atoms fall within 4.0 Å of the ligand's heavy atoms are rendered as capped sticks.

Scheme 1.^a

^aReagents and conditions: (a) $\text{H}_2\text{C}=\text{O}$ (37%), NaCNBH_3 , HOAc, MeOH/MeCN, r.t. 3 h; (b) conc. HCl, EtOH, , 2 h; (c) PhSO_2Cl , CH_2Cl_2 , 0 °C, 3 h.

15 **16****Chart 1.**

Structures of 5-HT (1), 5-HT₆ antagonists Ro 04-6790 (2), SB-258510 (3a), SB-271046 (3b), MS-245 (4a) and its des-methoxy analog 4b, and the agonist EMDT (5).

Table 1

Proposed modes of interaction for sulfonamide-containing 5-HT₆ antagonists.

Aryl ¹ -SO ₂ -N-Aryl-X- Amine				
Feature	Bromidge ¹⁴	Pullagurta <i>et al.</i> ¹⁷	Hirst <i>et al.</i> ¹⁶	Lopez-Rodriguez <i>et al.</i> ¹⁵
Amine	Asp106 [3.32]	Asp106 [3.32]	Asp106 [3.32]	Asp106 [3.32]
Aryl	Trp281 [6.48]	Phe284 [6.51] Phe285 [6.52]	Trp281 [6.48] Phe284 [6.51] Phe302 [7.35]	Val107 [3.33] Leu182 [e2] Trp281 [6.48] Phe284 [6.51] Ser111 [3.37] Thr196 [5.46] Ala157 [4.56] Ala192 [5.42] Phe285 [6.52]
SO₂	--	Leu182 [e2] Gln291 [e3]	Asn288 [6.55] Gln216 ^b	
Aryl¹	Phe277 [6.44]	Phe302 [7.35]	Phe188 [5.38]	

^a Other residues within 5 Å of docked molecules, in 0.5-Å distance increments, are listed in the Supporting Information section.

^b Both the rat and human receptor contain a glutamine moiety at position 218, but an arginine residue at position 216. Hence, it is uncertain which of the two amino acids was actually identified in this study.