CORRESPONDENCE

Impingement Syndrome of the Shoulder

by Dr. med. Christina Garving, Sascha Jakob, Dr. med. Isabel Bauer, Dr. med. Rudolph Nadjar, and Prof. Dr. med. Ulrich H. Brunner in issue 45/2017

If Possible Treat Without Surgery

The authors chose the title “impingement syndrome,” although internationally the term “subacromial pain syndrome” (SAPS) is regarded as more appropriate, because the anatomy and the pain barely correlate and imply a false causality (1).

In the article, one of the take-home messages is that the one-month prevalence of shoulder pain is between 16% and 30%, but the 30% refers to people older than 30. This means that the prevalence in the population in this study is only 16% (2). The statement that “Surgery is indicated if the symptoms fail to improve after 3 or more months of conservative treatment” is ambiguous and is backed up by only one, outdated, source from 1997. A more correct recommendation would be that surgery within the first 3 months is generally not indicated, and only in a few cases afterwards. The current Dutch guideline that the authors cite elsewhere (3) is clearer in this respect: SAPS should if at all possible be treated without surgery...

- If intensive non-surgical treatment does not bring about improvement and rehabilitation is not possible, bursectomy may be considered.
- No indication exists for operating on an asymptomatic rupture of the rotator cuff or calcific tendinitis.

Even after rupture of the rotator cuff, 90% of patients who have not had surgery have no problems or only negligible problems after 14 years (2). That is no worse than the long term results after surgery (4).

DOI: 10.3238/arztebl.2018.0342a

References

1. Garving C, Jakob S, Bauer I, Nadjar R, Brunner UH: Impingement syndrome of the shoulder. Dtsch Arztebl Int 2017; 114: 765–76.
2. Diercks R, Bron C, Dorrestijn O, et al.: Guideline for diagnosis and treatment of subacromial pain syndrome: a multidisciplinary review by the dutch orthopaedic association. Acta Orthop 2014; 85: 314–22.
3. Kijima H, Minagawa H, Nishi T, Kikuchi K, Shimada Y: Long-term follow-up of cases of rotator cuff tear treated conservatively. J Shoulder Elbow Surg 2012; 21: 491–4.
4. Jaeger M, Berndt T, Rühmann O, Lerch S: Patients with impingement syndrome with and without rotator cuff tears do well 20 years after arthroscopic subacromial decompression. Arthroscopy 2016; 32: 409–15.

Dr. med. Uwe Popert
Kassel
uwe.popert@web.de

Motion Sequence Disrupted

It is important that the summary of a selective literature search for the third most common musculoskeletal complaint in orthopedic practice is published in *Deutsches Ärzteblatt* (1). What is surprising and sobering is the fact that the focus of the article is on well-known

structural findings and surgical options, whereas the orthopedic aspect of structural functional causes is barely mentioned and is given too short shrift. The learning objectives are therefore not met.

How to explain the development of structural adjustments and defects in this “epidemic”? The human shoulder develops primarily as a pull-rotation system, in contrast to the hip, which is a push-rotation system (2). At the start of a shoulder movement, the humeral head in the glenoid needs to be pulled downwards and centered by using the humeral distractors (the long head of the biceps tendon and the latissimus dorsi muscle), before abduction of the glenohumeral joint becomes possible without pathological humero-acromial impingement and squeezing of the rotator cuff, which in the long term leads to subacromial adjustments and destructions. The element that is being moved is primarily the humeral head and not the acromion. Modern lifestyles and/or unilateral or incorrect training will weaken the humeral distractors vis-à-vis the cranializing and ventralizing muscle groups, and the physiologically necessary sequence of pull-rotation in the glenoid will be disrupted (3). Congenital or acquired instabilities also play a part. Expert sonography allows for functional analysis of the movement and stability of the shoulder long before structural remodeling processes will be visible on magnetic resonance imaging or changes are seen on the conventional radiograph (4).

Early, timely, and systematic instruction on optimizing movement patterns and behavioral habits can help prevent the development of the described structural defects (4, 2).

DOI: 10.3238/arztebl.2018.0342b

References

1. Garving C, Jakob S, Bauer I, Nadjar R, Brunner UH: Impingement syndrome of the shoulder. Dtsch Arztebl Int 2017; 114: 765–76.
2. Hien NM: Hüftdysplasie und Subakromialsyndrom – gemeinsame Aspekte zu Ursache und Behandlung. Orthopädische Praxis 2009; 45: 398–402.
3. Longo S, Corradi A, Michielon G, Sardaneli F, Sconfienza LM: Ultrasound evaluation of the subacromial space in healthy subjects performing three different positions of shoulder abduction in both loaded and unloaded conditions. Phys Ther Sport 2017; 23: 105–12.
4. Luding U: Funktionsanalyse der Abduktion im Gelenohumeralgelenk bei Supraspinatussyndrom anhand sonographischer Beobachtung. Inaugural Dissertation LMU München 1994.

Dr. med. Norbert M. Hien
81243 München
PraxisDrHien@t-online.de

Don't Forget Radiotherapy

We thank the authors for compiling what is known about impingement syndrome of the shoulder (1).

Regrettably, however, they did not mention that even low-dose radiotherapy can have a good analgetic effect. Furthermore, in individual case reports, periarticular calcifications regressed after radiotherapy (2). Retrospective studies have shown a reduction in pain or even absence of pain in 58–100% of patients. The results are better for early symptoms. Radiotherapy is recommended in the form of orthovoltage irradiation or by a linear accelerator with total reference doses of 3–6 Gy in 2–5 × weekly individual fractions of 0.5–1 Gy (2), which amounts to 1–3 weeks of radiotherapy.

Earlier randomized trials from 1952 and 1969, which did not establish any superiority of radiotherapy over placebo, should be regarded with caution because of low patient numbers, questionable endpoints, occasionally unusual indications, and unsatisfactorily short follow-up periods. A study published in 2014 on the painful shoulder syndrome (3), which compared individual doses of 0.5 Gy (total reference dose 3.0 Gy) and 1 Gy (total reference dose 6.0 Gy), showed a good response to radiotherapy on the one hand, but on the other hand it was found that the mentioned doses were isoeffective. Many institutions halved their doses afterwards. Studies comparing radiotherapy using a conventional dose with placebo (or a very low dose) are lacking for impingement syndrome. Gross et al. published a comparison with an alternative method (extracorporeal shockwave therapy, ESWT) (4); they showed equivalent results.

Local adverse reactions, such as skin irritations, have been observed very rarely. The risk of secondary tumors is extremely low when the appropriate technique is used and used sparingly in very young patients (<30 years).

DOI: 10.3238/arztebl.2018.0342c

References

1. Garving C, Jakob S, Bauer I, Nadjar R, Brunner UH: Impingement syndrome of the shoulder. *Dtsch Arztebl Int* 2017; 114: 765–76.
2. Ott OJ, Niewald M, Weitmann HD, et al.: DEGRO guidelines for the radiotherapy of non-malignant disorders. Part II: painful degenerative skeletal disorders. *Strahlenther Onkol* 2015; 191: 1–6.
3. Ott OJ, Hertel S, Gaipf US, Frey B, Schmidt M, Fietkau R: The Erlangen dose optimization trial for radiotherapy of benign painful shoulder syndrome. Long-term results. *Strahlenther Onkol* 2014; 190: 394–8.
4. Gross MW, Sattler A, Haake M, et al.: [The effectiveness of radiation treatment in comparison with extracorporeal shockwave therapy (ESWT) in supraspinatus tendon syndrome]. *Strahlenther Onkol* 2002; 178: 314–20.

On behalf of the German Working Group on Radiotherapy of Benign Diseases in the German Society of Radiation Oncology (DEGRO)

Prof. Dr. med. Marcus Niewald
66421 Homburg/Saar
marcus.niewald@uks.eu

Prof. Dr. med. Oliver Micke
Bielefeld

PD Dr. med. Ralph Mücke
Bad Kreuznach

In Reply:

We thank our correspondents for their responses to our article (1).

Regarding radiotherapy: Randomized controlled trials of the effect of radiotherapy in shoulder pain are

lacking, with the exception of the study reported by Plenck (2). A review article of the historical assessment describes how in the US the method was abandoned as early as in the 1960s because of too few studies (3). In the “critical study” by Plenck (2), patients with peritendinitis of the shoulder were randomized and blinded and received radiotherapy, with controls being shielded by a lead plate from radiation. After six weeks, no clinical or radiological differences were seen between groups. The studies by Ott et al (4, 5) did not differentiate any pathology, had no control group, and randomized two radiotherapy protocols. The evaluation of pain only, without clinical examination, was not significantly different. The DEGRO S2e guideline (4) refers only to the two studies by Ott and confirms for the radiotherapy protocol an evidence level of only 4, recommendation level C. An effect of the “pain radiotherapy” has not been confirmed in the literature and can at best be assumed. Any indication has to be classed as empirical.

Regarding the conservative treatment of impingement syndrome: We thank our correspondent for the emphasis on functional associations in the pathogenesis of shoulder pain. However, this does not release us from describing different etiologies and referring to terms and concepts that are firmly established in the international literature on the subject. In terms of biomechanics, we placed our emphasis on the shoulder joint’s force fit connection by muscle activity and its full range of motion, and separated out functional causes such as muscular dysbalances and mechanical “extrinsic” factors. We mentioned conservative approaches with evidence levels (for example, a program for stepwise shoulder training). The extent to which “optimizing movement patterns and behavioral habits” improve not only clinical symptoms but also help to “prevent the development of . . . structural defects” remains to be seen.

Regarding impingement syndrome: The Dutch guideline is the only existing one for the clinical syndrome of subacromial pain. It also defines the importance of the differential diagnostic clinical examination with specific tests as a first step and imaging after 6 weeks. The guideline also mentions structural pathologies—for example, bursitis, calcific tendinitis, or biceps lesions. Although the term “subacromial pain syndrome” avoids the association with the historical impingement theory, it does not spare us from the necessity of understanding the pathophysiological associations and treatment modalities of the individual symptoms, which again appear under an umbrella term—such as “periartthritis humeroscapularis” in the time of Duplay. The statement that “subacromial pain syndrome” should not be treated with surgery if possible, is similarly misleading if the underlying pathology is not known. In this context we wish to mention the differentiated treatment of rotator cuff lesions (6).

DOI: 10.3238/arztebl.2018.0343

References

1. Garving C, Jakob S, Bauer I, Nadjar R, Brunner UH: Impingement syndrome of the shoulder. *Dtsch Arztebl Int* 2017; 114: 765–76.
2. Plenck H P: Calcifying tendinitis of the shoulder; a critical study of the value of x-ray therapy. *Radiology* 1952; 59: 384–9.
3. Calabrese EJ, Dhawan G, Kapoor R: Use of X-rays to treat shoulder tendonitis/bursitis: a historical assessment. *Arch Toxicol* 2014; 88: 1503–17.
4. Ott OJ, Hertel S, Gaipf US, Frey B, Schmidt M, Fietkau R: The Erlangen dose optimization trial for radiotherapy of benign painful shoulder syndrome. Long-term results. *Strahlenther Onkol* 2014; 190:394–8.
5. Ott OJ, Niewald M, Weitmann HD, et al.: DEGRO guidelines for the radiotherapy of non-malignant disorders. Part II: painful degenerative skeletal disorders. *Strahlenther Onkol* 2015; 191: 1–6.
6. AWMF: Rotatorenmanschette. www.awmf.org/leitlinien/detail/II/033-041.html (last accessed on 28 March 2018).

On behalf of the authors

Prof. Dr. Ulrich H. Brunner

Abteilung für Unfall-, Schulter- und Handchirurgie,
Krankenhaus Agatharied GmbH, Hausham
ulrich.brunner@khagatharied.de

Conflict of interest statement

All authors declare that no conflict of interest exists.

CLINICAL SNAPSHOT

The Consequences of a “Harmless” Frontal Swelling

- a) The affected area after incomplete incisional biopsy in another institution.
b) The same site after four further procedures for the resection of dermatofibrosarcoma protuberans of the frontal region, with secondary granulation.

A 31-year-old man without antecedent illnesses and in good general condition was referred because of a painless, but growing and cosmetically disturbing swelling on the forehead, which he had first noticed some 9 months previously. It had reached the size of ca. 2 × 1 cm before subtotal excisional biopsy in another institution. The histopathological examination revealed a dermatofibrosarcoma protuberans of the facial skin; multiple further surgical procedures were needed before an R0 resection could be achieved. The resulting cutaneous defect was as large as the palm of a hand. Because of its size, the defect was allowed to heal by secondary intention in combination with a meshed split-thickness skin graft, yielding an adequate cosmetic result. This case underscores the importance of a differential diagnostic work-up even for seemingly harmless cosmetic flaws.

Dr. med. univ. Dr. med. dent. Florian J. Kupilas, M.Sc., D.A.L.M., Prof. Dr. med. Dr. med. dent. Johannes Kleinheinz,
Klinik für Mund-, Kiefer- und Gesichtschirurgie, Universitätsklinikum Münster, kupilas@uni-muenster.de

Conflict of interest statement: The authors state that they have no conflict of interest.

Cite this as: Kupilas FJ, Kleinheinz J: The consequences of a harmless frontal swelling. *Dtsch Arztebl Int* 2018; 115: 344.
DOI: 10.3238/arztebl.2018.0344

Translated from the original German by Ethan Taub, M.D.