Variable Tacrolimus Dosing After Protease Inhibitor-Based Antiretroviral Therapy Discontinuation in 2 HIV Patients Post-Kidney Transplantation

Journal of Pharmacy Technology
2018, Vol. 34(2) 86–88
© The Author(s) 2018
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/8755122517753684
journals.sagepub.com/home/pmt
SAGE

Patricia Pecora Fulco, PharmD¹ and Crystal Reiko Leibrand, PharmD²

To the Editor:

Recent data support the success of solid organ transplants in human immunodeficiency virus (HIV)-positive patients on suppressive antiretroviral therapy (ART).¹ Significant drug-drug interactions (DDIs) occur between calcineurin inhibitors (CNIs) and ART.^{2,3} In a recent study of 332 HIV-positive kidney transplant recipients, a 1.8- and 1.9-fold increased risk of allograft loss and death resulted, respectively, in 88 patients receiving protease inhibitor (PI)-based regimens compared with 244 patients on non-PI-based ART.⁴ If possible, eligible HIV-positive recipients should be converted to a non-PI-based regimen prior to or after kidney transplantation.

Tacrolimus is the CNI of choice in HIV-positive renal transplant recipients but has a narrow therapeutic index requiring close drug monitoring to avoid common adverse effects (nephrotoxicity and neurotoxicity) with supratherapeutic concentrations.^{2,5} CNI dose conversion after PI discontinuation is variable, with tacrolimus (TAC) dose escalation of 5 to 20 times the original dose.^{6,7} We report on 2 HIV-positive renal transplant recipients who were successfully switched from ritonavir (RTV)-boosted ART to dolutegravir or rilpivirine necessitating a 10- to 24-fold TAC dose escalation to maintain therapeutic concentrations.

Patient 1 is a 68-year-old male with HIV/hepatitis B (viral loads [VL] undetectable), deceased donor kidney transplant (DDKT), and hypertension. After 2 years of concurrent mycophenolate mofetil, prednisone, and TAC (0.2 mg weekly) (stable serum creatinine [SCr] 1.16–1.30 mg/dL) with fixed dose combination (FDC) abacavir/lamivudine and twice daily DRV/RTV/etravirine/raltegravir, ART was simplified to abacavir 600 mg/dolutegravir 50 mg/lamivudine 300 mg. TAC was empirically adjusted 10-fold with weekly concentrations remaining at goal (5–7 ng/mL; Figure 1). After dolutegravir transition, the SCr increased (2.28 mg/dL), resulting in an allograft biopsy demonstrating no acute rejection but chronic antibody-mediated changes with early transplant glomerulopathy. The SCr has remained stable for the following 2 years.

Patient 2 is a 52-year-old female with HIV (DRV/RTV ART with undetectable VL), hypertension, deep vein thrombosis, and recent DDKT receiving mycophenolic acid, prednisone, and TAC (0.5 mg weekly) (goal trough 6–8 ng/mL; Figure 1). Antiretroviral simplification was recommended with daily FDC abacavir/lamivudine and rilpivirine 25 mg (previous integrase resistance [140A, 148R, 203M] and dual tropic HIV). TAC dose titration stabilized at 6 mg twice daily (24-fold dose escalation). TAC weekly concentrations remained therapeutic with SCr improvement (1.23–1.01 mg/dL) and continued VL suppression at 4 months posttransplant.

A similar report describes a DDKT HIV-positive recipient receiving DRV/RTV/raltegravir.⁷ ART simplification to FDC dolutegravir/abacavir/lamivudine necessitated a 20-fold TAC dose increase (0.5 mg every 11th day to 5 mg twice daily) after PI discontinuation. The SCr subsequently rose (+0.4 mg/dL) with a negative biopsy. Dolutegravir's inhibition of organic cationic transporter-2 (OCT-2) in the renal tubule was suspected (similar to Patient 1). Rilpivirine also inhibits OCT-2 and a small SCr rise of 0.1 mg/dL is common but does not result in an alteration in the actual glomerular filtration rate. Additional caution should be given to substituting cobicistat-boosted ART in HIV-positive renal transplant recipients as a probable DDI with TAC exists and resulted in supratherapeutic TAC concentrations (111.2 ng/mL) in one reported case.^{8,9}

Recent data suggest hypertension as the primary etiology for the pretransplant chronic kidney disease in HIV-positive renal transplant patients as successful ART results in an almost normal life expectancy.⁴ Based on historical

¹Virginia Commonwealth University Health System, Richmond, VA, USA

²Virginia Commonwealth University, Richmond, VA, USA

Corresponding Author:

Patricia Pecora Fulco, Department of Pharmacy, Virginia Commonwealth University Health System, 401 N 12th Street, PO Box 980042, Richmond, VA 23298-0042, USA.

Email: patricia.fulco@vcuhealth.org

Figure 1. (A) Tacrolimus concentrations and (B) renal function after ART switch to either dolutegravir or rilpivirine therapy.

genotypes and in consultation with an HIV infectious disease specialist, HIV-positive candidates should be switched prior to planned renal transplant as post-allograft ART conversions will require weekly evaluations to maintain therapeutic CNI concentrations and HIV viral rebound (similar to the present cases). RTV or cobicistat-boosted regimens should be changed to ART devoid of cytochrome P450 CNI DDIs (dolutegravir, maraviroc, raltegravir, or rilpivirine or the soon to be approved bictegravir) to prolong allograft survival.³ If ARV resistance does not allow PI or cobicistat-boosted ART discontinuation, alternative immunosuppressive strategies (belatacept) devoid of CNI DDIs is recommended.³

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

1. Muller E, Barday Z, Mendelson M, Kahn D. HIV-positive-to-HIV-positive kidney transplantation—results at 3 to 5 years. *N Engl J Med*. 2015;372:613-620. doi:10.1056/NEJMoa1408896.
2. Primeggia J, Timponi JG Jr, Kumar PN. Pharmacological issues of antiretroviral agents and immunosuppressive regimens in HIV-infected solid organ transplant recipients. *Infect Dis Clin North Am*. 2013;27:473-486. doi:10.1016/j.idc.2013.02.011.
3. Azar MM, Malinis MF, Moss J, Formica RN, Villaneuva MS. Integrase strand transferase inhibitors: the preferred antiretroviral regimen in HIV-positive renal transplantation.

- Int J STD AIDS*. 2017;28:447-458. doi:10.1177/0956462416651528.
4. Sawinski D, Shelton BA, Mehta S, et al. Impact of protease-inhibitor-based anti-retroviral therapy on outcomes for HIV+ kidney transplant recipients. *Am J Transplant*. 2017;17:3114-3122. doi:10.1111/ajt.14419.
 5. Stock PG, Barin B, Murphy B, et al. Outcomes of kidney transplantation in HIV-infected recipients. *N Engl J Med*. 2010;363:2004-2014. doi:10.1056/NEJMoa1001197.
 6. Bickel M, Anadol E, Vogel M, et al. Daily dosing of tacrolimus in patients treated with HIV-1 therapy containing a ritonavir-boosted protease inhibitor or raltegravir. *J Antimicrob Chemother*. 2010;65:999-1004. doi:10.1093/jac/dkq054.
 7. Lee DH, Malat GE, Bias TE, Harhay MN, Ranganna K, Doyle AM. Serum creatinine elevation after switch to dolutegravir in a human immunodeficiency virus-positive kidney transplant recipient. *Transpl Infect Dis*. 2016;18:625-627. doi:10.1111/tid.12545.
 8. Han Z, Kane BM, Petty LA, Josephson MA, Sutor J, Pursell KJ. Cobicistat significantly increases tacrolimus serum concentrations in a renal transplant recipient with human immunodeficiency virus infection. *Pharmacotherapy*. 2016;36:e50-e53. doi:10.1002/phar.1752.
 9. Tseng A, Hughes CA, Wu J, Seet J, Phillips EK. Cobicistat versus ritonavir: similar pharmacokinetic enhancers but some important differences. *Ann Pharmacother*. 2017;51:1008-1022. doi:10.1177/1060028017717018.