Published in final edited form as:

J Antibiot (Tokyo). 2017 August ; 70(9): 962–966. doi:10.1038/ja.2017.67.

Antimicrobial sulfonamides clear latent Kaposi Sarcoma Herpesvirus infection and impair MDM2-p53 complex formation

Fabrizio Angius¹, Enrica Piras¹, Sabrina Uda¹, Clelia Madeddu², Roberto Serpe², Rachele Bigi³, Wuguo Chen³, Dirk P. Dittmer³, Raffaello Pompei¹, and Angela Ingianni¹

¹Department of Biomedical Sciences, University of Cagliari, Italy

²Department of Medical Sciences “Mario Aresu, University of Cagliari, Italy

³Department of Microbiology and Immunology, Lineberger Comprehensive Cancer Center, and Center for AIDS Research, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

Abstract

Kaposi Sarcoma Herpesvirus (KSHV), also known as Human Herpesvirus 8, is the causative agent of Kaposi sarcoma; this malignant angiosarcoma is usually treated with conventional anti-tumor agents, which can control disease evolution, but do not clear the latent KSHV episome that binds to cellular DNA. Some commercial antibacterial sulfonamides were tested for the ability to suppress latent KSHV. qPCR and cytofluorometry assays were used for detecting both viral DNA and the latency factor LANA in BC3 cells, respectively. The capacity of sulfonamides to impair MDM2-p53 complex formation was detected by an ELISA method. The analysis of variance was performed according to one-way ANOVA with Fisher as a post-hoc test.

Here we show that sulfonamide antibiotics are able to suppress the KSHV latent state in permanently-infected BC3 lymphoma-cells and interfere with the formation of the MDM2-p53 complex, which KSHV seemingly needs to support latency and to trigger tumor cell transformation. These findings detected a new molecular target for the activity of sulfonamides and offer a new potential perspective for treating KSHV-induced lymphoproliferative diseases.

Keywords

Kaposi sarcoma Herpesvirus; Sulfonamides; Antivirals; Oncogenic viruses; MDM2-p53; Kaposi Sarcoma

INTRODUCTION

The Kaposi Sarcoma Herpesvirus (KSHV) is a lymphotropic γ -herpesvirus known as the causative agent of the Kaposi sarcoma (KS) and other lymphoproliferative diseases.^{1,2} Like

Address correspondence to R. Pompei, Dept. of Biomedical Sciences, University of Cagliari, ViaPorcell 4, 09124, Cagliari, Italy, Tel. ++39.070.6758483; rpompei@unica.it.

CONFLICT OF INTEREST:

The authors declare that they have no conflict of interests.

all the species of herpesvirus, KSHV is also able to cause a lytic infection at first contact, and then to enter into a latent state, whereby the viral episome is bound to cellular DNA by means of the latency factor LANA.^{3–6}

At present, KS is treated as a malignant tumor with conventional chemotherapeutic agents and/or interferon-alpha.⁷ Such therapy is often able to control disease evolution, but does not cure the KSHV infection, since the virus can remain in a latent state as an episome bound to cellular DNA for its entire lifespan. In addition, conventional anti-herpes drugs, such as gancyclovir and foscarnet, are able to slightly suppress virus replication in the lytic phase, but do not clear the latent state.^{8,9}

Recent studies have demonstrated that latent herpesviruses, such as Epstein-Barr virus (EBV) and KSHV, can be inhibited by certain synthetic compounds, which are able to interfere with the interaction of the latency factor EBNA1 of EBV¹⁰ or with the ability of the KSHV latency protein LANA to bind to cellular DNA.¹¹

In this work we found that several sulfonamide antibacterials were able to suppress the KSHV latent state in permanently infected BC3 lymphoma cells. Studies on the mechanism of action revealed that they could interfere with the formation of the MDM2-p53 complex, which KSHV seemingly needs to support latency and to trigger tumor cell transformation.

MATERIALS AND METHODS

Cell culture

KSHV-positive BC3 cells were grown in RPMI-1640 medium with 10% fetal calf serum (Invitrogen, Paisley, UK) as previously described.^{12–14} THP-1 cells (a KSHV-negative macrophage-cell line) were purchased from Life Technologies (Paisley, UK). A pool-stock of HUVEC cells (primary endothelial human cells from Life Technologies) was grown in M200 medium (Gibco, UK) with LSGS supplement (Invitrogen). The sulfonamide compounds used in this study (see Table 1 and supplemental file) were commercial drugs purchased from Sigma-Aldrich (S. Louis, USA). Nutlin-3a (Sigma-Aldrich) and nimesulide (Sigma-Aldrich) were used as reference drugs. Compound cytotoxicity was evaluated by measuring the effect produced on cell morphology and replication as previously described.¹²

qPCR

Genomic DNA was isolated using an Easy-DNA kit (Invitrogen) and suspended in 40 µl of TE buffer (10 mM Tris-HCl, 1 mM EDTA). To detect KSHV DNA, 100 ng of the processed DNA were used as a template, and qPCR was performed on a Real-Time PCR System (StepOne; Applied Biosystems, Monza, Italy) as described before.¹⁴ The Custom TaqMan® Assay used consisted in KSHV Orf50 primers and probe (forward CAGCCATATGTAAGCTACTACAGCAA, reverse TGTAGAGCTTCAGTAGGGAAGAGGTT, and probe 6-FAM-CCTGGAGGCGACTCGTCTGCAATC-MGB).

Cytofluorometry assay

LANA expression was assayed by flow cytofluorometry in BC3 cells with a modification of the method described by Angius et al.¹² The cells were incubated at 37 °C and 5% CO₂ for 6 days in the presence of a 100 µM concentration of compounds. Subsequently, cells were washed twice with PBS and once with RPMI medium, pH 7.4, and incubated at 37°C 1 h with rat anti-LANA antibodies (Santa Cruz Biotech, Dallas, USA) and subsequently with rabbit anti-rat FITC-conjugated antibodies (Sigma-Aldrich). Following incubation, cells were washed with PBS and suddenly fixed with iced methanol for 10 min at RT. After fixation, cells were washed and harvested for measurements. Fluorescence was observed with a Perkin-Elmer flow cytofluorometry apparatus. All data were processed with BD FACSdiva 6.1.3 (Becton Dickinson, Mountain View, CA, USA). At least 10,000 cells were evaluated for each group and each experiment was carried out at least twice. The data are reported as the weighted mean value of fluorescence evaluated in positive cells normalized to the entire population.

MDM2-p53 complex inhibition assay

The interaction of sulfonamide compounds on the MDM2-p53 complex was performed according to the Böttger et al. method¹⁵ on high-binding microtiter-plates (Greiner Bio-One, Kremsmünster, Austria) coated by overnight incubation with 2 µg/ml p53 protein (human recombinant, Sigma) in PBS at 4°C. The plates were subsequently incubated with MDM2 (1.3 µg/ml) tagged with Glutathione-S-transferase (human GST-MDM2, Sigma) prepared in blocking buffer. The experiments were performed in two ways: i) pre-incubation: the compounds were added to the p53 coated-wells 30 min before adding the GST-MDM2; ii) post-incubation: the compounds were added to the wells 30 min after adding GST-MDM2 to the mixture. The plates were then incubated with rabbit anti-GST specific monoclonal antibody (Sigma-Aldrich), followed by incubation with an anti-rabbit horseradish peroxidase-conjugated secondary antibody (goat anti-rabbit IgG-HRP, Sigma). After incubation with Stabilized Chromogen (Invitrogen), the absorbance was measured with a microtiter plate reader (Model 680, Bio-Rad, Hercules, USA) and the relative luminescence units (RLU) were measured. The percentage of MDM2-p53 complex inhibition was calculated as the (RLU detected in the compound treated sample - RLU of DMSO control)/100. The analysis of variance was performed according to one-way ANOVA with Fisher as a post-hoc test. A significant difference was considered when $p < 0.05$.

RESULTS

Latent KSHV DNA was detected by qPCR (Figure 1) and the LANA antigen was assayed by a cytofluorometry method (Figure 2). After 6 days of cell culture in the presence of 100 µM drugs (SG3= 21.4 µg/ml; SA4= 17.2 µg/ml; ST5= 25.5 µg/ml; BC6= 25.3 µg/ml; see also the additional file), the cell number was counted and a maximum of 15-20% of difference in cell count was recorded in the various plates. We found that the KSHV DNA was cleared from most cells by some of the compounds tested: sulfoguanidine (SG3) inhibited viral DNA by 52±3%, while sulfanilamide (SA4), sulfamethoxazole (BC6) and sulfathiazole (ST5) showed antiviral activities of 97±2, 88±2 and 87±3% respectively. Nimesulide, known to reduce KSHV latent gene expression, disrupt p53-LANA-1 protein

complexes and activate the p53/p21 tumor-suppressor pathway,¹¹ was used as a reference control in our assays, demonstrating an antiviral effect of $55\pm3\%$ on KSHV DNA.

In this assay we observed an increasing anti-viral activity among the tested molecules from sulfaguanidine to sulfathiazole. As a matter of fact, a progressive increase of hydrophobicity from sulfaguanidine to sulfathiazole due to the presence of the heterocyclic moieties was observed, which suggests an increasing affinity to bind to the MDM2-p53 complex, as also indicated by Wang et al.²³

Figure 2 shows the inhibition of the LANA expression by the BC3 cells treated with the sulfonamide compounds at 100 μM concentration. After 6 days of treatment with the sulfonamides, the BC3 cells were separated into two populations: the bottom right sector of the frame indicates BC3+/LANA+, the upper right shows the cells BC3+/LANA-. In this case too, the drugs significantly inhibited HHV8 latent antigen expression; the final data were about $65\pm4\%$ of LANA inhibition for SG3 (33% of residual fluorescence, RF), $69\pm5\%$ for BC6 (32% of RF), $78\pm6\%$ for SA4 (21% of RF) and $85\pm3\%$ for ST5 (17% of RF). In these conditions nimesulide showed a score of $54\pm5\%$ (about 46% of RF). All the data were normalized to 10^6 cells.

The cytotoxic activity of the tested drugs was detected on BC3 (KSHV permanently infected lymphoblasts) and THP-1 (KSHV-free macrophages) cell lines, and also on HUVEC cells (primary human umbilical cord cells) after 6 days of culture. Sulfonamide drugs showed a CC_{50} around 100 $\mu\text{g}/\text{ml}$ on BC3 cells (Table 1), while the values for CC_{50} on THP-1 varied from 30 $\mu\text{g}/\text{ml}$ (ST5) to 125 $\mu\text{g}/\text{ml}$ (SG3, SA4, BC6); primary HUVEC cells appeared generally more resistant to drug cytotoxicity, with CC_{50} of 125 $\mu\text{g}/\text{ml}$ (BC6), 250 (SA4 and ST5), and 500 $\mu\text{g}/\text{ml}$ (SG3). The drug EC_{50} was detected in BC3 cells and the relative selectivity indexes (SI) were 4.76 for SG3, 12.5 for SA4, 16.6 for ST5 and 8.3 for BC6.

The activity of the sulfonamide drugs was tested on the MDM2-p53 complex before and after its formation. Figure 3 shows that most of the tested compounds significantly inhibited the formation of the MDM2-p53 complex when added to the test wells before its establishment. In comparison with nutlin-3a, used as a positive control, ST5 showed a slightly stronger activity (about 70% inhibition compared to 62% of nutlin-3a); SA4 and BC6 also inhibited complex formation by 56 and 50% respectively. Sulfaguanidine was the least active compound in this test, with a score of about 48%. Finally, both nutlin-3a and sulfonamide compounds exerted some activity in disrupting the MDM2-p53 complex after its formation: ST5, the most active drug, disrupted the complex by about 48% ($p<0.01$), compared to the around 28% inhibition exerted by nutlin-3a ($p<0.05$). All the other sulfonamides showed non-significant values in this assay, whilst the difference between pre- and post-treatment was significant ($p<0.05$) for each compound.

DISCUSSION

Sulfonamide antibiotics were the earliest selective anti-bacterial drugs to be discovered and used for treating microbial infections. Their capacity to antagonize folic acid synthesis made them one of the most frequently prescribed and widely used antibiotics in the 1930s and

1940s.¹⁶ Afterwards, their use and importance decreased, since more active and less toxic natural antibiotics were discovered. Recently, the worldwide diffusion of bacterial resistance has induced many researchers to re-evaluate sulfonamide antibiotics in the treatment of bacterial infections.¹⁷

In this work we show that some of the most common sulfonamide drugs are endowed with strong activity on latently KSHV-infected BC3 cells, thereby allowing them to play a possible potential role in clearing the KSHV-infected lymphoma cells. These compounds interfere with the formation of the MDM2-p53 complex, which has been demonstrated to also contain LANA in KSHV infected BC3 lymphoma cells.¹⁸ MDM2, as well as LANA, also binds to the N terminus of p53.²⁰ This interaction seems to be sensitive to non-toxic doses of sulfonamides. Hence, the nature of the complex must be such as to allow sulfonamides access to the p53 N terminus and the p53-MDM2 interaction site, after which the viral genome detaches from the cellular DNA and is destroyed by DNA-polymerases; as a consequence LANA, which is the specific factor of KSHV latency, also stops being transcribed and is cleared from cell nucleus and cytoplasm. Our findings fit with the Chen et al. model,²⁰ demonstrating that MDM2-p53 disruption via nutlin-3a will also release LANA from p53. Moreover sulfonamides disrupted purified MDM2-p53 complex *in vitro*, suggesting a direct interaction between these drugs and the MDM2-p53 complex, which may have the correct structure to bind LANA.

It is well known that p53 is a key protein for the regulation of cell differentiation.^{18,19} Interaction with the oncogene MDM2 impairs the p53 activity in normal cells, and over-expression of MDM2 can dysregulate cell replication, leading to an oncogenic cell transformation. KSHV sustains the MDM2-p53 complex by adding the LANA protein, thereby abolishing the p53 function, after which the virus remains in a latent state inside the cell as an episome.^{20,21} Both sulfonamides and nutlin-3a are able to reactivate p53 thus enhancing its role as a cell genome regulator.^{22,23} KSHV associated lymphomas and KS are unusual, since they contain wild-type p53, unlike the majority of non-viral cancers, in which p53 is mutated. This suggests that the MDM2-p53 complex is perhaps activated by KSHV at the protein levels or that some of p53's activities may indeed be beneficial to support latent persistence.^{24,25} The MDM2 gene encodes a nuclear E3 ubiquitin-ligase, which is also transcriptionally-regulated by p53. Sulfonamide antibiotics are able to disrupt this complex and reactivate p53, leading to cell differentiation or occasionally to cell apoptosis.²⁶ Moreover, KSHV also targets the p53-related p73 and disruption of this complex can also induce apoptosis.²⁷

As regards a possible structure-activity relationship, Wang et al.²³ demonstrated that the sulfonamide moiety can play an important role in the enhanced potency of the molecules in the MDM2-p53 complex, since the sulfonamide side chain acts as a directing group that orients the side chemical groups into the pocket of the complex. Li et al.¹⁰ also analyzed the interaction between EBNA1 (the latent factor of Epstein-Barr gamma-Herpesvirus) and the MDM2-p53 complex. They found that the sulfonyl motif of the tested sulfonamide derivatives mimics the interaction with the receptor. In particular, compounds with modifications that extend the hydrophobic motifs might enable compound analogues to reach deeper into the EBNA1-binding pocket. Chen et al.²⁰ have recently suggested that

under normal conditions, nutlin-3a disrupts purified MDM2-p53 complex binding by intercalating at the p53-MDM2 surface. They concluded that MDM2 contributes to LANA's interaction with p53 either directly or by changing p53 conformation to enable it to bind LANA. Thus, the MDM2-p53 complex is the key that allows LANA binding to and stabilization on cellular DNA. Disrupting MDM2-p53 automatically induces the termination of HHV8 latency. Therefore, the sulfonamide antibiotics used in this study seem to behave almost similarly to nutlin-3a, by binding to the MDM2-p53 complex, which is destabilized and releases latent HHV8 factors; in addition the different hydrophobicity of the tested sulfonamides can justify the observed increasing inhibitory activity from sulfaguanidine to sulfathiazole.

In conclusion the overall finding of this work is that some sulfonamide compounds, widely used as antibacterial drugs, can be considered a new potential tool for inactivating KSHV, and thus clearing latent infection specifically in KSHV infected cells. In addition, sulfonamides inhibit latent KSHV at low μM concentrations, which can easily be reached after standard sulfonamide therapy (sulfamethoxazole serum concentration up to 40-60 $\mu\text{g/ml}$ after a standard oral dose of 800 mg).²⁸ It would also be interesting to verify whether other herpesviruses could be inhibited in their latent state by these drugs. This finding could open up a very promising program of studies on both oncogenic and degenerative diseases, in which herpesvirus latency is suspected to be involved.²⁹

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Ms Sally Davies for helping in the preparation and correction of the manuscript. RP, AI and DD designed and wrote the paper; FA, EP, and SU performed RT-PCR and ELISA assays, CM, RS, RB and WC performed all the fluorescence and cytofluorometry tests. This work was supported by Fondazione Banco di Sardegna 2014-15 and US Public health service grant DE018304 to DPD.

References

1. Dourmishev LA, Dourmishev AL, Palmeri D, Schwartz RA, Lukac DM. Molecular Genetics of Kaposi's Sarcoma-Associated Herpesvirus (Human Herpesvirus 8) Epidemiology and Pathogenesis. *Microbiol Mol Biol Rev.* 2003; 67:175–212. [PubMed: 12794189]
2. Ganem D. KSHV and the pathogenesis of Kaposi sarcoma: listening to human biology and medicine. *J Clin Invest.* 2010; 120:939–949. [PubMed: 20364091]
3. Ablashi DV, Chatlynne LG, Whitman JE Jr, Cesarman E. Spectrum of Kaposi's sarcoma-associated herpesvirus, or human herpesvirus 8, diseases. *Clin Microbiol Rev.* 2002; 15:439–464. [PubMed: 12097251]
4. Chandran B. Early Events in Kaposi's Sarcoma-Associated Herpesvirus Infection of Target Cells. *J Virol.* 2010; 84:2188–2199. [PubMed: 19923183]
5. Uldrick TS, Whitby D. Update on KSHV epidemiology, Kaposi Sarcoma pathogenesis, and treatment of Kaposi Sarcoma. *Cancer Lett.* 2011; 305:150–162. [PubMed: 21377267]
6. Wen KW, Damania B. Kaposi sarcoma-associated herpesvirus (KSHV): Molecular biology and oncogenesis. *Cancer Lett.* 2010; 289:140–150. [PubMed: 19651473]

7. Friedrichs C, Neyts J, Gaspar G, De Clercq E, Wutzler P. Evaluation of antiviral activity against human herpesvirus 8 (HHV-8) and Epstein–Barr virus (EBV) by a quantitative real-time PCR assay. *Antivir Res.* 2004; 62:121–123. [PubMed: 15130535]
8. Caselli E, et al. Retinoic acid analogues inhibit human herpesvirus 8 replication. *Antivir Ther.* 2008; 13:199–209. [PubMed: 18505171]
9. Krug LT, Pozharskaya VP, Yu Y, Inoue N, Offermann MK. Inhibition of infection and replication of human herpesvirus 8 in microvascular endothelial cells by alpha interferon and phosphonoformic acid. *J Virol.* 2004; 78:8359–8371. [PubMed: 15254208]
10. Li N, et al. Discovery of Selective Inhibitors Against EBNA1 via High Throughput In Silico Virtual Screening. *PLoS One.* 2010; 5(4):e10126. [PubMed: 20405039]
11. Paul AG, Sharma-Walia N, Chandran B. Targeting KSHV/HHV-8 Latency with COX-2 Selective Inhibitor Nimesulide: A Potential Chemotherapeutic Modality for Primary Effusion Lymphoma. *PLoS One.* 2011; 6(9):e24379. [PubMed: 21980345]
12. Angius F, et al. High-density lipoprotein contribute to G0-G1/S transition in Swiss NIH/3T3 fibroblasts. *Sci Rep.* 2015 Dec 7.5:17812.doi: 10.1038/srep17812 [PubMed: 26640042]
13. Caselli E, et al. High prevalence of HHV8 infection and specific killer cell immunoglobulin-like receptors allotypes in Sardinian patients with type 2 diabetes mellitus. *J Med Virol.* 2014; 86:1745–1751. [PubMed: 24122895]
14. Dedoni S, Olinas MC, Ingiani A, Onali P. Type I interferons up-regulate the expression and signalling of p75 NTR/TrkA receptor complex in differentiated human SH-SY5Y neuroblastoma cells. *Neuropharmacology.* 2014; 79:321–334. [PubMed: 24333329]
15. Böttger A, Sparks A, Liu WL, Howard SF, Lane DP. Design of a synthetic Mdm2-binding mini protein that activates the p53 response in vivo. *Curr Bio.* 1997; 17:860–869.
16. Shah SS, Rivera G, Ashfaq M. Recent advances in medicinal chemistry of sulfonamides. Rational design as anti-tumoral, anti-bacterial and anti-inflammatory agents. *Mini Rev Med Chem.* 2013; 13:70–86. [PubMed: 22625411]
17. Haruki H, Grønlund-Pedersen M, Gorska KI, Pojer F, Johnsson K. Tetrahydrobiopterin Biosynthesis as an Off-Target of Sulfa Drugs. *Science.* 2013; 340:987–991. [PubMed: 23704574]
18. Kojima K. Mdm2 inhibitor Nutlin-3a induces p53-mediated apoptosis by transcription-dependent and transcription-independent mechanisms and may overcome Atm-mediated resistance to fludarabine in chronic lymphocytic leukemia. *Blood.* 2006; 108:993–1000. [PubMed: 16543464]
19. Sarek G, Ojala P. M p53 Reactivation Kills KSHV Lymphomas Efficiently In Vitro and In Vivo: New Hope for Treating Aggressive Viral Lymphomas. *Cell Cycle.* 2007; 6:2205–2209. [PubMed: 17890905]
20. Chen W, Hilton IB, Staudt MR, Burd CE, Dittmer DP. Distinct p53, p53:LANA, and LANA complexes in Kaposi's Sarcoma-associated Herpesvirus Lymphomas. *J Virol.* 2010; 84:3898–3908. [PubMed: 20130056]
21. Ohsaki E, Ueda K. Kaposi's Sarcoma-Associated Herpesvirus Genome Replication, Partitioning, and Maintenance in Latency. *Front Microbiol.* 2012; 3:7. [PubMed: 22291692]
22. Shen H, Maki CG. Pharmacologic activation of p53 by small-molecule MDM2 antagonists. *Curr Pharm Des.* 2011; 17:560–568. [PubMed: 21391906]
23. Wang W, et al. Design, synthesis and biological evaluation of novel 3,4,5-trisubstituted aminothiophenes as inhibitors of p53–MDM2 interaction. Part 1. *Bioorg Med Chem.* 2013; 21:2879–2885. [PubMed: 23601819]
24. Petre CE, Sin SH, Dittmer DP. Functional p53 signaling in Kaposi's sarcoma-associated herpesvirus lymphomas: implications for therapy. *J Virol.* 2007; 81(4):1912–1922. [PubMed: 17121789]
25. Sarek G, et al. Reactivation of the p53 pathway as a treatment modality for KSHV-induced lymphomas. *J Clin Invest.* 2007; 117(4):1019–1028. [PubMed: 17364023]
26. Wang H, Ma H, Ren S, Buolamwini JK, Yan CA. Small-Molecule Inhibitor of MDMX Activates p53 and Induces Apoptosis. *Mol Cancer Ther.* 2011; 10:69. [PubMed: 21075910]
27. Santag S, et al. Recruitment of the tumour suppressor protein p73 by Kaposi's Sarcoma Herpesvirus latent nuclear antigen contributes to the survival of primary effusion lymphoma cells. *Oncogene.* 2013; 32:3676–3685. [PubMed: 22964633]

28. Bruun JN, Ostby N, Bredesen JE, Kierulf P, Lunde PK. Sulfonamide and trimethoprim concentrations in human serum and skin blister fluid. *Antimicrob Agents Chemother.* 1981; 19:82–85. [PubMed: 7247364]
29. Tselis A. Evidence for viral etiology of multiple sclerosis. *Semin Neurol.* 2011; 31:307–316. [PubMed: 21964847]

Figure 1. Sulfonamide compounds clear latent KSHV DNA from BC3 cells

qPCR was applied to the orf50 KSHV gene for detecting viral DNA in latently infected BC3 cells. The drugs were used at 100 μ M concentration (SG3= 21.4 μ g/ml; SA4= 17.2 μ g/ml; ST5= 25.5 μ g/ml; BC6= 25.3 μ g/ml; nimesulide= 30.8 μ g/ml) and left on the cells for 6 days before performing the assay. Nimesulide was used as reference positive control. The data are reported as a percentage of the untreated BC3 control. Data were normalized to 10^6 cells and are expressed as the mean $\pm$ SE of three independent experiments.

Figure 2. Cytofluorometry assay for detecting the KSHV LANA latency antigen in BC3 cells
 The numbers in the frame indicate the percentage of residual fluorescence within the drug-treated latently infected BC3 cells and the data were normalized to 10^6 cells. All the drugs were used at the concentration of 100 μ M and left on the cell culture for 6 days. Each frame was set so that some sectors were identified: the bottom right sector indicates BC3+/LANA+ cells, the upper right shows the cells BC3+/LANA-; the bottom left is a negative control. All the sulfonamide drugs tested scored significant values as compared to the untreated BC3 control ($p < 0.05$).

Figure 3. Inhibitory activity of sulfonamide compounds on MDM2-p53 complex formation
 Empty columns (pre-treat): the compounds were added to the reaction mixture before the formation of the MDM2-p53 complex; black columns (post-treat): the drugs were added to the reaction after the formation of the MDM2-p53 complex. All the compounds were used at a concentration of 100 μ M (SG3= 21.4 μ g/ml; SA4= 17.2 μ g/ml; ST5= 25.5 μ g/ml; BC6= 25.3 μ g/ml; nutlin-3a= 58.1 μ g/ml). The values are reported as a percentage of the negative control. Nutlin-3a was used as a positive control. Data are expressed as mean $\pm$ SE of three independent experiments.

Cytotoxic activity of sulfonamide drugs on both human primary and tumor line cells and detection of EC₅₀ and SI on BC3 cells.

Table 1

Drugs	CC50 ^a μg/ml±SE			EC50 ^b μg/ml±SE	SI ^c (CC50/EC50)
	HUVEC	THP-1	BC3		
SG3	500±125	125±31	100±25	21±5	4.76
SA4	250±125	125±31	100±50	8±4	12.5
ST5	250±62	30±15	100±25	6±2	16.6
BC6	125±31	125±62	100±12	12±4	8.3

^aCC50: cytotoxic concentration that inhibits cell viability by 50%.

^bEC50: effective concentration was detected in BC3 cells by RT-PCR as inhibition of latent HHV8 DNA;

^cSI: selectivity index; Cells: BC3= HHV8 permanently infected lymphoblastic cell line; THP-1: macrophagic cell line; HUVEC= primary umbilical cord human cells. SG3: sulfaguanidine; SA4: sulfanilamide; ST5: sulfathiazole; BC6: sulfamethoxazole. Data are expressed as mean ± SE of three independent experiments.