Published in final edited form as:

J Geriatr Oncol. 2018 May ; 9(3): 194–203. doi:10.1016/j.jgo.2017.11.005.

Healthcare Experience among Older Cancer Survivors: Analysis of the SEER-CAHPS Dataset

Michael T. Halpern^{a,b}, Matthew P. Urato^a, Lisa M. Lines^a, Julia B. Cohen^a, Neeraj K. Arora^c, and Erin E. Kent^d

^aRTI International, 3040 East Cornwallis Road, Research Triangle Park, NC, 27709 USA and 307 Waverley Oaks Road, Suite 101, Waltham, MA, 02452 USA

^bTemple University, 1301 Cecil B. Moore Ave., Philadelphia PA, 19122 USA

^cPatient-Centered Outcomes Research Institute (PCORI), 1828 L Street, NW, Suite 900 Washington, DC, 20036 USA

^dNational Cancer Institute, 9609 Medical Center Drive, Rockville, MD 20850 USA

Abstract

Objective—Little is known about factors affecting medical care experiences of cancer survivors. This study examined experience of care among cancer survivors and assessed associations of survivors' characteristics with their experience.

Materials and Methods—We used a newly-developed, unique data resource, SEER-CAHPS (NCI's Surveillance Epidemiology and End Results [SEER] data linked to Medicare Consumer Assessment of Healthcare Providers and Systems [CAHPS] survey responses), to examine experiences of care among breast, colorectal, lung, and prostate cancer survivors age >66 years who completed CAHPS>1 year after cancer diagnosis and survived >1 year after survey completion. Experience of care was assessed by survivor-provided scores for overall care, health plan, physicians, customer service, doctor communication, and aspects of care. Multivariable logistic regression models assessed associations of survivors' sociodemographic and clinical characteristics with care experience.

Results—Among 19,455 cancer survivors with SEER-CAHPS data, higher self-reported general-health status was significantly associated with better care experiences for breast, colorectal, and prostate cancer survivors. In contrast, better mental-health status was associated with better care experience for lung cancer survivors. College-educated and Asian survivors were less likely to

Corresponding Author: Michael T. Halpern, MD, PhD, MPH, Dept. of Health Services Management and Policy, Temple University College of Public Health, 1301 Cecil B. Moore Ave, Ritter Annex 533, Philadelphia, PA 19122 USA, Phone: 1-215-204-5814, Fax: 1-215-204-5958, michael.halpern@temple.edu.

Author Contribution and Conflict of Interest

All authors participated in conception and design of this study, analysis and interpretation of data, manuscript writing, and approval of final article. The authors declare no conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

indicate high scores for care experiences. Few differences in survivors' experiences were observed by sex or years since diagnosis.

Conclusions—The SEER-CAHPS data resources allows assessment of factors influencing experience of cancer among U.S. cancer survivors. Higher self-reported health status was associated with better experiences of care; other survivors' characteristics also predicted care experience. Interventions to improve cancer survivors' health status, such as increased access to supportive care services, may improve experience of care.

Keywords

cancer; SEER; Medicare; survivors; patient satisfaction; CAHPS

Introduction

As of January 2016, there were approximately 15.5 million cancer survivors in the United States; this will grow to 20 million by 2026 [1]. Approximately three-quarters of cancer survivors are age 60 or older, and almost one-half are age 70 or older [1]. Cancer survivors have unique post-treatment needs, including surveillance for tumor recurrence and development of other cancers; treatment of chronic/late-occurring effects due to cancer or cancer treatment; and increased needs for preventive care and health promotion [2]. Cancer survivors also frequently experience needs for a range of psychosocial services [3]. Symptoms experienced among cancer survivors and their effects vary based on cancer treatment, stage at diagnosis, sociodemographic characteristics, and health status [4].

There is limited data on factors affecting the experience of care among cancer survivors, particularly older survivors [5]. The experience of survivorship care may differ substantially in older versus younger survivors. Older survivors may demonstrate better psychosocial adaptation to cancer than do younger survivors [6]. However, older cancer survivors may also experience more rapid decline in functional status compared with younger survivors or with age-matched individuals in the general population [7], and older age may be a risk factor for cancer treatment-related complications [8]. In addition, older survivors are more likely to have multiple chronic conditions, which may affect outcomes including quality of life and survival [9]. Almost 40% of older survivors have at least one on-going symptom due to cancer or cancer treatment [10]; older survivors with multiple chronic conditions are less resistant to decreases in physical functioning and are less likely to recover lost functional ability [11]. Overall, older survivors have unique needs, and their survivorship care should be tailored to these needs to enhance the experience of care [12].

SEER-CAHPS is a recently developed data resource linking patient-reported information from the Medicare Consumer Assessment of Healthcare Providers and Systems (M-CAHPS®) Survey with clinical information from NCI's Surveillance, Epidemiology, and End Results (SEER) Program [13, 14]. This unique resource is the first dataset permitting examination of sociodemographic and clinical characteristics and experiences of care among large U.S. populations of individuals diagnosed with cancer, including cancer survivors. The objective of this study is to examine the experience of care for older cancer survivors—

including experience with physicians, health plans, and other aspects of medical care—and assess associations of survivor characteristics with these experiences.

Materials and Methods

SEER-CAHPS contains information from Medicare beneficiaries who responded to the CAHPS survey and were diagnosed with their first recorded primary cancer while residing in SEER regions [13, 14]. We used data from Medicare beneficiaries diagnosed with non-metastatic (stage I-III) breast, colorectal, lung, or prostate cancer in SEER regions who completed the CAHPS survey from 2000–2011. Males with breast cancer were excluded (N=49). We included individuals who: (1) responded to an M-CAHPS survey at least once, at least one-year after cancer diagnosis (diagnosis dates 1988–2009); (2) survived for at least one-year after responding to the survey; and (3) were age 66 or older at survey completion (N=19,455). Details regarding sample selection, administration, and response rates for the M-CAHPS are presented elsewhere.[15–17]

Responses to four M-CAHPS global ratings were included in this study: overall care, health plan, primary physician, and specialist physician. Each measure was rated on a 0–10 scale, with zero representing the lowest rating. M-CAHPS responses also included patient reports on five composite, multi-item measures: *customer service*, *doctor communication*, *getting care quickly*, *getting needed care*, and *getting needed prescription drugs*. Each composite measure was scored 0–100. Appendix Table 1 presents descriptive statistics for each CAHPS measure.

Similar to previous CAHPS analysis,[18] because global ratings were very negatively skewed, with most responses at the extreme upper end of scales, we classified global ratings (scored 0–10) as indicating “high” experiences of care if responses were 9 or 10. The multi-item composite scores (scored 0–100) exhibited ceiling effects and few respondents provided scores between 90–100; we classified survivors’ reports on composite measures to indicate “high” care experiences for responses of 100. Previous studies have also dichotomized CAHPS ratings or reports, defining the upper scores as “high” [19–21]. Use of the term “high” to classify upper-end scores does not imply that M-CAHPS participants judged their experience of care as high compared with other experiences of care. Details regarding these measures are presented elsewhere [22, 23]. Additional information on M-CAHPS is at <https://cahps.ahrq.gov/surveys-guidance/hp/about/Medicare-CAHPS-HP-Survey.html>.

We examined associations of survivor characteristics with high scores for experience of care using multivariable logistic regression analyses, with separate models for each of the four global and five composite measures. Dependent variables were specified as high/not high for each measure. Sociodemographic independent variables (with reference categories listed first) were: sex (female/male, for colorectal and lung cancer survivors only); race/ethnicity (Non-Hispanic White, Non-Hispanic Black, Non-Hispanic Asian, Hispanic, or other); education (less than high school, high school, some college, or college graduate); Medicare plan type (managed-care vs. fee-for-service); and age at completion of CAHPS survey (66–74, 75–79, 80–84, or 85+).

Survivors' health status was controlled for using self-reported general health status (GHS) and mental health status (MHS) on a 5-point scale (poor, fair, good, very good, or excellent). In regression analyses, excellent and very good health status were combined due to infrequent excellent ratings for GHS. Previous studies of factors associated with CAHPS ratings generally use self-rated health status (GHS and MHS) to control for patient comorbidities. Studies using other survey instruments to capture information from cancer survivors on self-rated quality of care have also found that health status is a more relevant measure than comorbidities for this population [24].

In addition, we controlled for years since cancer diagnosis (1–2, more than 2 to 5, more than 5 years from cancer diagnosis to completion of CAHPS survey); stage at cancer diagnosis (1 vs. 2 vs. 3); geographic location (state of residence at diagnosis); number of primary cancers at diagnosis (single vs. multiple); and survey year (2000–2011). Regression analysis results for years since cancer diagnosis and stage at diagnosis are presented in Appendix Tables 2 and 3. All analyses were performed on unweighted data using SAS 9.4 (SAS Institute, Cary NC). Statistical significance was assessed at $p < 0.05$. As this study focused on hypothesis generation (i.e., identifying factors associated with experience of care among older cancer survivors), statistical significance was not corrected for multiple comparisons [25]. This study was classified as exempt by the RTI Institutional Review Board.

Results

Characteristics of Study Population

Table 1 presents characteristics of the study population by cancer type. Colorectal cancer survivors were the oldest group; prostate cancer survivors were the youngest and included the highest proportions of college graduates and Black and Hispanic individuals. Medicare fee-for-service (vs. managed care) status was similar across all groups.

The distribution of general health status (GHS) and mental health status (MHS) category among the study population are also presented in Table 1. For all cancer types, the proportion of survivors reporting excellent/very good GHS was less than the proportion reporting excellent/very good MHS. This was particularly evident for lung cancer survivors; 21.5% of this groups reported excellent/very good GHS while 58.5% indicated excellent/very good MHS. Pearson correlations coefficients between GHS and MHS by cancer type were: breast, 0.45; colorectal, 0.49; lung, 0.44; and prostate, 0.49 (data not shown).

Table 1 also presents the proportion of individuals in each survivor group indicating high scores for each CAHPS measure. Few large differences were observed across cancer types for a specific measure. The measure with the lowest proportion of high scores was Customer Service (51.9%), while the measures with the greatest proportion indicating high scores were Getting Needed Prescription Drugs (77.2%), Specialist Physician (73.3%), and Primary Physician (72.9%).

Association of GHS with High Scores for Experience of Care

Figure 1 presents results from multivariable regression analyses examining associations of very good/excellent GHS with high scores for survivor's experience. The top graph (Figure

1a) presents associations for global ratings; the bottom graph (Figure 1b) presents associations for composite measures. Among all four survivor groups, those with very good/excellent general health status were significantly more likely to provide high ratings for all four global measures, except for specialist physician ratings among lung cancer survivors (Figure 1a). Breast, colorectal, and prostate cancer survivors with very good/excellent general health status were also significantly more likely to indicate high scores for experience with almost all composite measures (Figure 1b). However, among lung cancer survivors, very good/excellent GHS was significantly associated with high scores for only one composite measure (getting needed care).

Association of MHS with High Scores for Experience of Care

Figure 2 presents results from multivariable regression analyses examining associations of very good/excellent MHS with high scores for experience of care. The top graph (Figure 2a) presents associations for global ratings, while the bottom graph (Figure 2b) presents associations for composite measures. Significant associations of MHS with the global ratings (Figure 2a) were inconsistent for breast, colorectal, and prostate cancer survivors. However, among lung cancer survivors, all global ratings except health plan were significantly associated with very good/excellent MHS. Lung cancer survivors with very good/excellent MHS were six to seven times more likely to provide high ratings for overall care, primary physician, and specialist physician than were those with poor MHS. The magnitudes of the associations between lung cancer survivors' MHS and likelihood of high CAHPS ratings were larger than any other odds ratios in this study.

While very good/excellent GHS was associated with high composite measure scores among breast, colorectal, and prostate cancer survivors (Figure 1b), there were no significant associations between very good/excellent MHS and high composite scores for colorectal and prostate cancer survivors (Figure 2b). Very good/excellent MHS was significantly associated with high experience of care for two composite measures among breast cancer survivors and three measures among lung cancer survivors. As with the global measures (Figure 2a), odds ratios for associations of very good/excellent MHS among lung cancer survivors with high CAHPS composite measure scores were notably large.

Association of Age with High Scores for Experience of Care

Table 2 presents results from multivariable regression analyses examining the association of survivor characteristics with provision of high scores for the global ratings, while Table 3 presents similar regression analysis results for the composite measures. Separate results are present by cancer site. Among breast, colorectal, and prostate cancer survivors, older age was associated with increased likelihood of high health plan rating (Table 2); older colorectal survivors were also more likely to indicated high ratings for specialist physician. In contrast, among prostate cancer survivors, older age was associated with decreased likelihood of high ratings for customer service and doctor communications (Table 3). There were no significant associations between age and high ratings among lung cancer survivors.

Association of Education and Sex with High Scores for Experience of Care

College graduate breast cancer survivors were significantly less likely to provide high ratings for all global measures and for doctor communications and getting needed care. Highly educated colorectal and prostate cancer survivors were also less likely to provide high ratings for health plan. Education was not significantly associated with high scores among lung cancer survivors.

Sex was included as a covariate in models of colorectal and lung cancer survivors. Among both these groups, men (vs. women) were significantly less likely to provide high ratings for two of the nine measures, health plan or primary physician.

Association of Race/Ethnicity with High Scores for Experience of Care

Asian survivors (compared with non-Hispanic White survivors) were less likely to indicate high scores for most composite measures among breast, colorectal, and prostate cancer survivors (Table 3); for two composite measures (getting care quickly and getting needed prescription drugs) among lung cancer survivors (Table 3); and for two global measures (overall care and specialist physician) among breast cancer survivors (Table 2). Compared to non-Hispanic Whites, Black breast cancer survivors were more likely to indicate high scores for experience with health plan and customer service, while Black colorectal cancer survivors were more likely to provide high ratings for their primary physician and Black prostate cancer survivors were more likely to provide high scores for health plan, customer service, and getting needed care. Results among Hispanic survivors were mixed. Hispanic breast cancer survivors were more likely to provide high ratings for health plan and getting needed care. Hispanic colorectal cancer survivors were more likely to provide high ratings for overall care and health plan. However, Hispanic colorectal cancer survivors (and Hispanic prostate cancer survivors) were less likely to provide high scores for getting care quickly and getting needed drugs (Table 3).

Association of Medicare Plan Type with High Scores for Experience of Care

Among breast and prostate cancer survivors, having fee-for-service Medicare (vs. managed care) was associated with decreased likelihood of scoring customer service highly, but increased likelihood of high scores for getting needed prescription drugs. Breast cancer survivors with fee-for-service Medicare were also more likely to indicate high scores for access to care, getting care quickly and getting needed care.

Discussion

This is the first study to use data from SEER-CAHPS, a newly developed data resource, to examine associations of experience with health plans, physicians, and medical care with characteristics of older cancer survivors. In this population, self-reported health status is strongly associated with experience of care. Overall, we found that more than 50% of older cancer survivors provided high scores for each CAHPS measure. However, breast, colorectal, lung, and prostate cancer survivors with very good/excellent GHS were significantly more likely to provide high scores for the majority of global rating measures included in the study; breast, colorectal, and prostate cancer survivors with very good/

excellent GHS were also more likely to indicate high scores for most composite measures. In contrast, higher MHS among breast, colorectal, and prostate cancer survivors had few significant associations with higher scores, while lung cancer survivors with very good/excellent MHS were significantly more likely to indicate high scores for the majority of their global and composite measures. The cause of the difference between lung cancer survivors and the other survivor groups is uncertain. Only 20% of lung cancer survivors reported very good/excellent GHS versus approximately 30% of breast and colorectal cancer survivors and 38% of prostate cancer survivors (Table 1). As health status is a subjective measure, it is possible that even those lung cancer survivors reporting very good/excellent GHS experienced greater health limitations than did survivors of other types of cancer. Among survivors with lower GHS, MHS may assume more importance in their experience of medical care. In addition, lung cancer survivors may experience a greater decline over time in GHS, which has been associated with decreased satisfaction [26]. Due to the cross-sectional nature of the CAHPS data, we are unable to assess changes in survivors' experience or health status over time.

Previous studies have also reported links between self-rated health status and outcomes among individuals with cancer. High health status was associated with increased satisfaction and increased survival among cancer survivors [24, 27–29], while being off work due to health reasons, anxiety and depression, and perceived stress were associated with decreased satisfaction [30, 31]. Among overweight or obese older cancer survivors, increased general health status or mental health status was associated with increased resistance to declines in physical functioning [11]. While older survivors are more likely to have multiple chronic conditions, older survivors are less likely to report being in excellent or very good health compared with age-matched individuals without cancer even after controlling for comorbidities [32].

Our findings and those of earlier studies highlight the importance of supportive care services in survivorship care. These services can improve physical/mental health, quality of life, and satisfaction among individuals with cancer [33–36]. For example, eliminating surgical side effects and comorbidities among breast cancer survivors helped to prevent reductions in self-rated health [37]. In addition, supportive care services can assist with comorbidities experienced by survivors; these comorbidities may be predictive of functional declines among older survivors [38]. Older adults with cancer are interested in supportive care services to improve their health-related quality of life [39], and recent recommendations advocate earlier integration of supportive/palliative care into standard oncology care, which may improve survivors' experience of care [35, 40, 41]. Inclusion of health promotion activities, a key component of survivorship care, may also be particularly important for older survivors. Improvements in modifiable risk factors including obesity, physical activity, and diet are associated with increased functional status and quality of life among older cancer survivors, which may improve experience of care [7, 42]. For example, older female survivors who are physically-active have similar general health, physical functioning, and other quality of life domains compared with age-matched women who never had cancer, while physically-inactive survivors have significantly decreased quality of life [42]. Exercise interventions in older survivors can decrease symptoms and may increase survival [6].

The significant associations between self-rated health status and experience of care also highlights the importance of assessing and managing long-term and late effects of cancer and cancer treatment among survivors [43]. Systematic review have documented that survivors may experience cancer-related symptoms for multiple years following diagnosis [44–46]. Therefore, interventions to improve the health status of older survivors may have long-term impacts on the experience of care.

In the present study, associations of older survivors' characteristics with experience of care varied substantially by cancer type and category of experience being assessed. Previous studies examining associations of survivors' experience of or satisfaction with care have also shown varied results. Several studies have reported that satisfaction among survivors is not significantly associated with age, sex, ethnicity, or education [24, 30, 31], although others have reported that more highly educated and African American survivors have lower satisfaction [47]. Studies have also found that female and older survivors and those with better health status had more positive assessments of the quality of their care, and satisfaction varied substantially by cancer type [24, 47–49].

Similar to previous CAHPS analyses from broader populations [50, 51], our study found that Asian survivors were less likely to indicate high scores for experience of care, particularly for the composite measures, than were non-Hispanic Whites; this difference is not likely due to lack of measurement equivalence [50]. In contrast, Hispanic colorectal cancer survivors were significantly more likely to provide high ratings for health plan and overall care, but significantly less likely to report high scores for getting care quickly and getting needed prescription drugs. It is unclear why Hispanic colorectal cancer survivors would be more likely to rate their health plan and overall care highly but less likely to report high scores for other aspect of care. These responses appear contradictory, as the ability to get care quickly and get needed prescription drugs would presumably influence survivors' experience of overall care and health plan. While these findings are limited by the accuracy of the SEER and Medicare race/ethnicity variables, they suggest that cancer survivorship care programs need to consider and address potential distinctions in experience based on cultural differences and patient preferences.

Among breast cancer survivors, those with fee-for-service Medicare were more likely to indicate high scores for three composites (getting care quickly, getting needed care, and getting needed prescription drugs) but less likely for customer service. The decreased likelihood of high scores for customer service among fee-for-service Medicare survivors may reflect the lack of a unified health care plan; that is, in contrast to Medicare Advantage, fee-for-service does not have a single entity or organization responsible for overall care for a beneficiary. Despite the increased likelihood of high scores for customer services among survivors with Medicare Advantage, the significantly lower likelihood of high scores for three other composites suggest that these managed care plans need to improve access to and delivery of survivorship services.

To our knowledge, no published study has compared CAHPS responses between Medicare beneficiaries diagnosed with cancer and those not diagnosed with cancer. Thiels et al. examined the impact of a cancer diagnosis on CAHPS scores for individuals (Medicare

beneficiaries and those with other insurance) who were hospitalized [52]. In this population, patients with a cancer diagnosis were significantly more likely to have high Hospital CAHPS (HCAHPS) scores in pain management and communication with physician.

Limitations to our study stem primarily from the cross-sectional study design of the CAHPS survey that does not allow us to explore changes in survivors' experience for individual respondents over time. The study population was limited to individuals living in states that participate in the SEER program who completed the CAHPS survey; results may not be generalizable to all cancer survivors. A strong ceiling effect was observed, with more than 50% of survivors indicating high ratings for each CAHPS measure. Although validations of CAHPS ratings are not available, CAHPS surveys have been widely used for almost 20 years to assess patient-reported perceptions of care and associations of patient demographic, clinical, and insurance-related factors with these ratings [53, 54]. In addition, while GHS and MHS are not able to be compared or validated with other measures of survivors' health status, self-reported GHS and MHS are widely recognized as appropriate predictors of morbidity and mortality [55–57]. Furthermore, completion of CAHPS is not uniform across the SEER-Medicare population; for example, individuals in Medicare Advantage plans are more likely to have the opportunity to complete this survey. Therefore, the study population may not be representative of the overall SEER-Medicare population. Also, while information on cancer recurrence would be useful for this study, SEER does not include this information and use of Medicare claims would miss a majority of individuals experiencing recurrence [58].

Despite these limitations, this study provides important information about experiences of care for older cancer survivors using a new resource, SEER-CAHPS. Among these individuals, experience with health plans, physicians, and aspects of medical care are associated with sociodemographic, insurance, and clinical characteristics. These findings provide guidance for developing patient-centered programs to improve experience with care among older cancer survivors. Survivorship care for older individuals is affected by factors including functional and cognitive impairments, multiple comorbidities, distinct goals of care, and increased needs for social support [59]. Subsequent research is needed to examine programs addressing potentially modifiable factors, including obesity, physical activity, diet, GHS and MHS, on older survivors' experience with care and to enhance the tailoring of survivorship programs to individuals' characteristics. In particular, evaluations of tailored, patient-centered programs and policies addressing older survivors' health status concerns, risk factors, and unmet needs are needed to assess the importance of and potential for supportive care and lifestyle interventions to improve health and experiences of care.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was funded by the National Cancer Institute, contract number HHSN261201000166U and HHSN261201500132U. All statements in this report, including its findings and conclusions, are solely those of the

authors and do not necessarily represent the views of the Patient-Centered Outcomes Research Institute (PCORI), its Board of Governors or Methodology Committee, or the National Cancer Institute.

References

1. Miller KD, Siegel RL, Lin CC, Mariotto AB, Kramer JL, Rowland JH, et al. Cancer treatment and survivorship statistics, 2016. *CA: a cancer journal for clinicians*. 2016; 66:271–89. [PubMed: 27253694]
2. Halpern MT, Viswanathan M, Evans TS, Birken SA, Basch E, Mayer DK. Models of Cancer Survivorship Care: Overview and Summary of Current Evidence. *Journal of oncology practice / American Society of Clinical Oncology*. 2014; 11:e19–27.
3. Landier W. Survivorship care: essential components and models of delivery. *Oncology (Williston Park, NY)*. 2009; 23:46–53.
4. LeMasters T, Madhavan S, Sambamoorthi U, Kurian S. A population-based study comparing HRQoL among breast, prostate, and colorectal cancer survivors to propensity score matched controls, by cancer type, and gender. *Psycho-oncology*. 2013; 22:2270–82. [PubMed: 23606210]
5. Halpern MT, McCabe MS, Burg MA. The Cancer Survivorship Journey: Models of Care, Disparities, Barriers, and Future Directions. American Society of Clinical Oncology educational book / ASCO American Society of Clinical Oncology Meeting. 2016; 35:231–9.
6. Rowland JH, Bellizzi KM. Cancer survivorship issues: life after treatment and implications for an aging population. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2014; 32:2662–8. [PubMed: 25071099]
7. Kenzik KM, Morey MC, Cohen HJ, Sloane R, Demark-Wahnefried W. Symptoms, weight loss, and physical function in a lifestyle intervention study of older cancer survivors. *Journal of geriatric oncology*. 2015; 6:424–32. [PubMed: 26362355]
8. Bellizzi KM, Mustian KM, Palesh OG, Diefenbach M. Cancer Survivorship and Aging: Moving the Science Forward. *Cancer*. 2008; 113:3530–9. [PubMed: 19058147]
9. Flannery M, Mohile SG, Dale W, Arora NK, Azar L, Breslau ES, et al. Interventions to improve the quality of life and survivorship of older adults with cancer: The funding landscape at NIH, ACS and PCORI. *Journal of geriatric oncology*. 2016; 7:225–33. [PubMed: 27197917]
10. Avis NE, Deimling GT. Cancer survivorship and aging. *Cancer*. 2008; 113:3519–29. [PubMed: 19058151]
11. Duan-Porter W, Cohen HJ, Demark-Wahnefried W, Sloane R, Pendergast JF, Snyder DC, et al. Physical resilience of older cancer survivors: An emerging concept. *Journal of geriatric oncology*. 2016; 7:471–8. [PubMed: 27478133]
12. Guerard EJ, Nightingale G, Bellizzi K, Burhenn P, Rosko A, Artz AS, et al. Survivorship care for older adults with cancer: U13 conference report. *Journal of geriatric oncology*. 2016; 7:305–12. [PubMed: 27424802]
13. Chawla N, Urato M, Ambs A, Schussler N, Hays RD, Clauser SB, et al. Unveiling SEER-CAHPS(R): a new data resource for quality of care research. *Journal of general internal medicine*. 2015; 30:641–50. [PubMed: 25586868]
14. National Cancer Institute. SEER-CAHPS Linked Data Resource. 2017. <https://healthcaredelivery.cancer.gov/seer-cahps/>
15. Centers for Medicare & Medicaid Services. Quality Assurance Protocols & Technical Specifications Version 7.0. Medicare Advantage and Prescription Drug Plan CAHPS® Survey. 2016. http://www.ma-pdpcahps.org/globalassets/ma-pdp/quality-assurance/final_ma-pdp-cahps-qapts-v7-final-508-7-10-17.pdf
16. Centers for Medicare & Medicaid Services. Implementation of Medicare CAHPS® Fee-for-Service Survey—Final Report for Year 1. https://www.cms.gov/Research-Statistics-Data-and-Systems/Research/ConsumerResearch/Downloads/01_FFSCAHPSYr1SYN.pdf
17. Hays, R. The SEER and Medicare CAHPS Linked Dataset. 2015. <https://www.ahrq.gov/sites/default/files/wysiwyg/cahps/news-and-events/events/ahrq-conference-2015/cahps-preconf-hays.pdf>

18. Jha AK, Orav EJ, Zheng J, Epstein AM. Patients' perception of hospital care in the United States. *The New England journal of medicine*. 2008; 359:1921–31. [PubMed: 18971493]
19. Schmocker RK, Cherney Stafford LM, Siy AB, Levenson GE, Winslow ER. Understanding the determinants of patient satisfaction with surgical care using the Consumer Assessment of Healthcare Providers and Systems surgical care survey (S-CAHPS). *Surgery*. 2015; 158:1724–33. [PubMed: 26195107]
20. Tevis SE, Kennedy GD. Patient satisfaction: does surgical volume matter? *The Journal of surgical research*. 2015; 196:124–9. [PubMed: 25796107]
21. Zweifler J, Hughes S, Lopez RA. Controlling for race/ethnicity: a comparison of California commercial health plans CAHPS scores to NCBD benchmarks. *International Journal for Equity in Health*. 2010; 9:4. [PubMed: 20181011]
22. Damiano PC, Elliott M, Tyler MC, Hays RD. Differential use of the CAHPS® 0–10 global rating scale by Medicaid and commercial populations. *Health Serv Outcomes Res Methodol*. 2004; 5:193–205.
23. Hargraves JL, Hays RD, Cleary PD. Psychometric properties of the Consumer Assessment of Health Plans Study (CAHPS) 2.0 adult core survey. *Health services research*. 2003; 38:1509–27. [PubMed: 14727785]
24. Arora NK, Reeve BB, Hays RD, Clauser SB, Oakley-Girvan I. Assessment of quality of cancer-related follow-up care from the cancer survivor's perspective. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2011; 29:1280–9. [PubMed: 21357781]
25. Feise RJ. Do multiple outcome measures require p-value adjustment? *BMC medical research methodology*. 2002; 2:8. [PubMed: 12069695]
26. Nguyen TV, Anota A, Bredart A, Monnier A, Bosset JF, Mercier M. A longitudinal analysis of patient satisfaction with care and quality of life in ambulatory oncology based on the OUT-PATSAT35 questionnaire. *BMC cancer*. 2014; 14:42. [PubMed: 24460858]
27. Gupta D, Patel K, Lis CG. Self-rated health supersedes patient satisfaction with service quality as a predictor of survival in prostate cancer. *Health and quality of life outcomes*. 2015; 13:137. [PubMed: 26337960]
28. Hoffman RM, Hunt WC, Gilliland FD, Stephenson RA, Potosky AL. Patient satisfaction with treatment decisions for clinically localized prostate carcinoma. Results from the Prostate Cancer Outcomes Study. *Cancer*. 2003; 97:1653–62. [PubMed: 12655522]
29. Lis CG, Patel K, Gupta D. The Relationship between Patient Satisfaction with Service Quality and Survival in Non-Small Cell Lung Cancer - Is Self-Rated Health a Potential Confounder? *PloS one*. 2015; 10:e0134617. [PubMed: 26230934]
30. Amler S, Sauerland MC, Deiters C, Buchner T, Schumacher A. Factors influencing life satisfaction in acute myeloid leukemia survivors following allogeneic stem cell transplantation: a cross-sectional study. *Health and quality of life outcomes*. 2015; 13:28. [PubMed: 25888906]
31. Zlatar ZZ, Meier EA, Montross Thomas LP, Palmer BW, Glorioso D, Marquine MJ, et al. Life satisfaction and its correlates among older cancer survivors: critical role of psychosocial factors. *Psycho-oncology*. 2015; 24:241–4. [PubMed: 24917214]
32. Keating NL, Norredam M, Landrum MB, Huskamp HA, Meara E. Physical and mental health status of older long-term cancer survivors. *Journal of the American Geriatrics Society*. 2005; 53:2145–52. [PubMed: 16398900]
33. Bauman JR, Temel JS. The integration of early palliative care with oncology care: the time has come for a new tradition. *Journal of the National Comprehensive Cancer Network : JNCCN*. 2014; 12:1763–71. quiz 71. [PubMed: 25505216]
34. Bruera E, Yennurajalingam S. Palliative care in advanced cancer patients: how and when? *The oncologist*. 2012; 17:267–73. [PubMed: 22252934]
35. Glare PA. Early implementation of palliative care can improve patient outcomes. *Journal of the National Comprehensive Cancer Network : JNCCN*. 2013; 11(Suppl 1):S3–9.
36. Nakajima N, Abe Y. Concurrent Specialized Palliative Care Upon Initiation of First-Line Chemotherapy for Cancer Progression: Is It Early Enough? *The American journal of hospice & palliative care*. 2016; 33:340–5. [PubMed: 25433067]

37. Schootman M, Deshpande AD, Pruitt S, Aft R, Jeffe DB. Estimated effects of potential interventions to prevent decreases in self-rated health among breast cancer survivors. *Annals of epidemiology*. 2012; 22:79–86. [PubMed: 22226030]
38. Garman KS, Pieper CF, Seo P, Cohen HJ. Function in elderly cancer survivors depends on comorbidities. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2003; 58:M1119–24.
39. Pergolotti M, Deal AM, Williams GR, Bryant AL, Bensen JT, Muss HB, et al. Activities, function, and health-related quality of life (HRQOL) of older adults with cancer. *Journal of geriatric oncology*. 2017; 8:249–54. [PubMed: 28285980]
40. Levy MH, Adolph MD, Back A, Block S, Codada SN, Dalal S, et al. Palliative care. *Journal of the National Comprehensive Cancer Network : JNCCN*. 2012; 10:1284–309. [PubMed: 23054879]
41. Shin J, Temel J. Integrating palliative care: when and how? Current opinion in pulmonary medicine. 2013; 19:344–9. [PubMed: 23715290]
42. Blair CK, Robien K, Inoue-Choi M, Rahn W, Lazovich D. Physical inactivity and risk of poor quality of life among elderly cancer survivors compared to women without cancer: the Iowa Women's Health Study. *Journal of cancer survivorship : research and practice*. 2016; 10:103–12. [PubMed: 26008207]
43. Runowicz CD, Leach CR, Henry NL, Henry KS, Mackey HT, Cowens-Alvarado RL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2016; 34:611–35. [PubMed: 26644543]
44. Harrington CB, Hansen JA, Moskowitz M, Todd BL, Feuerstein M. It's not over when it's over: long-term symptoms in cancer survivors--a systematic review. *International journal of psychiatry in medicine*. 2010; 40:163–81. [PubMed: 20848873]
45. Jansen L, Koch L, Brenner H, Arndt V. Quality of life among long-term (5 years) colorectal cancer survivors Systematic review. *Eur J Cancer*. 2010:46.
46. Maass SW, Roorda C, Berendsen AJ, Verhaak PF, de Bock GH. The prevalence of long-term symptoms of depression and anxiety after breast cancer treatment: A systematic review. *Maturitas*. 2015; 82:100–8. [PubMed: 25998574]
47. Mariano C, Hanson LC, Deal AM, Yang H, Bensen J, Hendrix L, et al. Healthcare satisfaction in older and younger patients with cancer. *Journal of geriatric oncology*. 2016; 7:32–8. [PubMed: 26774226]
48. Ayanian JZ, Zaslavsky AM, Guadagnoli E, Fuchs CS, Yost KJ, Creech CM, et al. Patients' perceptions of quality of care for colorectal cancer by race, ethnicity, and language. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2005; 23:6576–86. [PubMed: 16116149]
49. Charalambous A. Variations in patient satisfaction with care for breast, lung, head and neck and prostate cancers in different cancer care settings. *European journal of oncology nursing : the official journal of European Oncology Nursing Society*. 2013; 17:588–95. [PubMed: 23453567]
50. Hays RD, Chawla N, Kent EE, Arora NK. Measurement equivalence of the Consumer Assessment of Healthcare Providers and Systems (CAHPS(R)) Medicare survey items between Whites and Asians. *Qual Life Res*. 2016
51. Morales LS, Elliott MN, Weech-Maldonado R, Spritzer KL, Hays RD. Differences in CAHPS adult survey reports and ratings by race and ethnicity: an analysis of the National CAHPS benchmarking data 1.0. *Health services research*. 2001; 36:595–617. [PubMed: 11482591]
52. Thiels CA, Hanson KT, Yost KJ, Zielinski MD, Habermann EB, Cima RR. Effect of Hospital Case Mix on the Hospital Consumer Assessment of Healthcare Providers and Systems Star Scores: Are All Stars the Same? *Annals of surgery*. 2016; 264:666–73. [PubMed: 27355268]
53. Anhang Price R, Elliott MN, Zaslavsky AM, Hays RD, Lehrman WG, Rybowski L, et al. Examining the role of patient experience surveys in measuring health care quality. *Medical care research and review : MCRR*. 2014; 71:522–54. [PubMed: 25027409]
54. Crofton C, Lubalin JS, Darby C. Consumer Assessment of Health Plans Study (CAHPS). Foreword. *Medical care*. 1999; 37:Ms1–9.

55. Fayers PM, Sprangers MA. Understanding self-rated health. *Lancet*. 2002; 359:187–8. [PubMed: 11812551]
56. Lee Y. The predictive value of self assessed general, physical, and mental health on functional decline and mortality in older adults. *Journal of epidemiology and community health*. 2000; 54:123–9. [PubMed: 10715745]
57. Miilunpalo S, Vuori I, Oja P, Pasanen M, Urponen H. Self-rated health status as a health measure: the predictive value of self-reported health status on the use of physician services and on mortality in the working-age population. *Journal of clinical epidemiology*. 1997; 50:517–28. [PubMed: 9180644]
58. Warren JL, Mariotto A, Melbert D, Schrag D, Doria-Rose P, Penson D, et al. Sensitivity of Medicare Claims to Identify Cancer Recurrence in Elderly Colorectal and Breast Cancer Patients. *Medical care*. 2016; 54:e47–54. [PubMed: 24374419]
59. Committee on Improving the Quality of Cancer Care IoM. Cancer Care in Older Adults. In: Levit, L.Balogh, E.Nass, S., Ganz, PA., editors. *Delivering High-Quality Cancer Care: Charting a New Course for a System in Crisis*. Washington, DC: National Academies Press (US); 2013.

(1a) Global Ratings

(1b) Composite Measures

Fig. 1.

Mean odds ratio and 95% confidence intervals of associations of very good/excellent general health status (GHS) with high scores for global ratings (2a) and composite measures (2b). Odds ratios determined by multivariable logistic regressions controlling for all variables listed in Table 1 as well as MHS, state of SEER registry, year of CAHPS survey, and presence of multiple cancers. OC=overall care, HP=health plan, PP=primary physician, SP=specialist physician (SP), Cust Serv=customer service, MD Comm=doctor communications, Get Rx=get needed prescription drugs

(2a) Global Ratings

(2b) Composite Measures

Fig. 2.

Mean odds ratio and 95% confidence intervals of associations of very good/excellent mental health status (MHS) with high scores for global ratings (3a) and composite measures (3b). Odds ratios determined by multivariable logistic regressions controlling for all variables listed in Table 1 as well as GHS, state of SEER registry, year of CAHPS survey, and presence of multiple cancers. OC=overall care, HP=health plan, PP=primary physician, SP=specialist physician (SP), Cust Serv=customer service, MD Comm=doctor communications, Get Rx=get needed prescription drugs

Table 1Characteristics of Study Population by Cancer Site^a

Percent of Population	Breast (n=8,261)	Colorectal (n=4,231)	Lung (n=1,096)	Prostate (n=5,867)	All (n=19,455)
Male	0.0%	46.8%	46.6%	100.0%	43.0%
Age: 66-74	49.2%	38.1%	48.8%	53.6%	48.1%
75-79	22.0%	23.9%	26.4%	25.1%	23.6%
80-84	16.7%	20.2%	16.3%	14.5%	16.8%
85+	12.1%	17.8%	8.5%	6.9%	11.6%
Education:<High School	16.2%	24.3%	23.2%	17.2%	18.7%
High School Grad	34.9%	33.0%	36.0%	25.2%	31.6%
Some College	27.1%	22.7%	24.3%	23.5%	24.9%
College Grad	21.8%	20.0%	16.5%	34.1%	24.8%
Race/ethnicity: Non-Hispanic White	78.5%	76.9%	81.8%	74.4%	77.1%
Non-Hispanic Black	5.8%	6.6%	5.5%	10.2%	7.3%
Non-Hispanic Asian	6.4%	7.2%	6.4%	5.1%	6.2%
Hispanic	6.0%	6.4%	3.7%	7.4%	6.4%
Other	3.2%	2.8%	2.6%	2.9%	3.0%
Fee-for-service Medicare	45.6%	46.8%	50.5%	46.5%	46.4%
AJCC Stage: 1	63.2%	38.0%	72.3%	4.0%	40.3%
2	32.7%	37.2%	7.4%	73.5%	44.5%
3	4.2%	24.9%	20.4%	22.5%	15.1%
Years Since Diagnosis: 0-2	10.6%	13.9%	23.4%	16.2%	13.7%
2-5	28.3%	31.1%	36.8%	41.0%	33.2%
5+	61.1%	55.1%	39.8%	42.8%	53.1%
Year of CAHPS completion: ^b 2000	1.4%	1.8%	1.9%	1.1%	1.4%
2001	5.2%	6.8%	5.3%	3.3%	5.0%
2002	5.5%	7.4%	5.7%	3.2%	5.2%
2003	7.3%	7.4%	5.9%	4.5%	6.4%
2004	7.4%	7.9%	8.1%	4.0%	6.5%

Percent of Population	Breast (n=8,261)	Colorectal (n=4,231)	Lung (n=1,096)	Prostate (n=5,867)	All (n=19,455)
2006	3.4%	3.2%	3.0%	2.2%	3.0%
2007	11.3%	11.1%	11.3%	10.1%	10.9%
2008	13.5%	13.1%	13.1%	14.6%	13.7%
2009	16.4%	14.7%	15.5%	18.9%	16.7%
2010	15.6%	14.6%	14.7%	19.1%	16.4%
2011	13.0%	11.9%	15.4%	19.0%	14.7%
GHS: Poor	4.24%	5.55%	10.31%	3.70%	4.70%
Fair	21.87%	23.97%	32.94%	18.24%	21.86%
Good	41.97%	39.99%	35.22%	38.47%	40.10%
Very Good/Excellent	31.92%	30.49%	21.53%	39.59%	33.34%
MHS: Poor	1.21%	1.68%	1.82%	1.40%	1.40%
Fair	7.46%	9.36%	10.58%	6.08%	7.63%
Good	26.44%	26.78%	29.11%	23.71%	25.84%
Very Good/Excellent	64.90%	62.18%	58.49%	68.81%	65.12%
Proportion with High Scores for each Measure					
Overall Care	66.1%	65.9%	60.8%	60.6%	64.1%
Health Plan	61.2%	62.8%	60.6%	58.4%	60.7%
Primary Physician	74.3%	71.7%	72.9%	71.8%	72.9%
Specialist Physician	74.3%	74.5%	73.9%	70.9%	73.3%
Customer Service	52.2%	53.9%	50.0%	50.4%	51.9%
Doctor Communications	56.7%	56.4%	55.7%	58.1%	57.0%
Get Care Quickly	60.3%	61.5%	61.3%	59.9%	60.5%
Get Needed Care	64.7%	68.5%	65.2%	65.4%	65.8%
Get Needed Prescription Drugs	77.3%	78.0%	77.0%	76.6%	77.2%

^a AJCC: American Joint Committee on Cancer; GHS: general health status; MHS: mental health status.

^b The Medicare CAHPS survey was not administered in 2005.

Table 2

Odds ratios (95% CI) from Multivariable Logistic Regression Analyses for Associations with High Global Ratings^a

	High Rating for Health Plan	High Rating Overall	High Rating for Primary Physician	High Rating for Specialist Physician
Breast Cancer Survivors				
N included in regression	7,896	7,011	6,883	5,071
Age (reference: 66–74)				
75–79	1.421 (1.26–1.603)	1.16 (1.015–1.325)	1.144 (0.991–1.321)	1.139 (0.964–1.346)
80–84	1.588 (1.386–1.819)	1.218 (1.049–1.414)	1.188 (1.013–1.394)	1.197 (0.986–1.453)
85-up	1.485 (1.273–1.733)	1.152 (0.973–1.364)	1.18 (0.982–1.418)	0.828 (0.665–1.029)
Education (reference: < High School Grad)				
High School Grad	0.515 (0.436–0.608)	0.742 (0.619–0.89)	0.622 (0.51–0.759)	0.603 (0.475–0.765)
Some College	0.808 (0.696–0.938)	0.997 (0.847–1.174)	0.763 (0.639–0.911)	0.869 (0.695–1.086)
College Grad	0.627 (0.536–0.734)	0.829 (0.698–0.984)	0.699 (0.58–0.844)	0.78 (0.62–0.982)
Race (reference: non-Hispanic White)				
Non-Hispanic Asian	0.9 (0.722–1.124)	0.491 (0.384–0.629)	0.781 (0.6–1.016)	0.595 (0.438–0.807)
Non-Hispanic Black	1.262 (1.022–1.559)	0.919 (0.733–1.152)	1.016 (0.792–1.303)	0.993 (0.726–1.359)
Hispanic	1.333 (1.072–1.657)	0.945 (0.746–1.197)	1.037 (0.803–1.34)	1.233 (0.892–1.705)
Other	1.089 (0.824–1.439)	0.858 (0.627–1.175)	0.93 (0.665–1.3)	0.981 (0.651–1.476)
Insurance (reference: Managed Care)				
Fee-For-Service	1.03 (0.932–1.137)	1.084 (0.972–1.208)	1.026 (0.91–1.155)	1.096 (0.955–1.258)
Colorectal Cancer Survivors				
N included in regression	4,060	3,450	3,432	2,352
Age (reference: 66–74)				
75–79	1.137 (0.959–1.348)	1.093 (0.901–1.326)	0.948 (0.777–1.157)	0.998 (0.78–1.277)
80–84	1.339 (1.113–1.61)	1.033 (0.841–1.27)	0.978 (0.789–1.212)	1.066 (0.815–1.393)
85-up	1.417 (1.164–1.725)	1.024 (0.823–1.274)	1.021 (0.814–1.281)	1.417 (1.044–1.923)
Education (reference: < High School Grad)				
High School Grad	0.621 (0.501–0.77)	0.845 (0.667–1.072)	0.783 (0.611–1.004)	0.699 (0.51–0.958)
Some College	0.927 (0.769–1.117)	1.156 (0.939–1.424)	1.041 (0.84–1.288)	0.82 (0.617–1.09)

	High Rating for Health Plan	High Rating Overall	High Rating for Primary Physician	High Rating for Specialist Physician
College Grad	0.751 (0.612–0.922)	1.212 (0.963–1.525)	1.016 (0.801–1.288)	0.975 (0.713–1.334)
Race (reference: non-Hispanic White)				
Non-Hispanic Asian	0.863 (0.632–1.178)	0.82 (0.58–1.161)	0.825 (0.579–1.173)	0.664 (0.427–1.034)
Non-Hispanic Black	1.197 (0.902–1.589)	0.864 (0.634–1.177)	1.64 (1.14–2.36)	0.857 (0.545–1.349)
Hispanic	1.742 (1.283–2.366)	1.495 (1.056–2.116)	1.358 (0.962–1.918)	1.134 (0.734–1.754)
Other	1.03 (0.678–1.564)	1.172 (0.745–1.844)	1.224 (0.746–2.008)	0.907 (0.491–1.674)
Insurance (reference: Managed Care)				
Fee-For-Service	1.077 (0.936–1.239)	1.112 (0.951–1.3)	1.036 (0.879–1.221)	1.123 (0.915–1.379)
Gender (reference: female)				
Male	0.825 (0.721–0.943)	0.864 (0.743–1.004)	0.823 (0.704–0.963)	0.987 (0.81–1.201)
Lung Cancer Survivors				
N included in regression	1,055	935	943	709
Age (reference: 66–74)				
75–79	1.371 (0.99–1.898)	1.483 (1.045–2.106)	1.685 (1.141–2.489)	0.98 (0.631–1.522)
80–84	1.13 (0.769–1.661)	1.155 (0.766–1.742)	1.733 (1.095–2.743)	0.691 (0.407–1.173)
85-up	0.956 (0.581–1.574)	1.242 (0.722–2.137)	1.751 (0.96–3.192)	0.713 (0.376–1.354)
Education (reference: < High School Grad)				
High School Grad	0.505 (0.323–0.79)	1.028 (0.634–1.665)	1.15 (0.684–1.931)	1.1 (0.601–2.015)
Some College	1.025 (0.709–1.482)	1.391 (0.935–2.07)	1.396 (0.915–2.128)	1.018 (0.603–1.718)
College Grad	0.724 (0.486–1.08)	1.004 (0.654–1.54)	1.187 (0.749–1.881)	0.742 (0.429–1.286)
Race (reference: non-Hispanic White)				
Non-Hispanic Asian	1.458 (0.765–2.779)	0.709 (0.364–1.382)	0.872 (0.436–1.743)	0.638 (0.289–1.405)
Non-Hispanic Black	0.97 (0.54–1.744)	1.377 (0.709–2.677)	1.513 (0.715–3.202)	1.873 (0.696–5.04)
Hispanic	0.822 (0.41–1.649)	0.617 (0.286–1.33)	1.536 (0.614–3.842)	0.85 (0.315–2.296)
Other	0.834 (0.358–1.939)	1.649 (0.582–4.675)	1.049 (0.371–2.962)	0.829 (0.244–2.812)
Insurance (reference: Managed Care)				
Fee-For-Service	1.331 (0.996–1.778)	1.242 (0.913–1.689)	1.175 (0.834–1.654)	1.102 (0.743–1.634)
Gender (reference: female)				
Male	0.722 (0.551–0.947)	0.778 (0.582–1.041)	0.656 (0.476–0.904)	0.798 (0.549–1.16)

	High Rating for Health Plan	High Rating Overall	High Rating for Primary Physician	High Rating for Specialist Physician
Prostate Cancer Survivors				
N included in regression	5,642	5,001	4,807	3,680
Age (reference: 66–74)				
75–79	1.269 (1.112–1.448)	1.054 (0.914–1.217)	1.059 (0.904–1.24)	0.964 (0.808–1.152)
80–84	1.627 (1.377–1.923)	1.061 (0.888–1.267)	0.974 (0.805–1.179)	0.959 (0.77–1.194)
85–up	1.453 (1.157–1.823)	1.003 (0.789–1.274)	1.09 (0.837–1.419)	0.945 (0.707–1.264)
Education (reference: < High School Grad)				
High School Grad	0.618 (0.516–0.74)	0.913 (0.752–1.11)	0.876 (0.709–1.082)	0.797 (0.618–1.027)
Some College	0.963 (0.805–1.151)	1.129 (0.929–1.372)	0.949 (0.77–1.17)	1.187 (0.909–1.548)
College Grad	0.794 (0.66–0.956)	1.013 (0.829–1.238)	0.895 (0.721–1.113)	0.795 (0.612–1.033)
Race (reference: non-Hispanic White)				
Non-Hispanic Asian	0.88 (0.661–1.172)	0.826 (0.606–1.126)	0.917 (0.658–1.277)	0.718 (0.497–1.038)
Non-Hispanic Black	1.314 (1.077–1.601)	0.769 (0.622–0.95)	0.913 (0.724–1.151)	0.934 (0.689–1.267)
Hispanic	1.103 (0.873–1.392)	0.938 (0.728–1.209)	0.903 (0.691–1.179)	1.194 (0.856–1.665)
Other	1.298 (0.918–1.837)	1.078 (0.746–1.556)	1.219 (0.802–1.852)	1.225 (0.762–1.968)
Insurance (reference: Managed Care)				
Fee-For-Service	1.008 (0.9–1.128)	1.032 (0.914–1.165)	0.972 (0.849–1.112)	1.062 (0.912–1.237)

^aIn addition to independent variables listed in this Table, regressions controlled for GHS, MHS, years from diagnosis to survey completion, AJCC stage at diagnosis, state of SEER registry, year of CAHPS survey, and presence of multiple cancers. Bold indicates significance at $p < .05$.

Table 3

Odds ratios (95% CI) from Multivariable Logistic Regression Analyses for Association with High Composite Scores^a

	High Scores for Customer Service	High Scores for Doctor Communications	High Scores for Getting Care Quickly	High Scores for Getting Needed Care	High Scores for Getting Needed Prescription Drugs
Breast Cancer Survivors					
N included in regression	2,410	6,692	6,703	6,783	7,452
Age (reference: 66–74)					
75–79	1.166 (0.944–1.439)	0.964 (0.848–1.097)	0.962 (0.846–1.094)	1.098 (0.961–1.254)	1.095 (0.948–1.264)
80–84	1.169 (0.922–1.482)	0.872 (0.757–1.005)	1.017 (0.88–1.176)	1.166 (1.003–1.356)	1.299 (1.101–1.533)
85-up	1.152 (0.852–1.557)	0.859 (0.729–1.012)	0.924 (0.783–1.09)	1.003 (0.842–1.195)	1.106 (0.923–1.325)
Education (reference: < High School Grad)					
High School Grad	0.725 (0.539–0.975)	0.645 (0.54–0.77)	0.768 (0.642–0.918)	0.664 (0.552–0.799)	0.857 (0.709–1.036)
Some College	1.042 (0.794–1.368)	0.827 (0.707–0.968)	1.001 (0.852–1.176)	1.05 (0.886–1.244)	1.275 (1.074–1.513)
College Grad	0.87 (0.656–1.155)	0.774 (0.655–0.914)	0.87 (0.735–1.031)	0.749 (0.628–0.893)	0.961 (0.803–1.149)
Race (reference: non-Hispanic White)					
Non-Hispanic Asian	0.884 (0.596–1.311)	0.725 (0.566–0.927)	0.518 (0.404–0.665)	0.77 (0.602–0.984)	0.594 (0.46–0.768)
Non-Hispanic Black	1.73 (1.191–2.512)	1.069 (0.858–1.332)	0.85 (0.68–1.063)	1.236 (0.971–1.574)	1.028 (0.81–1.304)
Hispanic	0.963 (0.681–1.364)	1.233 (0.978–1.555)	1.007 (0.795–1.275)	1.334 (1.039–1.711)	0.947 (0.742–1.208)
Other	1.509 (0.905–2.515)	0.874 (0.646–1.183)	0.633 (0.468–0.856)	0.888 (0.649–1.215)	0.758 (0.552–1.042)
Insurance (reference: Managed Care)					
Fee-For-Service	0.799 (0.655–0.974)	1.05 (0.943–1.168)	1.111 (1–1.235)	1.234 (1.106–1.378)	1.149 (1.022–1.293)
Colorectal Cancer Survivors					
N included in regression	1,131	3,333	3,266	3,373	3,643
Age (reference: 66–74)					
75–79	0.986 (0.709–1.37)	0.963 (0.8–1.159)	0.992 (0.82–1.199)	0.985 (0.808–1.202)	0.959 (0.775–1.186)
80–84	0.82 (0.581–1.159)	0.908 (0.743–1.109)	0.916 (0.749–1.12)	0.988 (0.799–1.223)	0.937 (0.748–1.173)
85-up	1.096 (0.746–1.61)	0.98 (0.792–1.212)	1.181 (0.947–1.472)	1.024 (0.815–1.287)	1.158 (0.907–1.479)
Education (reference: < High School Grad)					
High School Grad	0.721 (0.475–1.094)	0.778 (0.616–0.983)	0.797 (0.629–1.011)	0.667 (0.522–0.854)	0.819 (0.634–1.058)

	High Scores for Customer Service	High Scores for Doctor Communications	High Scores for Getting Care Quickly	High Scores for Getting Needed Care	High Scores for Getting Needed Prescription Drugs
Some College	1.389 (0.958–2.015)	1.058 (0.867–1.292)	0.945 (0.766–1.165)	0.994 (0.797–1.239)	1.389 (1.107–1.743)
College Grad	1.073 (0.721–1.596)	0.996 (0.799–1.243)	1.091 (0.865–1.375)	0.814 (0.641–1.034)	0.94 (0.736–1.202)
Race (reference: non-Hispanic White)					
Non-Hispanic Asian	0.534 (0.311–0.917)	0.57 (0.405–0.801)	0.504 (0.356–0.714)	0.372 (0.263–0.528)	0.364 (0.259–0.512)
Non-Hispanic Black	1.244 (0.735–2.107)	1.304 (0.958–1.775)	0.755 (0.549–1.037)	1.036 (0.74–1.45)	0.914 (0.655–1.277)
Hispanic	0.92 (0.553–1.531)	1.095 (0.798–1.501)	0.693 (0.505–0.951)	1.088 (0.774–1.529)	0.689 (0.493–0.961)
Other	0.482 (0.236–0.981)	1.11 (0.712–1.73)	0.721 (0.467–1.113)	0.754 (0.473–1.203)	0.615 (0.384–0.985)
Insurance (reference: Managed Care)					
Fee-For-Service	0.745 (0.552–1.006)	1.068 (0.917–1.244)	1.022 (0.876–1.192)	1.104 (0.938–1.299)	1.085 (0.913–1.289)
Gender (reference: female)					
Male	1.025 (0.793–1.326)	0.894 (0.773–1.035)	0.899 (0.775–1.043)	0.934 (0.8–1.092)	1.132 (0.958–1.337)
Lung Cancer Survivors					
N included in regression	312	910	907	918	987
Age (reference: 66–74)					
75–79	1.237 (0.66–2.319)	1.165 (0.824–1.648)	1.34 (0.951–1.888)	1.237 (0.867–1.764)	1.078 (0.726–1.6)
80–84	1.555 (0.672–3.597)	1.006 (0.668–1.514)	1.177 (0.782–1.773)	1.358 (0.868–2.124)	1.421 (0.872–2.317)
85-up	0.567 (0.211–1.523)	1.198 (0.712–2.017)	1.576 (0.908–2.737)	0.88 (0.516–1.499)	1.296 (0.702–2.392)
Education (reference: < High School Grad)					
High School Grad	0.456 (0.186–1.117)	0.975 (0.604–1.574)	0.738 (0.46–1.182)	0.788 (0.483–1.284)	0.944 (0.554–1.611)
Some College	0.567 (0.27–1.192)	1.016 (0.688–1.499)	1.074 (0.724–1.592)	0.945 (0.625–1.43)	1.473 (0.952–2.279)
College Grad	0.885 (0.399–1.96)	0.859 (0.563–1.311)	1.046 (0.685–1.598)	0.739 (0.475–1.149)	1.266 (0.786–2.04)
Race (reference: non-Hispanic White)					
Non-Hispanic Asian	0.453 (0.115–1.783)	0.563 (0.289–1.096)	0.511 (0.261–0.999)	0.599 (0.315–1.139)	0.31 (0.156–0.616)
Non-Hispanic Black	0.731 (0.268–1.995)	1.381 (0.701–2.719)	0.719 (0.371–1.393)	1.05 (0.505–2.183)	1.176 (0.563–2.458)
Hispanic	0.196 (0.05–0.758)	0.959 (0.452–2.033)	0.906 (0.424–1.934)	0.457 (0.213–0.982)	0.796 (0.336–1.883)
Other	4.154 (0.569–30.333)	0.928 (0.369–2.336)	1.668 (0.566–4.919)	0.621 (0.227–1.695)	1.064 (0.359–3.151)
Insurance (reference: Managed Care)					
Fee-For-Service	0.793 (0.423–1.487)	0.959 (0.705–1.303)	0.902 (0.662–1.228)	1.086 (0.791–1.49)	1.357 (0.953–1.934)

	High Scores for Customer Service	High Scores for Doctor Communications	High Scores for Getting Care Quickly	High Scores for Getting Needed Care	High Scores for Getting Needed Prescription Drugs
Gender (reference: female)					
Male	0.703 (0.414–1.193)	0.851 (0.637–1.137)	0.972 (0.727–1.301)	1.007 (0.747–1.358)	0.971 (0.696–1.353)
Prostate Cancer Survivors					
N included in regression	1,583	4,684	4,771	4,756	5,137
Age (reference: 66–74)					
75–79	0.679 (0.529–0.872)	0.844 (0.73–0.976)	0.954 (0.826–1.102)	0.941 (0.809–1.094)	1.035 (0.881–1.217)
80–84	0.847 (0.611–1.175)	0.777 (0.651–0.928)	0.865 (0.725–1.032)	1.016 (0.844–1.222)	1.025 (0.84–1.251)
85-up	0.599 (0.391–0.918)	0.73 (0.573–0.93)	0.797 (0.624–1.016)	0.888 (0.693–1.137)	1.079 (0.821–1.419)
Education (reference: < High School Grad)					
High School Grad	0.669 (0.48–0.933)	0.854 (0.702–1.04)	1.059 (0.867–1.293)	0.926 (0.749–1.145)	0.9 (0.726–1.115)
Some College	0.782 (0.557–1.099)	0.864 (0.71–1.052)	1.01 (0.827–1.234)	1.246 (1.003–1.547)	1.037 (0.837–1.285)
College Grad	0.785 (0.553–1.115)	0.827 (0.675–1.013)	0.988 (0.805–1.213)	0.97 (0.78–1.206)	1.011 (0.81–1.263)
Race (reference: non-Hispanic White)					
Non-Hispanic Asian	0.868 (0.503–1.497)	0.584 (0.424–0.803)	0.631 (0.457–0.872)	0.676 (0.487–0.939)	0.608 (0.435–0.848)
Non-Hispanic Black	1.584 (1.072–2.34)	1.066 (0.857–1.326)	0.983 (0.786–1.228)	1.283 (1.001–1.645)	0.998 (0.789–1.261)
Hispanic	1.174 (0.789–1.746)	0.775 (0.602–0.998)	0.72 (0.561–0.926)	0.937 (0.72–1.219)	0.741 (0.57–0.963)
Other	0.999 (0.513–1.943)	0.823 (0.57–1.188)	0.737 (0.513–1.058)	1.121 (0.763–1.648)	0.949 (0.638–1.412)
Insurance (reference: Managed Care)					
Fee-For-Service	0.665 (0.519–0.853)	1.02 (0.901–1.155)	1.066 (0.943–1.204)	1.125 (0.989–1.28)	1.17 (1.02–1.342)

^aIn addition to independent variables listed in this Table, regressions controlled for GHS, MHS, years from diagnosis to survey completion, AJCC stage at diagnosis, state of SEER registry, year of CAHPS survey, and presence of multiple cancers. Bold indicates significance at p<.05.