Published in final edited form as:

Tob Control. 2018 October ; 27(E2): e136–e142. doi:10.1136/tobaccocontrol-2017-053972.

Negative affect, message reactance, and perceived risk: How do pictorial cigarette pack warnings change quit intentions?

Marissa G. Hall^{1,2}, Paschal Sheeran^{3,1}, Seth M. Noar^{4,1}, Marcella H. Boynton^{2,1}, Kurt M. Ribisl^{2,1}, Humberto Parada Jr.⁵, Trent O. Johnson⁶, and Noel T. Brewer^{2,1}

¹Lineberger Comprehensive Cancer Center, University of North Carolina

²Department of Health Behavior, Gillings School of Global Public Health, University of North Carolina

³Department of Psychology & Neuroscience, University of North Carolina

⁴School of Media and Journalism, University of North Carolina

⁵Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina

⁶Stanford Prevention Research Center, Stanford University School of Medicine

Abstract

Objective—Pictorial warnings on cigarette packs increase motivation to quit smoking. We sought to examine the potential mediating role of negative affect, message reactance (i.e., an oppositional reaction to a message), and perceived risk in shaping quit intentions.

Methods—In 2014 and 2015, we randomly assigned 2,149 adult US smokers to receive either pictorial warnings or text-only warnings applied to their cigarette packs for four weeks. Analyses used structural equation modeling with bootstrapped standard errors to test our theorized mediational model.

Findings—Pictorial warnings increased negative affect, message reactance, and quit intentions (all $p < .001$), but not perceived risk (i.e., perceived likelihood and severity of harms of smoking). Negative affect mediated the impact of pictorial warnings on quit intentions (mediated effect = .16, $p < .001$). Message reactance weakened the impact of pictorial warnings on quit intentions,

Corresponding author: Marissa G. Hall, PhD, Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, 324 Rosenau Hall CB7440, Chapel Hill, NC 27599 USA, Phone: +1 336-972-3365, Fax: +1 919-966-2921, mghall@unc.edu.

Contributors

MGH analyzed the data and drafted the manuscript, with oversight and input from NTB. NTB, SMN, KMR, and MGH designed the study and developed the measures. NTB, SMN, KMR, MGH, and TOJ implemented and oversaw data collection. PS, MHB, HP, and NTB provided input on the statistical analyses. All authors critically revised the manuscript.

Competing Interest

Drs. Brewer and Ribisl have served as paid expert consultants in litigation against tobacco companies. The other authors declare no conflicts of interest.

Patient Consent

Obtained.

Ethics Approval

The University of North Carolina Institutional Review Board approved the procedures for this trial.

although the effect was small (mediated effect=-.04, $p<.001$). Although pictorial warnings did not directly influence perceived risk, the model showed additional small mediation effects on quit intentions through negative affect and its *positive* association with perceived risk (mediated effect=.02, $p<.001$), as well as reactance and its *negative* association with perceived risk (mediated effect=-.01, $p<.001$).

Conclusions—Pictorial cigarette pack warnings increased quit intentions by increasing negative affect. Message reactance partially attenuated this increase in intentions. The opposing associations of negative affect and reactance on perceived risk may explain why pictorial warnings did not lead to observable changes in perceived risk.

Keywords

Reactance; risk appraisals; risk perceptions; perceived risk; emotion; negative affect; tobacco control; pictorial warnings; graphic warnings

Introduction

Tobacco use is the leading cause of preventable morbidity and mortality worldwide, causing nearly six million deaths each year.[1] The World Health Organization Framework Convention on Tobacco Control calls for its signatory countries to implement large pictorial images on cigarette packs based on evidence that pictorial warnings are more effective than text-only warnings.[2] Indeed, systematic reviews and a recent randomized controlled trial demonstrate that pictorial warnings elicit stronger quit intentions[3, 4] and increase subsequent cessation behavior, compared to text-only warnings.[3, 5] Characterizing how pictorial warnings strengthen quit intentions can help policymakers design more effective warnings.

The 2009 Family Smoking Prevention and Tobacco Control Act requires pictorial warnings on cigarette packs in the US.[6] However, a 2012 lawsuit brought forth by the tobacco industry stalled implementation of this facet of the law.[7] In the court case, the warnings were criticized for being “unabashed attempts to evoke emotion.”[7] Indeed, research has shown that pictorial warnings elicit fear and other negative emotions.[4, 8–12] These negative emotions may serve an important information processing role by making the warning content accessible and salient, thereby more effectively influencing outcomes such as risk perceptions and quit intentions.[8, 9] Emotional responses are typically formed with greater confidence and more quickly than cognitive appraisals, and are therefore likely to precede cognitive responses such as perceived risk.[13] Affect may also help people process risk information.[14–18] Understanding the role of negative affective responses to pictorial warnings may be relevant to future legal battles in the US and may also inform the creation of strong warnings in countries that are implementing new pictorial warning policies or changing their existing pictorial warnings.

In addition to negative affect, pictorial warnings may elicit *message reactance*, defined as cognitive and emotional resistance to a health message in response to a perceived threat to one’s freedom.[19–21] Several studies have found that pictorial warnings elicit greater reactance than text-only warnings.[22–25] However, the extent to which reactance weakens

the impact of pictorial warnings on outcomes such as quit intentions remains an important research gap. Moreover, research is needed to understand whether negative emotions and message reactance are associated with changes in perceived risk.

In the current study, we sought to determine the psychological mechanisms by which pictorial cigarette pack warnings elicit stronger quit intentions using experimental data. Specifically, we aimed to understand whether pictorial warnings heightened negative affect and reactance, and whether, in turn, negative affect and reactance were associated with quit intentions via perceived risk (i.e., perceived likelihood of harm from smoking and perceived severity of harm from smoking).

Methods

Participants

From September 2014 to August 2015, we recruited[26] a convenience sample of adult smokers in North Carolina and California, US to participate in a trial comparing the impact of pictorial and text versus text-only warnings.[3] Participants were age 18 or older, proficient in English, and current smokers, defined as having smoked at least 100 cigarettes during their lifetime and now smoking every day or some days. Exclusion criteria included pregnancy, current enrollment in a smoking cessation trial, smoking only roll-your-own cigarettes, smoking fewer than seven cigarettes per week, and living in the same household as another trial participant. Details regarding recruitment, design, and methods have previously been reported.[3]

Procedures

In our trial, smokers received warnings on their own cigarette packs for four weeks using a protocol developed by our team.[3, 27] Participants brought in an eight-day supply of cigarettes weekly. At baseline, we randomly assigned participants to receive one of eight warnings: four pictorial warnings with text, and four text-only controls in a two-arm trial testing pictorial vs. text-only warnings. The four pictorial warnings contained text required by the Tobacco Control Act and a picture to illustrate a health harm of smoking selected from the FDA's originally-proposed set of images (Figure 1).[28] We chose these four warning images because they performed well in a previous internet study and avoided many of the criticisms in the lawsuits (e.g., using a cartoon or a rare health harm of smoking).[29, 30] We removed the FDA quitline number from the images, which was a source of contention in litigation against the warnings.[7, 31] Staff applied the pictorial warning labels to the top half of the front and back panels of participants' cigarette packs, in accordance with the FDA requirements.[6] The four text-only control warnings used the US Surgeon General's warning statements that have been required on the side of cigarette packs since 1985. Staff applied the text-only warning labels on the side of the packs covering the existing US Surgeon General's warnings. We applied the new warning labels on top of the existing warnings to control for the effect of putting a label on smokers' packs. Randomization created groups that did not differ on demographics assessed (all $p > .05$).[3] For this reason, we did not adjust analyses for demographics. Trial participants were diverse,

including a substantial number of sexual minority, African American, low-education, and low-income smokers (Table 1).

Participants completed two computer surveys at the first visit (i.e., baseline and immediately after seeing their assigned warning, which was immediate post-test), and one survey at each visit thereafter for four weeks. Participants received a cash incentive at the end of each visit, totaling up to \$185 in North Carolina and \$15 higher in California due to the higher cost of living. At the end of the final follow-up appointment, participants received information about local smoking cessation programs. The University of North Carolina Institutional Review Board approved the procedures for this trial.

Measures

The baseline survey and the week 2 follow-up survey assessed perceived likelihood of harm from smoking and perceived severity of harm from smoking. We measured negative affect elicited by the warning at immediate post-test.[32] We originally planned to examine fear alone as a mediator given its importance in the Extended Parallel Process Model,[33] but sensitivity analyses revealed that fear and the other negative affect items exhibited a nearly identical pattern of associations, and confirmatory factor analysis supported treating negative affect as a single latent factor. The immediate post-test survey also assessed message reactance using the Brief Reactance to Health Warnings Scale.[25] This measure captures the three main components of reactance: anger toward the warning, perceived threat to freedom, and counterarguing against the warning.[19, 20, 24] Finally, we measured quit intentions at baseline, immediate post-test, and all follow-up surveys.[34] Item wording, response scales, and factor loadings for all measures appear in Table 2.

Data Analysis

Analyses used Stata/SE version 14.1 and *Mplus* version 7.4 with two-tailed tests and a critical alpha of .05. Analyses used measures of negative affect and message reactance taken at immediate post-test, and measures of perceived likelihood and perceived severity taken at week 2 follow-up (Table 2). The outcome was quit intentions at week 4 follow-up.

First, we examined associations among the constructs using bivariate correlations; this analysis used mean scores for each construct. Next, we examined the association between the individual items and their latent constructs using a measurement model that treated the mediators and outcome as latent variables. The model specification process pointed toward modeling perceived risk as a second-order latent variable that was a function of perceived likelihood and perceived severity. Then, we examined the impact of pictorial warnings on quit intentions in an unadjusted structural model. Finally, we examined the simultaneous influence of the hypothesized mediator and suppressor variables using a final, adjusted structural model. Suppression occurs when the direct and mediated effects have opposite signs, in this case demonstrating that the mediator detracts from the effectiveness of pictorial warnings.[35] In contrast, a direct and mediated effect with the same sign signals mediation, indicating that the mediator contributes to the effectiveness of pictorial warnings.[35] Candidate mediator/suppressor variables were negative affect, message reactance, and perceived risk. The model tested theoretically-driven predictions[4, 15, 33, 36] about the

indirect effects of pictorial warnings on quit intentions first via negative affect and message reactance, and then through perceived risk.

Intent-to-treat analyses included all participants randomized, using the last observation available.[37] Participants who had quit smoking at week 4 did not answer the quit intentions items, so we used the last observation available for them as well. When data remained missing, the models employed full information maximum likelihood estimation, an approach commonly recommended for structural equation models that makes use of all available observations.[38–40] We report results as standardized path coefficients (β s). Mediation analyses used bootstrapped 95% confidence intervals (CI) with 1,000 repetitions, as this approach does not assume that indirect effects are normally distributed.[41] We evaluated several indicators of acceptable model fit, including the root mean square error of approximation (RMSEA<.08),[42] the Tucker-Lewis Index (TLI>.95),[43] and the Bentler Comparative Fit Index (CFI>.95).[44]

Results

The measurement model fit the data well (RMSEA=.040 [90% CI=.036, .043], CFI=.997, TLI=.997; Table 2). Factor loadings for the indicator variables were all statistically significant and ranged from .75 to .98. Bivariate correlations between the model constructs ranged from -.24 to .38 (Table 3). Pictorial warnings increased negative affect, message reactance, and quit intentions (all $p<.001$, Table 3). Pictorial warnings did not increase perceived risk ($p=.65$, Table 3).

The final structural model exhibited excellent fit (RMSEA=.038 [90% CI=.034, .041], CFI=.997, TLI=.997; Figure 2; Supplementary Table 1). Pictorial warnings generated higher quit intentions than text-only warnings ($\beta=.08$, $p<.001$). As expected, negative affect mediated the impact of pictorial warnings on quit intentions, such that pictorial warning exposure increased negative affect ($\beta=.40$, $p<.001$; Figure 2) which, in turn, was associated with greater quit intentions ($\beta=.41$, $p<.001$; mediated effect=.16, $p<.001$; Table 4). Negative affect was also associated with greater perceived risk ($\beta=.30$, $p<.001$), and perceived risk was associated with greater quit intentions (mediated effect through negative affect and perceived risk=.02, $p<.001$).

Pictorial warnings also increased message reactance ($\beta=.25$, $p<.001$), and message reactance was associated with lower quit intentions ($\beta=-.17$, $p<.001$; mediated effect=-.04, $p<.001$, Table 4). Message reactance was associated with lower perceived risk ($\beta=-.36$, $p<.001$), which was in turn associated with greater quit intentions (mediated effect through reactance and perceived risk=-.01, $p<.001$).

Discussion

Pictorial cigarette pack warnings elicited stronger quit intentions than text-only warnings. Negative affect (which included fear, guilt, disgust, anxiety, and sadness) was a key driver of the effect of pictorial warnings on quit intentions. Compared to text-only warnings, pictorial warnings elicited more negative affect, which was associated with greater quit intentions. We also found in adjusted analyses that negative affect was associated with greater perceived

risk of harm from smoking. Reactance, however, was associated with lower perceived risk. Perceived risk was associated with stronger quit intentions, building on prior work showing associations between risk perceptions and intentions and behavior.[45–48] The opposing associations of negative affect and reactance with perceived risk may offer insight into the well-established finding that pictorial cigarette pack warnings do not cause observable changes in risk perceptions.[4, 8, 9, 49]

Our study suggests that the emotion evoked by the warnings may be a precursor to beneficial changes in quit intentions, building on prior research about emotional reactions to pictorial warnings.[4, 8–12] This is an important point that the US courts did not acknowledge when criticizing pictorial warnings simply for evoking emotion.[7] A recent legal analysis of the pictorial warning lawsuits in the US concluded that the “warnings do not bypass reason simply by reaching for the heart,” and that the emotions evoked by the warnings should not deem them unconstitutional.[50] Indeed, we found pictorial warnings were effective *because* of the emotions they elicited.

Some have argued that discrete negative emotions fall into distinct categories and therefore play unique roles in shaping intentions and behaviors.[51–53] The Extended Parallel Process Model argues that fear is the most important affective motivator of behavior change.[33] However, we found support for a constructionist view of emotion[52] because – in the context of pictorial cigarette pack warnings – fear, guilt, disgust, anxiety, and sadness were highly correlated and functioned quite similarly. However, in contrast with negative affect, anger (a facet of reactance) was associated with *lower* quit intentions. This finding builds on previous research indicating that anger and fear produce opposite effects on intentions and behavior.[54–56] In the context of pictorial warnings, the distinctions between fear, guilt, disgust, anxiety, and sadness may not be meaningful, but it is crucial to distinguish between anger and other types of negative affect. Message reactance, which includes anger, works against the warnings’ purpose to motivate quitting, while increasing other types of negative affect advances this purpose.

As predicted, message reactance weakened the effect of pictorial warnings on quit intentions, although the magnitude of the mediated effect was small. Message reactance was also associated with lower perceived risk. Previous studies have found that pictorial warnings cause greater reactance than text-only warnings,[4, 22–24] but few studies have examined whether message reactance or other forms of defensive processing are associated with deleterious consequences, such as lower quit intentions.[24, 57, 58] Our experimental test of the suppression effects of message reactance following repeated exposure to pictorial warnings adds to this body of research. However, given research demonstrating the effectiveness of pictorial warnings, it would be unwise to conclude that pictorial warnings are counterproductive simply because they produce reactance, as others have argued.[23] In fact, the best warnings may be those that elicit strong reactions of all kinds, including reactance. Reactance had a small effect in the model and its influence did not undo the positive effects of warnings. Measuring message reactance can help to better identify the types of individuals who are resistant to the warnings and therefore may benefit from alternative interventions. Moreover, message reactance could be a particularly useful as a way of vetting candidate warnings in the early stages of message development and testing.

Given a choice between two warnings that elicit similar levels of negative affect, message designers might choose the warning that elicits less reactance.

Study strengths include the use of an experimental design, a large and diverse sample of smokers who received the warnings on the cigarette packs they used every day, and the 4 week data collection period that allowed us to establish the temporality of mediated effects. However, the trial examined the potential effect of adding pictorial warnings to cigarette packs as well as implementing other label formatting changes required by the 2009 Tobacco Control Act compared with the present text-only warnings in the United States. While the trial aimed to compare the current warning policy to the new one in the Act, the use of this research design leaves open the possibility that the observed effects on mediators and intentions may be due to the combination of adding pictures and other changes (e.g., location, size, and content). The generalizability of these findings to different contexts (e.g., outside the US) or over a longer period of time has yet to be established. Finally, some of the mediated pathways were based on observational data, limiting our ability to assume causal associations between the model factors.

Conclusions

Understanding how pictorial warnings exert their influence can help researchers and policymakers design more effective warnings. Pictorial warnings elicited stronger quit intentions in the present study. However, message reactance partially suppressed pictorial warnings' effect on quit intentions. We found that increased negative affect was a key mechanism by which pictorial warnings changed quit intentions. Negative affect was also associated with stronger perceived risk, which was in turn, associated with stronger quit intentions. Thus, negative affect is a vital and productive result of pictorial warning exposure.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the research participants for taking part in our trial. We also thank Laura Bach, staff at the Pacific Institute for Research and Evaluation, and staff at Ewald and Wasserman Research Consultants LLC for helping to carry out the trial. Finally, we thank Cathy Zimmer for statistical consulting.

Funding

Research reported in this publication was supported by P30CA016086-38S2 from the National Cancer Institute and the FDA Center for Tobacco Products (CTP). F31CA196037 and T32-CA057726 from the National Cancer Institute of the National Institutes of Health and P50CA180907 from the National Cancer Institute and the FDA Center for Tobacco Products supported MGH's time writing the paper. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or the Food and Drug Administration.

References

1. World Health Organization. WHO report on the global tobacco epidemic, 2013: Enforcing bans on tobacco advertising, promotion and sponsorship. Geneva, Switzerland: World Health Organization; 2013.

2. World Health Organization. Framework Convention on Tobacco Control. 2013
3. Brewer NT, Hall MG, Noar SM, et al. Effect of pictorial cigarette pack warnings on changes in smoking behavior: A randomized clinical trial. *JAMA Intern Med.* 2016; 176(7):905–912. [PubMed: 27273839]
4. Noar SM, Hall MG, Francis DB, et al. Pictorial cigarette pack warnings: A meta-analysis of experimental studies. *Tob Control.* 2016; 25(3):341–354. [PubMed: 25948713]
5. Noar SM, Francis DB, Bridges C, et al. The impact of strengthening cigarette pack warnings: Systematic review of longitudinal observational studies. *Soc Sci Med.* 2016; 164:118–129. [PubMed: 27423739]
6. United States Public Laws. Family smoking prevention and tobacco control act of 2009. 111th Congress, 1st Session Public Law 111-31[H.R. 1256]. 2009
7. R.J. Reynolds Tobacco Company vs United States Food and Drug Administration. Civil Case No. 11-1482(RJL). United States District Court for the District of Columbia; 2011.
8. Emery LF, Romer D, Sheerin KM, et al. Affective and cognitive mediators of the impact of cigarette warning labels. *Nicotine Tob Res.* 2014; 16(3):263–269. [PubMed: 23946325]
9. Evans AT, Peters E, Strasser AA, et al. Graphic warning labels elicit affective and thoughtful responses from smokers: Results of a randomized clinical trial. *PLoS One.* 2015; 10(12):e0142879. [PubMed: 26672982]
10. Romer D, Ferguson SG, Strasser AA, et al. Effects of pictorial warning labels for cigarettes and quit-efficacy on emotional responses, smoking satisfaction, and cigarette consumption. *Ann Behav Med.* 2017
11. Netemeyer RG, Burton S, Andrews JC, et al. Graphic health warnings on cigarette packages: the role of emotions in affecting adolescent smoking consideration and secondhand smoke beliefs. *Journal of Public Policy & Marketing.* 2016; 35(1):124–143.
12. Wang AL, Lowen SB, Romer D, et al. Emotional reaction facilitates the brain and behavioural impact of graphic cigarette warning labels in smokers. *Tob Control.* 2015; 24(3):225–232. [PubMed: 25564288]
13. Zajonc RB. Feeling and thinking: Preferences need no inferences. *Am Psychol.* 1980; 35(2):151–175.
14. Schwarz N, Clore GL. Mood, misattribution, and judgments of well-being: Informative and directive functions of affective states. *J Pers Soc Psychol.* 1983; 45(3):513–523.
15. Slovic, Finucane ML, Peters E, et al. The affect heuristic. *Eur J Oper Res.* 2007; 177(3):1333–1352.
16. Slovic, Finucane ML, Peters E, et al. Risk as analysis and risk as feelings: Some thoughts about affect, reason, risk, and rationality. *Risk Anal.* 2004; 24(2):311–322. [PubMed: 15078302]
17. Evans AT, Peters E, Shoben AB, et al. Cigarette graphic warning labels increase both risk perceptions and smoking myth endorsement. *Psychol Health.* 2017:1–22. [PubMed: 27616462]
18. Kiviniemi MT, Ellis EM, Hall MG, et al. Mediation, moderation, and context: Understanding complex relations among cognition, affect, and health behaviour. *Psychol Health.* 2017:1–19. [PubMed: 27616462]
19. Brehm JW. A theory of psychological reactance. New York, NY: Academic Press; 1966.
20. Brehm JW, Brehm SS. Psychological reactance: A theory of freedom and control. New York, NY: Academic Press; 1981.
21. Fitzsimons GJ, Lehmann DR. Reactance to recommendations: When unsolicited advice yields contrary responses. *Marketing Science.* 2004; 23(1):82–94.
22. Erceg-Hurn DM, Steed LG. Does exposure to cigarette health warnings elicit psychological reactance in smokers? *J Appl Soc Psychol.* 2011; 41(1):219–237.
23. LaVoie NR, Quick BL, Riles JM, et al. Are graphic cigarette warning labels an effective message strategy? A test of psychological reactance theory and source appraisal. *Communic Res.* 2015 Advance online publication.
24. Hall MG, Sheeran P, Noar SM, et al. Reactance to health warnings scale: Development and validation. *Ann Behav Med.* 2016; 50(5):736–750. [PubMed: 27333895]

25. Hall MG, Sheeran P, Noar SM, et al. A brief measure of reactance to health warnings. *J Behav Med.* 2017; 40(3):520–529. [PubMed: 28120228]
26. Brodar KE, Hall MG, Butler EN, et al. Recruiting diverse smokers: Enrollment yields and cost. *International journal of environmental research and public health.* 2016; 13(12)
27. Brewer NT, Hall MG, Lee JG, et al. Testing warning messages on smokers' cigarette packages: a standardised protocol. *Tob Control.* 2015; 25(2):153–159. [PubMed: 25564282]
28. Nonnemaker J, Farrelly M, Kamyab K. , et al. Experimental study of graphic cigarette warning labels: Final results report. Research Triangle Park NC: RTI International; 2010.
29. Cameron LD, Pepper JK, Brewer NT. Responses of young adults to graphic warning labels for cigarette packages. *Tob Control.* 2015; 24(e1):e14–22. [PubMed: 23624558]
30. Berman ML, Byron JB, Hemmerich N, et al. Communicating tobacco product information to the public. *Food Drug Law Journal.* In Press.
31. Kraemer JD, Baig SA. Analysis of legal and scientific issues in court challenges to graphic tobacco warnings. *Am J Prev Med.* 2013; 45(3):334–342. [PubMed: 23953361]
32. Nonnemaker J, Choiniere CJ, Farrelly MC, et al. Reactions to graphic health warnings in the United States. *Health Educ Res.* 2015; 30(1):46–56. [PubMed: 24974219]
33. Witte K. Putting the fear back into fear appeals: The extended parallel process model. *Commun Monogr.* 1992; 59(4):329–349.
34. Klein WM, Zajac LE, Monin MM. Worry as a moderator of the association between risk perceptions and quitting intentions in young adult and adult smokers. *Ann Behav Med.* 2009; 38(3):256–261. [PubMed: 20049660]
35. MacKinnon DP, Krull JL, Lockwood CM. Equivalence of the mediation, confounding and suppression effect. *Prev Sci.* 2000; 1(4):173–181. [PubMed: 11523746]
36. Rogers RW. A protection motivation theory of fear appeals and attitude change. *J Psychol.* 1975; 91(1):93–114. [PubMed: 28136248]
37. Brody T. Clinical trials: study design, endpoints and biomarkers, drug safety, and FDA and ICH guidelines. London, UK: Academic Press; 2011.
38. Peters CL, Enders C. A primer for the estimation of structural equation models in the presence of missing data: Maximum likelihood algorithms. *Journal of Targeting, Measurement and Analysis for Marketing.* 2002; 11(1):81–95.
39. Bollen KA. Structural equations with latent variables. New York, NY: Wiley; 1989.
40. Kline RB. Principles and practice of structural equation modeling. New York, NY: Guilford Press; 2011.
41. Hayes AF. Beyond Baron and Kenny: Statistical mediation analysis in the new millennium. *Commun Monogr.* 2009; 76(4):408–420.
42. Steiger JH. Structural model evaluation and modification: An interval estimation approach. *Multivariate Behav Res.* 1990; 25(2):173–180. [PubMed: 26794479]
43. Tucker LR, Lewis C. A reliability coefficient for maximum likelihood factor analysis. *Psychometrika.* 1973; 38(1):1–10.
44. Bentler PM. Comparative fit indexes in structural models. *Psychol Bull.* 1990; 107(2):238–246. [PubMed: 2320703]
45. Sheeran P, Harris PR, Epton T. Does heightening risk appraisals change people's intentions and behavior? A meta-analysis of experimental studies. *Psychol Bull.* 2014; 140(2):511–543. [PubMed: 23731175]
46. Costello MJ, Logel C, Fong GT, et al. Perceived risk and quitting behaviors: results from the ITC 4-country survey. *Am J Health Behav.* 2012; 36(5):681–692. [PubMed: 22584095]
47. Brewer NT, Chapman GB, Gibbons FX, et al. Meta-analysis of the relationship between risk perception and health behavior: The example of vaccination. *Health Psychol.* 2007; 26(2):136–145. [PubMed: 17385964]
48. Floyd DL, Prentice-Dunn S, Rogers RW. A meta-analysis of research on protection motivation theory. *J Appl Soc Psychol.* 2000; 30(2):407–429.
49. Peebles K, Hall MG, Pepper JK, et al. Adolescents' responses to pictorial warnings on their parents' cigarette packs. *J Adolesc Health.* 2016; 59(6):635–641. [PubMed: 27646499]

50. Goodman EP. Visual gut punch: persuasion, emotion, and the constitutional meaning of graphic disclosure. *Cornell Law Rev.* 2013; 99(3):513–570.
51. Barrett LF. Are Emotions Natural Kinds? *Perspect Psychol Sci.* 2006; 1(1):28–58. [PubMed: 26151184]
52. Lindquist KA, Siegel EH, Quigley KS, et al. The hundred-year emotion war: are emotions natural kinds or psychological constructions? Comment on Lench, Flores, and Bench (2011). *Psychol Bull.* 2013; 139(1):255–263. [PubMed: 23294094]
53. Keltner D, Ellsworth PC, Edwards K. Beyond simple pessimism: effects of sadness and anger on social perception. *J Pers Soc Psychol.* 1993; 64(5):740–752. [PubMed: 8505705]
54. Lerner, Gonzalez, Small, et al. Effects of fear and anger on perceived risks of terrorism a national field experiment. *Psychol Sci.* 2003; 14(2):144–150. [PubMed: 12661676]
55. Lerner, Keltner. Fear, anger, and risk. *J Pers Soc Psychol.* 2001; 81(1):146–159. [PubMed: 11474720]
56. Dabbs JM Jr, Leventhal H. Effects of varying the recommendations in a fear-arousing communication. *J Pers Soc Psychol.* 1966; 4(5):525–531. [PubMed: 5972084]
57. Cho YJ, Thrasher JF, Swayampakala K, et al. Does reactance against cigarette warning labels matter? Warning label responses and downstream smoking cessation amongst adult smokers in Australia, Canada, Mexico and the United States. *PLoS One.* 2016; 11(7):e0159245. [PubMed: 27411100]
58. Yong HH, Borland R, Thrasher JF, et al. Mediatonal pathways of the impact of cigarette warning labels on quit attempts. *Health Psychol.* 2014; 33(11):1410–1420. [PubMed: 24977309]

What This Paper Adds

- In a four-week trial with smokers, we found that pictorial warnings increased quit intentions.
- Negative affect was a key driver of the association between pictorial warnings and quit intentions. Message reactance weakened this association, although pictorial warnings were more effective than text-only warnings on the whole.
- Negative affect and reactance had opposite and offsetting associations with perceived risk, which may explain why pictorial cigarette pack warnings do not change perceived risk.

Figure 1.
Intervention warnings (left) and control warnings (right)

Figure 2.

Structural equation model assessing the impact of pictorial warnings on quit intentions ($n=2,149$)

Note. To simplify presentation, the figure omits factor loadings, non-significant pathways, and residuals. RMSEA=.038 [90% CI=.034, .041], CFI=.997, TLI=.997.

* $p<.05$

Table 1Participant characteristics (*n*=2,149)

	<u>Text-only warnings (<i>n</i>=1,078)</u>		<u>Pictorial warnings (<i>n</i>=1,071)</u>	
	<i>n</i>	(%)	<i>n</i>	(%)
Demographics				
Age				
18–24 years	171	(16.1)	152	(14.5)
25–39 years	377	(35.5)	398	(37.9)
40–54 years	338	(31.8)	304	(29.0)
55+ years	176	(16.6)	195	(18.6)
Mean (SD) years	39.7	(13.4)	39.8	(13.7)
Gender				
Female	548	(51.2)	512	(48.2)
Male	507	(47.4)	532	(50.0)
Transgender	15	(1.4)	19	(1.8)
Gay, lesbian, or bisexual	173	(16.3)	195	(18.8)
Hispanic	92	(8.6)	89	(8.5)
Race				
Black or African American	484	(45.8)	510	(48.9)
White	393	(37.2)	358	(34.3)
Other/multiracial	134	(12.7)	117	(11.2)
Asian	28	(2.7)	42	(4.0)
American Indian or Alaska Native	7	(0.6)	11	(1.0)
Native Hawaiian or other Pacific Islander	11	(1.0)	6	(0.6)
Education				
High school graduate or less	333	(31.1)	344	(32.5)
Some college	519	(48.5)	502	(47.4)
College graduate	156	(14.6)	156	(14.7)
Graduate degree	63	(5.9)	58	(5.5)
Household income, annual				
\$0–\$24,999	566	(53.3)	589	(55.8)
\$25,000–\$49,999	272	(25.6)	266	(25.2)
\$50,000–\$74,999	110	(10.3)	92	(8.7)
\$75,000+	115	(10.8)	109	(10.3)
Low income (< 150% of Federal Poverty Level)				
No	506	(47.0)	477	(44.8)
Yes	570	(53.0)	589	(55.2)
Trial site				
California	594	(55.1)	592	(55.3)
North Carolina	484	(44.9)	479	(44.7)
Cigarettes smoked per day, mean (SD)	8.8	(6.6)	8.7	(7.3)
Smoking frequency				

	Text-only warnings (n=1,078)		Pictorial warnings (n=1,071)	
	n	(%)	n	(%)
Non-daily	211	(19.6)	207	(19.3)
Daily	866	(80.4)	864	(80.7)
Mediators and outcome at baseline, mean (SD)				
Perceived likelihood of harm from smoking	3.3	(0.9)	3.3	(0.9)
Perceived severity of harm from smoking	3.7	(0.6)	3.7	(0.6)
Quit intentions	2.2	(0.9)	2.3	(0.9)

Note. Study characteristics, mediators and outcomes at baseline did not differ by trial arm.[3] Missing demographic data ranged from 0.7% to 2.2%. The baseline surveys could not assess negative affect or message reactance as participants had not yet seen the warnings.

Table 2Latent variables used in the measurement and structural equation models ($n=2,149$)

Latent variable (Timepoint used in analysis)	Indicator item wording [response scale]	Factor loading
Negative affect (Immediate post-test)	How much did the warning on your cigarette packs make you feel... Scared? [not at all (coded as 1), a little (2), somewhat (3), very (4), extremely (5)]	.93
	Regretful? [not at all (1), a little (2), somewhat (3), very (4), extremely (5)]	.91
	On edge? [not at all (1), a little (2), somewhat (3), very (4), extremely (5)]	.88
	Disgusted? [not at all (1), a little (2), somewhat (3), very (4), extremely (5)]	.88
	Sad? [not at all (1), a little (2), somewhat (3), very (4), extremely (5)]	.86
Message reactance (Immediate post-test)	Please say how much you agree or disagree with each statement below about the warning we put on your packs. The health effect on this warning is overblown. [strongly disagree (1), somewhat disagree (2), neither agree nor disagree (3), somewhat agree (4), strongly agree (5)]	.82
	This warning is trying to manipulate me. [strongly disagree (1), somewhat disagree (2), neither agree nor disagree (3), somewhat agree (4), strongly agree (5)]	.77
	This warning annoys me. [strongly disagree (1), somewhat disagree (2), neither agree nor disagree (3), somewhat agree (4), strongly agree (5)]	.75
Perceived risk ¹ (Week 2)	Perceived likelihood of harm from smoking ²	1.00
	What is the chance that you will one day get <u>heart disease</u> if you continue to smoke cigarettes? [no chance (1), low chance (2), moderate chance (3), high chance (4), certain (5)]	.94
	What is the chance that you will one day get <u>cancer</u> if you continue to smoke cigarettes? [no chance (1), low chance (2), moderate chance (3), high chance (4), certain (5)]	.93
	What is the chance that you will one day get a <u>permanent breathing problem</u> if you continue to smoke cigarettes? [no chance (1), low chance (2), moderate chance (3), high chance (4), certain (5)]	.87
	Perceived severity of harm from smoking	.56
	How much would getting a <u>heart disease</u> because of smoking affect your life? [not at all (1), a little (2), a moderate amount (3), a lot (4)]	.97
	How much would getting a <u>cancer</u> because of smoking affect your life? [not at all (1), a little (2), a moderate amount (3), a lot (4)]	.97
	How much would getting a <u>permanent breathing problem</u> because of smoking affect your life? [not at all (1), a little (2), a moderate amount (3), a lot (4)]	.97
Quit intentions (Week 4)	How much do you plan to quit smoking in the next month? [not at all (1), a little (2), somewhat (3), very much (4)]	.98
	How interested are you in quitting smoking in the next month? [not at all interested (1), a little interested (2), somewhat interested (3), very interested (4)]	.95
	How likely are you to quit smoking in the next month? [not at all likely (1), a little likely (2), somewhat likely (3), very likely (4)]	.94

Note. Table reports standardized factor loadings.

¹ Modeled as a second-order latent factor

² For purposes of estimation, we fixed the value of the loading for perceived likelihood at 1

Table 3
Means and bivariate correlations among variables in the multiple mediation model ($n=2,093$)

	Mean (SD)	1	2	3	4	5
Pictorial warnings (1)	—	—	—	—	—	—
Negative affect (2)	2.25 (1.21)	.38**	—	—	—	—
Message reactance (3)	2.08 (.89)	.22**	.03	—	—	—
Perceived risk (4)	3.55 (.63)	.01	.20**	-.24**	—	—
Quit intentions (5)	2.57 (1.07)	.08**	.35**	-.15**	.23**	—

Note. Analyses excluded 56 participants with missing data on at least one of the variables. Pictorial warnings coded as 1 (vs. text-only, coded as 0). Mediators measured at immediate post-test (negative affect, message reactance) or the week 2 follow-up survey (perceived risk). Quit intentions measured at week 4.

* $p<.05$,
** $p<.001$

Table 4Mediation of association between pictorial warning exposure and quit intentions ($n=2,149$)

Mediation path	Mediated effect	95% CI
Pictorial warning → negative affect → quit intentions	.16	(.14, .19)
Pictorial warning → negative affect → perceived risk → quit intentions	.02	(.01, .02)
Pictorial warning → reactance → quit intentions	-.04	(-.06, -.02)
Pictorial warning → reactance → perceived risk → quit intentions	-.01	(-.02, -.01)

Note. Table reports standardized path coefficients for mediated effects. Mediators measured at immediate post-test (negative affect, message reactance) or the week 2 follow-up survey (perceived risk). Quit intentions measured at week 4.