Published in final edited form as:

Angew Chem Int Ed Engl. 2018 February 12; 57(7): 1939–1943. doi:10.1002/anie.201712145.

SuFEx Chemistry of Thionyl Tetrafluoride (SO_F₄) with Organolithium Nucleophiles: Synthesis of Sulfonimidoyl Fluorides, Sulfoximines, Sulfonimidamides and Sulfonimidates

Dr. Bing Gao^[a], Dr. Suhua Li^[a], Dr. Peng Wu^[b], Dr. John E. Moses^[c], and Dr. K. Barry Sharpless^[a]

^[a]Department of Chemistry, The Scripps Research Institute., 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

^[b]Department of Molecular Medicine, The Scripps Research Institute 10550 North Torrey Pines Road, La Jolla, CA, 92037, USA

^[c]La Trobe Institute For Molecular Science, La Trobe University, Bundoora, Melbourne, Victoria 3083, Australia

Abstract

Thionyl tetrafluoride (SO_F₄) is a valuable connective gas for sulfur fluoride exchange (SuFEx) click chemistry, that enables multidimensional linkages to be created via sulfur–oxygen (S–O) and sulfur–nitrogen (S–N) bonds. In this report, we expand the available SuFEx chemistry of SO_F₄ to include organolithium nucleophiles, and demonstrate, for the first time, the controlled projection of sulfur–carbon links at the sulfur center of SO_F₄-derived iminosulfur oxydifluorides (R¹–N=SO_F₂). This protocol enables rapid and modular access to sulfonimidoyl fluorides [R¹–N=SOFR²], another array of versatile SuFEx connectors with readily tunable reactivity of the S–F handle. Divergent connections derived from these valuable sulfonimidoyl fluoride units are also demonstrated, including the synthesis of sulfoximines, sulfonimidamides, and sulfonimidates.

Grahical Abstract

SuFEx with Sulfur–Carbon Link: Thionyl tetrafluoride (SO_F₄) is a connective gas that enables multidimensional linkages to be created through sulfur fluoride exchange (SuFEx). The installation of sulfur–carbon bond(s) at the sulfur center of SO_F₄-derived iminosulfur oxydifluorides is realized in the reaction with organolithium nucleophiles, affording sulfonimidoyl fluorides as versatile SuFEx-able connectors. The synthetic convenience of sulfonimidoyl fluorides is further demonstrated in the preparation of sulfoximines, sulfonimidamides and sulfonimidates.

Correspondence to: K. Barry Sharpless.

Conflict of Interest

A patent application covering this work has been filed by The Scripps Research Institute (TSRI, PCT Patent Application No. PCT/US2017/063746).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201xxxxxx>.

Keywords

Click Chemistry; SuFEx reaction; SOF_4 ; Organolithium; Sulfonimidoyl Fluoride

The sulfur fluoride exchange (SuFEx) family of click reactions enable the modular, reliable and practical formation of sulfur linked connections via connective hubs such as sulfonyl fluoride (SO_2F_2),^[1] ethenesulfonyl fluoride (ESF),^[2] and thionyl tetrafluoride (SOF_4).^[3] Sulfur fluoride derivatives are uniquely suited for click chemistry owing to their inherent thermodynamic and redox stability,^[4] yet, given the right conditions have alacrity for certain transformations through a cooperative engagement of catalyst, reagent and solvent.^[1] The power of SuFEx click chemistry has been demonstrated in applications, including the development of synthetic methods,^[5] the preparation of polymers,^[6] and in medicinal chemistry programs.^[7] Most intriguingly, there is growing evidence to suggest that S-F handles (e.g. fluorosulfates and sulfonyl fluorides) possess context-dependent protein reactivity that could be selective covalent (or non-covalent) enzyme inhibitors or chemical probes.^[1, 7]

We recently demonstrated thionyl tetrafluoride (SOF_4) gas as the first hub for creating multidimensional SuFEx connections.^[3a] In a typical process, the treatment of the trigonal bipyramidal SOF_4 with primary amines affords tetrahedral iminosulfur oxydifluorides (Figure 1A, J). The later has two further SuFEx-able S-F handles, which readily undergo sequential exchange with amines/anilines or derivatives of phenols. A diverse array of sulfur fluoride compounds and their downstream derivatives obtained from the SOF_4 assembly have dramatically enriched the SuFEx combinatorial library.

The SuFEx connections created from the SOF_4 hub to date are limited to S-O and S-N links involving naturally abundant nucleophiles (amines and phenols). Connections enabled through the S-C link(s) that would engage carbon nucleophiles, on the contrary, have not been explored. To further expand the scope of SuFEx connectivity and functionality, the S-C bond(s) construction via selective cleavage of S-F bond(s) at the sulfur center of SOF_4 or its

derivatives is worthwhile. Because it would enable divergent access to sulfur fluoride derivatives with novel skeletons and reactivity.

Herein, we report the controlled projection of S–C link(s) from iminosulfur oxydifluorides (**I**) upon reaction with organolithiums (Figure 1B). This protocol allows the synthesis of sulfonimidoyl fluorides in two steps from SOF₄ (Figure 1B, *II*). First reported by Johnson in 1983, sulfonimidoyl fluorides were conventionally prepared by the treatment of sulfonimidoyl chlorides with a fluoride source (Figure 1C).^[8] Because of the harsh reaction conditions and a lengthy *de novo* four-step process, however, only a confined number of derivatives were available.^[1, 9] Our protocol is complementary, but has significant advantages due to its straightforward, rapid and modular operation.^[10] The synthetic convenience for creating diverse connections from sulfonimidoyl fluorides, through either S–C or S–N/O links, is further demonstrated in the preparation of sulfoximines,^[11] sulfonimidamides,^[12] and sulfonimides.^[13] These compounds are known for their valuable utilities in synthetic transformations and medicinal chemistry.^[11–13]

The reaction between phenyliminosulfur difluoride (Ph–N=SF₂, S^{IV}) and phenyllithium was one of the rare examples of sulfur fluorides to react with organometallic reagents, but only affording a mixture of triphenylsulfilimine (Ph–N=S(Ph)₂) and diphenyl sulfide (PhSPh).^[14] In another case, the reduction of sulfur(VI) to lower-valent adducts was also observed during the nucleophilic substitution of sulfonyl fluoride derivatives (R–SO₂F, S^{VI}) with Grignard reagents.^[15] Fortunately, by using aryllithiums at low temperature, we could dramatically diminish the over-substitution and sulfur-reduction side paths in the reaction with iminosulfur oxydifluorides (**I**). The optimized conditions, 1.35 equivalents of phenyllithium in anhydrous cyclopentyl methyl ether (CPME) at –78 °C for 5 mins, allows mono-substitution of difluorides **I** into the corresponding sulfonimidoyl fluorides **II** in excellent yields (Table 1, see supporting information for condition optimization).^[16]

In general, aniline-derived difluorides **I** exhibited good selectivity for mono-substitution (**II-1–II-9**), while strong electron-withdrawing substituents (R¹), such as *p*-toluenesulfonyl group (**II-10**), tended to induce over-substitution. It is noteworthy that iminosulfur oxydifluorides (–N=SO₂F) are more electrophilic than fluorosulfates (–OSO₂F) in **II-4–II-6** that could be selectively functionalized by phenyllithium.^[3a] Such a differentiation implies it is possible for divergent functionalization of compounds containing multiple S–F handles.^[17] On the other hand, aliphatic primary amine-derived difluorides are relatively less electrophilic compared to the aniline derivatives, therefore exhibit even enhanced selectivity for mono-substitution (**II-11–II-20**).^[18] Functional groups that might be liable to the oxidative chlorination conditions in Johnson's route, such as methylthio motif (**II-7**), alkyl motif (**II-8**), olefin (**II-9**, **II-18**) and ferrocenyl group (**II-14**), are well tolerated under our conditions. Bromide and azide motifs are also compatible, thereby allowing for further functionalization (**II-3**, **II-15**). Finally, the synthesis of bis(sulfonimidoyl fluoride) **II-21** is showcased with moderate yield.

A variety of aryllithiums, prepared from halogen-lithium exchange or direct deprotonation,^[19] were investigated in the reaction with difluoride **I-11**.^[20] Especially, various heteroaryllithiums afforded the mono-substituted products in good to excellent yield (Table

1, **II-21–II-33**). Vinylic and alkynylic lithiums were examined, but no substitution was observed, probably due to their poor nucleophilicity. These results are also in good accordance with examples **II-8** and **II-9**, where terminal alkyl and olefin motifs would not interfere with the substitution reactions with phenyllithium. Alkylolithium reagents were also investigated in the reaction with **I-2**. Primary and secondary alkylolithium, such as *n*-butyllithium and *s*-butyllithium, mainly afford di-substituted products (sulfoximines **III-2**, **III-3**) rather than mono-substituted ones (**II-34**, **II-35**). In contrast, tertiary alkylolithiums *t*-butyllithium led exclusively to mono-substituted product **II-36**, with only trace amount of di-substituted sulfoximine **III-4**. This is probably due to a combined effect of steric hindrance and basicity of the nucleophiles and the products.^[21]

With a reliable process to sulfonimidoyl fluorides **II**, we next studied the formation of a second S–C link with the remaining S–F handle for the preparation of sulfoximines. THF was used as solvent to enhance the reactivity of organolithium nucleophiles for the more demanding second substitution.^[19] As illustrated in Table 2, the treatment of **II** with 2.0 equivalents of aryllithiums or alkylolithiums afforded the corresponding sulfoximines **III** in excellent yields. Importantly, we also identified an interesting trend that sulfonimidoyl fluoride motif in **II-6** is slightly more electrophilic than fluorosulfate motif to react with phenyllithium (determined based on the ratio of **III-5** and **III-6**). Finally, a Lewis acid induced S–F bond cleavage and subsequent S–C bond formation also gave sulfoximine **III-14** from sulfonimidoyl fluoride **II-3**.^[8]

Aside from creating sequential S–C linkages, the SuFEx reaction with heteroatom linkages has proved a robust approach for diverse functionalization through S–F handles. Sulfonimides and sulfonimidamides hitherto have mainly been prepared from sulfonimidoyl chlorides, which are liable to hydrolysis and redox side reactions.^[22, 23] In Table 3, we demonstrate the modification of the natural products Capsaicin, Estrone and (+)- δ -Tocopherol, through DBU-promoted formation of S–O links with sulfonimidoyl fluorides, giving the corresponding sulfonimides in excellent yields and with good functional group tolerance. As highlighted earlier in this manuscript, an *N*-alkyl substituent (**IV-1**) generally makes the S-center less electrophilic in comparison with an *N*-aryl substituent (**IV-2**, **IV-3**), and therefore requires excess DBU to get full conversion from the corresponding sulfonimidoyl fluorides. Again, DBU also serves as a promising base in the reaction with secondary amines (showcased in example **V-1**). For the reaction with primary amine and anilines, we found that deprotonation of the amino group with *n*-butyllithium prior to the nucleophilic substitution is necessary (**V-2–V-4**). These conditions are representative and generally applicable to a wide range of substrates.

To summarize, we have developed a new protocol for the controlled installation of S–C link(s) at the sulfur center of iminosulfur oxydifluorides ($R^1-N=SOF_2$), by sequential exchange of S–F bonds. As part of the fast modular assembly idea for the construction of multidimensional connections from the SOF_4 hub, this transformation might not be perfect, but is highly worthwhile. It enables a practical route for the rapid synthesis of sulfonimidoyl fluorides. Sulfonimidoyl fluorides themselves serve as versatile tetrahedral connectors with tunable reactivity of the S–F handles. Through the subsequent reaction with diverse nucleophiles, they readily afford sulfoximines, sulfonimides and sulfonimidamides that are

conventionally prepared through tedious routes. The current process is complementary to the established SuFEx chemistry featuring S–O/N links, yet offers enriched diversity to the family of sulfur fluorides and their derivatives that are valuable scaffolds for synthetic transformations and medicinal utilities.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the NIH (P50 GM103368, R01 GM117145) for financial support for this project. S.L. was partially supported by the Skaggs Institute for Chemical Biology. JEM thanks the Australian Research Council for a Future Fellowship (Project ID: FT170100156).

References

- Dong JJ, Krasnova L, Finn MG, Sharpless KB. *Angew Chem Int Ed.* 2014; 53:9430. *Angew Chem.* 2014; 126:9584.
- a) Krutak JJ, Burpitt RD, Moore WH, Hyatt JA. *J Org Chem.* 1979; 44:3847. b) Zheng QH, Dong JJ, Sharpless KB. *J Org Chem.* 2016; 81:11360. [PubMed: 27764941] c) Qin HL, Zheng QH, Bare GAL, Wu P, Sharpless KB. *Angew Chem Int Ed.* 2016; 55:14155. *Angew Chem.* 2016; 128:14361. d) Chinthakindi PK, Govender KB, Kurnar AS, Kruger HG, Govender T, Naicker T, Arvidsson PI. *Org Lett.* 2017; 19:480. [PubMed: 28075600] e) Zha GF, Zheng QH, Leng J, Wu P, Qin HL, Sharpless KB. *Angew Chem Int Ed.* 2017; 56:4849. *Angew Chem.* 2017; 129:4927.
- a) Li SH, Wu P, Moses JE, Sharpless KB. *Angew Chem Int Ed.* 2017; 56:2903. *Angew Chem.* 2017; 129:2949. b) Cramer R, Coffman DD. *J Org Chem.* 1961; 26:4010.
- a) Steinkopf W. *J Prakt Chem.* 1927; 117:1. b) Steinkopf W, Jaeger P. *J Prakt Chem.* 1930; 128:63. c) Petranek J. *J Chromatogr A.* 1961; 5:254.
- a) Kende AS, Mendoza JS. *J Org Chem.* 1990; 55:1125. b) Hyatt JA, White AW. *Synthesis.* 1984:214. c) Chen WT, Dong JJ, Li SH, Liu Y, Wang YJ, Yoon L, Wu P, Sharpless KB, Kelly JW. *Angew Chem Int Ed.* 2016; 55:1835. *Angew Chem.* 2016; 128:1867. d) Zhang EX, Tang JZ, Li SH, Wu P, Moses JE, Sharpless KB. *Chem Eur J.* 2016; 22:5692. [PubMed: 26990693] e) Ren GR, Zheng QH, Wang H. *Org Lett.* 2017; 19:1582. [PubMed: 28332844] f) Dondoni A, Marra A. *Org Biomol Chem.* 2017; 15:1549. [PubMed: 28116403]
- a) Dong JJ, Sharpless KB, Kwisnek L, Oakdale JS, Fokin VV. *Angew Chem Int Ed.* 2014; 53:9466. *Angew Chem.* 2014; 126:9620. b) Gao B, Zhang L, Zheng Q, Zhou F, Klivansky LM, Lu J, Liu Y, Dong J, Wu P, Sharpless KB. *Nat Chem.* 2017; 9:1083. [PubMed: 29064495] c) Oakdale JS, Kwisnek L, Fokin VV. *Macromolecules.* 2016; 49:4473. d) Wang H, Zhou F, Ren GR, Zheng QH, Chen HL, Gao B, Klivansky L, Liu Y, Wu B, Xu QF, Lu JM, Sharpless KB, Wu P. *Angew Chem Int Ed.* 2017; 56:11203. *Angew Chem.* 2017; 129:11355.
- a) Fahrney DE, Gold AM. *J Am Chem Soc.* 1963; 85:997. b) Baker BR. *Acc Chem Res.* 1969; 2:129. c) Narayanan A, Jones LH. *Chem Sci.* 2015; 6:2650. [PubMed: 28706662] d) Chen W, Dong J, Plate L, Mortenson DE, Brighty GJ, Li S, Liu Y, Galmozzi A, Lee PS, Hulce JJ, Cravatt BF, Saez E, Powers ET, Wilson IA, Sharpless KB, Kelly JW. *J Am Chem Soc.* 2016; 138:7353. [PubMed: 27191344] e) Fadeyi OO, Hoth LR, Choi C, Feng X, Gopalsamy A, Hett EC, Kyne RE, Robinson RP, Jones LH. *ACS Chem Biol.* 2017; 12:2015. [PubMed: 28718624] f) Baranczak A, Liu Y, Connelly S, Du WGH, Greiner ER, Genereux JC, Wiseman RL, Eisele YS, Bradbury NC, Dong JJ, Noodleman L, Sharpless KB, Wilson IA, Encalada SE, Kelly JW. *J Am Chem Soc.* 2015; 137:7404. [PubMed: 26051248]
- Johnson CR, Bis KG, Cantillo JH, Meanwell NA, Reinhard MFD, Zeller JR, Vonk GP. *J Org Chem.* 1983; 48:1.
- a) Vanleusen D, Vanleusen AM. *Recl Trav Chim Pay-Bas.* 1984; 103:41. b) Kowalczyk R, Edmunds AJF, Hall RG, Bolm C. *Org Lett.* 2011; 13:768. [PubMed: 21235264] c) Kolesnik NP, Rozhenko

- AB, Kinzhybalo V, Lis T, Shermolovich YG. *J Fluorine Chem.* 2014; 160:16.d) Garlyauskayte RY, Bezdudny AV, Michot C, Armand M, Yagupolskii YL, Yagupolskii LM. *J Chem Soc Perkin Trans 1.* 2002;1887.e) Reggelin M, Weinberger H. *Tetrahedron Lett.* 1992; 33:6959.
10. As analogues of sulfonyl fluorides, the reactivity of the sulfonimidoyl fluorides is readily tunable with our protocol (through manipulating R^1 and/or R^2). That would be valuable for future investigation of sulfonimidoyl fluorides as covalent pharmacophores (see ref. 1).
11. a) Lücking U. *Angew Chem Int Ed.* 2013; 52:9399. *Angew Chem.* 2013; 125:9570.b) Frings M, Bolm C, Blum A, Gnam C. *Eur J Med Chem.* 2017; 126:225. [PubMed: 27821325] c) Reggelin M, Zur C. *Synthesis.* 2000;1.d) Okamura H, Bolm C. *Chem Lett.* 2004; 33:482.e) Shen X, Hu JB. *Eur J Org Chem.* 2014;4437.f) Bizet V, Kowalczyk R, Bolm C. *Chem Soc Rev.* 2014; 43:2426. [PubMed: 24549291] g) Bizet V, Hendriks CMM, Bolm C. *Chem Soc Rev.* 2015; 44:3378. [PubMed: 25941981] h) Zenzola M, Doran R, Degennaro L, Luisi R, Bull JA. *Angew Chem Int Ed.* 2016; 55:7203. *Angew Chem.* 2016; 128:7319.i) Bizet V, Buglioni L, Bolm C. *Angew Chem Int Ed.* 2014; 53:5639. *Angew Chem.* 2014; 126:5745.j) Yu H, Li Z, Bolm C. *Angew Chem Int Ed.* 2018; 57:324. *Angew Chem.* 2018; 130:330.
12. a) Chinthakindi PK, Naicker T, Thota N, Govender T, Kruger HG, Arvidsson PI. *Angew Chem Int Ed.* 2017; 56:4100. *Angew Chem.* 2017; 129:4160.b) Pemberton N, Graden H, Evertsson E, Bratt E, Lepistö M, Johannesson P, Svensson PH. *ACS Med Chem Lett.* 2012; 3:574. [PubMed: 24900513] c) Gnam C, Jeanguenat A, Dutton AC, Grimm C, Kloer DP, Crossthwaite AJ. *Bioorg Med Chem Lett.* 2012; 22:3800. [PubMed: 22552196]
13. a) Roy AK. *J Am Chem Soc.* 1993; 115:2598.b) Leca D, Fensterbank L, Lacote E, Malacria M. *Org Lett.* 2002; 4:4093. [PubMed: 12423094] c) Felim A, Toussaint A, Phillips CR, Leca D, Vagstad A, Fensterbank L, Lacote E, Malacria M. *Org Lett.* 2006; 8:337. [PubMed: 16408909]
14. Smith WC, Tullock CW, Smith RD, Engelhardt VA. *J Am Chem Soc.* 1960; 82:551.
15. a) Gilman H, Fothergill RE. *J Am Chem Soc.* 1929; 51:3501.b) Shirota Y, Nagai T, Tokura N. *Tetrahedron.* 1967; 23:639.
16. Although 5 minutes is given as a standard reaction time herein, the transformations are extremely rapid that could get instant conversion upon the addition of lithium reagents.
17. Hanley PS, Ober MS, Krasovskiy AL, Whiteker GT, Kruper WJ. *ACS Catal.* 2015; 5:5041.
18. *N*-alkyl substituents could be converted to *N*-acyl substituents, see: Lamers P, Priebbenow DL, Bolm C. *Eur J Org Chem.* 2015:5594.
19. Clayden, J. *Organolithiums: Selectivity for Synthesis.* Elsevier Science & Technology Books; 2002.
20. *p*-Methoxybenzyl group is removable at the late stage of transformations: Sirvent JA, Bierer D, Webster R, Lücking U. *Synthesis.* 2017:1024.
21. Please see supporting information for summary of the reactions between I-2 and alkylolithiums (Section 5, Page 29).
22. Johnson CR, Jonsson EU, Bacon CC. *J Org Chem.* 1979; 44:2055.
23. There are alternative protocols for the preparation of sulfonimidamides, see selected examples: Wen J, Cheng HC, Dong SX, Bolm C. *Chem Eur J.* 2016; 22:5547. [PubMed: 26892735] Izzo F, Schafer M, Stockman R, Lücking U. *Chem Eur J.* 2017; 23:15189. [PubMed: 28833686] Richards-Taylor CS, Martinez-Lamenca C, Leenaerts JE, Trabanco AA, Oehlrich D. *J Org Chem.* 2017; 82:9898. [PubMed: 28809121] Davies TQ, Hall A, Willis MC. *Angew Chem Int Ed.* 2017; 56:14937. *Angew Chem.* 2017; 129:15133.

A, Previous work: "S-O" and "S-N" Links at the SOF₄ Hub (2017, Sharpless; 1961, Cramer & Coffman)

B, This work: "S-C" Link(s) at the SOF₄-derived Iminosulfur Oxydifluoride Hubs

C, Previous Work: Conventional Route to Sulfonimidoyl Fluorides (1983, Johnson)

Figure 1.

(A) Sulfur-heteroatom links enabled connections at the SOF₄ hub; (B) Sulfur-carbon link(s) enabled connections at the SOF₄ hub; (C) Conventional route to sulfonimidoyl fluorides.

Table 1

Synthesis of Sulfonylimidoyl Fluorides from SO_2F_2 ^[a]

$\text{O}=\text{S}(\text{F})_2 \xrightarrow[\text{Et}_3\text{N (2.0 equiv), CH}_3\text{CN, RT, < 30 min}]{\text{R}^1\text{-NH}_2} \text{R}^1\text{-N}(\text{F})\text{SO}_2\text{F} \text{ (I)} \xrightarrow[\text{CPME (0.1 M), -78 }^\circ\text{C, 5 min}]{\text{R}^2\text{-Li (1.35 equiv)}} \text{R}^1\text{-N}(\text{F})\text{SO}_2\text{R}^2 \text{ (II)}$
II-1 (±), 81% (0.5 g scale)	II-2 (±), 70%	II-3 (±), 74% (1.5 g scale)	II-4 (±), 65%	II-5 (±), 72%	II-6 (±), 71% (0.6 g scale)
II-7 (±), 87%	II-8 (±), 64% ^[b]	II-9 (±), 87%	II-10 (±), 10% ^[c]	II-11 (±), 82% (2.2 g scale)	II-12 (±), 66%
II-13 (±), 79% (1.2 g scale)	II-14 (±), 89%	II-15, 85% d.r. = 1.0 : 2.4	II-16, 87% d.r. = 1.0 : 2.1	II-17, 90% d.r. = 1.0 : 0.6	II-18, 83% d.r. = 1.0 : 1.4
II-19, 91% d.r. = 1.0 : 1.1	II-20, 93% d.r. = 1.0 : 4.0	II-21 (±), 47% ^[d]	II-22 (±), 81% ^[e]	II-23 (±), 90% ^[f]
II-24 (±), 81% ^[f]	II-25 (±), 69% ^[g]	II-26 (±), 45% ^[f]	II-27 (±), 17% ^[f]	II-28 (±), 67% ^[f]
II-29 (±), 42%	II-30 (±), 53% ^[e]	II-31 (±), 70% ^[e]	II-32 (±), 93% ^[e]	II-33 (±), 42% ^[e]	

^[a] See supporting information (SI) for the preparation of **I**; For the substitution reaction: **I** (0.2 mmol), ArLi (0.27 mmol), cyclopentyl methyl ether (CPME, 2.0 mL), -78°C , 5 mins; then quenched with $\text{CH}_3\text{CO}_2\text{H}$ (1.0 M in MeOH) at -78°C . Isolated yields from **I** to **II** were reported.

^[b] 2.4 equiv of PhLi used.

^[c] Di-substituted product **III-1** was isolated in 54% yield.

^[d] 2.7 equiv of PhLi used.

^[e] ArLi was prepared from direct C–H lithiation, see SI.

^[f] ArLi were prepared from aryl bromides (ArBr) via halogen-lithium exchange and THF were used as solvent in most cases, see SI.

^[g] ArLi was prepared from aryl iodide (ArI) via halogen-lithium exchange, see SI.

Table 2

Sulfoximines from Sulfonimidoyl Fluorides.^[a] ^[a] **II** (0.2 mmol), R²-Li (0.4 mmol), THF (2.0 mL), -78 °C, 10 mins.^[b] **II** (0.2 mmol), Ph(TMSO)=CH₂ (0.26 mmol), TiCl₄ (0.2 mmol), DCM (2.0 mL), -78 °C, 30 mins.

Table 3

Sulfonimidates and Sulfonimidamides from Sulfonimidoyl Fluorides.

^[a] **II** (0.2 mmol), R³OTBS (0.2 mmol), DBU (20 mol% to 2.0 equiv), CH₃CN (2.0 mL), 60 °C, 10 h.

^[b] **II** (0.2 mmol), Pyrrolidine (0.4 mmol), DBU (0.4 mmol), CH₃CN (2.0 mL), 60 °C, 24 h.

^[c] R³NH₂ (0.4 mmol), *n*-BuLi (0.4 mmol), THF (2.0 mL), -70 °C, 15 mins, then **II** (0.2 mmol), 0 °C to RT.