Andrew A. Monte, M.D.*

University of Colorado
Aurora, Colorado

and

Rocky Mountain Poison & Drug Center
Denver, Colorado

Hao Sun, M.D.*

University of Colorado
Aurora, Colorado

Anna Malin Rapp-Olsson, M.S.

Rocky Mountain Poison & Drug Center
Denver, Colorado

Fahim Mohamed, Ph.D.

University of Peradeniya
Peradeniya, Sri Lanka

and

University of Sydney
Sydney, Australia

Indika Gawarammana, Ph.D.

University of Peradeniya
Peradeniya, Sri Lanka

Nicholas A. Buckley, M.D.

University of Peradeniya
Peradeniya, Sri Lanka

and

University of Sydney
Sydney, Australia

Christopher M. Evans, Ph.D.

Ivana V. Yang, Ph.D.
David A. Schwartz, M.D.

University of Colorado
Aurora, Colorado

*A.A.M. and H.S. share co-first authorship on this work.

References

- Gawarammana IB, Buckley NA. Medical management of paraquat ingestion. *Br J Clin Pharmacol* 2011;72:745–757.
- Eddleston M, Gunnell D, Karunaratne A, de Silva D, Sheriff MH, Buckley NA. Epidemiology of intentional self-poisoning in rural Sri Lanka. *Br J Psychiatry* 2005;187:583–584.
- Gunawardena G, Roberts DM, Buckley NA. Randomized control trial of immunosuppression in paraquat poisoning. *Crit Care Med* 2007;35:330–331, author reply 331.
- Roy MG, Livraghi-Butrico A, Fletcher AA, McElwee MM, Evans SE, Boerner RM, et al. Muc5b is required for airway defence. *Nature* 2014;505:412–416.
- Nakano Y, Yang IV, Walts AD, Watson AM, Helling BA, Fletcher AA, et al. Muc5b promoter variant rs35705950 affects muc5b expression in the distal airways in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2016;193:464–466.
- Seibold MA, Wise AL, Speer MC, Steele MP, Brown KK, Loyd JE, et al. A common MUC5B promoter polymorphism and pulmonary fibrosis. *N Engl J Med* 2011;364:1503–1512.
- Peljo AL, Zhang Y, Fingerlin TE, Ma SF, Garcia JG, Richards TJ, et al. Association between the MUC5B promoter polymorphism and survival in patients with idiopathic pulmonary fibrosis. *JAMA* 2013;309:2232–2239.
- Senarathna L, Eddleston M, Wilks MF, Woollen BH, Tomenson JA, Roberts DM, et al. Prediction of outcome after paraquat poisoning by measurement of the plasma paraquat concentration. *QJM* 2009;102:251–259.
- Wunnapuk K, Mohammed F, Gawarammana I, Liu X, Verbeeck RK, Buckley NA, et al. Prediction of paraquat exposure and toxicity in clinically ill poisoned patients: a model based approach. *Br J Clin Pharmacol* 2014;78:855–866.
- Buckley NA. Pulse corticosteroids and cyclophosphamide in paraquat poisoning. *Am J Respir Crit Care Med* 2001;163:585.

Copyright © 2018 by the American Thoracic Society

Downregulation of MicroRNA-126 Augments DNA Damage Response in Cigarette Smokers and Patients with Chronic Obstructive Pulmonary Disease

To the Editor:

Increased DNA damage response (DDR) contributes to the pathophysiology of aging disorders, including chronic obstructive pulmonary disease (COPD), cardiovascular disease (CVD), and cancer (1, 2). DNA damage resulting from cigarette smoke-induced oxidative stress activates the DDR signaling pathway, controlled mainly by the ATM (ataxia-telangiectasia mutated) protein kinase. Accumulation of DNA damage promotes cellular senescence and disrupts tissue homeostasis (2). Using circulating endothelial precursors called blood outgrowth endothelial cells (BOEC) or endothelial colony-forming cells, we have previously demonstrated increased DDR, endothelial senescence, and dysfunction in smokers and patients with COPD (3), supporting the concept of accelerated aging of the endothelium as contributing to CVD in these groups.

MicroRNAs are important regulators of almost every cellular process, including DNA repair. MicroRNA-126 (miR-126; also known as miR-126-3p), a microRNA enriched in endothelial cells, plays a key role in angiogenesis and vascular homeostasis (4). miR-126 is reduced in patients susceptible to developing or with CVD (5–7) and in several types of cancer, including lung cancer (8). The link between miR-126 and DDR signaling in COPD is currently unknown. Therefore, we investigated whether miR-126 is reduced by cigarette smoke *in vivo* and in endothelial and lung epithelial cells from patients with COPD, as well as its link to DDR activation.

Methods

To investigate DDR resulting from cigarette smoke *in vivo*, male C57BL/6 mice were challenged for 28 days with cigarette smoke or ambient air, as previously described (9). BOEC were isolated from human blood samples, and human primary bronchial epithelial cells were isolated from lung tissue as described (3, 10). Informed consent was obtained from all individuals (Table 1). Data have been derived from subjects and animals used in previous reports (3, 9, 10). Human umbilical vein endothelial cells were transfected with miRCURY

This work was supported by a Wellcome Trust Programme Grant (093080/Z/10/Z). K.E.P. is financially supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Imperial College Healthcare NHS Trust and Imperial College London. A.Z. and R.D.S. were financially supported by the British Heart Foundation (BHF). M.M. is a BHF Chair Holder, and his research is supported by the BHF, NIHR Biomedical Research Centre based at Guy's and St. Thomas' NHS Foundation Trust and King's College London, and Fondation Leducq Transatlantic Network of Excellence. A.M.R.'s research is supported by a BHF programme grant and the BHF Centre for Vascular Regeneration (Rm/30158).

Author Contributions: Conception and design: K.E.P., A.M.R., and P.J.B.; data analysis and interpretation: K.E.P., A.Z., J.R.B., M.A.B., R.D.S., M.G.B., L.E.D., M.M., A.M.R., and P.J.B.; experimental performance: K.E.P., A.Z., and J.R.B.; and writing: K.E.P., A.M.R., and P.J.B.

Originally Published in Press as DOI: 10.1164/rccm.201706-1304LE on July 28, 2017

Table 1. Clinical Characteristics of Volunteers

	Healthy Nonsmokers	Healthy Smokers	COPD
Blood outgrowth endothelial cells			
Number, <i>n</i>	7	6	5 (4 moderate; 1 severe)*
Sex, M/F	4/3	4/2	5/0
Age, yr	54 ± 4	57 ± 4	69 ± 3 [†]
Smoking, pack-years	0	37 ± 10	68 ± 38
FEV ₁ , % predicted	99.1 ± 6.4	93.6 ± 3.8	61.4 ± 5.8 ^{‡,§}
FEV ₁ /FVC, % predicted	77.9 ± 0.8	71.6 ± 3.6	53 ± 5.2 ^{§,}
Lung epithelial cells			
Number, <i>n</i>	6		6 (3 moderate; 1 severe; 2 very severe)*
Sex, M/F	0/6		5/1
Age, yr	63 ± 6		66 ± 2
Smoking, pack-years	0		47 ± 5
FEV ₁ , % predicted	97.2 ± 5.6		47.2 ± 10.5
FEV ₁ /FVC, % predicted	75.2 ± 3.4		42.02 ± 9.2

Definition of abbreviation: COPD = chronic obstructive pulmonary disease. Values are expressed as means ± SEM unless otherwise indicated. FEV₁ and FEV₁/FVC ratio are post-bronchodilator for subjects with COPD, smokers, or nonsmokers.

*Staging of COPD is according to the Global Initiative for Chronic Obstructive Lung Disease criteria.

[†]*P* < 0.05 (comparison between nonsmokers and COPD).

[‡]*P* < 0.001 (comparison between nonsmokers and COPD).

[§]*P* < 0.01 (comparison between smokers and COPD).

^{||}*P* < 0.01 (comparison between nonsmokers and COPD).

locked-nucleic-acid microRNA inhibitor hsa-miR-126-3p (Exiqon; #4109148-001) or negative control (#199006-001) at 50 nM. RNA was extracted using the mirVana Isolation Kit (Ambion) or the RNeasy kit (Qiagen). miR-126-3p levels, and the small nucleolar RNAs RNU44 and snoRNA135 as reference genes, were measured by TaqMan assays (#477887, #4427975, #4427975; Life Technologies) (7). Levels of ATM, SPRED-1 (Sprouty-related EVH1 domain containing 1), and PIK3R2 (phosphoinositol-3 kinase regulatory subunit 2; also known as p85-β) normalized to GAPDH mRNA were measured by quantitative real-time PCR. Primer sequences are available on request. Immunofluorescence was performed with the phospho-(serine/threonine) ATM/ataxia telangiectasia and Rad3-related protein (ATR) substrate antibody (Cell Signaling Technology, New England Biolabs). Nuclei were visualized using deep red anthraquinone 5 (DRAQ5) (Biostatus Limited). The following antibodies were used for Western blotting: ATM (Cell Signaling Technology), phospho-ATM-(Ser1981) (Millipore), and α-tubulin (Sigma-Aldrich Co. Ltd). Statistical analysis was performed with GraphPad Prism 7. Data are expressed as mean ± SEM. Comparisons between groups were performed with one-way ANOVA, Student's *t* tests, or Mann-Whitney *U* test, according to normality of the data. Correlations were estimated with Spearman correlation coefficient (*r*).

Results

The ATM protein kinase is rapidly phosphorylated and activated in response to DNA double-strand breaks and regulates the DDR (2). ATM phosphorylation was significantly increased in BOEC from patients with COPD compared with nonsmokers (Figure 1A), suggesting activation of ATM signaling pathway in COPD. We next investigated whether ATM activation is linked to miR-126 dysregulation. miR-126 levels were reduced in BOEC from smokers and patients with COPD compared with nonsmokers (Figure 1B), suggesting cigarette smoke downregulates miR-126 expression in the endothelium and could contribute to the aberrant activation of DDR in these groups.

Subsequently, we examined whether miR-126 regulates ATM protein kinase and DDR. As previously shown (4), inhibition of miR-126 function in human umbilical vein endothelial cells increased the expression of PIK3R2 and SPRED1 (Figure 1C). Under the same conditions, ATM mRNA and protein levels were increased (Figure 1C), suggesting miR-126 controls DDR by repressing ATM expression.

We then used a mouse model of subchronic exposure to cigarette smoke to confirm our findings *in vivo*. We found that miR-126 levels were reduced in lungs from mice exposed to cigarette smoke (Figure 1D). Within the same samples, there was also increased ATM expression and activity (Figure 1E-F), in line with our *ex vivo* findings in patients' endothelial cells.

As miR-126 was dysregulated in whole-lung tissue in our *in vivo* model, we investigated whether miR-126 expression is affected in lung epithelial cells in COPD. Interestingly, we found reduced miR-126 levels in lung epithelial cells from patients with COPD (all current or ex-smokers) compared with nonsmokers (Figure 1G). In addition, miR-126 levels in lung epithelial cells negatively correlated with smoking history assessed as pack-years (Spearman's *r* = −0.73; *P* < 0.01) and positively correlated with disease severity measured as FEV₁% predicted (Spearman's *r* = 0.57; *P* < 0.05), suggesting miR-126 expression is downregulated with extended exposure to cigarette smoke and increased severity of lung disease.

These results identify a novel miR-126-dependent pathway controlling DDR caused by cigarette smoke, where downregulation of miR-126 enhances ATM activation and thereby promotes tissue aging and dysfunction.

Discussion

In this study, we demonstrate that chronic exposure to cigarette smoke causes reduced expression of miR-126 and increases the DDR. Using a combined *in vivo* and *ex vivo* approach, namely, mice exposed to cigarette smoke and endothelial and lung epithelial cells from patients with COPD, we show that exposure to cigarette smoke downregulates miR-126 expression.

miR-126 is critical for endothelial function, and miR-126 supplementation has been proposed therapeutically in *in vivo* models of vascular injury and pulmonary hypertension (5, 6). We have previously shown that BOEC from both smokers and patients with COPD exhibit increased DDR, senescence, and dysfunction (3); these molecular defects could contribute to CVD, a major cause of mortality in these individuals. Here we demonstrate that miR-126 is downregulated in BOEC and in lung epithelial cells from patients with COPD, highlighting miR-126 as a possible contributor to endothelial and lung tissue dysfunction, and thus a potential therapeutic target for COPD.

Figure 1. MicroRNA-126 (miR-126) is downregulated and enhances ATM (ataxia-telangiectasia mutated) activation in smokers and patients with chronic obstructive pulmonary disease (COPD) and in an *in vivo* model of subchronic cigarette smoke exposure. (A) Activation of ATM was assessed by measuring phosphorylation of the ATM/ATR substrate by immunofluorescence staining ($n = 3$). Deep red anthraquinone 5 (DRAQ5) was used for nuclear staining. The number of nuclear foci per nucleus was quantified with Volocity software in at least five optic fields (z-stack images). An increased number of cells with focal nuclear staining of phosphorylated ATM/ATR (p-ATM/ATR) was observed in patients with COPD compared with nonsmokers. Arrows show nuclei with increased number of nuclear foci (scale bars, 20 μ m). (B) We measured miR-126 by TaqMan real-time PCR in blood outgrowth endothelial cells from nonsmokers ($n = 7$), healthy smokers ($n = 6$), and patients with COPD ($n = 5$). RNU44 was used for normalization. (C) Human umbilical vein endothelial cells were treated with specific anti-miR-126 inhibitor versus control for 72 hours (baseline conditions, no growth factors). mRNA levels for PIK3R2 (phosphoinositide-3 kinase regulatory subunit 2; also known as p85- β) and SPRED1 (Sprouty-related EVH1 domain containing 1) (positive controls for miR-126 inhibition) and for ATM were measured by real-time PCR. ATM protein levels were quantified by Western blotting. GAPDH and α -tubulin were measured for normalization. (D–F) Male C57BL/6 mice were challenged for 28 days with cigarette smoke (for 1 hour twice daily, 4 hours apart) or ambient air. (D) miR-126 was measured by TaqMan real-time PCR in lung tissue from mice exposed to cigarette smoke ($n = 6$) versus control mice lung tissue ($n = 5$). Small nucleolar RNA135 (snoRNA135) was measured for normalization. (E and F) ATM mRNA levels were measured by real-time PCR and activation of ATM (phospho-ATM), and ATM protein levels were measured by Western blotting in lung tissue from mice exposed to cigarette smoke ($n = 6$) versus control mice ($n = 6$). GAPDH and α -tubulin were measured for normalization. (G) miR-126 was measured by TaqMan real-time PCR in lung epithelial cells from nonsmokers ($n = 6$) and patients with COPD ($n = 6$). RNU44 was used for normalization. Data presented are means $\pm$ SEM. * $P < 0.05$; ** $P < 0.01$. ATR = ataxia telangiectasia and Rad3-related protein.

We also show that miR-126 controls the DDR by repressing ATM protein kinase activity in endothelial cells. This is important, as it highlights a novel pathogenetic pathway of miR-126 in age-related diseases, including cancer. Further studies will show whether restoration of miR-126 has a beneficial effect on lung and vascular function in COPD by regulating the aberrant activation of ATM and downstream molecular pathways that promote accelerated lung and vascular aging. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Acknowledgment: The authors thank Prof. James Scott, Prof. Dorian Haskard, and Prof. Justin Mason for their constructive criticism. They also thank Peter Fenwick and Sally Meah for collecting patients' samples.

Koralia E. Paschalaki, M.D., Ph.D.
Imperial College London
London, United Kingdom

Anna Zampetaki, Ph.D.
King's College London British Heart Foundation Centre of Research
Excellence
London, United Kingdom

Jonathan R. Baker, Ph.D.
Mark A. Birrell, Ph.D.
Richard D. Starke, Ph.D.
Maria G. Belvisi, Ph.D.
Louise E. Donnelly, Ph.D.
Imperial College London
London, United Kingdom

Manuel Mayr, M.D., Ph.D.
King's College London British Heart Foundation Centre of Research
Excellence
London, United Kingdom

Anna M. Randi, M.D., Ph.D.*
Peter J. Barnes, M.D., D.Sc.*
Imperial College London
London, United Kingdom

*These authors share senior authorship.

ORCID ID: 0000-0002-4114-6140 (K.E.P.).

References

1. Aoshiba K, Zhou F, Tsuji T, Nagai A. DNA damage as a molecular link in the pathogenesis of COPD in smokers. *Eur Respir J* 2012;39:1368–1376.
2. Childs BG, Durik M, Baker DJ, van Deursen JM. Cellular senescence in aging and age-related disease: from mechanisms to therapy. *Nat Med* 2015;21:1424–1435.
3. Paschalaki KE, Starke RD, Hu Y, Mercado N, Margariti A, Gorgoulis VG, Randi AM, Barnes PJ. Dysfunction of endothelial progenitor cells from smokers and chronic obstructive pulmonary disease patients due to increased DNA damage and senescence. *Stem Cells* 2013;31:2813–2826.
4. Fish JE, Santoro MM, Morton SU, Yu S, Yeh RF, Wythe JD, Ivey KN, Bruneau BG, Stainier DY, Srivastava D. miR-126 regulates angiogenic signaling and vascular integrity. *Dev Cell* 2008;15:272–284.
5. Jansen F, Yang X, Hoelscher M, Cattelan A, Schmitz T, Proebsting S, Wenzel D, Vosen S, Franklin BS, Fleischmann BK, et al. Endothelial microparticle-mediated transfer of MicroRNA-126 promotes vascular endothelial cell repair via SPRED1 and is abrogated in glucose-damaged endothelial microparticles. *Circulation* 2013;128:2026–2038.
6. Potus F, Ruffenach G, Dahou A, Thebault C, Breuils-Bonnet S, Tremblay É, Nadeau V, Paradis R, Graydon C, Wong R, et al. Downregulation of microRNA-126 contributes to the failing right ventricle in pulmonary arterial hypertension. *Circulation* 2015;132:932–943.
7. Zampetaki A, Kiechl S, Drozdov I, Willeit P, Mayr U, Prokopi M, Mayr A, Weger S, Oberhollenzer F, Bonora E, et al. Plasma microRNA profiling reveals loss of endothelial miR-126 and other microRNAs in type 2 diabetes. *Circ Res* 2010;107:810–817.
8. Ebrahimi F, Gopalan V, Smith RA, Lam AK. miR-126 in human cancers: clinical roles and current perspectives. *Exp Mol Pathol* 2014;96:98–107.
9. Eltom S, Stevenson CS, Rastrick J, Dale N, Raemdonck K, Wong S, Catley MC, Belvisi MG, Birrell MA. P2X7 receptor and caspase 1 activation are central to airway inflammation observed after exposure to tobacco smoke. *PLoS One* 2011;6:e24097.
10. Baker JR, Vuppusetty C, Colley T, Papaioannou AI, Fenwick P, Donnelly L, Ito K, Barnes PJ. Oxidative stress dependent microRNA-34a activation via PI3Kα reduces the expression of sirtuin-1 and sirtuin-6 in epithelial cells. *Sci Rep* 2016;6:35871.

Copyright © 2018 by the American Thoracic Society

X Chromosome–encoded MicroRNAs Are Functionally Increased in Cystic Fibrosis Monocytes

To the Editor:

Cystic fibrosis (CF) is an autosomal recessive disease. In addition to primary *CFTR* (CF conductance regulator) gene defects, other factors contribute to CF lung disease, including altered microRNA (miRNA) expression. miRNAs are short regulatory RNAs responsible for posttranscriptional inhibition of protein expression. Various miRNA expression profiling studies have demonstrated altered miRNA expression profiles in people with CF and in *in vitro* CF cell models (reviewed in [1]). Researchers in functional studies have assessed the biological consequences of some of the altered miRNAs. Whether the expression of miRNA in peripheral blood monocytes, and specifically miRNAs encoded on the X chromosome, is altered in p.Phe508del homozygous individuals is not known.

The human genome comprises 22 pairs of autosomes and a pair of sex chromosomes, XX in females and XY in males. X chromosome inactivation achieves dosage compensation between the sexes, whereas X upregulation achieves a balanced expression between the X chromosome and autosomes. These mechanisms can be inconsistent, leading to variability in gene expression between sexes, individuals, cell types, and developmental stages (2, 3). The X chromosome has a high density of miRNAs compared with autosomes, encoding approximately 7% of the total miRNAs in the genome (miRBase, release 21). The contribution of sex chromosomes to CF is virtually unknown.

In the present study, we examined the expression patterns of 86 X-linked miRNAs in monocytes of males and females with and without CF using PCR arrays. Twenty-four individuals were recruited into this study; 12 were stable p.Phe508del *CFTR* homozygotes confirmed by genotyping, and 12 were non-CF control subjects (6 females and 6 males per cohort) with mean $\pm$ SD ages of 23.5 ± 5.1 years and 27.3 ± 3.6 years, respectively. FEV₁

Funded by the Health Research Board in Ireland (grant PHD/2007/11), the European Respiratory Society GSK Rare Disease Award 2013, and Cystic Fibrosis Foundation Therapeutics (GREENE15XXO).

Originally Published in Press as DOI: 10.1164/rccm.201707-1417LE on August 10, 2017