Published in final edited form as:

Methods Enzymol. 2018 ; 605: 335–349. doi:10.1016/bs.mie.2018.02.011.

Recent Advances in the Genetic Manipulation of *Methylosinus trichosporium* OB3b

Soo Y. Ro and Amy C. Rosenzweig¹

Northwestern University, Evanston, IL, United States

Abstract

Methanotrophic bacteria utilize methane as their sole carbon and energy source. Studies of the model Type II methanotroph *Methylosinus trichosporium* OB3b have provided insight into multiple aspects of methanotrophy, including methane assimilation, copper accumulation, and metal-dependent gene expression. Development of genetic tools for chromosomal editing was crucial for advancing these studies. Recent interest in methanotroph metabolic engineering has led to new protocols for genetic manipulation of methanotrophs that are effective and simple to use. We have incorporated these newer molecular tools into existing protocols for *Ms. trichosporium* OB3b. The modifications include additional shuttle and replicative plasmids as well as improved gene delivery and genotyping. The methods described here render gene editing in *Ms. trichosporium* OB3b efficient and accessible.

1. INTRODUCTION

Methanotrophic bacteria are unique in their ability to utilize methane, a potent greenhouse gas, as their sole carbon source (Hanson & Hanson, 1996). In the first step of their metabolic pathway, methane is oxidized to methanol, which can be utilized as a cheap carbon feedstock for industrial chemical processes (Lawton & Rosenzweig, 2016). For this reason, there have been extensive efforts to understand methanotrophy in multiple species of bacteria. Most aerobic methanotrophs are classified as Type I (Gammaproteobacteria) or Type II (Alphaproteobacteria) based on their use of either the ribulose monophosphate or serine cycles for carbon assimilation, respectively, as well as their phospholipid compositions, cell morphologies, and membrane arrangements (Hanson & Hanson, 1996; Semrau, DiSpirito, & Yoon, 2010). Methane is oxidized to methanol by methane monooxygenase enzymes (MMOs), followed by methanol conversion to formaldehyde by methanol dehydrogenases (MDHs). There are two types of MMO, a soluble and cytoplasmic iron-containing enzyme (sMMO) and a “particulate” and inner membrane-bound copper-containing enzyme (pMMO) (Ross & Rosenzweig, 2016). With the exception of the genera *Methylocella* and *Methyloferula*, aerobic methanotrophs express pMMO under most conditions (Semrau et al., 2010; Vorobev et al., 2011). Under copper-starved conditions, some Type I and Type II methanotrophs can switch to sMMO expression (Semrau et al., 2010), and some *Methylosinus* and *Methylocystis* species also produce the copper chelator methanobactin (Dassama, Kenney, & Rosenzweig, 2017).

¹Corresponding author: amy@northwestern.edu.

Methylosinus trichosporium OB3b is the best studied Type II methanotroph (Stein et al., 2010). This bacterium can express both sMMO and pMMO, produces methanobactin (Dassama et al., 2017), and can express two distinct MDHs, the calcium-binding MxaF and the lanthanide-binding XoxF (Haque, Kalidass, et al., 2015). Genetic tools developed previously for *Ms. trichosporium* OB3b have been used to mutate or disrupt the genes encoding several important proteins. Recent studies have focused on sMMO (Borodina, Nichol, Dumont, Smith, & Murrell, 2007; Lock, Nichol, Murrell, & Smith, 2017; Semrau et al., 2013; Smith & Murrell, 2011; Smith, Slade, Burton, Murrell, & Dalton, 2002), proteins in the extended *pmo* operon (Gu, Haque, & Semrau, 2017), methanobactin transport and biosynthetic proteins (Dassama, Kenney, Ro, Zielazinski, & Rosenzweig, 2016; Gu, Baral, DiSpirito, & Semrau, 2017; Gu et al., 2016; Gu, Haque, et al., 2017), and genes related to MDH (Haque, Gu, DiSpirito, & Semrau, 2015).

In recent years, new genetic manipulation tools have been developed for two Type I haloalkaliphilic methanotrophs, *Methylobacterium alcaliphilum* 20Z (Kalyuzhnaya et al., 2008; Ojala, Beck, & Kalyuzhnaya, 2011) and *Methylobacterium buryatense* 5G (Puri et al., 2015; Yan, Chu, Puri, Fu, & Lidstrom, 2016). These protocols have been used to engineer methanotrophs for biofuel production (de la Torre et al., 2015; Henard et al., 2016; Kalyuzhnaya, Puri, & Lidstrom, 2015) and have also enabled studies of lanthanide-dependent methanol oxidation (Chu & Lidstrom, 2016), ectoine biosynthesis (Mustakhimov et al., 2009), and fatty acid biosynthesis (Demidenko, Akberdin, Allemann, Allen, & Kalyuzhnaya, 2017). This chapter describes the incorporation of these newer techniques developed for Type I methanotrophs into the traditional mutagenesis protocol for *Ms. trichosporium* OB3b (Murrell, 1994; Lloyd, Finch, Dalton, & Murrell, 1999; Smith & Murrell, 2011). By simplifying cloning, gene delivery via conjugation or electroporation, and DNA isolation for genotyping, this hybrid method will facilitate future investigations and engineering of *Ms. trichosporium* OB3b.

2. MS. TRICHOSPORIUM OB3B GROWTH CONDITIONS

Ms. trichosporium OB3b strains are grown in nitrate minimal salt (NMS) media (Hakemian et al., 2008) (Table 1). Cultures are grown under either low copper ($0.1 \mu\text{M CuSO}_4$) or high copper ($10\text{--}25 \mu\text{M CuSO}_4$) conditions in liquid cultures and agar plates. Many genes of interest in methanotrophs are copper regulated, and it may be necessary to alter the copper concentration to ensure viability and to avoid selection against the genetic modifications. Similar considerations may be appropriate for genes regulated by other environmental factors.

Initially, cells at an OD_{600} of 0.1 are inoculated into 50 mL cultures in 250-mL Erlenmeyer flasks sealed with rubber septa. The flasks are then incubated at 30°C with shaking at 200 rpm, and oxygen and methane levels are maintained via daily sparging with a 1:3 methane-to-air ratio at 1 L/min for 5 min. An OD_{600} of ~ 1.5 is typically reached after 1 week of growth. Agar plates are incubated in GasPak plate incubation chambers (BD) at 30°C , and gas exchange is facilitated by use of a vacuum to create negative pressure inside the chamber followed by gas flow at a 1:1 methane-to-air ratio at 1 L/min for 3 min. Colonies on plates typically appear in 1–2 weeks. Antibiotics such as kanamycin ($25 \mu\text{g/mL}$) and gentamicin (5

µg/mL) are used to select for mutants, and naladixic acid (10 µg/mL) is used to remove donor *Escherichia coli* strains as described later. The bacterial strains used in this protocol are listed in Table 2.

3. PLASMIDS FOR MUTAGENESIS

For chromosomal site-directed mutagenesis or gene deletion, a broad host range vector pk18*mobsacB* (Schäfer et al., 1994) has been used in several methanotrophs, including *Ms. trichosporium* OB3b (Borodina et al., 2007). However, it is sometimes difficult to insert methanotroph genes, including those in the *pmo* operon, into this plasmid due to leaky expression at high copy number and subsequent toxicity of the product proteins in *E. coli* (Ojala et al., 2011). We have employed the following two strategies to circumvent this issue. First, we use the chemically competent *E. coli* strain CopyCutter EPI400 (Lucigen), designed to control plasmid copy number, to increase the success rate for cloning of methanotroph genes. However, the plasmid must then be incorporated into the *E. coli* donor strain S17-1 (Simon, Priefer, & Pühler, 1983) for subsequent conjugation. Transformation into S17-1 can also fail due to leaky expression. As a second, alternative strategy, we have constructed a modified version of pk18*mobsacB* under p15a copy control (Selzer, Som, Itoh, & Tomizawa, 1983) to generate the plasmid pk18*mobsacB*_p15a (Fig. 1; Table 3). This shuttle vector has improved the efficiency of cloning methanotroph genes, such as those in the *pmo* operon (Ojala et al., 2011).

Replicative plasmids are useful tools for protein overexpression or probing promoters. We tested a replicative plasmid compatible with Type I methanotrophs for function in *Ms. trichosporium* OB3b. The Lidstrom laboratory has constructed a replicative plasmid pAWP89 that can be maintained in *M. buryatense* 5G (Puri et al., 2015). We have determined that this plasmid can be maintained in *Ms. trichosporium* OB3b (Fig. 2). For the purposes of overexpressing toxic methanotroph genes, we have replaced the origin of replication in pAWP89 with p15a to create pAWP89_p15a (Fig. 1; Table 3).

4. GENETIC MANIPULATION VIA CONJUGATION

The most common and efficient method for genetic manipulation in methanotrophs requires conjugation (Lloyd et al., 1999; Murrell, 1994; Smith & Murrell, 2011). Typically after conjugation, the shuttle plasmid is incorporated into the chromosome of the recipient strain, followed by counterselection to ensure double recombination and removal of the vector backbone from the chromosome (Lalioi & Heath, 2001). In *Ms. trichosporium* OB3b, the established conjugation method involves the addition of liquid cultures of both methanotroph and *E. coli* cells onto a 0.2-µm sterile filter paper for mating (Smith & Murrell, 2011). This protocol requires the use and subsequent dismantling of a plastic filter unit or a Pyrex microfiltration glass assembly, followed by cell recovery from filter paper, which can increase the risk of contamination. A simpler conjugation protocol has been developed for the Type I methanotrophs *M. buryatense* 5G and *M. alcaliphilum* 20Z (Ojala et al., 2011; Puri et al., 2015). We have modified this protocol, which requires only cells grown on agar plates and disposable cell spreaders, for conjugation in *Ms. trichosporium* OB3b (Fig. 3):

1. The vector *pk18mobsacB* or *pk18mobsacB_p15a* (carrying an insert consisting of 5' and 3' DNA regions flanking the site of gene insertion, deletion, or mutagenesis, as well as any sequence to be inserted or altered) is transformed into the donor strain *E. coli* S17-1 via either heat shock or electroporation (Simon et al., 1983).
2. One day prior to bacterial mating, the *E. coli* S17-1 cells are streaked onto LB agar plates to obtain a bacterial lawn.
3. The wild-type recipient strain *Ms. trichosporium* OB3b is streaked onto an NMS agar medium plate (Table 1) and given 1 week to grow, forming at least a partial bacterial lawn.
4. The donor and recipient strains are harvested from plates and evenly plated and spread onto the same mating agar plate (Table 1) with a cell ratio of 2:1 donor-to-recipient strain. Specifically, a sterile cell spreader is dragged once across the plate containing the recipient methanotroph cells, and then is immediately used to inoculate the mating plate (Table 1). This is repeated for the donor (*E. coli*) strain, and the donor cells are plated directly on top of the recipient cells in the mating plate.
5. These plates are then incubated at a 1:1 methane-to-air ratio for 48 h at 30°C in a GasPak chamber.
6. A fraction of the mated cells, just enough to be visible on the bottom of the sterile cell spreader, are scraped off the mating plate and directly plated onto a selection agar plate (Table 1) containing kanamycin (25 µg/mL). The selection plate is incubated under the same conditions as the mating plate.
7. Single transconjugant colonies that appear are picked with a sterile loop and separately streaked onto selection plates containing kanamycin (25 µg/mL) as well as naladixic acid (10 µg/mL) to eliminate remaining cells from the donor *E. coli* strain.
8. Single transconjugant colonies on selection plates are picked using a sterile loop and streaked onto counterselection agar plates (Table 1) containing 2.5% sucrose and appropriate antibiotics. Single colonies from these counterselection plates are then picked and streaked once more to obtain sufficient biomass for genotype screening as described in Section 6.1.

5. GENE MANIPULATION VIA ELECTROPORATION

In situations where cloning or conjugation fails, electroporation can be employed, as reported for *Methylocella silvestris* BL2, *Methylocystis* sp. strain SC2, and *M. buryatense* 5G (Baani & Liesack, 2008; Crombie & Murrell, 2011; Yan et al., 2016). Linear DNA fragments produced via PCR can be also electroporated into *Ms. trichosporium* OB3b and integrated into the chromosome (Dassama et al., 2016). Thus far, intact plasmids purified from *E. coli* cloning strains have not been successfully incorporated into *Ms. trichosporium* OB3b via this method.

5.1 Construction of Linear DNA Fragment

Homologous recombination of a linear DNA fragment is a technique used for gene disruption, insertion, and modification (Orr-Weaver, Szostak, & Rothstein, 1981; Sharan, Thomason, Kuznetsov, & Court, 2009). One common use of this technique is gene disruption via insertion of a linear DNA fragment containing an antibiotic resistance cassette into the middle of the target gene. Construction of the resulting linear DNA fragment can be performed via several methods, including fusion PCR (Shevchuk et al., 2004). In that method, ~500 bp DNA regions flanking the gene of interest and a gentamicin resistance cassette are amplified by PCR. Primers are designed to add 25 bp regions onto the 5' and 3' ends of the gentamicin resistance cassette that overlap with the DNA regions flanking the gene of interest. The three PCR products are then joined together through two rounds of fusion PCR (Fig. 4). PCR products for each round of amplification are isolated and purified through gel extraction. In addition, it is necessary to perform a final cleanup of the PCR product using a PCR Purification Kit (Qiagen) or equivalent technique prior to sequencing and electroporation.

5.2 Electrocompetent Cell Preparation

50 mL cultures of *Ms. trichosporium* OB3b cells are grown in NMS medium (0.1 μ M CuSO₄) as described in Section 2. Cells are harvested at an OD₆₀₀ of 1.0, centrifuged at 7000 $\times$ g at 4°C for 10 min, and washed twice with 20 mL of 10% sterile glycerol at 4°C before a final 1 mL resuspension in 10% glycerol at 4°C. Fresh electrocompetent cells are recommended for every electroporation round.

5.3 Electroporation

1. 100–500 ng of the linear DNA fragment is added to 100 μ L electrocompetent cells in sterile microcentrifuge tubes kept on ice.
2. The mixture is then transferred into a prechilled sterile electroporation cuvette with a 0.2-cm gap width (Bio-Rad). The cuvette is inserted into an electroporation system (Bio-Rad MicroPulser) and subjected to one pulse at 2.2 kV for 5 ms at 1 Å.
3. After electroporation, the cell mixture is added to 5 mL of NMS liquid medium (without antibiotics) in borosilicate culture tubes sealed with rubber septa.
4. The cultures are incubated overnight at 30°C and 200 rpm after gas sparging at a 1:3 methane-to-air ratio.
5. After 24 h of growth, the cells (the outgrowth culture) are centrifuged at 7000 $\times$ g and 4°C for 10 min, and are then resuspended in 500 μ L standard NMS liquid medium.
6. 100–300 μ L of the outgrowth culture is then spread onto NMS selection plates containing gentamicin (5 μ g/mL).

6. ANALYSIS OF *MS. TRICHOSPORIUM* OB3B MUTANTS

6.1 DNA Extraction From *Ms. trichosporium* OB3b Cells on Agar Plates

DNA extraction from *Ms. trichosporium* OB3b is less efficient than DNA extraction from many Type I or Type II methanotrophic bacteria. Existing methods use the neutral lysis/CsCl method or a DNeasy Blood Tissue Kit (Qiagen) for DNA extractions from liquid cultures (Gu et al., 2016; Smith & Murrell, 2011). However, growing liquid cultures to genotype multiple colonies is time-consuming. Instead, an improved method has been developed to extract DNA from cells growing on agar plates using the MasterPure Complete DNA Purification Kit (Epicentre, Cat. No. MC85200). This method allows for higher-throughput of genotype screening and provides increased yields of DNA from smaller amounts of cells. The MasterPure Complete DNA Purification Kit protocol has been modified for *Ms. trichosporium* OB3b:

1. Add 20 μ L Proteinase K (NEB, 800 U/mL) into 300 μ L $2 \times$ tissue and cell lysis solution for each sample.
2. Drag a sterile inoculating loop across the agar plate culture until the loop is covered with cells (Fig. 5).
3. Swirl the inoculating loop in the lysis solution and resuspend fully by pipetting.
4. Incubate the sample at 65°C overnight at 500 rpm.
5. Add 5 μ L RNase A (Epicentre, 5 μ g/ μ L). Incubate for 1 h at 37°C without shaking.
6. Place the samples on ice for 5 min.
7. Add 150 μ L MPC protein precipitation reagent and vortex.
8. Centrifuge the sample at 10,000 $\times g$ at 4°C for 10 min to pellet cell debris.
9. Transfer the supernatant to a new tube and add 500 μ L 100% isopropanol on ice. Manually invert the sample 30 times.
10. Centrifuge the sample at 4°C for 30 min at 20,000 $\times g$ to pellet DNA.
11. Wash the DNA pellet with 750 μ L 70% ethanol on ice. Centrifuge at 20,000 $\times g$ at 4°C for 5 min, remove the ethanol, and repeat the wash step once more.
12. Let the pellet dry at 37°C until residual ethanol is gone.
13. Resuspend the pellet with 50 μ L TE buffer and incubate overnight at 4°C.
14. DNA yield can vary from 10 to 60 μ g. Additional cleanup steps using silica spin columns (Zymo) may be required for uses other than simple Sanger sequencing.

6.2 Genotyping Mutants

To confirm the genotype of *Ms. trichosporium* OB3b variants, PCR amplification of the altered DNA region is performed. Due to the high genomic GC content of Type II methanotrophs, PCR amplification is more successful under reaction conditions optimized for high-GC samples, such as the $2 \times$ iProof GC Master Mix (Bio-Rad). 100 ng of *Ms.*

trichosporium OB3b genomic DNA is an appropriate starting point for a 50- μ L PCR reaction. Gradient or touchdown PCR at annealing temperatures of 55–65°C can be employed to identify conditions for specific amplification. PCR cleanup (QIAquick PCR Purification Kit, Qiagen) will significantly improve sequencing results.

7. SUMMARY

The modifications in plasmid design, construct delivery, and genotyping described here allow for simpler and more efficient genetic manipulation of *Ms. trichosporium* OB3b, at least until CRISPR or other gene editing methods are developed (Tian et al., 2017). These advances will expedite metabolic engineering efforts in this methanotrophic strain and enhance studies of proteins involved in the key processes of methanotrophy.

Acknowledgments

Methanotroph work in the Rosenzweig laboratory is supported by NIH Grant GM118035 and DOE Grant DE-SC0016284. We thank Dr. Grace Kenney for sharing her expertise on *Ms. trichosporium* OB3b.

References

- Baani M, Liesack W. Two isozymes of particulate methane monooxygenase with different methane oxidation kinetics are found in *Methylocystis* sp. strain SC2. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105:10203–10208. [PubMed: 18632585]
- Borodina E, Nichol T, Dumont MG, Smith TJ, Murrell JC. Mutagenesis of the ‘leucine gate’ to explore the basis of catalytic versatility in soluble methane monooxygenase. *Applied and Environmental Microbiology*. 2007; 73:6460–6467. [PubMed: 17704278]
- Chu F, Lidstrom ME. XoxF acts as the predominant methanol dehydrogenase in the Type I methanotroph *Methylobacterium buryatense*. *Journal of Bacteriology*. 2016; 198:1317–1325. [PubMed: 26858104]
- Crombie, A., Murrell, JC. Development of a system for genetic manipulation of the facultative methanotroph *Methylocella silvestris* BL2. In: Rosenzweig, AC., Ragsdale, SW., editors. *Methods in enzymology*: Vol. 495. *Methods in methane metabolism, part B: Methanotrophy*. Amsterdam: Elsevier; 2011. p. 119–133.
- Dassama LMK, Kenney GE, Ro SY, Zielazinski EL, Rosenzweig AC. Methanobactin transport machinery. *Proceedings of the National Academy of Sciences of the United States of America*. 2016; 113:13027–13032. [PubMed: 27807137]
- Dassama LMK, Kenney GE, Rosenzweig AC. Methanobactins: From genome to function. *Metallomics*. 2017; 9:7–20. [PubMed: 27905614]
- de la Torre A, Metivier A, Chu F, Laurens LML, Beck DAC, Pienkos PT, et al. Genome-scale metabolic reconstructions and theoretical investigation of methane conversion in *Methylobacterium buryatense* strain 5G(B1). *Microbial Cell Factories*. 2015; 14:188. [PubMed: 26607880]
- Demidenko A, Akberdin IR, Allemann M, Allen EE, Kalyuzhnaya MG. Fatty acid biosynthesis pathways in *Methylobacterium buryatense* 5G(B1). *Frontiers in Microbiology*. 2017; 7:2167. [PubMed: 28119683]
- Gu W, Baral BS, DiSpirito AA, Semrau JD. An aminotransferase is responsible for the deamination of the N-terminal leucine and required for formation of oxazolone ring A in methanobactin of *Methylosinus trichosporium* OB3b. *Applied and Environmental Microbiology*. 2017; 83:e02616–16.
- Gu W, Haque MFU, Baral BS, Turpin EA, Bandow NL, Kremmer E, et al. A TonB-dependent transporter is responsible for methanobactin uptake by *Methylosinus trichosporium* OB3b. *Applied and Environmental Microbiology*. 2016; 82:1917–1923. [PubMed: 26773085]

- Gu W, Haque MFU, Semrau JD. Characterization of the role of copCD in copper uptake and the 'copper-switch' in *Methylosinus trichosporium* OB3b. FEMS Microbiology Letters. 2017; 364 <https://doi.org/10.1093/femsle/fnx094>.
- Hakemian AS, Kondapalli KC, Telser J, Hoffman BM, Stemmler TL, Rosenzweig AC. The metal centers of particulate methane monooxygenase from *Methylosinus trichosporium* OB3b. Biochemistry. 2008; 47:6793–6801. [PubMed: 18540635]
- Hanson RS, Hanson TE. Methanotrophic bacteria. Microbiological Reviews. 1996; 60:439–471. [PubMed: 8801441]
- Haque MFU, Gu W, DiSpirito AA, Semrau JD. Marker exchange mutagenesis of mxaF, encoding the large subunit of the Mxa methanol dehydrogenase, in *Methylosinus trichosporium* OB3b. Applied and Environmental Microbiology. 2015; 82:1549–1555. [PubMed: 26712545]
- Haque MFU, Kalidass B, Bandow N, Turpin EA, DiSpirito AA, Semrau JD. Cerium regulates expression of alternative methanol dehydrogenases in *Methylosinus trichosporium* OB3b. Applied and Environmental Microbiology. 2015; 81:7546–7552. [PubMed: 26296730]
- Henard CA, Smith H, Dowe N, Kalyuzhnaya MG, Pienkos PT, Guarnieri MT. Bioconversion of methane to lactate by an obligate methanotrophic bacterium. Scientific Reports. 2016; 6:21585. [PubMed: 26902345]
- Kalyuzhnaya MG, Khmelenina V, Eshinimaev B, Sorokin D, Fuse H, Lidstrom M, et al. Classification of halo(alkali)philic and halo(alkali)tolerant methanotrophs provisionally assigned to the genera *Methylomicrobium* and *Methylobacter* and emended description of the genus *Methylomicrobium*. International Journal of Systematic and Evolutionary Microbiology. 2008; 58:591–596. [PubMed: 18319461]
- Kalyuzhnaya MG, Puri AW, Lidstrom ME. Metabolic engineering in methanotrophic bacteria. Metabolic Engineering. 2015; 29:142–152. [PubMed: 25825038]
- Lalio M, Heath J. A new method for generating point mutations in bacterial artificial chromosomes by homologous recombination in *Escherichia coli*. Nucleic Acids Research. 2001; 29:e14. [PubMed: 11160916]
- Lawton TJ, Rosenzweig AC. Methane-oxidizing enzymes: An upstream problem in biological gas-to-liquids conversion. Journal of the American Chemical Society. 2016; 138:9327–9340. [PubMed: 27366961]
- Lloyd JS, Finch R, Dalton H, Murrell JC. Homologous expression of soluble methane monooxygenase genes in *Methylosinus trichosporium* OB3b. Microbiology. 1999; 145:461–470. [PubMed: 10075428]
- Lock M, Nichol T, Murrell JC, Smith TJ. Mutagenesis and expression of methane monooxygenase to alter regioselectivity with aromatic substrates. FEMS Microbiology Letters. 2017; 364 <https://doi.org/10.1093/femsle/fnx137>.
- Murrell JC. Molecular genetics of methane oxidation. Biodegradation. 1994; 5:145–159. [PubMed: 7765830]
- Mustakhimov II, Reshetnikov AS, Glukhov AS, Khmelenina VN, Kalyuzhnaya MG, Trotsenko YA. Identification and characterization of EctR1, a new transcriptional regulator of the ectoine biosynthesis genes in the halotolerant methanotroph *Methylomicrobium alcaliphilum* 20Z. Journal of Bacteriology. 2009; 192:410–417. [PubMed: 19897647]
- Ojala, DS., Beck, DAC., Kalyuzhnaya, MG. Genetic systems for moderately halo(alkali)philic bacteria of the genus *Methylomicrobium*. In: Rosenzweig, AC., Ragsdale, SW., editors. Methods in enzymology: Vol. 495. Methods in methane metabolism, part B: Methanotrophy. Amsterdam: Elsevier; 2011. p. 99–118.
- Orr-Weaver TL, Szostak JW, Rothstein RJ. Yeast transformation: A model system for the study of recombination. Proceedings of the National Academy of Sciences of the United States of America. 1981; 78:6354–6358. [PubMed: 6273866]
- Puri AW, Owen S, Chu F, Chavkin T, Beck DAC, Kalyuzhnaya MG, et al. Genetic tools for the industrially promising methanotroph *Methylomicrobium buryatense*. Applied and Environmental Microbiology. 2015; 81:1775–1781. [PubMed: 25548049]
- Ross MO, Rosenzweig AC. A tale of two methane monooxygenases. Journal of Biological Inorganic Chemistry. 2016; 22:307–319. [PubMed: 27878395]

- Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: Selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene*. 1994; 145:69–73. [PubMed: 8045426]
- Selzer G, Som T, Itoh T, Tomizawa J. The origin of replication of plasmid p15A and comparative studies on the nucleotide sequences around the origin of related plasmids. *Cell*. 1983; 32:119–129. [PubMed: 6186390]
- Semrau JD, DiSpirito AA, Yoon S. Methanotrophs and copper. *FEMS Microbiology Reviews*. 2010; 34:496–531. [PubMed: 20236329]
- Semrau JD, Jagadevan S, DiSpirito AA, Khalifa A, Scanlan J, Bergman BH, et al. Methanobactin and MmoD work in concert to act as the ‘copper-switch’ in methanotrophs. *Environmental Microbiology*. 2013; 15:3077–3086. [PubMed: 23682956]
- Sharan SK, Thomason LC, Kuznetsov SG, Court DL. Recombineering: A homologous recombination-based method of genetic engineering. *Nature Protocols*. 2009; 4:206–223. [PubMed: 19180090]
- Shevchuk NA, Bryksin AV, Nusinovich YA, Cabello FC, Sutherland M, Ladisch S. Construction of long DNA molecules using long PCR-based fusion of several fragments simultaneously. *Nucleic Acids Research*. 2004; 32:e19. [PubMed: 14739232]
- Simon R, Priefer U, Pühler A. A broad host range mobilization system for in vivo genetic engineering: Transposon mutagenesis in gram negative bacteria. *Nature Biotechnology*. 1983; 1:784–791.
- Smith, TJ., Murrell, JC. Mutagenesis of soluble methane monooxygenase. In: Rosenzweig, AC., Ragsdale, SW., editors. *Methods in enzymology: Vol. 495. Methods in methane metabolism, part B: Methanotrophy*. Amsterdam: Elsevier; 2011. p. 135-147.
- Smith TJ, Slade SE, Burton NP, Murrell JC, Dalton H. Improved system for protein engineering of the hydroxylase component of soluble methane monooxygenase. *Applied and Environmental Microbiology*. 2002; 68:5265–5273. [PubMed: 12406713]
- Stein LY, Yoon S, Semrau JD, DiSpirito AA, Crombie A, Murrell JC, et al. Genome sequence of the obligate methanotroph *Methylosinus trichosporium* strain OB3b. *Journal of Bacteriology*. 2010; 192:6497–6498. [PubMed: 20952571]
- Tian P, Wang J, Shen X, Rey JF, Yuan Q, Yan Y. Fundamental CRISPR-Cas9 tools and current applications in microbial systems. *Synthetic and Systems Biotechnology*. 2017; 2:219–225. [PubMed: 29318202]
- Vorobev AV, Baani M, Doronina NV, Brady AL, Liesack W, Dunfield PF, et al. *Methyloferula stellata* gen. nov., sp. nov., an acidophilic, obligately methanotrophic bacterium that possesses only a soluble methane monooxygenase. *International Journal of Systematic and Evolutionary Microbiology*. 2011; 61:2456–2463. [PubMed: 21097638]
- Yan X, Chu F, Puri AW, Fu Y, Lidstrom ME. Electroporation-based genetic manipulation in Type I methanotrophs. *Applied and Environmental Microbiology*. 2016; 82:2062–2069. [PubMed: 26801578]

Fig. 1.
Plasmid maps of *pk18mobsacB_p15a* and *pAWP89_p15a*.

Fig. 2.
Ms. trichosporium OB3b transformed with pAWP89_p15a (A) compared to wild-type *Ms. trichosporium* OB3b (B) on 25 µg/mL kanamycin NMS agar selection plates.

Fig. 3.

Gene deletion via conjugation in *Ms. trichosporium* OB3b. Following conjugation, the introduced plasmid recombines into the targeted site in the chromosome during the first crossover. Transconjugant selection followed by sucrose counterselection induces a second crossover event to remove the plasmid backbone from the chromosome and produce the mutant with the gentamicin resistance cassette (*gen^R*) disrupting *gene A*.

Fig. 4.

Construction of a gene-disrupting linear DNA fragment using fusion PCR. (1) PCR amplification of *gen^R* and the 5' and 3' DNA regions flanking gene targeted for disruption. (2) The 5' DNA fragment and *gen^R* are used as templates for the first round of fusion PCR. (3) The resulting PCR product and the 3' DNA fragment are used as templates for the second round of PCR to produce the final gene-disrupting linear DNA construct. The linear DNA construct is (4) electroporated into wild-type *Ms. trichosporium* OB3b followed by (5) selection for the *Ms. trichosporium* OB3b *gene A* mutant. Primer binding sites are represented by *half arrows*.

Fig. 5.
Amount of *Ms. trichosporium* OB3b cells required for DNA extraction shown on a 1- μ L inoculating loop.

Table 1*Ms. trichosporium* OB3b Growth Conditions

	Growth	Mating	Selection	Counterselection
<i>NMS medium (1 L)</i>				
NaNO ₃	0.85 g	0.85 g	0.85 g	0.85 g
K ₂ SO ₄	0.17 g	0.17 g	0.17 g	0.17 g
MgSO ₄ ·7H ₂ O	0.037 g	0.037 g	0.037 g	0.037 g
CaCl ₂ ·2H ₂ O	0.01 g	0.01 g	0.01 g	0.01 g
40 mM FeSO ₄ ·7H ₂ O	1 mL	1 mL	1 mL	1 mL
100 mM CuSO ₄ ·5H ₂ O	Varies	Varies	Varies	Varies
Agar (plates only)	15 g	15 g	15 g	15 g
LB (Difco)	—	1.2 g	—	—
Phosphate buffer solution	10 mL	10 mL	10 mL	10 mL
Trace elements solution	2 mL	2 mL	2 mL	2 mL
50% sucrose	—	—	—	50 mL
<i>Phosphate solution (g/L)</i>				
Na ₂ HPO ₄ ·7H ₂ O	48.06			
KH ₂ PO ₄	23.4			
<i>Trace elements (g/L)</i>				
ZnSO ₄ ·7H ₂ O, 0.288; MnCl ₂ ·4H ₂ O, 0.16; H ₃ BO ₃ , 0.06; Na ₂ MoO ₄ ·2H ₂ O, 0.048; CoCl ₂ ·6H ₂ O, 0.048; KI, 0.083				

Table 2**Bacterial Strains**

Strain	Growth Medium
<i>Ms. trichosporium</i> OB3b	NMS
<i>E. coli</i> S17-1	LB (Difco)
<i>E. coli</i> CopyCutter EPI400	LB (Difco)

Table 3

Plasmids

Plasmid Name	Source/Reference
pk18 <i>mobsacB</i>	ATCC 87097 (Schäfer et al., 1994)
pk18 <i>mobsacB</i> _p15a	This study
pAWP89	Addgene 61264 (Puri et al., 2015)
pAWP89_p15a	This study