Published in final edited form as:

Neuropsychologia. 2008 March 07; 46(4): 1060–1068. doi:10.1016/j.neuropsychologia.2007.11.004.

Movement interference in autism spectrum disorder

E Gowen¹, J Stanley², and R.C Miall³

¹Faculty of Life Sciences, University of Manchester, UK ²School of Medicine, University of Otago, Wellington, New Zealand ³School of Psychology, University of Birmingham, UK

Abstract

Movement interference occurs when concurrently observing and executing incompatible actions and is believed to be due to co-activation of conflicting populations of mirror neurons. It has also been suggested that mirror neurons contribute towards the imitation of observed actions. However, the exact neural substrate of imitation may depend on task demands: a processing route for goal-directed meaningful actions may be distinct from one for non-goal directed actions. A more controversial role proposed for these neurons is in theory of mind processing, along with the subsequent suggestion that impairment in the mirror neuron circuit can contribute to autism spectrum disorder (ASD) where individuals have theory of mind deficits. We have therefore examined movement interference in 9 ASD participants and 9 matched controls while performing actions congruent and incongruent with observed meaningless arm movements. We hypothesised that if the mirror neuron system was impaired, reduced interference should be observed in the ASD group. However, control and ASD participants demonstrated an equivalent interference effect in an interpersonal condition, with greater movement variability in the incongruent compared to the congruent condition. A component of movement interference which is independent of congruency did differ between groups: ASD participants made generally more variable movements for the interpersonal task than for biological dot motion task, while the reverse was true for the control participants. We interpret these results as evidence that the ASD participant group either rely to a greater extent on the goal directed imitation pathway, supporting claims that they have a specific deficit of the non-goal directed imitation pathway or exhibit reduced visuomotor integration.

Keywords

Imitation; autism; mirror neurons; facilitation

Introduction

The role of the mirror neuron system in action observation and imitation

In recent years it has become well established that observation of human movement influences the observer's own motor system. For example, concurrent observation and execution of a compatible action (e.g. lifting the index finger while observing index finger elevation) results in a facilitation in reaction time, whereas during incompatible observation/action combinations (e.g., lifting the index finger while observing finger depression) interference in movement initiation is indicated by increased reaction times (Brass, Bekkering, Wohlschlaeger and Prinz, 2000; Brass, Bekkering and Prinz, 2001; Craighero, Bello, Fadiga and Rizzolatti, 2002; Vogt, Taylor and Hopkins, 2003; see Vogt and Thomaschke, 2007 for a recent review of the perception-action system). The neural basis of such effects has been clarified with the discovery of mirror neurons, present in the pars opercularis of the ventral premotor cortex (BA 44) and inferior parietal lobe (Buccino *et al* 2001; Grafton, Arbib, Fadiga and Rizzolatti, 1996; Grezes, Armony, Rowe and Passingham, 2003; Iacoboni *et al.*, 1999; Rizzolatti, Fadiga, Gallese and Fogassi, 1996). Mirror neurons discharge during both observation and execution of an action, leading to the theory that facilitation and interference effects may result from co-activation of compatible or incompatible sets of mirror neurons (Blakemore and Frith, 2005).

Owing to the combined observation/execution properties of mirror neurons, Iacoboni and others have described them as core components of an imitation network (Iacoboni *et al.*, 1999; Iacoboni, 2005; Iacoboni and Dapretto, 2006; Rizzolatti, Fogassi and Gallese, 2001). However, recent evidence from neuroimaging and neuropsychological studies indicates that there are at least two different neural pathways underlying imitation, dependent on the semantic content of the action to be imitated. One is a goal directed, transitive (meaningful) route, the other a non-goal directed, intransitive (meaningless) route. There is some disagreement at present as to the exact neural areas involved in each route. One hypothesis of imitation proposes that transitive actions are associated with the inferior temporal gyrus and hippocampus and that intransitive actions have greater association with visuospatial areas such as the superior temporal lobe, parieto-occipital and occipitotemporal junctions (Rumiati *et al* 2005; Tessari, Canessa, Ukmar and Rumiati, 2007). Focussing particularly on the mirror neuron system, Molnar-Szakacs, Iacoboni, Koski and Mazziotta (2005) proposed a functional and anatomical split within the pars opercularis of the inferior frontal gyrus whereby the ventral region processes details relating to action goals and the dorsal region guides detailed observation-execution matching in conjunction with the superior temporal sulcus (STS) and inferior parietal regions. Similarly, activation of ventral premotor mirror neurons has also been associated to a greater extent with goal directed actions, whereas intransitive actions preferentially activated superior and inferior parietal lobes (Decety *et al.*, 1997; Grezes, Costes and Decety, 1998; Koski *et al.*, 2002). However, as an alternative to this hypothesis, several authors have suggested a model whereby goals are represented in the intraparietal sulcus, whereas the inferior frontal gyrus is more important in encoding the specific properties of an action (Hamilton, in press; Hamilton and Grafton, 1996; Rizzolatti and Craighero, 2004).

The role of the mirror neuron system in theory of mind

Several authors have recently suggested that mirror neurons may contribute towards action understanding, action prediction and to Theory of Mind (ToM) processes by mentally reproducing or “simulating” the goals and outcomes of others’ actions (Frith and Frith, 1999; Gallese and Goldman, 1998; Hamilton and Grafton, 2006; Iacoboni *et al* 1999; 2005; Jeannerod, 2001; Rizzolatti, Fogassi and Gallese, 2001). ToM is the ability to understand other people in terms of their intentions, beliefs and desires. Consequently, deficits in the mirror neuron system have been proposed to underlie some of the behavioural characteristics present in autistic spectrum disorder (ASD) where affected individuals fail to acquire a ToM. A number of neuroimaging studies have reported altered mirror neuron activity in ASD (Bernier, Dawson, Webb and Murias, 2007; Dapretto *et al* 2006; Hadjikhani, Joseph, Snyder and Tager-Flusberg, 2006; Oberman *et al.*, 2005; Nishitani, Avikainen and Hari, 2004; Williams *et al.*, 2006). These data support the hypothesis that many of the social impairments present in ASD may be attributed to an underlying mirror neuron system deficit.

Imitation and ASD

If mirror neuron activity is abnormal in those with ASD, one would expect to see altered movement facilitation and interference during observation of actions, as well as other effects such as impaired action understanding, prediction and imitation. However, evidence on this topic is contradictory which may be a consequence of task demands such as the effector used (face vs. hand), or the transitive versus intransitive nature of the imitation. For example, Théoret *et al* (2005) reported reduced motor cortical modulation using transcranial magnetic stimulation (TMS) during observation of intransitive finger movements in ASD participants. Imitation tasks that use meaningless gestures evoke impaired performance in those with ASD (McIntosh, Reichmann-Decker, Winkelman and Wilbarger, 2006; Williams, Whiten and Singh, 2004), whereas those that use goal directed actions demonstrate intact ability (Hamilton, Brindley and Frith, 2007; Hobson and Lee, 1999; McIntosh *et al* 2006; Williams, Whiten and Singh, 2004). In addition, recent work indicates that adults with high functioning autism or Asperger syndrome are able to represent the actions of others’ (Sebanz *et al.*, 2005). Consequently, the contribution of mirror neuron dysfunction to behavioural characteristics in ASD is likely to be more complex than first suggested. On balance, one possible interpretation of these studies is that during imitation tasks, individuals with ASD have differential impairment of those circuits involved in goal-less directed imitation. A model to this effect has been put forward by Hamilton (in press), suggesting that individuals with ASD have an intact goal directed pathway mediated by occipital-parietal connections but a deficient pathway for the mimicry of meaningless actions mediated by occipital-frontal connections.

In light of this conflicting literature and limited behavioural data concerning facilitation and interference in ASD, the current work examined movement interference in a group of ASD individuals using a previously reported paradigm (Kilner, Paulignan and Blakemore, 2003). In this task, participants make either horizontal or vertical intransitive and continuous arm movements in time with the movements of an experimenter so that the two peoples’ movements are either congruent (i.e. both moving in the same plane) or incongruent (i.e.

participant moving their arm in plane perpendicular to that of experimenter). Finger tip movement variability (as measured in the orthogonal plane) is greater in the orthogonal plane for incongruent than for congruent conditions (Kilner, Paulignan and Blakemore, 2003; Stanley, Gowen and Miall, 2007) and is termed the “interference effect”. We predict that if the mirror neuron system was abnormal in ASD, particularly for the processing of observed meaningless actions, then the ASD participant group would demonstrate a smaller or null interference effect compared to a matched control group.

We have recently extended the interpersonal interference task by examining interference during presentation of a small dot projected onto a screen that could have either a biological (pre-recorded human arm movement) or non-biological (computer generated, constant-velocity sinusoid) motion profile (Stanley, Gowen and Miall, 2007). Our data demonstrated that the interference effect was present if participants were instructed that both dot motion stimuli were human generated and also uncovered a component of movement variability that was independent of congruency and agency instruction, but was driven by the higher variability of the biological as opposed to non-biological dot-motion profiles (Stanley, Gowen and Miall, 2007). We attributed this to a bottom-up effect of stimulus content on the ongoing arm movement. This congruency-independent or “bottom up” interference effect may be of particular relevance for the intransitive route of action processing that is concerned with the specifics of movement detail. This is supported by data indicating that during imitation of goal-less actions more emphasis appears to be placed on effector selection and movement execution, whereas goal-directed actions are imitated correctly in respect to the goal while details regarding the movement may be ignored (Bekkering, Wohlschläger and Gattis 2000; Franz, Ford and Werner, 2007). Moreover, adults performing more difficult imitation tasks frequently imitate the goal of an action rather than perform an exact replication (Tessari and Rumiati, 2004; Wohlschläger, Gattis and Bekkering, 2003). Consequently, the effect of stimulus variability may provide an indication of the degree of visuomotor coupling between the observed and executed movements. If ASD individuals rely more on abstract, goal directed simulation, and are less able to simulate goal-less intransitive actions, then their arm trajectories should be less affected by the bottom-up increased variability in the biological dot-motion profile. Therefore, in the current study we explored the effect of stimulus presentation on our ASD participants, using these biological and not biological dot stimuli, as well as the interpersonal task.

In summary, based on the assumption that the mirror neuron system is dysfunctional in ASD, we predict less congruency-dependent motor interference (“incongruent interference”) in that group than in controls, as well as less congruency-independent (“bottom-up”) influence of the stimulus

Methods

Participants

Participants were 12 ASD individuals (6 males) and 12 sex, age and IQ matched healthy controls. The data of three male participants from the ASD group was excluded due one being left-handed, one being unable to maintain the correct pace of arm movement and one exhibiting consistently greater movement variability across all tasks ($> 2SD$).

Neuropsychological assessment of these latter two participants using the tasks documented below did not differ from the rest of the group. ASD participants had been given a diagnosis by outside clinical assessment (DSM IV criteria, American Psychiatric Association, 1994 or ICD-10, WHO 1992) of either high functioning autism (n=2) or Aspergers' syndrome (n=7). None of the participants reported a diagnosis of developmental coordination disorder. Average age ($\pm$ standard deviation) of the 9 ASD and control participants was 33.9 ± 13.2 and 32.0 ± 11.8 respectively. Average IQ ($\pm$ standard deviation) measured using the abbreviated WAIS was 117.6 ± 20.5 and 115.7 ± 14.1 for the ASD and control participants respectively. Age ($t = 0.22$ $p = 0.83$) and IQ ($t = 0.83$; $p = 0.96$) did not significantly differ between the two groups. All participants demonstrated binocular Snellens visual acuity of better than 6/9. Each participant gave written informed consent to participate and the study was approved by a local ethical committee, being performed in accordance with ethical standards laid down in the 1964 Declaration of Helsinki.

Assessment of ToM

To measure their mentalizing ability, all subjects performed false belief tasks, the results of which are shown in Table 1 in order of increasing ability. Four of these (Sally-Ann task (Wimmer and Perner, 1983), smarties, coat/ice cream story, (Perner and Wimmer, 1985)) were scored as pass or fail. Responses to a penny hiding task (Baron-Cohen, 1992) were given a % score out of 6 repetitions and responses to Happe stories (Happe, 1994) were given a % score out of a total of 16. Each ASD participant was given a total score across these ToM tasks (Table 1). Mean autism-spectrum quotient (AQ) scores were significantly higher for the ASD group compared to the control group (ASD: 36.7 ± 6 , minimum 34; Control: 11.6 ± 5 , maximum 18; $t = 9.8$; $p < 0.001$). It has been suggested that a score of 32+ distinguishes individuals who have clinically significant levels of autistic traits (Baron-Cohen, Wheelwright, Skinner, Martin and Clubely, 2001).

Movement Recording and Signal Processing

Fingertip position was recorded using the Optotrak 3020 active marker system (Northern Digital Instruments, Inc.). An infrared sensor was attached to a plastic thimble on the participant's right hand index finger, and sensor position recorded at 250 Hz, with 0.01 mm spatial resolution; only data from the horizontal x-plane and the vertical z-plane were used in the analysis. The plane of instructed movement (e.g., the x-plane during a horizontal movement, and the z-plane in a vertical movement) is referred to hereafter as the "dominant" plane; the orthogonal plane is referred to as the "error" plane. The experimenter's fingertip position was also recorded for those trials where the participant moved in time with the experimenter, but is not reported here.

Following data acquisition, fingertip position data were filtered with a 20 Hz Butterworth filter (all analysis conducted in Matlab, Mathworks Inc.) prior to movement scoring. Each trial's data were split into single movement segments (e.g., from extreme left to extreme right was one segment, and returning from right to left another segment). Endpoints for each segment were detected by finding the datapoints where velocity in the dominant movement plane crossed zero.

To quantify interference, the standard deviation of fingertip position (mm) was calculated within the error plane for each movement segment. The mean of these deviation scores was then calculated across all movement segments to give a single average of error-plane deviation for each trial, for each participant. As noted by Stanley *et al.* (2007), using SD of error plane variations appears preferable to using variance (as in the original Kilner, Paulignan and Blakemore (2003) study) as the non-linear nature of the variance measure produces group/condition means that are inappropriately influenced by outlying data, with consequences for subsequent analysis.

Stimuli

The visual stimulus was a 1-cm diameter white dot, presented on a black background projected onto a white screen 1.9 m from the participant. The centre of the projector screen was at a height of 1.55 m; the projector refresh rate was 60 Hz. The metronome for self-pacing movements was a sequence of tones presented at 1 Hz over headphones.

The biological dot-motion stimulus was pre-recorded data of the experimenter moving his arm in time with the audio metronome. The presented trajectories therefore included variation in the plane orthogonal to the dominant movement direction (e.g., there were fluctuations in vertical position during the presented horizontal movements). Data were scaled so that 50 cm in the recorded data corresponded to 50 cm on the projection screen. Movement frequency and the amplitude of the individual movement segments varied naturally over the course of the recording (also true for trials where the participant observed the experimenter moving). The non-biological dot-motion stimulus was of a constant speed (50 cm/s) with fixed amplitude of 50 cm on the projection screen, and a fixed frequency of 0.5 Hz. The dot instantly changed direction when reaching either end of its range. For this stimulus, position orthogonal to the main plane of movement direction was invariable. The difference in the motion between the two dot trajectories was clearly visible to the participants. Both the biological and non-biological dot-motion stimuli were identical to those used by Stanley, Gowen and Miall (2007) where greater detail of the velocity profiles can be found.

Procedure

Testing dot-motion stimuli—Participants were instructed to perform movements of about 50 cm amplitude. Initially, each participant performed two practice trials (30s duration) in time with the auditory metronome – one block of horizontal arm movements, followed by one block of vertical arm movements, while fixating a stationary circular target. Movements were timed so that the endpoints of each movement coincided with the tone.

Participants then performed the dot-motion trials that were blocked by motion profile of the observed dot stimulus (biological or non-biological), with the order of presentation counterbalanced across participants. Participants were informed that both dot stimuli were human generated (i.e. pre-recorded arm movements) and were asked to move in phase with the dot movement. At the start of each trial, the participant was instructed as to the plane in which to move his or her arm, as well as the dot-motion plane. Recordings began 2 to 3 seconds after the participant was moving in phase with the displayed dot stimulus, and lasted

for 30 seconds. Four trials, consisting of the different combinations of congruent and incongruent arm/dot-motion conditions for the two movement planes were conducted for each dot-motion type, with order again counterbalanced across participants.

Testing interpersonal interference—Trial structure was identical to the dot-motion task, except that participants moved in phase with a gender-matched experimenter, who stood 190 cm away (toe-to-toe distance). The experimenter performed arm movements in time with the auditory metronome with eyes closed. The interpersonal task was always conducted last to preclude the possibility of carry over effects contaminating the dot-motion trial data (see discussion of this issue in Stanley, Gowen and Miall, 2007).

Data Analysis—Data for fingertip position standard deviation in the error plane were analyzed across both dot-motion and interpersonal movement task using a $3 \times 2 \times 2 \times 2$ ANOVA design with factors of task (biological dot-motion, non-biological dot-motion, interpersonal), direction of participant's performed movement (horizontal, vertical), congruency of observed action, and subject group (ASD, control). For significant main effects and two-way interactions, t-tests were performed between the appropriate levels of the factors involved, using a Bonferroni α -correction for multiple tests. Correlations between interference effects and ASD measures (ToM and ASQ scores) were conducted using Pearson's correlation.

Results

The results can be seen in Figs 1a-b and 2a-b, for the biological, non-biological and interpersonal conditions. From the omnibus ANOVA, main effects of congruency ($F(1,16) = 18.90, p < 0.001$) and task ($F(2,32) = 7.78, p = 0.002$) were observed, along with interactions for task $\times$ congruency ($F(2,32) = 7.58, p = 0.02$) and task $\times$ subject group ($F(2,32) = 3.64, p = 0.001$). Importantly, there was no overall significant difference between the two participant groups ($F(1,16) = 0.015, p = 0.9$) and no interactions between group and congruency ($F(1,16) = 0.294, p = 0.595$). No other interactions were significant ($F(1,16) = 1.99, p = 0.17$).

Main effects of congruency and task

As expected, the main effect of congruency signified greater interference for the incongruent (mean $\pm$ SE = 11.53 ± 0.49) compared to congruent trials (mean $\pm$ SE = 8.23 ± 0.65) (Figs 1a-b). Concerning the main effect of task (on average over congruency, direction and group), error plane deviation in the biological dot profile (mean $\pm$ SE = 10.83 ± 0.71) and interpersonal task (mean $\pm$ SE = 10.4 ± 0.45) was significantly greater than in the non-biological dot profile task (mean $\pm$ SE = 8.45 ± 0.5) ($t(17) = 3.41; p = 0.003; t(17) = 3.1; p = 0.007$ respectively; Bonferroni corrected critical $\alpha = .0167$, for three tests). Collapsed across both groups, error plane deviation did not vary between the biological dot-motion and interpersonal tasks ($t(17) = 0.65; p = 0.523$).

Task $\times$ congruency interaction: Greater incongruent interference in the interpersonal task

In regards to the task $\times$ congruency interaction, interference effects (Incongruent trial standard deviation - Congruent trial standard deviation) were calculated for each task, and

three paired *t*-tests (Bonferroni corrected critical $\alpha = .0167$) compared these interference effects across tasks (on average over the ASD and control groups). The interpersonal task produced a greater interference effect than both the biological dot task ($t(17) = 3.61, p = .002$) and the non-biological dot task ($t(17) = 3.27, p = .005$). Interference effects in the biological and non-biological dot tasks were not significantly different ($t(17) = 0.72, p = .48$). The amount of incongruent interference for each task can be seen in Fig. 2a.

Task x group interaction: Bottom up interference differs between the biological dot motion and interpersonal tasks according to participant group

In order to follow up the task x group interaction, difference scores were calculated for the three possible comparisons between the task conditions (biological dot, non-biological dot, and interpersonal tasks). These difference scores (Table 2) provided information on which motion type produced relatively higher error-plane deviations, independent of congruency (bottom-up interference). The unpaired *t*-test comparing the two groups on the difference between the biological dot task and the interpersonal task was significant, $t(16) = 2.69, p = .0161$ (Bonferroni corrected critical $\alpha = .0167$, for three tests). For control participants, fingertip position deviation was higher in the biological dot task than in the interpersonal task (Table 2); for ASD participants, the reverse was true with lower scores of fingertip position deviation in the biological dot task relative to the interpersonal task. This pattern can be observed in Fig. 2b. Comparisons of the relative differences between biological/non-biological and non-biological/interpersonal task suggested no significant pattern of differences between the groups (see Table 2).

Summary

In summary, greater error plane deviation was observed during incongruent compared to congruent trials for both participant groups indicating that the ASD group also displayed incongruent interference. Less overall error plane deviation was observed in the non-biological motion condition than in the biological dot-motion or in the interpersonal conditions for both groups. Incongruent interference was greater for the interpersonal task across both subject groups. Compared to the ASD participants, control participants showed a greater degree of bottom-up, congruency-independent movement interference in the biological task relative to the interpersonal task.

Comparisons between ToM/ASQ scores and interference effects

The ASQ scores for each ASD participant were compared to the amount of incongruent interference for each task (biological, non-biological and interpersonal) using Pearson's correlation. Interference was measured as the difference in error plane variance between congruent and incongruent conditions. This was also performed for a total interference score calculated by taking the average across all 3 tasks. No significant correlations were observed for any comparison ($r = 0.43; p = 0.25$). Correlations were also performed between incongruent interference and ToM score. Once again, no significant correlations were observed for any comparison ($r = 0.32; p = 0.41$), although a larger sample size is required to fully explore this issue.

Discussion

Contrary to our first prediction, the ASD participants demonstrated incongruent interference effects in the interpersonal task, equivalent to the control group. Furthermore, both ASD and control participants demonstrated less error plane deviation in the non-biological dot-motion task compared to the interpersonal tasks. However, the two participant groups demonstrated a different pattern of bottom-up interference across the three tasks: the ASD group exhibited greater error plane deviation during the interpersonal task, while the control group produced greater error plane deviation during the biological dot motion task. Overall, these results suggest that in our group of ASD individuals, the mechanisms underlying concurrent action observation and execution in a simple, apparently meaningless task were sufficient to produce equivalent incongruent interference effects as in the control participants. If co-activation of mirror neurons during the interpersonal task condition is indeed responsible for the interference effect, this would further imply that the ASD group exhibited behaviourally equivalent mirror neuron function. Nevertheless, the different pattern of bottom-up interference across the three tasks (the significant task x group interaction) does suggest potential differences in function between the two groups, which we will return to later on.

ASD individuals demonstrate congruency-dependent interference on the interpersonal task

The finding of an equivalent incongruent interference effect for both ASD and control participants on the interpersonal task is inconsistent with the hypothesis of a global mirror neuron dysfunction in the ASD group. We suggest three possible interpretations for these findings (1) Mirror neuron activity is not responsible for the interference effect (2) The mirror neuron system was intact in our group of high functioning ASD participants (3) Certain components of the mirror neuron system retain function in ASD individuals.

Firstly, the incongruent interference may be a consequence of shared representational resources other than co-activation of mirror neurons, in common with more general spatial stimulus-response compatibility effects that occur with non-biological stimuli (Cho and Proctor, 2003). Indeed, facilitation and interference during stimulus-response compatibility tasks using biological stimuli can arise due to the stimulus and response sets being spatially congruent or incongruent respectively (Brass, Bekkering, Wohlschlaeger and Prinz, 2000; Stürmer, Aschersleben and Prinz, 2000). Both spatial compatibility and automatic imitation (the latter thought to depend on the mirror neuron system) have been shown to influence the amount of interference and facilitation during imitation (Bertenthal, Longo and Kosobud, 2006). Although experiments have not yet been performed to directly test for mirror neuron activity during the task we used here, evidence from our previous work (Stanley, Gowen and Miall, 2007) using the biological and non-biological dot-motion stimuli speaks to this: Interference was not observed during the two dot-motion conditions when participants believed them to be computer generated. If general spatial compatibility effects were responsible for interference one would expect interference to be present consistently for the interpersonal and dot-motion tasks. However, future work is required to directly test this suggestion.

Secondly, in regards to the high functioning ability of our participants, we believe that it will be important to assess a wider range of participants in future studies. However, those neuroimaging studies that have reported abnormal mirror neuron functioning in ASD have used participant groups equivalent to ours (Bernier *et al.*, 2007; Dapretto *et al* 2006; Nishitani, Avikainen and Hari, 2004; Oberman *et al* 2005; Williams *et al* 2006). Therefore, according to these studies interference should be reduced even in high functioning individuals.

Thirdly, components of the mirror system may still retain some degree of function in both children and adults with ASD, as indicated by findings of their intact abilities to imitate and represent others' actions (Hamilton, Brindley and Frith, 2007; Hobson and Lee, 1999; McIntosh *et al* 2006; Sebanz *et al.*, 2005; Williams, Whiten and Singh, 2004). Indeed, it has previously been suggested that individuals with ASD may have a selective impairment of the goal-less directed route of imitation, with unimpaired goal directed imitation (Hamilton, in press). Although the repetitive arm action in our task appears meaningless, participants may have successfully performed the task by attributing a goal to it, or to each movement end point, hence depending more on those mirror neuron systems involved in processing transitive actions, and thus causing an interference effect.

The use of different processing strategies by individuals with ASD could help to explain the conflict between neuroimaging and behavioural findings. Those papers that have examined mirror neuron activity in the manual domain have employed simple, meaningless tasks (Oberman *et al.*, 2005; Williams *et al* 2006). So, if ASD participants were performing the imitation task in a different (but behaviourally correct) manner i.e. using a goal directed pathway, functional activity would be expected to be different to the control group. Future studies should attempt to specifically examine transitive versus intransitive actions using a combination of both imaging and behavioural data.

ASD and control participants demonstrate different patterns of congruency-independent interference

As with our previous work (Stanley, Gowen and Miall, 2007; also see Bouquet *et al* 2007), the biological dot-motion condition created greater arm movement variability in the control participants regardless of congruency than did the non-biological dot-motion. We previously interpreted this as due to bottom-up processing of the stimulus properties as there was greater variability present in the biological as opposed to non-biological dot stimulus and because the effect was independent of agency instruction and congruency (Stanley, Gowen and Miall, 2007). In addition, we further suggested that this congruency-independent bottom-up interference has similarities with imitation of intransitive actions and hence with processing of the specific details of movement. This differential effect of stimulus properties on action execution is also highlighted in studies of imitation style. Bekkering, Wohlschläger and Gattis (2000) demonstrated that transitive actions are imitated correctly in respect to the goal, but details regarding the movement may be ignored, whereas in intransitive action imitation, greater attention is paid to movement execution. In addition, during observation of goal directed movements, observer's eye movements are predictive rather than linked reactively to the observed movement (Flanagan and Johansson, 2003), implying that the goal

is identified in advance of movement execution. Furthermore, as imitation of meaningless actions becomes more familiar, functional brain activation switches from visual areas (V3, V5, MT) to more frontal and parietal areas, indicating less focus on the visual stimulus once the outcome or goal has been established (Grezes, Costes and Decety, 1999).

The results of our current work support this interpretation as incongruent (top-down) interference was greater for the interpersonal task as opposed to both types of dot motion task across both participant groups, suggesting that goal attribution occurred to a greater extent in the interpersonal task. However, the pattern of bottom-up interference in the biological dot and interpersonal tasks differed between the ASD participants and the control group. These findings parallel previous observations that children with ASD are less precise in their imitation of action (Hobson and Lee, 1999; Vanvuchelen, Roeyers and Weerdt, 2007). The pattern suggests that attention to and subsequent processing of the visual stimuli differed between both tasks and subject groups. Perhaps due to the meaningless nature of the biological dot, visual attention of this stimulus was higher in the control participants than for the interpersonal arm movement. In contrast, visual processing in ASD participants may have been biased towards the interpersonal task where apparent goals may have been more obvious.

Alternatively, our results may be explained by reduced visuomotor integration in the ASD participants so that the visual properties of the observed dot motion were less efficiently integrated into the executed movement. In addition, some visuomotor differences between ASD and control participants may be evident in the higher baseline variability in the ASD group, as indicated by their greater (albeit non-significant) error plane variability in both non-biological and interpersonal tasks. This makes the biological dot motion findings potentially more revealing. Abnormal sensory motor integration deficits in ASD have previously been suggested from studies that examine visually guided actions and dorsal stream processing (Gowen and Miall, 2005; Mari et al., 2003; Piek and Dyck, 2006; Spencer et al., 2000). There is growing evidence that processing within occipital visual areas and the STS and subsequent connections with the inferior frontal and temporal areas may be impaired in those with ASD (Castelli, Frith, Happe and Frith, 2002; Just, Cherkassky, Keller and Minshew, 2004; Villalobos *et al.*, 2005; Williams *et al.*, 2006; Zilbovicius *et al.* 2006) which may result in altered visually guided movements and less accurate imitation.

Finally, there exists a well established literature documenting motor difficulties in ASD (Dewey, Cantell and Crawford, 2007; Glazebrook, Elliot and Lyons, 2006; Gowen and Miall, 2005, 2007; Green et al., 2002; Miyahara et al., 1997; Hallet et al., 1993; Rinehart, Bradshaw, Brereton and Tonge, 2001; Ming, Brimacombe and Wagner, 2007). As mentioned above, the ASD participants appeared to show slightly more error plane variability for all tasks except the biological dot motion task, which suggests the presence of motor deficits within our ASD population. However, this error plane variability was not significantly different between the two participant groups and any concomitant motor deficits were not severe, which is supported by the absence of developmental coordination disorder comorbidity in our group. So while visuomotor integration differences may explain the pattern of bottom-up interference, all nine ASD participants showed a top-down incongruent

interference effect, indicating that this observation is a robust finding irrespective of any motor abnormalities.

Conclusions

Our results demonstrate an equivalent incongruent interference effect in both ASD and control participants in all versions of the task. However, ASD participants demonstrated greater error plane deviation during the interpersonal task where incongruent interference was maximal, whereas control participants exhibited greater error plane deviation in the biological dot-motion condition, indicating differences in way the two groups performed the tasks. We suggest two possible reasons for this difference. Firstly, in line with recent work (Hamilton, in press), visual processing in the ASD participants may have been biased towards stimuli that were more obviously goal related, such as the arm movement compared to the dot motion, whereas control participants showed the reverse tendency. Secondly, altered visuomotor integration in the ASD participants may have reduced the impact of bottom-up stimulus variability.

Acknowledgments

We would like to thank Jonathan Winter for his technical guidance. This work is supported by the Wellcome Trust.

References

- Baron-Cohen S. Out of sight out of mind? Another look at deception in autism. *Journal of child psychology psychiatry*. 1992; 33:1141–1155. [PubMed: 1400697]
- Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E. The autism-spectrum quotient (AQ): evidence from ASD syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*. 2001; 31:5–17. [PubMed: 11439754]
- Bekkering H, Wohlschlagel A, Gattis M. Imitation of gestures in children is goal-directed. *Q J Exp Psychol A*. 2000; 53:153–164. [PubMed: 10718068]
- Bernier R, Dawson G, Webb S, Murias M. EEG mu rhythm and imitation impairments in individuals with autism spectrum disorder. *Brain Cogn*. 2007
- Bertenthal BI, Longo MR, Kosobud A. Imitative response tendencies following observation of intransitive actions. *J Exp Psychol Hum Percept Perform*. 2006; 32:210–225. [PubMed: 16634666]
- Blakemore SJ, Frith C. The role of motor contagion in the prediction of action. *Neuropsychologia*. 2005; 43:260–267. [PubMed: 15707910]
- Brass M, Bekkering H, Prinz W. Movement observation affects movement execution in a simple response task. *Acta Psychol (Amst)*. 2001; 106:3–22. [PubMed: 11256338]
- Brass M, Bekkering H, Wohlschlagel A, Prinz W. Compatibility between observed and executed finger movements: comparing symbolic, spatial, and imitative cues. *Brain Cogn*. 2000; 44:124–143. [PubMed: 11041986]
- Buccino G, Binkofski F, Fink GR, Fadiga L, Fogassi L, Gallese V, Seitz RJ, Zilles K, Rizzolatti G, Freund HJ. Action observation activates premotor and parietal areas in a somatotopic manner: an fMRI study. *Eur J Neurosci*. 2001; 13:400–404. [PubMed: 11168545]
- Castelli F, Frith C, Happe F, Frith U. Autism, Asperger syndrome and brain mechanisms for the attribution of mental states to animated shapes. *Brain*. 2002; 125:1839–1849. [PubMed: 12135974]
- Cho YS, Proctor RW. Stimulus and response representations underlying orthogonal stimulus-response compatibility effects. *Psychon Bull Rev*. 2003; 10:45–73. [PubMed: 12747491]
- Cisek P, Kalaska JF. Neural correlates of mental rehearsal in dorsal premotor cortex. *Nature*. 2004; 431:993–996. [PubMed: 15496925]

- Craigheo L, Bello A, Fadiga L, Rizzolatti G. Hand action preparation influences the responses to hand pictures. *Neuropsychologia*. 2002; 40:492–502. [PubMed: 11749979]
- Dapretto M, Davies MS, Pfeifer JH, Scott AA, Sigman M, Bookheimer SY, Iacoboni M. Understanding emotions in others: mirror neuron dysfunction in children with autism spectrum disorders. *Nat Neurosci*. 2006; 9:28–30. [PubMed: 16327784]
- Decety J, Grezes J, Costes N, Perani D, Jeannerod M, Procyk E, Grassi F, Fazio F. Brain activity during observation of actions. Influence of action content and subject's strategy. *Brain*. 1997; 120(Pt 10):1763–1777. [PubMed: 9365369]
- Dewey D, Cantell M, Crawford SG. Motor and gestural performance in children with autism spectrum disorders, developmental coordination disorder, and/or attention deficit hyperactivity disorder. *J Int Neuropsychol Soc*. 2007; 13:246–256. [PubMed: 17286882]
- Flanagan JR, Johansson RS. Action plans used in action observation. *Nature*. 2003; 424:769–771. [PubMed: 12917683]
- Franz EA, Ford S, Werner S. Brain and cognitive processes of imitation in bimanual situations: Making inferences about mirror neuron systems. *Brain Res*. 2007; 1145:138–149. [PubMed: 17349983]
- Frith CD, Frith U. Interacting minds--a biological basis. *Science*. 1999; 286:1692–1695. [PubMed: 10576727]
- Gallese V, Goldman A. Mirror neurons and the simulation theory of mind-reading. *Trends Cogn Sci*. 1998; 2:493–501. [PubMed: 21227300]
- Glazebrook CM, Elliott D, Lyons J. A kinematic analysis of how young adults with and without autism plan and control goal-directed movements. *Motor Control*. 2006; 10:244–264. [PubMed: 17106133]
- Gowen E, Miall RC. Behavioural aspects of cerebellar function in adults with Asperger syndrome. *Cerebellum*. 2005; 4:279–289. [PubMed: 16321884]
- Gowen E, Miall RC. The cerebellum and motor dysfunction in neuropsychiatric disorders. *The Cerebellum*. 2007; 6(3):268–279. [PubMed: 17786823]
- Grafton ST, Arbib MA, Fadiga L, Rizzolatti G. Localization of grasp representations in humans by positron emission tomography. 2. Observation compared with imagination. *Exp Brain Res*. 1996; 112:103–111. [PubMed: 8951412]
- Green D, Baird G, Barnett AL, Henderson L, Huber J, Henderson SE. The severity and nature of motor impairment in Asperger's syndrome: a comparison with specific developmental disorder of motor function. *J Child Psychol Psychiatry*. 2002; 43:655–668. [PubMed: 12120861]
- Grezes J, Armony JL, Rowe J, Passingham RE. Activations related to "mirror" and "canonical" neurones in the human brain: an fMRI study. *Neuroimage*. 2003; 18:928–937. [PubMed: 12725768]
- Grezes J, Costes N, Decety J. Top down effect of strategy on the perception of human biological movement: A PET investigation. *cognitive neuropsychology*. 1998; 15:553–582. [PubMed: 22448838]
- Grezes J, Costes N, Decety J. The effects of learning and intention on the neural network involved in the perception of meaningless actions. *Brain*. 1999; 122(Pt 10):1875–1887. [PubMed: 10506090]
- Hadjikhani N, Joseph RM, Snyder J, Tager-Flusberg H. Abnormal activation of the social brain during face perception in autism. *Hum Brain Mapp*. 2007; 28:441–449. [PubMed: 17133386]
- Hallett M, Lebedowska MK, Thomas SL, Stanhope SJ, Denckla MB, Rumsey J. Locomotion of autistic adults. *Arch Neurol*. 1993; 50:1304–1308. [PubMed: 8257307]
- Hamilton AF. Emulation and mimicry for social interaction: A theoretical approach to imitation in autism. *Q J Exp Psychol*. in press.
- Hamilton AF, Brindley RM, Frith U. Imitation and action understanding in autistic spectrum disorders: how valid is the hypothesis of a deficit in the mirror neuron system? *Neuropsychologia*. 2007; 45:1859–1868. [PubMed: 17234218]
- Hamilton AF, Grafton ST. Goal representation in human anterior intraparietal sulcus. *J Neurosci*. 2006; 26:1133–1137. [PubMed: 16436599]
- Hobson RP, Lee A. Imitation and identification in autism. *J Child Psychol Psychiatry*. 1999; 40:649–659. [PubMed: 10357170]

- Happe FG. An advanced test of theory of mind: understanding of story characters' thoughts and feelings by able autistic, mentally handicapped, and normal children and adults. *Journal of Autism and Developmental Disorders*. 1994; 24:129–154. [PubMed: 8040158]
- Iacoboni M. Neural mechanisms of imitation. *Curr Opin Neurobiol*. 2005; 15:632–637. [PubMed: 16271461]
- Iacoboni M, Dapretto M. The mirror neuron system and the consequences of its dysfunction. *Nat Rev Neurosci*. 2006; 7:942–951. [PubMed: 17115076]
- Iacoboni M, Molnar-Szakacs I, Gallese V, Buccino G, Mazziotta JC, Rizzolatti G. Grasping the intentions of others with one's own mirror neuron system. *PLoS Biol*. 2005; 3:e79. [PubMed: 15736981]
- Iacoboni M, Woods RP, Brass M, Bekkering H, Mazziotta JC, Rizzolatti G. Cortical mechanisms of human imitation. *Science*. 1999; 286:2526–2528. [PubMed: 10617472]
- Jeannerod M. Neural simulation of action: a unifying mechanism for motor cognition. *Neuroimage*. 2001; 14:S103–S109. [PubMed: 11373140]
- Just MA, Cherkassky VL, Keller TA, Minshew NJ. Cortical activation and synchronization during sentence comprehension in high-functioning autism: evidence of underconnectivity. *Brain*. 2004; 127:1811–1821. [PubMed: 15215213]
- Kilner JM, Paulignan Y, Blakemore SJ. An interference effect of observed biological movement on action. *Curr Biol*. 2003; 13:522–525. [PubMed: 12646137]
- Koski L, Wohlschlaeger A, Bekkering H, Woods RP, Dubeau MC, Mazziotta JC, Iacoboni M. Modulation of motor and premotor activity during imitation of target-directed actions. *Cereb Cortex*. 2002; 12:847–855. [PubMed: 12122033]
- Mari M, Castiello U, Marks D, Marraffa C, Prior M. The reach-to-grasp movement in children with autism spectrum disorder. *Philos Trans R Soc Lond B Biol Sci*. 2003; 358:393–403. [PubMed: 12639336]
- McIntosh DN, Reichmann-Decker A, Winkielman P, Wilbarger JL. When the social mirror breaks: deficits in automatic, but not voluntary, mimicry of emotional facial expressions in autism. *Dev Sci*. 2006; 9:295–302. [PubMed: 16669800]
- Ming X, Brimacombe M, Wagner GC. Prevalence of motor impairment in autism spectrum disorders. *Brain Dev*. 2007; 29:565–570. [PubMed: 17467940]
- Miyahara M, Tsujii M, Hori M, Nakanishi K, Kageyama H, Sugiyama T. Brief report: motor incoordination in children with Asperger syndrome and learning disabilities. *J Autism Dev Disord*. 1997; 27:595–603. [PubMed: 9403374]
- Molnar-Szakacs I, Iacoboni M, Koski L, Mazziotta JC. Functional segregation within pars opercularis of the inferior frontal gyrus: evidence from fMRI studies of imitation and action observation. *Cereb Cortex*. 2005; 15:986–994. [PubMed: 15513929]
- Nishitani N, Avikainen S, Hari R. Abnormal imitation-related cortical activation sequences in Asperger's syndrome. *Ann Neurol*. 2004; 55:558–562. [PubMed: 15048895]
- Oberman LM, Hubbard EM, McCleery JP, Altschuler EL, Ramachandran VS, Pineda JA. EEG evidence for mirror neuron dysfunction in autism spectrum disorders. *Brain Res Cogn Brain Res*. 2005; 24:190–198. [PubMed: 15993757]
- Perner J, Wimmer H. “John thinks that Mary thinks that...”: attribution of second-order beliefs by 5- to 10- year old children. *Journal of Experimental Child Psychology*. 1985; 39:437–471.
- Piek JP, Dyck MJ. Sensory-motor deficits in children with developmental coordination disorder, attention deficit hyperactivity disorder and autistic disorder. *Hum Mov Sci*. 2004; 23:475–488. [PubMed: 15541530]
- Rinehart NJ, Bradshaw JL, Brereton AV, Tonge BJ. Movement preparation in high-functioning autism and Asperger disorder: a serial choice reaction time task involving motor reprogramming. *J Autism Dev Disord*. 2001; 31:79–88. [PubMed: 11439757]
- Rizzolatti G, Craighero L. The mirror-neuron system. *Annu Rev Neurosci*. 2004; 27:169–192. [PubMed: 15217330]
- Rizzolatti G, Fadiga L, Gallese V, Fogassi L. Premotor cortex and the recognition of motor actions. *Brain Res Cogn Brain Res*. 1996; 3:131–141. [PubMed: 8713554]

- Rizzolatti G, Fogassi L, Gallese V. Neurophysiological mechanisms underlying the understanding and imitation of action. *Nat Rev Neurosci.* 2001; 2:661–670. [PubMed: 11533734]
- Rumiati RI, Weiss PH, Tessari A, Assmus A, Zilles K, Herzog H, Fink GR. Common and differential neural mechanisms supporting imitation of meaningful and meaningless actions. *J Cogn Neurosci.* 2005; 17:1420–1431. [PubMed: 16197695]
- Saygin AP, Wilson SM, Hagler DJ Jr, Bates E, Sereno MI. Point-light biological motion perception activates human premotor cortex. *J Neurosci.* 2004; 24:6181–6188. [PubMed: 15240810]
- Sebanz N, Knoblich G, Stumpf L, Prinz W. Far from action blind: Representations of others actions in individuals with autism. *Cognitive Neuropsychology.* 2005; 22:433–454. [PubMed: 21038260]
- Spencer J, O'Brien J, Riggs K, Braddick O, Atkinson J, Wattam-Bell J. Motion processing in autism: evidence for a dorsal stream deficiency. *Neuroreport.* 2000; 11:2765–2767. [PubMed: 10976959]
- Stanley J, Gowen E, Miall RC. Effects of agency on movement interference during observation of a moving dot stimulus. *J Exp Psychol Hum Percept Perform.* in press.
- Sturmer B, Aschersleben G, Prinz W. Correspondence effects with manual gestures and postures: a study of imitation. *J Exp Psychol Hum Percept Perform.* 2000; 26:1746–1759. [PubMed: 11129371]
- Tessari A, Canessa N, Ukmar M, Rumiati RI. Neuropsychological evidence for a strategic control of multiple routes in imitation. *Brain.* 2007; 130:1111–1126. [PubMed: 17293356]
- Tessari A, Rumiati RI. The strategic control of multiple routes in imitation of actions. *J Exp Psychol Hum Percept Perform.* 2004; 30:1107–1116. [PubMed: 15584818]
- Theoret H, Halligan E, Kobayashi M, Fregni F, Tager-Flusberg H, Pascual-Leone A. Impaired motor facilitation during action observation in individuals with autism spectrum disorder. *Curr Biol.* 2005; 15:R84–R85. [PubMed: 15694294]
- Vanvuchelen M, Roeyers H, De Weerd W. Nature of motor imitation problems in school-aged males with autism: how congruent are the error types? *Dev Med Child Neurol.* 2007; 49:6–12. [PubMed: 17209969]
- Villalobos ME, Mizuno A, Dahl BC, Kemmotsu N, Muller RA. Reduced functional connectivity between V1 and inferior frontal cortex associated with visuomotor performance in autism. *Neuroimage.* 2005; 25:916–925. [PubMed: 15808991]
- Vogt S, Taylor P, Hopkins B. Visuomotor priming by pictures of hand postures: perspective matters. *Neuropsychologia.* 2003; 41:941–951. [PubMed: 12667530]
- Williams JWG, Gilchrist J, Perrett G, Murray A, Whiten A. Neural mechanisms of imitation and "mirror neuron" functioning in autism spectrum disorder. *Neuropsychologia.* 2006; 44:610–621. [PubMed: 16140346]
- Williams JH, Whiten A, Singh T. A systematic review of action imitation in autistic spectrum disorder. *J Autism Dev Disord.* 2004; 34:285–299. [PubMed: 15264497]
- Wimmer H, Perner J. Beliefs about beliefs – representation and constraining function of wrong beliefs in young children's understanding of deception. *Cognition.* 1983; 13:103–128. [PubMed: 6681741]
- Wohlschlager A, Gattis M, Bekkering H. Action generation and action perception in imitation: an instance of the ideomotor principle. *Philos Trans R Soc Lond B Biol Sci.* 2003; 358:501–515. [PubMed: 12689376]
- Zilbovicius M, Meresse I, Chabane N, Brunelle F, Samson Y, Boddaert N. Autism, the superior temporal sulcus and social perception. *Trends Neurosci.* 2006; 29:359–366. [PubMed: 16806505]

Fig 1a-b.

Error plane standard deviation (mm) in the ASD (a) and control group (b) for congruent (white bars) and incongruent trials (black bars) during biological dot-motion trials, non-biological dot-motion trials and interpersonal trials. Hatched bar indicates error plane standard deviation for practice trials. Data is collapsed across direction. Standard error bars are shown.

Fig 2a.

Congruency dependent (a) and congruency independent (b) interference (mm) in the ASD (white bars) and control group (black bars) during biological dot-motion trials, non-biological dot-motion trials and interpersonal trials. Congruency dependent data was calculated by subtracting error plane deviation in the incongruent plane from the congruent plane for each participant then taking the average. Congruency independent data was calculated by summing congruent and incongruent error plane deviation for each participant, then taking the average. Data is collapsed across direction. Standard error bars are shown.

Table 1

Scores for the 9 ASD participants on mentalizing tasks in increasing order of ability. Controls have been grouped together and averaged for each separate task and “total” reflects the total of these averages. 1=pass, 0 = fail. Penny hiding and Happe stories were given a % score out of a total of 6 and 16 respectively. Final two columns display IQ and AQ score respectively. ToM = theory of mind. Dash indicates tasks not performed.

Subject	Smarties	Sally-Ann	Coat story	Ice-cream story	Penny hiding	ToM Happe stories	Total	IQ	AQ scores
8	1	1	0	-	-	0.63	2.63	123	41
7	1	0	0	1	0.67	0.81	3.48	94	40
9	1	1	0	0	1	0.56	3.56	108	45
3	1	1	1	1	0	0.94	4.94	132	35
6	1	1	1	1	0.33	0.75	5.08	131	35
1	1	1	1	1	0.67	0.63	5.3	139	43
4	1	1	1	1	0.671	0.88	5.55	78	35
5	1	1	1	1	1	0.69	5.69	119	41
2	1	1	1	1	1	0.88	5.88	134	34
Controls (N=9)	1	1	1	1	1	0.94	5.94	115.7 ± 14.1 Range: 98-136	11.6 ± 5.0 Range: 5-18

Table 2
Differences in error-plane deviation scores by task, for the ASD and Control groups. T-tests compare difference scores between ASD and Control groups.

Difference score	ASD		Control		Between group <i>t</i> -tests	
	Mean	SEM	Mean	SEM	<i>t</i> (16)	<i>p</i>
Biological - Non-Biological	1.32	0.81	3.45	1.07	1.59	0.13
Non-biological - Interpersonal	-2.54	1.00	-1.26	0.70	1.04	0.313
Biological - Interpersonal	-1.22	0.76	2.19	1.02	2.69	0.016