CASE REPORT

Aortopulmonary window with pulmonary atresia with ventricular septal defect with D-transposition of great arteries: extremely rare anomaly

Sachin Sondhi,¹ PC Negi,² Rajesh Sharma,¹ Ayushi Mehta³

¹Department of Cardiology, Indira Gandhi Medical College, Shimla, Himachal Pradesh, India

²Head of Department of cardiology, Indira Gandhi Medical College, Shimla, Himachal Pradesh, India

³Department of Anaesthesia, Indira Gandhi Medical College, Shimla, Himachal Pradesh, India

Correspondence to
Dr Sachin Sondhi,
ssachin119@gmail.com

Accepted 27 April 2018

SUMMARY

Aortopulmonary window (APW) is rare a congenital heart disease accounting for 0.1%–0.2% of all congenital heart defects. The 35% of the APW has been associated with wide variety of other structural heart diseases such as ventricular septal defect, persistent ductus arteriosus, arch anomalies and coronary artery anomalies. To the best of our knowledge, only six cases of APW with pulmonary atresia with ventricular septal defect has been described in the literature. It resembles the type 1 truncus arteriosus, and differentiation from this condition is important prior to surgical correction. We present a case of 14-year-old girl child; she was diagnosed with APW with pulmonary atresia with ventricular septal defect and D transposition of great arteries with the help of echocardiography, cardiac catheterisation and cardiac CT.

BACKGROUND

The aortopulmonary window (APW) with pulmonary atresia with ventricular septal defect is extremely rare congenital anomaly. To the best of our knowledge, only six of such cases have been reported in the literature until.^{1–4} Although the patent ductus arteriosus or major aortopulmonary collaterals are the major route of blood supply to the lungs in case of pulmonary

Figure 2 ECG showing right axis deviation with clockwise rotation (note small q waves in lead 2, lead 3 and lead aVF), right ventricular hypertrophy (prominent R in V1), sudden transition of R in V1 to S in V2 (suggestive of ventricular septal defect) and R in V6 is 10 mm and S is 7 mm (suggestive of preserved left ventricular forces).

atresia. This case illustrates that the APW can be the only route of blood supply to the lungs in patients of pulmonary atresia.

CASE PRESENTATION

A 14-year-old girl child presented with complaints of palpitations, exertional breathlessness since childhood. There was history of recurrent chest infections and failure to gain weight in past. She denied any history of cyanotic spells, orthopnoea, paroxysmal nocturnal dyspnoea, swelling feet, angina and syncope. Her natal and postnatal history was

Figure 1 Chest X-ray showing situs solitus, levocardia, boot-shaped heart with right ventricle type of apex (blue star), atretic end of main pulmonary artery (yellow arrow), dilated aorta (white arrow) and borderline pulmonary vascularity traced up to middle one-third of lung fields.

Figure 3 Parasternal long axis view showing non-restrictive subaortic ventricular septal defect with atretic pulmonary valve (marked by #) and aortic valve (marked by *). Note aorta arising from right ventricle (RV) and pulmonary artery from left ventricle (LV).

To cite: Sondhi S, Negi PC, Sharma R, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224401

Figure 4 Echocardiogram showing atretic pulmonary valve (blue arrow), aortopulmonary window (marked by yellow *), dextro transposition of great arteries as aorta from anatomical right ventricle (RV) through aortic valve and blind pulmonary end towards left ventricle (LV). Note adequate size of LV and RV.

Figure 5 Echocardiogram showing flow from left ventricle (LV) through large subaortic ventricular septal defect into aorta. No flow from LV to pulmonary artery because of atretic pulmonary valvular end (marked by white arrow).

uneventful, and there was no family history of congenital heart disease in the other siblings. She had not been previously investigated. On general physical examination, she had short stature as her

Video 1 Right ventricle angiography lateral view.

height was 133 cm (less than third percentile for her age), her body weight was 31 kg (less than third percentile for her age), hyperterolism, low set ears, flat nose bridge, short and smooth philtrum, vitals showed blood pressure of 108/54 mm Hg and pulse 98 beats per minute regular and collapsing in nature, baseline oxygen saturation of 79% and she had mild central cyanosis, grade 1 clubbing. She had primary amenorrhoea and poorly developed secondary sexual characters. On cardiovascular examination, jugular venous pressure was not raised, apical impulse not palpable, grade 2 parasternal heave, S2 is single and accentuated with aortic ejection click and no murmur was heard.

INVESTIGATIONS

The chest X-ray revealed situs solitus, levocardia, cardiothoracic ratio 0.5, right ventricular (RV) type apex, dilated ascending aorta, blunt atretic end of main pulmonary artery (MPA) and borderline lung vascularity (figure 1). ECG showed right axis deviation, clockwise rotation, right ventricular hypertrophy (prominent R wave in V1 with sudden transition

Figures 6 Right ventricle(RV) angiography (coarsely trabeculated chamber) on lateral view showed (A) filling of left ventricle (LV) through ventricular septal defect marked by arrow; (B) atretic pulmonary end (marked by curved arrow) towards LV and (C) dilated aorta arising from RV.

Video 2 Right ventricle angiography anteroposterior view.

from R in V1 to S in V2), in V6 R wave is 10 mm and S wave is 7 mm suggestive of preserved left ventricular forces ([figure 2](#)).

Echocardiogram showed situs solitus, atrioventricular concordance, ventriculoarterial discordance, inferior vena cava opening into right atrium, all pulmonary veins opening into left atrium, left ventricle (LV) and RV are of adequate size, aorta arising from anterior chamber (RV), blind pulmonary end with atretic pulmonary valve towards RV; these findings suggestive of D transposition of great arteries, non-restrictive subaortic ventricular septal defect (VSD) with left to right shunt and pulmonary artery arising from aorta through APW ([figures 3–5](#)).

On cardiac catheterisation, on RV, lateral view angiography ([video 1](#)) showed filling of LV through ventricle septal defect ([figure 6A](#)), atretic pulmonary valvular end towards LV ([figure 6B](#)) and dilated aorta arising from RV ([figure 6B](#)).

On RV angiography with anteroposterior view ([video 2](#)) showed atretic pulmonary valvular end from LV ([Figure 7A](#)), aorta arising from RV ([figure 7B](#)) and main pulmonary artery arising from aorta through APW ([figure 7C](#)).

Figures 7 Right ventricle (RV) angiography on anteroposterior view showed (A) filling of left ventricle (LV) through ventricular septal defect and blind pulmonary end (marked by vertical arrow) towards LV; (B) aorta arising from RV and (C) filling of pulmonary artery from aorta through aortopulmonary window (marked by curved arrow).

Video 3 Left ventricle angiography on left anterior oblique cranial view.

On LV, angiography ([video 3](#)) showed atretic pulmonary end towards LV ([figure 8A](#)), initially filling of RV through VSD and then filling of aorta ([figure 8B](#)) and filling of MPA from ascending aorta through APW ([figure 8C](#)).

On aortic root angiography ([videos 4 and 5](#)) by pigtail catheter showed normal origin of coronaries from aortic sinuses and origin of pulmonary artery from ascending aorta ([figure 9A](#)). Then, we selectively entered into main pulmonary artery from aortic side and on giving contrast, the main pulmonary artery was dividing into left and right pulmonary artery ([figure 9B,C](#)). On descending aortogram, there was no evidence of major aortopulmonary collaterals ([video 6](#)).

The pressure in pulmonary artery (127/86/104) was same as aortic pressure (136/91/108). The systemic saturation in aorta was 77% and for pulmonary artery was also 77%. Pulmonary vascular resistance (PVR) was 24 wood units. On giving 100% oxygen for 10 min systemic saturation was 87% and PVR was 14 wood units ([tables 1 and 2](#)).

Cardiac CT scan confirmed the diagnosis of pulmonary atresia with APW ([figure 10](#)).

Figures 8 Left ventricle (LV) angiography revealed (A) atretic pulmonary end (marked by arrow); (B) filling of right ventricle (RV) through ventricular septal defect and then filling of aorta and (C) filling of main pulmonary artery from aorta.

Video 4 Aortic root angiography on left anterior oblique cranial view.

Video 5 Main pulmonary artery selectively entered from aortic side through aortopulmonary window in left anterior oblique cranial view.

OUTCOME AND FOLLOW-UP

The diagnosis of APW, pulmonary atresia, VSD and dextro transposition of great arteries was made. Mean pulmonary artery pressures remained same after pulmonary vasodilator testing. Although there was fall in PVR from 24 to 14 wood units after vasodilator testing, the PVR and PVR/systemic vascular resistance ratio was still higher than the values taken for operability. So patient was discharged on 20 mg of sildenafil thrice a day.

DISCUSSION

The APW with pulmonary atresia is an extremely rare congenital anomaly. To the best of our knowledge, only six of such cases have been reported in the literature until.¹⁻⁴ APW with pulmonary atresia with ventricular septal defect is a rare cardiac malformation first described by Faulkner *et al*¹ in 1974. He described a 2-month-old infant with APW, large VSD, membranous pulmonary atresia, anomalous origin of the left coronary artery from the pulmonary artery and a right aortic arch.¹ Then in 1981 Shore *et al*² described three cases

(3 month child, 18 year adult, 23 year adult) of APW with pulmonary atresia with VSD. In 2004, Kuhn *et al*³ diagnosed this entity by prenatal echocardiography; this case underwent primary surgical correction at age of 4 weeks due to intractable heart failure. The sixth case was diagnosed by Shah *et al*⁴ in 2012 in 9 days old neonate and surgically corrected by barbero-marcial technique.

Figures 9 Aortic root angiography on left anterior oblique cranial view showed (A) origin of main pulmonary artery (MPA) from ascending aorta; (B) MPA selectively entered with pigtail catheter from aortic side through aortopulmonary defect and (C) on giving contrast into MPA in right anterior oblique cranial view, it is dividing into left pulmonary artery (LPA) and right pulmonary artery (RPA).

Video 6 Descending aortogram. No evidence of major aortopulmonary collateral arteries.

Table 1 Cardiac catheterisation data before and after giving 100% oxygen for 10 min. PCWP (Pulmonary capillary wedge pressure)

Chamber	Oxygen saturation (preoxygenation)	Oxygen saturation (postoxygenation)	Pressures (preoxygenation)	Pressures (postoxygenation)
Superior vena cava	60			
Right atrium	62	68	11/5/8	12/7/9
Inferior vena cava	64			
MvO ₂	61			
Right ventricle	62	72	144/10	140/12
Pulmonary artery (through aortopulmonary window)	77	87	127/86/104	118/85/100
PCWP(mean)	97	99	13	13
Left ventricle	91	97	146/14	142/14
Ascending aorta	77	87	136/91/109	136/90/109
Femoral artery	77	87	154/70/104	150/68/100

Table 2 Calculation of pulmonary blood flow (PBF, Q_p), systemic blood flow (SBF, Q_s), pulmonary vascular resistance (PVR) and systemic vascular resistance (SVR)

	Pre Oxygenation	Post Oxygenation
PBF (Q _p)	3.82	6.38
SBF (Q _s)	5.10	4
PBF/SBF (Q _p /Q _s)	0.71	1.74
PVR	24	14
SVR	20	25
PVR/SVR	1.2	0.56

Note: VO₂ 125 mL/min/m², BSA 1 and haemoglobin 12 g%.

This is the unusual differential diagnosis of common arterial trunk.⁴ Common arterial trunk is defined as a single arterial vessel that arises from the base of the heart and gives origin

to the coronary, pulmonary and systemic arteries. APW has features in common with truncus arteriosus so that some authors include cases of window within general grouping of truncus arteriosus.⁵ In pulmonary atresia, there is a single great vessel seen usually overriding the interventricular septum with a large subarterial VSD. The great vessel (aorta) gives rise to coronary arteries and head and neck vessels and continues as the descending aorta. Usually, there is a posterior descending artery (PDA) supplying the branch pulmonary arteries, and in the absence of PDA, there are large major aortopulmonary collaterals. Our case demonstrated that despite PDA and major aortopulmonary collateral arteries, APW is another way of supply to the pulmonary arteries in pulmonary atresia. Differentiation from truncus arteriosus is an important aspect. In the current case, blind atretic pulmonary valvular end visible on echocardiography and confirmed on cardiac catheterisation gave a clue to the diagnosis. Closure of APW along with

Figure 10 Cardiac CT (transverse and coronal section) showed origin of aorta from right ventricle and origin of main pulmonary artery from aorta through aortopulmonary window.

Learning points

- This case highlights the fact that aortopulmonary window can also be a sole supply to branch pulmonary arteries in cases of pulmonary atresia.
- It emphasises the importance of looking for a 'stump' proximal to branch pulmonary arteries in a suspected common arterial trunk on echocardiography and on cardiac catheterisation.
- Early diagnosis of this rare entity is important, as early surgical repair will prevent the development of irreversible pulmonary hypertension.

Rastelli procedure may be performed in this type of scenario which involves routing of LV to aorta, patch closure of ventricular septal defect and placement of conduit between RV and pulmonary artery.⁶

Contributors All the authors were involved in the management of the patient. Echocardiography was done by PN. Cardiac catheterisation was performed by SS, RS and AM. SS and AM wrote the manuscript. Manuscript was approved by all the authors.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Faulkner SL, Oldham RR, Atwood GF, *et al.* Aortopulmonary window, ventricular septal defect, and membranous pulmonary atresia with a diagnosis of truncus arteriosus. *Chest* 1974;65:351–353.
- 2 Shore DF, Ho SY, Anderson RH, *et al.* Aortopulmonary septal defect coexisting with ventricular septal defect and pulmonary atresia. *Ann Thorac Surg* 1983;35:132–7.
- 3 Kuehn A, Oberhoffer R, Vogt M, *et al.* Aortopulmonary window with ventricular septal defect and pulmonary atresia: prenatal diagnosis and successful early surgical correction. *Ultrasound Obstet Gynecol* 2004;24:793–6.
- 4 Shah S, Kavya SR, Saraf R, *et al.* Unusual differential diagnosis of common arterial trunk. *Pediatr Cardiol* 2012;33:1180–2.
- 5 Calder L, Van Praagh R, Van Praagh S, *et al.* Truncus arteriosus communis. *Am Heart J* 1976;92:93.
- 6 Morell VO, Jacobs JP, Quintessenza JA. Surgical management of transposition with ventricular septal defect and obstruction to the left ventricular outflow tract. *Cardiol Young* 2005;15(Suppl 1):102–5.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow