Mr. Bowman said he came as a practitioner to speak to practitioners, and had therefore avoided going into anything like abstract or physiological reasoning. In one case he had seen the connection referred to by Dr. Richardson; but in that case there was a large amount of sugar, and the specific gravity was not higher than 1025. It was a case of chronic glaucoma—the sight of one eye was entirely gone and but little remained in the other. He performed the operation of iridectomy, and nothing could be more satisfactory than the recovery of the patient. With regard to the blood conditions, there might be varied morbid conditions of the blood not marked by albuminuria, or by anything gouty, rheumatic, or diabetic in the system. There was much more to be discovered on the subject by future observers, and there might be states of blood found to be intimately connected with glaucoma; but he thought the symptoms were rather referable to a local condition of the blood vessels and the nervous coats of the blood vessels in the eye. With regard to the *rationale* of the operation, he followed his friend Von Graefe. He at first thought that the result was produced by opening a freer communication between the back and front of the iris, thus allowing the fluids from behind to percolate more freely into the aqueous humour. He was not prepared to agree with Von Graefe's early supposition, that it might depend upon a removal of a portion of the iris, the iris being considered as the secreting agent of the aqueous humour, and that thus there might be a diminished secretion of the aqueous humour in the eye. He did not think it depended upon the quantity of iris removed; but upon the exactitude with which the marginal part of the iris was turned away from the ciliary bodies. With regard to the cause of the tension, he believed that in chronic cases it was an enlargement of the vitreous chamber of the eye, and in many cases a partial consolidation of the substance of the vitreous humour, not necessarily connected with any serum, or fibrous, or cloudy flocculi.

Employment of Glycerine in the Treatment of Diseases of the Eye. By W. ABBOTTS SMITH, M. D.

Notwithstanding that glycerine is largely used as a bland, emollient application to the skin, in several affections of that delicate membrane, its importance as a remedial agent in certain diseases of the conjunctiva, appears to be considerably underrated, although several marked points of analogy exist between the two structures.

Having been induced some time since by the perusal of a paper published in the *Bulletin de Therapeutique*, by M. Foucher, upon this subject, I have given a fair trial to the topical employment of glycerine, and the results have been such as to warrant me in recommending its use in some of the forms of ophthalmia, and other superficial affections of the eye.

The application of *pure* glycerine, uncombined, is decidedly advantageous for the prevention and removal of the hard crusts produced by irritation which are occasionally closely adherent to the eye-lids in ophthalmia, which disorder they consequently tend to perpetuate.

In ectropion, and in cases of partial loss of the eye-lashes, glycerine forms an agreeable and efficient application; and it will also be found similarly useful in epiphora, and in any other superficial diseases of the eye which are dependent upon diminished secretion by the Meibomian glands. Among these may be especial mentioned the peculiar condition of the conjunctiva to which Von Ammon gave the name of *Xerosis Conjunctivæ*, in reference to the dry, cuticular appearance of the eye-ball, and which is most commonly met with in persons of scrofulous diathesis, who have long suffered from chronic ophthalmia. The free play of the eye-lids upon the front surface of the eye-ball is preserved by the topical employment of the glycerine; and the tendency to symblepharon, or to

an uncomfortable temporary adhesion of the eyelids, which occurs after sleep, is counteracted.

Unctuous applications are always preferable to watery solutions in ophthalmia, both because they are not so readily washed away by the tears, and because they more closely resemble the natural greasy secretion of the Meibomian glands, by which the conjunctival surface is lubricated in the healthy state; and the circumstance that glycerine alone, when employed locally, is highly beneficial in the ophthalmic affections already referred to, also points to that substance as a valuable vehicle in the preparation of collyria.

M. Foucher enumerates, in his paper in the *Bulletin de Therapeutique*, a long list of medicaments with which he has combined glycerine in the treatment of ophthalmia in numerous cases which had come under his care at the *Hopital de Enfants*. The substances which I have found to be most usefully employed in this form are the sulphate of zinc, the sulphate of copper, and laudanum; but M. Foucher also combines glycerine with the perchloride of iron, with borax, with tincture of iodine, and other agents.

It should be here stated that the nitrate of silver, owing to the chemical action which ensues upon its coming in contact with organic matter, cannot be used conjointly with glycerine; and it further remains to be mentioned that the glycerine should be well purified, in which condition it is perfectly natural, does not possess any odor or taste, and does not produce irritation when it is applied to the eye.

The Submaxillary Ganglion.

M. Claude Bernard, in a late communication to the Academy of Sciences, treated of the special function of the submaxillary ganglion. The results of his experiments prove that the tongue is connected with the submaxillary gland by two kinds of nervous arcs, which are, in a measure, concentric; the one of the greatest extent passing through the brain, the other, far shorter, passing through the submaxillary ganglion. There results from these two nervous connections two kinds of reflex action destined to affect the submaxillary gland. The first, which traverses the brain, is *intelligent*, and is excited more particularly by the gustatory faculty of the tongue; the second, which is unintelligent, is transmitted by the submaxillary ganglion, and seems to be provoked more especially by the dry or moist condition of the buccolingual mucous membrane. The submaxillary ganglion not only possesses the faculty of developing reflex action, which may, by its mediation, reach the gland without passing through the encephalic centre, but it also exerts a special influence upon the intermittent character of the salivary secretion. Everyone is aware of the fact, that the flow of the saliva takes place in an intermittent form, and that only when an exciting cause, reflex or direct, calls forth the activity of the gland, and that the secretion ceases when the exciting cause is withdrawn. M. Bernard has observed that after the section of the submaxillary ganglion, the lingual and chorda tympani nerves still remaining intact, the secretion of the submaxillary gland becomes continuous, although it may be augmented by the application of strong-tasted stimulants to the surface of the tongue. Another fact of interest brought to light by the same experiment is, that when separated from the encephalic centre, the ganglion after a certain time loses its controlling influence over the secretive powers of the gland, which instead of ceasing its function, remains in a condition of permanent activity.—From which it results that the submaxillary ganglion is at one and the same time independent of, and subordinate to, the encephalic centre.