Published in final edited form as:

Curr Protoc Chem Biol. 2018 June ; 10(2): e38. doi:10.1002/cpch.38.

Characterizing Protein Kinase Substrate Specificity Using the Proteomic Peptide Library (ProPeL) Approach

Joshua M. Lubner¹, Jeremy L. Balsbaugh², George M. Church³, Michael F. Chou³, and Daniel Schwartz¹

¹University of Connecticut, Department of Physiology and Neurobiology, Storrs, Connecticut

²University of Connecticut, Proteomics & Metabolomics Facility, Center for Open Research Resources & Equipment, Storrs, Connecticut

³Harvard Medical School, Department of Genetics, Boston, Massachusetts and Wyss Institute for Biologically Inspired Engineering, Boston, Massachusetts

Abstract

Characterizing protein kinase substrate specificity motifs represents a powerful step in elucidating kinase-signaling cascades. The protocol described here uses a bacterial system to evaluate kinase specificity motifs *in vivo*, without the need for radioactive ATP. The human kinase of interest is cloned into a heterologous bacterial expression vector and allowed to phosphorylate *E. coli* proteins *in vivo*, consistent with its endogenous substrate preferences. The cells are lysed, and the bacterial proteins are digested into peptides and phosphoenriched using bulk TiO₂. The pooled phosphopeptides are identified by tandem mass spectrometry, and bioinformatically analyzed using the pLogo visualization tool. The ProPeL approach allows for detailed characterization of wildtype kinase specificity motifs, identification of specificity drift due to kinase mutations, and evaluation of kinase residue structure-function relationships.

Keywords

protein kinase; phosphorylation motif; kinase specificity; mass spectrometry; proteomic peptide library; ProPeL; pLogo

Key References

Chou et al., 2012

Original proof-of-concept publication for ProPeL

Lubner et al., 2017

Publication using the current ProPeL workflow

Internet Resources

<http://schwartzlab.uconn.edu/pepextend>

PeptidExtender web tool, used in this protocol to map tryptic phosphopeptides to the reference *E. coli* proteome and create modification-centered 15mers.

<http://plogo.uconn.edu/>

Motif visualization tool.

<http://motif-x.med.harvard.edu>

Motif discovery tool.

INTRODUCTION

Protein kinases are enzymes that catalyze the covalent addition of phosphate to specific amino acids within protein substrates as posttranslational modifications. Such changes can often alter the biological function of the target protein, so understanding the relationship between kinases and their substrates can provide important biological insights. However, traditional co-immunoprecipitation methods do not work to identify these transient interactions, so alternative approaches including the one described here have been developed.

Kinases discriminate their substrates, in part, by recognizing short linear patterns of amino acids or “motifs” that surround the phosphoacceptor residue (Pinna and Ruzzene, 1996; Ubersax and Ferrell Jr, 2007), and the identification of these motifs has proven to be a powerful tool for substrate hypothesis generation (Obenauer et al., 2003; Miller et al., 2008). This unit presents a non-radioactive, bacterial approach for querying protein kinase substrate specificity *in vivo* termed ProPeL (for Proteomic Peptide Library, Chou et al., 2012). In this method, a human kinase is expressed in *E. coli* cells (which have a very low background level of endogenous phosphorylation). The bacterial proteome functions as a substrate library for the human kinase to phosphorylate *in vivo*, consistent with its distinct specificity. In this way, the *E. coli* acts as a living mini reaction vessel, facilitating thousands of simultaneous *in vivo* phosphorylation events and generating thousands of kinase-specific phosphorylation sites that are isolated and identified by liquid chromatography tandem mass spectrometry (LC-MS/MS). Using our lab’s suite of computational tools, we can extract and visualize kinase specificity motifs, and make high-confidence predictions of downstream targets. ProPeL can also be used to evaluate the influence of disease-associated mutations on kinase substrate specificity (Lubner et al., 2017).

The Basic Protocol describes the overall ProPeL workflow, which is represented in Figure 1. The major steps include expression of the kinase of interest and *in vivo* phosphorylation of bacterial proteins, tryptic digestion, phosphopeptide enrichment, phosphopeptide identification by LC-MS/MS, and computational analysis. Prior to carrying out ProPeL, the kinase of interest must be cloned into an appropriate bacterial expression vector (Strategic Planning), and *in vivo* activity may need to be optimized (Troubleshooting). In the event that the kinase cannot be expressed in an active form in *E. coli*, it is possible to perform an *in vitro* version of ProPeL using recombinant (or endogenously purified) kinase (Alternate Protocol).

STRATEGIC PLANNING

At the start of a new ProPeL project, it is important to design the correct kinase insert, and choose an appropriate bacterial expression vector. A successful ProPeL experiment necessarily requires the expression of a soluble, constitutively active protein kinase. This may require expressing a truncation that omits inhibitory sequences (such as the C-terminal PKC inhibitory tail) or mimics caspase cleavage (as is required for activation of MST3). In other instances, it is as simple as expressing the naked catalytic subunit, as is the case for PKA. Plasmid design is also an important variable that can enhance kinase solubility (see

Critical Parameters). Although the background serine/threonine/tyrosine phosphorylation in *E. coli* is only around 0.4% (Macek et al., 2008; Soares et al., 2013; Hansen et al., 2013), it is still advisable to create a kinase-dead mutant as a negative control. This is most easily achieved by mutating the catalytic aspartate to an asparagine (within the HRD motif) or the invariant lysine to an alanine (VAIK motif), as these residues are essential for catalysis (Hanks et al., 1988; Gibbs and Zoller, 1991). The best indicator of a successful ProPeL result is the demonstration of strong *in vivo* phosphorylation of bacterial proteins. Therefore, it is critical to optimize expression conditions prior to mass spectrometry sample preparation (see Critical Parameters and Troubleshooting).

While we have found success with several phylogenetically distant kinases using the standard *in vivo* ProPeL approach, there are nevertheless instances of kinases that are challenging for the system. Kinases with highly complex activation requirements (such as involvement in multiple activation cascades or requirements for large protein scaffolding structures), cytoplasmic conditions that are unsustainable for *E. coli* growth, or kinases that are toxic to *E. coli* through their activity would be poor targets for ProPeL. Similarly, a kinase that is part of a cascade, such as the MAPK kinases, will be unsuitable for *in vivo* ProPeL. While it is possible to recapitulate activating cascades by co-expressing kinases in *E. coli* (Khokhlatchev et al., 1997), the greater the complexity of the cascade, the more difficult it is to determine which individual phosphorylation sites should be attributed to each individual kinase. However, those kinases are suitable candidates for the *in vitro* ProPeL approach, provided they can be successfully purified in the active state (see Alternate Protocol).

BASIC PROTOCOL

E. coli Kinase Expression and Mass Spectrometry Sample Preparation

This approach expresses an active protein kinase in *E. coli*, facilitating the *in vivo* phosphorylation of bacterial proteins. Following expression, the cells are harvested, lysed, and evaluated for kinase activity. Bacterial proteins are tryptically digested, and enriched for phosphopeptides using TiO₂. The resulting samples are ready for sequence identification by LC-MS/MS.

Materials

Appropriate *E. coli* cell strain (see *Critical Parameters*)
Bacterial expression vector with appropriate kinase insert
LB plates and liquid broth, and appropriate antibiotic (see *Reagents and Solutions*)
Isopropyl β -D-1-thiogalactopyranoside (IPTG) (Promega cat. no. V3951 or V3955)
Tris Base (Sigma Aldrich cat. no. T6066)
Sodium Chloride (Fisher Scientific cat. no. S671-500)
Urea (Fisher Scientific cat. no. BP169-500)

Halt Protease Inhibitor Cocktail, EDTA-Free (Thermo Fisher Scientific cat. no. PI87785)

Halt Phosphatase Inhibitor Cocktail (Thermo Fisher Scientific cat. no. PI78420)

Disposable Sterile Syringe with Luer Lok, 5 mL (Fisher Scientific cat. no. 14-829-45)

Sterile Syringe Filter, 0.22 μ m (Fisher Scientific cat. no. SLGL0250S)

BCA Assay Kit (Thermo Fisher Scientific cat. no. PI23225)

SDS-PAGE Gel, Laemmli Loading Buffer, and Running Buffers (see *Reagents and Solutions*)

Pro-Q Diamond Phosphoprotein Gel Stain (Fisher Scientific cat. no. P33300)

Pro-Q Diamond Phosphoprotein Destain Solution (see *Reagents and Solutions*)

PeppermintStick Ladder (Fisher Scientific cat. no. P27167)

All Blue Protein Standards, *optional* (Bio-Rad cat. no. 1610373)

GelCode Blue (Thermo Fisher Scientific cat. no. PI24592)

Sodium Acetate, ACS grade (Fisher Scientific cat. no. S210-500)

Dithiothreitol (DTT) (Fisher Scientific cat. no. BP172)

Iodoacetamide, mass spectrometry grade (Sigma Aldrich cat. no. I1149)

Calcium Chloride (Fisher Scientific cat. no. AC349615000)

Trypsin, Sequencing Grade Modified (Promega cat. no. V5111 or V5117)

Potassium Phosphate Monobasic (Fisher Scientific cat. no. BP362-500)

Water (double distilled or Ultrapure)

Water, HPLC or LC/MS grade (Fisher Scientific cat. no. W6-1)

Acetonitrile, ACS grade (Fisher Scientific cat. no. A21-4)

Acetonitrile, LC/MS grade (Fisher Scientific cat. no. A955-1)

Methanol, ACS grade (Fisher Scientific cat. no. A412-4)

Methanol, LC/MS grade (Fisher Scientific cat. no. A456-500)

Acetic Acid, Glacial ACS grade (Fisher Scientific cat. no. A38SI-212)

Acetic Acid, LC/MS grade (Fisher Scientific cat. no. A11350)

Chloroform, HPLC grade (Fisher Scientific cat. no. C607SK-4)

Trifluoroacetic acid, LC/MS grade (Fisher Scientific cat. no. A116-50)

Formic acid, LC/MS grade (Fisher Scientific cat. no. A117-50)

Lactic acid (Sigma Aldrich cat. no. 69785-1L)

tC18 SEP-Pak cartridges (Waters cat. no. WAT054925)

Titansphere TiO₂ 5 µm beads (GL Sciences, cat. no. 1400B500)
Empore SPE Disks C18 (Sigma-Aldrich cat. no. 66883-U)
Shaking bacterial culture incubator
Probe sonicator
Refrigerated centrifuge
Electrophoresis chamber
Gel imager
Water bath
Kel-F hub (KF), point style 3, gauge 16 needle (Hamilton Company cat. no. 90516)
Plunger assembly N, RN, LT, LTN for model 1702 (Hamilton Company cat. no. 1122-01)
Room temperature 15 mL conical tube shaker

Kinase Expression and *In Vivo* Phosphorylation of Bacterial Proteins

This protocol assumes that the appropriate kinase coding sequence has been cloned into a bacterial expression vector, and transformed into an appropriate *E. coli* cell strain. The following steps are for standard protein expression using an IPTG-inducible vector. Optimal protein expression conditions need to be determined empirically, and the expression steps should be adjusted accordingly.

- 1 Using a sterile pipette tip, streak a fresh plate from a bacterial glycerol stock. Incubate plate upside down at 37°C overnight.
- 2 Inoculate a well-isolated colony in 5 mL LB media (+ appropriate antibiotic) and grow at 37°C and 250 rpm overnight.
- 3 Inoculate 100 mL LB media (+ appropriate antibiotic) with 2 mL of overnight culture in a 500 mL Erlenmeyer flask.
- 4 Grow bacteria to an OD₆₀₀ of 0.4–0.6 (mid-log) and induce with 0.5 mM IPTG. Incubate at 37°C and 250 rpm for 3–24 hours.

This is an optimization point for kinase expression and phosphorylation. See Critical Parameters and Troubleshooting.

- 5 [OPTIONAL] Collect a 1 mL aliquot in a separate tube and store at 4°C for later analysis by Pro-Q Diamond.
- 6 Pellet cells by centrifugation at 6,000 g for 15 minutes at 4°C.

Pellet may be stored at –80°C, but it is best to proceed promptly.

Lysis and Evaluating the Success of *In Vivo* Phosphorylation

When preparing sample for LC-MS/MS, use all LC/MS-grade solvents and Eppendorf brand microcentrifuge tubes for sample preparation. If evaluating an aliquot during the kinase

expression optimization phase, ACS-grade reagents are acceptable. ACS-grade solvents may be used for all SDS-PAGE steps.

- 7 Prepare Lysis Buffer (see *Reagents and Solutions*), add at 2 mL/g of wet pellet, and resuspend by pipet mixing.
- 8 Lyse cells by sonication, using 15-second pulses on 15% power, until solution is no longer opaque.

To prevent cells over-heating, keep tubes on ice (between and during sonications) with at least 1 minute rest between pulses. The solution will be colored, but should be clear.
- 9 Centrifuge solution at 20,000 g for 30 minutes at 4°C. Transfer clarified supernatant to a fresh tube and discard pellet. If necessary, repeat centrifugation to further clarify.
- 10 Filter lysate with a disposable syringe and 0.22 µm filter attachment to further remove cellular debris.
- 11 Quantify samples by BCA assay (or by NanoDrop using the protein A280 measurement if evaluating an aliquot during the optimization phase for kinase expression).

Note that a NanoDrop A280 measurement is less accurate for quantifying protein concentration, and tends to overestimate protein concentration in crude cell lysate by a factor of 3–4X relative to a BCA assay. Accordingly, additional sample should be loaded when using NanoDrop readings. Using a NanoDrop is acceptable for optimization and gel evaluation, but when preparing a sample for mass spectrometry a BCA assay is critical for accurate protein quantification.
- 12 For each sample, separate 25 µg (or 75 µg if using NanoDrop A280 measurement) by SDS-PAGE, with 2 µL PeppermintStick Phosphoprotein ladder (and 5 µL All Blue Protein Standards, *optional*).

Stain gel

Analyze with Pro-Q Diamond stain as described below, according to manufacturer's instructions. All incubations should be carried out on a rocker at room temperature.

- 13 Immerse gel in 100 mL fix solution (see *Reagents and Solutions*) and incubate for 30 minutes. Discard the fix solution and add fresh 100 mL fix solution. Incubate for at least 30 minutes.

This is a pause point, as gel can be left in fix solution overnight.
- 14 Discard the fix solution and wash with 100 mL ultrapure water. Incubate for 10 minutes, discard, and repeat twice for a total of 3 water washes.

All subsequent incubations must be done in the dark, as Pro-Q Diamond is light sensitive.

- 15 Add 60 mL Pro-Q Diamond stain and incubate 90 minutes.
- 16 Discard stain and add 90 mL Pro-Q Diamond destain solution (see *Reagents and Solutions*), incubating for 30 minutes. Discard destain and repeat twice more for a total of 3 destain washes. Rinse with ultrapure water for 5 minutes, discard and repeat water wash once.
- 17 Visualize on appropriate imager (Typhoon, ChemiDoc etc.) using the following wavelengths: Ex: 555 nm, Em: 580 nm. Adjust the signal such that only the two phosphoprotein bands (23.k kDa and 40 kDa) on the PeppermintStick ladder are clearly visible (18 kDa band may be faintly visible).

To control for loading differences (which can change the level of background signal), it is important to perform a total protein stain. We use GelCode Blue according to manufacturer's instructions, but other stains (such as Coomassie staining) are acceptable.

- 18 Add 20 mL GelCode Blue and incubate on a rocker at room temperature for at least an hour. Preferably, leave gel in GelCode Blue overnight for clearest signal.
- 19 Destain using ultrapure water. For best results, change water several times until background signal has been completely removed. Gel may be left in water overnight.
- 20 Image using the Coomassie setting (and white light conversion screen).

If autophosphorylation of the kinase of interest is evident and/or there is a marked increase in phosphorylation of proteins throughout the gel relative to the negative control (see *Troubleshooting* and Figure 2), proceed with the remaining steps for LC MS/MS sample preparation. Otherwise, optimize kinase expression/activity (see *Critical Parameters and Troubleshooting*).

Protein Reduction, Alkylation and Tryptic Digestion

The following protocol is for the preparation of sample from 10 mg crude lysate (as quantified by BCA assay). Sample volumes can be scaled as needed. We have observed excellent results starting with as low as 1 mg crude lysate, but this is contingent on the activity of the target kinase within the *E. coli*.

Methanol/chloroform extract 10 mg of crude protein lysate as follows:

- 21 Add 4X sample volume methanol, vortex, add 1X sample volume chloroform, vortex, and add 3X sample volume water and vortex.
- 22 Spin sample at 14,000 g for 10 minutes at 4°C.

There will be a lower organic layer, a middle protein disc and a top, aqueous layer. If separation is insufficient, increase centrifugation time, but DO NOT increase centrifuge speed. Remove top aqueous layer without disturbing protein disc.
- 23 Add 4X methanol, vortex and spin at 14,000 g for 10 minutes at 4°C. Remove as much methanol as possible and air dry (approximately 10 minutes).

Over-drying pellet will make it very difficult to resuspend. For best results, proceed immediately to the next step.

- 24** Resuspend pellet in 1 mL Lysis Buffer for a protein concentration of 10 mg/mL. Pipet mixing can be aided by gently vortexing the sample and using heat for short durations (do not exceed 50°C). Sample can also be allowed to resolubilize overnight at 4°C.

- 25** Add DTT from a fresh 0.5 M stock (10.1 µL of 0.5 M DTT) to a final concentration of 5 mM DTT. Incubate at 56°C for 25 minutes.

During this step, the protein is unfolded by heat and DTT denaturation. Avoid temperatures above 60°C which can cause urea-based carbamylation of lysines and protein N-termini.

- 26** Allow mixture to cool to room temperature, and add iodoacetamide from a fresh 0.5 M stock to a final concentration of 14 mM iodoacetamide (29.1 µL of 0.5 M iodoacetamide). Incubate for 30 minutes at room temperature in the dark.

During this step, the exposed free cysteine residues are alkylated to prevent disulfide bond formation. Iodoacetamide is light sensitive. Store the 0.5 M iodoacetamide stock solution in the dark, and carry out the alkylation and quenching steps in the dark.

- 27** Quench alkylation by adding an additional 5 mM DTT from a 0.5 M stock (10.5 µL of 0.5 M DTT). Incubate 15 minutes at room temperature in the dark.

- 28** Transfer the sample to a 15 mL conical tube, and dilute the sample 1:5 by adding 4.2 mL of 25 mM Tris-HCl (pH 8.2), to reduce urea concentration to 1.6 M.

- 29** Add CaCl₂ from a 0.1 M stock to a final concentration of 1 mM (53 µL of 0.1 M CaCl₂).

- 30** Add 100 µg trypsin (for an enzyme:substrate ratio of 1:100), and incubate at 37°C for 16 hours.

- 31** Allow the digest to cool to room temperature, and stop digestion by acidification with 25 µL trifluoroacetic acid to 0.4% (v/v). Verify that pH < 2.5, otherwise add additional trifluoroacetic acid. Centrifuge at 2,500 g for 10 minutes at room temperature, and transfer supernatant to a fresh tube.

Peptides may be stored for a short period of time at 4°C, but it is best to proceed to desalting immediately.

Peptide Desalting

Peptides must be desalted prior to TiO₂ phosphoenrichment. When using 200 mg tC18 SEP-Pak cartridges, each cartridge capacity is 5%, so up to 10 mg can be purified on a single 200 mg cartridge. Other capacities can be used for different sample volumes, with the wash steps below scaled proportionally by bed volume. Sample processing can be accomplished by gravity flow, or the use of a vacuum manifold.

- 32 Prepare the following solutions (see *Reagents and Solutions*):
- Desalting Wash Solution A, Desalting Wash Solution B, Desalting Elution Solution.
- 33 Condition the cartridge using 6 mL of acetonitrile followed by 2 mL of Desalting Elution Solution.
- 34 Equilibrate with 6 mL of Desalting Wash Solution A.
- 35 Load sample from step 31.
- 36 Wash/desalt with 6 mL of Desalting Wash Solution A.
- 37 Wash with 500 μ L Desalting Wash Solution B.
- 38 Elute into a clean 1.5 mL tube with 1 mL Desalting Elution Solution.
- 39 Measure protein concentration by NanoDrop A280 (or other protein estimation method) and determine total quantity of protein in sample. Aliquot 4 mg of peptides into a clean 1.5 mL tube.
- 40 Snap-freeze with liquid nitrogen and speed vac until the sample is a white (sometimes yellowish) pellet.
- Avoid over-drying pellet, as this will make it very difficult to resuspend. Samples can be stored at -20°C if TiO_2 phosphoenrichment cannot be performed immediately, but for best results proceed immediately to the next step.

Combined TiO_2 Phosphoenrichment and Terminal Desalting

Phosphopeptides will be isolated by bulk TiO_2 enrichment. Combining the TiO_2 and terminal desalting steps helps to prevent loss and/or carryover of beads. Commercial desalting tips may be purchased (Thermo Fisher Scientific cat. no. PI89873), however we routinely prepare StageTips in-house using 5 C18 plugs packed gently into a 200 μ L tip, according to Rappsilber et al., 2007. StageTips should be prepared during the TiO_2 incubation.

- 41 Prepare the following solutions (see *Reagents and Solutions*):
- TiO_2 Binding Solution, TiO_2 Elution Solution B, Desalting Wash Solution C, Desalting Elution Solution.
- 42 Add 1 mL TiO_2 Binding Solution to dried down peptides and allow to resolubilize for > 30 min. Transfer to 15 mL conical tube and add an additional 2.6 mL TiO_2 Binding Solution (final volume 3.6 mL TiO_2 Binding Solution).
- Due to the lactic acid, the TiO_2 Binding Solution can result in gloves becoming sticky. Gloves exposed to TiO_2 binding solution may stick to tubes.
- 43 Condition enough TiO_2 beads to allow for a 1:1 ratio of beads to peptides based on post-SEP-Pak NanoDrop reading (i.e. 4 mg beads for every 4 mg peptides).

- 44** Condition the beads by washing in 50X bead volume of TiO₂ Binding Solution (50 µL TiO₂ Binding Solution per 1 mg beads), centrifuge at 600 g for 30 seconds. Remove supernatant and repeat TiO₂ Binding Solution conditioning step.
- 45** Resuspend beads in the appropriate volume of TiO₂ Binding Solution to obtain a bead concentration of 10 µg/µL (e.g. resuspend 4 mg of beads in 400 µL of Binding Solution).
- 46** Add 400 µL TiO₂ beads to 4 mg of resolubilized peptides from step 42. Final peptide concentration is 1 mg/mL.

The TiO₂ beads settle rapidly. To avoid adding an incorrect volume of beads, resuspend the TiO₂ bead slurry by pipet mixing immediately before dispensing to each peptide sample.

- 47** To bind phosphopeptides, incubate in conical tube shaker at maximum speed and at room temperature for 1 hour.
- 48** Pellet beads by centrifugation at 600 g for 30 seconds, and remove supernatant. Be careful not to remove beads. Binding buffer supernatant and all subsequent washes may be saved as “non-phosphopeptides” for analysis if desired.
- 49** Wash beads with 1 mL TiO₂ Binding Solution.
- 50** Pellet beads and remove supernatant. Repeat wash two more times with fresh TiO₂ Binding Solution for a total of 3 washes.
- 51** Resuspend beads in 200 µL TiO₂ Binding Solution.

Perform all subsequent steps by centrifuging at 2,000 g for the minimum time required to pass the liquid through the StageTip (~30 sec/50 µL). In order to make sure StageTips do not over-dry, it may be necessary to centrifuge some StageTips longer than others.

- 52** Condition StageTips with 50 µL methanol.
- 53** Pre-clear StageTips by washing with 50 µL Desalting Elution Solution, and then equilibrate by washing twice with 50 µL TiO₂ Binding Solution.
- 54** Load the TiO₂ bead slurry onto the top of the StageTip and centrifuge, saving flow-through with any residual “non-phosphopeptides” if desired.
- 55** Wash the combined StageTips/TiO₂ column twice with 150 µL TiO₂ Binding Solution. This can be added to the flow-through if desired.
- 56** Equilibrate the combined StageTips/TiO₂ column with 100 µL Desalting Wash Solution C.
- 57** Elute phosphopeptides with 150 µL TiO₂ Elution Solution. Repeat once.
- At this stage, phosphopeptides will be retained on the C18 disc.
- 58** Wash with 100 µL Desalting Wash Buffer C.
- 59** Elute phosphopeptides off the disc with 100 µL Desalting Elution Solution.

- 60** Dry the eluent in speed vac, and store at -20°C until analysis by LC-MS/MS (see Support Protocol 1 for mass spectrometry acquisition and analysis procedures).

Phosphopeptide List Creation and Filtering

Prior to motif visualization, mass spectrometry data must be filtered for appropriate sequences and converted into a set of phosphorylation-centered 15mers. This is accomplished by removing undesirable peptide matches within the Phospho(STY) modification-specific file, and bioinformatically determining the *in vivo* 15-residue context centered around the phosphoacceptor.

- 61** Within the Phospho(STY) modification-specific file, remove any peptides that match to a reverse database, or are identified as a contaminant.

- 62** If additionally searching against the human proteome, remove any autophosphorylation sites on the kinase of interest.

While these sites are often of biological interest, autophosphorylation sites (particularly those that occur in *cis*) frequently do not conform to consensus motifs and therefore should be removed prior to motif analysis.

- 63** Filter phosphopeptides to only retain high-confidence sites with a localization probability of ≥ 0.9 . This is most easily accomplished by a text find-and-replace where “(1)” and “(0.9)” are replaced with “*”.

Note that this localization probability value is based on the software detailed in Support Protocol 1. If a different search algorithm is used (or parameters are changed), this value may need to be determined empirically.

Each site must be converted to a modification-centered 15mer, therefore any phosphorylation site that is too close to a protein terminus to be extended to a centered 15mer is discarded. This is the same procedure as in *motif-x* analyses (Schwartz and Gygi, 2005; Chou and Schwartz, 2011). We have created a web tool, PeptidExtender, which accomplishes this task, and is freely accessible at <http://schwartzlab.uconn.edu/pepextend>.

- 64** Paste the peptide sequences into the input box in the top left corner of the PeptidExtender webpage. This will cause “modification markers” to populate. Select “*” as the modification marker, select “right of modified residue” for position, and enter a target sequence width of 15. Select the “*E. coli*” proteome as the extension database. Click “extend peptides!” to create a list of unique phosphorylation-centered 15mers.

PeptidExtender automatically filters out non-selected potential markers (i.e. non-amino acid characters such as numbers and brackets), deletes redundant sequences, and removes any sequence that fails to generate a full 15mer. The output from PeptidExtender is correctly formatted to be

directly pasted into pLogo, although additional negative-control subtraction is necessary prior to motif visualization.

It is critical to remove endogenous *E. coli* phosphorylation sites from the foreground data set. We have curated a master negative control list (Lubner et al., 2017), which was generated by pooling phosphopeptides previously identified in negative control experiments (Chou et al., 2012), previously identified endogenous *E. coli* phosphorylation sites (Macek et al., 2008; Soares et al., 2013), and phosphorylation sites identified in empty vector and kinase dead negative control experiments. This list is available as Supplementary Table S1. Any additional sites identified in endogenous *E. coli* and negative control experiments should be added. Phosphorylation sites on this master negative control list must be removed from each target kinase data set to generate a final list of kinase-specific phosphorylation sites. We typically make use of the webtool Venny (Oliveros, J.C., 2007), which can be freely accessed at <http://bioinfogp.cnb.csic.es/tools/venny/>. After control subtraction, phosphorylation site lists from all runs can be merged within each kinase variant (redundant sites will be automatically removed by our software). These final data sets can be used for motif visualization by pLogo (see Support Protocol 2) or motif analysis by *motif-x* (Schwartz and Gygi, 2005; Chou and Schwartz, 2011).

ALTERNATE PROTOCOL

In Vitro Kinase Reaction

As mentioned above, there are several challenges associated with heterologous kinase expression in *E. coli*. However, many of the positive features of ProPeL are retained in an *in vitro* version of the protocol. Conceptually, the issues of kinase expression and/or activation are solved through purification of recombinant or endogenous kinase from an alternative source, and activation with the required co-factors. By adding recombinant kinase to *E. coli* lysate in a traditional *in vitro* kinase reaction, the target kinase is still able to phosphorylate bacterial proteins, which can be isolated as in the standard ProPeL workflow. This format still allows the high signal to noise ratio and direct link afforded by the low endogenous *E. coli* serine/threonine/tyrosine phosphorylation, and reactions still take place with full-length protein substrates. Identification by LC-MS/MS is the same, allowing high throughput identification, and the preservation of intra-motif correlations due to the proteomic context of each phosphopeptide.

There are a few different options for producing recombinant kinase. The target kinase can still be expressed in *E. coli*, and then affinity purified. This would be most useful in the case of re-constituting a kinase cascade, where the individual kinases can be purified separately by using distinct affinity tags. Alternatively, other host systems such as yeast, insect, or human cells may be employed to produce recombinant kinase. Finally, the use of cell-free protein systems allow for production of posttranslationally modified kinases without the need for cell culture (Oza et al., 2015). Optimal kinase reaction conditions need to be determined empirically.

Additional Materials

Untransformed or empty vector *E. coli* bacterial stock

Recombinant Kinase (purchase or purify in-house)
Appropriate Kinase Reaction Buffer (determine empirically)
Adenosine 5'-triphosphate (ATP) (Fisher Scientific cat. no. BP413-25)

Preparation of Bacterial Substrate Library

For the *in vitro* version of ProPeL, bacterial proteins still function as a substrate library. Using either an untransformed bacterial stock or an empty vector stock, cells are grown and harvested similarly to the Basic Protocol, but lysed in kinase reaction buffer.

- 1 Using a sterile pipette tip, streak a fresh plate from a bacterial glycerol stock. Incubate plate upside down at 37°C overnight.
- 2 Inoculate a well-isolated colony in 5 mL LB media (+ appropriate antibiotic) and grow at 37°C and 250 rpm overnight.
- 3 Inoculate 100 mL LB media (+ appropriate antibiotic) with 2 mL of overnight culture in a 500 mL Erlenmeyer flask.
- 4 [OPTIONAL] Grow bacteria to an OD₆₀₀ of 0.4–0.6 (mid-log) and induce with 0.5 mM IPTG.
- 5 Incubate at 37°C and 250 rpm for 3–24 hours.
- 6 Pellet cells by centrifugation at 6,000 g for 15 minutes at 4°C.
- 7 Pellet may be stored at –80°C, but it is best to proceed promptly.

When preparing sample for LC-MS/MS, use all LC/MS grade solvents and Eppendorf brand microcentrifuge tubes for sample preparation. If evaluating an aliquot during the kinase expression optimization phase, ACS grade reagents are acceptable. ACS grade solvents may be used for all SDS-PAGE steps.

- 8 Prepare Kinase Reaction Buffer, add at 2 mL/g of wet pellet, and resuspend by pipet mixing.
- 9 Lyse cells by sonication, using 15-second pulses on 15% power, until solution is no longer opaque.

To prevent cells over-heating, tubes should be kept on ice (in between and during sonication) with at least 1 minute rest between pulses.
Solution will be discolored, but should be clear.
- 10 Clarify solution by centrifugation at 20,000 g for 30 minutes at 4°C. Transfer clarified supernatant to a fresh tube and discard pellet. If necessary, repeat centrifugation to further clarify.
- 11 Filter lysate with a disposable syringe and 0.22 µm filter attachment to further remove cellular debris.
- 12 Quantify samples by BCA assay.

***In Vitro* Kinase Reaction and Evaluation of Kinase Activity**

In the Alternate Protocol, recombinant kinase is incubated with bacterial lysate in an *in vitro* kinase reaction. Optimal reaction conditions should be determined empirically using small-scale reactions before proceeding to the full 10 mg reaction. After this section, sample preparation for LC-MS/MS is resumed as in the Basic Protocol.

- 13** Aliquot 10 mg bacterial lysate to a 1.5 mL tube.
- 14** Add appropriate volume recombinant kinase and 1X kinase reaction buffer, and incubate 3 hours at 30°C.

This is an optimization point. Appropriate buffer conditions, kinase:substrate ratio, incubation duration and temperature may need to be determined empirically.
- 15** For each sample, separate 25 µg (or 75 µg if using NanoDrop A280 measurement) by SDS-PAGE, with 2 µL PeppermintStick Phosphoprotein ladder (and 5 µL All Blue Protein Standards, *optional*).
- 16** Analyze with Pro-Q Diamond stain, according to manufacturer's instructions. All incubations should be carried out on a rocker.
- 17** Immerse gel in 100 mL fix solution (see *Reagents and Solutions*) and incubate for 30 minutes. Discard the fix solution and add fresh 100 mL fix solution. Incubate for at least 30 minutes.

This is a pause point, as gel can be left in fix solution overnight.
- 18** Discard the fix solution and wash with 100 mL ultrapure water. Incubate for 10 minutes, discard, and repeat twice for a total of 3 water washes.

All subsequent incubations must be done in the dark, as Pro-Q Diamond is light sensitive.
- 19** Add 60 mL Pro-Q Diamond stain and incubate 90 minutes.
- 20** Discard stain and add 90 mL Pro-Q Diamond destain solution (see *Reagents and Solutions*), incubating for 30 minutes. Discard destain and repeat twice more for a total of 3 destain washes. Rinse with ultrapure water for 5 minutes, discard and repeat water wash once.
- 21** Visualize on appropriate imager (Typhoon, ChemiDoc etc.) using the following wavelengths: Ex: 555 nm, Em: 580 nm. Adjust the signal such that only the two phosphoprotein bands (23.k kDa and 40 kDa) on the PeppermintStick ladder are clearly visible (18 kDa band may be faintly visible).

To control for loading differences (which can change the level of background signal), it is important to perform a total protein stain. We use GelCode Blue according to manufacturer's instructions, but other stains (such as Coomassie staining) are acceptable.

- 22** Add 20 mL GelCode Blue and incubate on a rocker at room temperature for at least an hour. Preferably, leave gel in GelCode Blue overnight for clearest signal.

- 23 Destain using ultrapure water. For best results, change water several times until background signal has been completely removed. Gel may be left in water overnight.
- 24 Image using the Coomassie setting (and white light conversion screen).

If phosphorylation is acceptable, proceed with the remaining steps for LC-MS/MS sample preparation in the Basic Protocol, beginning with ***Protein Reduction, Alkylation and Tryptic Digestion***. Otherwise, optimize kinase activity before proceeding.

SUPPORT PROTOCOL 1

Phosphopeptide Identification by Liquid Chromatography-Tandem Mass Spectrometry

The following protocol represents our current instrumentation, and is provided as a reference.

Instrumentation

Q Exactive HF Orbitrap mass spectrometer (Thermo Scientific)

Ultimate 3000 RSLC (Thermo Scientific)

250 mm nanoEase M/Z peptide BEH C18 column

130Å, 1.7 µm particle size, 75 µm i.d (Waters cat. no. 186008795)

10 µm silica PicoTip emitter (New Objective, Inc. cat. no. FS360-20-10-N-20-C12)

Liquid Chromatography and Mass Spectrometry Analysis

Dried and enriched phosphopeptides are resuspended in 40 µL 0.1% formic acid in water. Peptide identification is achieved using electrospray ionization (ESI) and nanoLC-MS/MS on a Q Exactive HF Orbitrap mass spectrometer (QE-HF) coupled to an Ultimate 3000 RSLC operated in nanoflow mode. A 250 mm nanoEase M/Z peptide BEH C18 column is fitted to a 10 µm silica PicoTip emitter to permit ESI directly into the QE-HF inlet. For all samples, 1 µL is loaded and subject to a 150 min, 300 nL/min linear reversed phase gradient (Solvent A: 0.1% formic acid in water, Solvent B: 0.1% formic acid in acetonitrile) as follows: initial 4% solvent B hold for 10 min, increase to 30% solvent B over 90 min, increase to 90% solvent B over 20 min, 90% solvent B hold for 10 min, then decrease to 4% solvent B over 2 min, followed by a re-equilibration period for 18 min. Column temperature is set to 50°C for the entire gradient. The QE-HF is operated in positive ion mode with a spray voltage of 1.5 kV. The capillary temperature is set to 250°C and all source gas flows are turned off. A Top 15 data-dependent (dd) MS/MS method is used that implements the following parameters for full MS scans: 1 microscan, 60,000 resolution at 200 m/z, 1e6 AGC target, 60 ms max ion time, and 300 to 1800 m/z mass range. MS/MS scans are acquired with the following parameters: 15,000 resolution at 20 m/z, 1e5 AGC target, 40 ms max IT time, 2.0 m/z isolation window, 0 m/z isolation offset, 200 to 2000 m/z mass range, 27 normalized collision energy, peptide match “preferred,” exclude isotopes “on,” a 30 s dynamic exclusion window, and charge state exclusion set to exclude +1 and >+8 ions. All spectral data are collected in profile mode. A QC analysis of tryptic BSA peptides (optional)

is analyzed between each ProPeL injection to minimize sample carryover and gauge instrument performance stability. Similar instrument parameters are used but a shorter, 60 min gradient is performed in place of the 120 min gradient previously described.

Phosphopeptide Identification and Site Localization

Raw files are searched with the Andromeda search engine and MaxQuant (Cox and Mann, 2008) against the UniProt *E. coli* strain B/BL21-DE3 proteome database (Proteome ID UP000002032). A reversed *E. coli* protein sequence database is automatically generated by the software and searched concurrently. All searches include the following parameters: 4.5 and 20 ppm mass tolerances for precursor and fragment ions, respectively, trypsin enzyme specificity, up to 2 missed cleavages, fixed carbamidomethyl C modification, variable phosphorylation of serine/threonine/tyrosine, oxidation of methionine, and acetylation of protein N-termini. Minimum peptide length is set to 5 and the contaminant database is included. All results are filtered at a 1% false discovery rate at the peptide spectrum match, protein and site levels. All other parameters are kept at MaxQuant default settings (version 1.6.0.1 at time of publication).

SUPPORT PROTOCOL 2

Motif Visualization with pLogo

pLogos (O'Shea et al., 2013) depict residues proportional to the log-odds of their binomial probabilities with respect to a given background. In a pLogo, the most statistically significant residues appear closest to the x-axis, with residues above the x-axis indicating overrepresentation and those below the x-axis indicating underrepresentation. Given the existence of one or more different residues at a given substrate position, it is possible to compute conditional probabilities of all remaining amino acids and positions to determine significant positions *given* specific residues at specific positions. We refer to this as “fixing” a given residue at a given position, which allows the exploration of correlated or uncorrelated residues across positions in the kinase specificity motif. Fixed positions within the pLogo (e.g., the central position) are depicted on a grey background, and red horizontal lines denote the $p = 0.05$ significance threshold (after Bonferroni correction). pLogos can be scaled for clarity. For each pLogo, the foreground data is the list of phosphorylation-centered 15mers, with negative control sites removed. The *E. coli* background data set is generated by pLogo through alignment of all unique phosphoacceptor-centered 15mers in the *E. coli* proteome. Below are basic instructions for generating pLogos:

1. Access the pLogo website (and register for an account if desired): <http://plogo.uconn.edu>
2. Paste desired foreground data set of aligned 15mers into the box on the left of the page.
3. Select “Protein” and then “e. coli k12” from the available backgrounds on the right of the page.
4. [OPTIONAL] If logged into a personal account, the user may add a job name.
5. Click the “generate pLogo” button in the center of the page.

6. Residues can be fixed or unfixed by clicking on them; however, residues that do not achieve statistical significance may not be fixed. Alternatively, users may fix significant residues by checking the corresponding box in the “statistics” tab to the left of the pLogo.
7. The zoom can be changed by clicking the “customize” tab, and either hitting the +/- buttons, or entering a value. Clicking “renormalize” will rescale the pLogo back to its default size.

For additional functionality and explanations, see:

O’Shea JP, Chou MF, Quader SA, Ryan JK, Church GM, & Schwartz D. (2013). pLogo: A probabilistic approach to visualizing sequence motifs. *Nat Methods* 10, 1211–1212.

REAGENTS AND SOLUTIONS

LB (Lysogeny Broth) Solid Plates (500 mL)

5 g Tryptone (Sigma-Aldrich cat. no. 61044-1KG)

2.5 g Yeast Extract (Sigma-Aldrich cat. no. 09182-1KG-F)

5 g NaCl

7.5 g Agar (Fisher Scientific cat. no. BP1423-500)

Water (double distilled or Ultrapure)

Add dry reagents in 1 L Pyrex media bottle, and add water to final volume 500 mL and mix thoroughly. It is not necessary to adjust the pH. Autoclave on liquid cycle, then cool autoclaved media in a water bath set to 55°C. Once cooled, add appropriate antibiotic if desired, swirly gently or rotating end-over-end to mix. Pour media to ~0.5” depth into sterile petri dishes. Prop lid (to minimize dust) and allow plates to cool in sterile area. Once cool, store plates in plastic sleeve at 4°C up to 3 months. Discard any plates with visible growth.

If adding antibiotic, wait until liquid media has cooled to 55°C. Adding antibiotic while the liquid media is still hot may cause the antibiotic to degrade.

LB (Lysogeny Broth) Liquid Media (1 L)

10 g Tryptone (Sigma-Aldrich cat. no. 61044-1KG)

5 g Yeast Extract (Sigma-Aldrich cat. no. 09182-1KG-F)

10 g NaCl

Water (double distilled or Ultrapure)

Add dry reagents in 1 L Pyrex media bottle, and add water to approximately 950 mL and mix thoroughly. Adjust pH to 7.5 with NaOH, then add water to bring to 1 L final volume. Split between 2 Pyrex media bottles (<600 mL per bottle) and autoclave on

liquid cycle. Once cooled, combine bottles and store at 4°C up to 6 months. Discard media if it appears cloudy.

If adding antibiotic, wait until liquid media has cooled to 55°C. Adding antibiotic while the liquid media is still hot may cause the antibiotic to degrade.

Lysis Buffer (10 mL)

500 µL 1 M Tris-HCl (pH 8.2)

750 µL 1 M NaCl

4.80 g Urea

100 µL 100X Protease Inhibitor

100 µL 100X Phosphatase Inhibitor

100 µL 100 mM phenylmethylsulfonyl fluoride

Water (HPLC/MS Grade)

Add Tris-HCl, NaCl, and urea to a clean falcon tube. Add water to approximately 9.5 mL and mix thoroughly. Add Protease and Phosphatase Inhibitors, and PMSF immediately prior to use. Add water to bring to 10 mL final volume, and store at room temperature.

Make buffer fresh daily.

Laemmli Loading Buffer, 6X (10 mL)

1.2 g Sodium Dodecyl Sulfate (SDS) (Fisher Scientific cat. no. BP166-500)

6 mg Bromophenol Blue (Fisher Scientific cat. no. 403160050)

4.7 mL Glycerol (Promega cat. no. H5433)

1.2 mL Tris-HCl (pH 6.8)

0.93 g Dithiothreitol (DTT)

2.1 mL Water (double distilled or Ultrapure)

Add all reagents (except DTT) to a 15 mL conical tube and mix thoroughly. Solution may be heated to 37°C to aid solubilization (which may take a period of several hours). Once fully dissolved, add DTT and mix thoroughly. Aliquot and store at -20°C for up to 1 year.

SDS-PAGE Running Buffer, 10X

30 g Tris Base

144 g Glycine

10 g SDS

Water (double distilled or Ultrapure)

Add reagents and water to a final volume of 1 L, and mix thoroughly. It is not necessary to adjust the pH (which should be pH 8.3), and can be stored at room temperature for up to 1 year.

SDS-PAGE Gel

12% Separating Gel

3.4 mL Water (double distilled or Ultrapure)

4 mL Bis/Acrylamide (37.5:1) (Bio-Rad cat. no. 1610158)

2 mL 1.87 M Tris-HCl (pH 8.9)

100 µL 10% SDS Solution

5 µL TEMED (Fisher Scientific cat. no. BP150-20)

0.5 mL Ammonium Persulfate (15 mg/mL solution) (Fisher Scientific cat. no. BP179-100)

3.5% Stacking Gel

0.826 mL Water (double distilled or Ultrapure)

0.232 mL Bis/Acrylamide

0.4 mL 0.312 M Tris-HCl (pH 6.7)

20 µL 10% SDS Solution

1 µL TEMED

0.533 mL Ammonium Persulfate (15 mg/mL solution)

Gel Casting Stand and Clamps (Bio-Rad cat. no. 1658050)

Glass Short Plates (Bio-Rad cat. no. 1653308)

Glass Spacer Plates 1.5 mm (Bio-Rad cat. no. 1653312)

10-well comb (Bio-Rad cat. no. 1653365) or 15-well comb (Bio-Rad cat. no. 1653366)

Assemble casting stand and plates. Mix *separating gel* reagents gently (avoid introducing bubbles), and pour separating gel until the level reaches approximately 1" below the top of the cassette. Layer isobutanol (use a 1:1 ratio of isobutanol:water, and use only the top, less dense layer) on top of the separating gel to prevent bubbles and level the gel. Allow gel to polymerize for approximately 30 minutes. Pour off isobutanol and rinse with water. Mix *stacking gel* reagents gently (avoid introducing bubbles), and pour stacking gel to the top of the cassette. Insert an appropriate comb and allow gel to polymerize for approximately 30 minutes. Gels can be wrapped in plastic wrap and stored at 4°C for several days.

Fix Solution (1 L)

700 mL Water (double distilled or Ultrapure)

500 mL Methanol

100 mL Glacial Acetic Acid

Add liquid reagents in 1 L Pyrex media bottle, mix thoroughly, and store at room temperature up to 1 year.

Pro-Q Diamond Destain Solution (1 L)

750 mL Water (double distilled or Ultrapure)

200 mL Acetonitrile

100 mL 1 M Sodium Acetate (pH 4.0)

Add liquid reagents in 1 L Pyrex media bottle, mix thoroughly, and store at room temperature up to 1 year.

Desalting Wash Solution A (100 mL)

99.9 mL Water (HPLC/MS Grade)

100 μ L Trifluoroacetic Acid (HPLC/MS Grade)

Add liquid reagents in 100 mL Pyrex media bottle, mix thoroughly, and store at room temperature.

Desalting Wash Solution B (50 mL)

49.75 mL Water (HPLC/MS Grade)

250 μ L Acetic Acid (HPLC/MS Grade)

Add liquid reagents in 50 mL Pyrex media bottle, mix thoroughly, and store at room temperature.

Desalting Wash Solution C (50 mL)

49.5 mL Water (HPLC/MS Grade)

500 μ L Formic Acid (HPLC/MS Grade)

Add liquid reagents in 50 mL Pyrex media bottle, mix thoroughly, and store at room temperature.

Desalting Elution Solution (50 mL)

24.75 mL Water (HPLC/MS Grade)

25 mL Acetonitrile (HPLC/MS Grade)

250 μ L Acetic Acid (HPLC/MS Grade)

Add liquid reagents in 50 mL Pyrex media bottle, mix thoroughly, and store at room temperature.

TiO₂ Binding Solution (100 mL)

33.44 mL Water (HPLC/MS Grade)

50 mL Acetonitrile (HPLC/MS Grade)

16.56 mL Lactic Acid (90%, HPLC/MS Grade)

Add liquid reagents in 100 mL Pyrex media bottle, mix thoroughly, and store at room temperature.

The lactic acid makes the solution very sticky. Gloves exposed to TiO₂ binding solution may stick to tubes.

TiO₂ Elution Solution (50 mL)

0.34 g KH₂PO₄

Water (HPLC/MS Grade)

Add dry reagents in 50 mL Pyrex media bottle. Add water to approximately 45 mL and mix thoroughly. Adjust pH to 10 with NaOH. Add water to bring to 50 mL final volume, mix thoroughly, and store at room temperature.

COMMENTARY**Background Information**

There are several existing strategies for determining kinase specificity motifs. The simplest approach is to align experimentally determined phosphorylation sites for a given kinase on the phosphoacceptor, and extract motifs bioinformatically. However, due to the complex, overlapping nature of kinase signaling cascades, it has been difficult to unambiguously pair identified phosphorylation sites with their upstream kinase. As a result of this bottleneck, several strategies have been developed for determining protein kinase substrate specificity motifs. Early success came from the use of an oriented peptide library, in a strategy pioneered by Cantley and colleagues (Songyang et al., 1994). This method utilized a synthetic peptide library of 2.5 billion distinct sequences, which all contained a single phosphoacceptor in the center of the peptide flanked by 4 variable positions upstream and downstream, and linker residues at the termini. The kinase of interest was incubated with the library and ³²P-ATP in an *in vitro* kinase reaction. Phosphopeptides were then separated by ferric iminodiacetic acid, and sequenced by Edman degradation. Observed residue frequencies in each position were compared to their respective abundance (obtained by sequencing the peptide library) to identify important determinants. While successful, this approach has several drawbacks. Notably, identification by Edman degradation is laborious and time-consuming, and technical limitations prevent querying tryptophan or cysteine (which interfere with sequencing due to oxidation), or additional phosphoacceptors (which would render the phosphorylation site ambiguous). In order to eliminate these challenges, the technique was refined by Turk and colleagues to a matrix format. Peptide libraries are arranged in a grid, with a single fixed residue/position for each well (representing all the possible amino acids in each position), while the other positions are randomized. After incubation with the kinase and ³²P-ATP, the reactions are spotted onto a membrane, and

phosphorylation preferences are read by measuring radioactive incorporation at different fixed residue positions (Hutti et al., 2004). While these methods provide a wealth of data, the combinatorial libraries are prohibitively expensive for most labs. Additionally, they require the use of radioactivity and large amounts of recombinant kinase. Importantly, these approaches use peptide substrates (which are less physiologically relevant than protein substrates), and there is no ability to discern correlation between motif positions.

The next milestone came with the utilization of depleted cell lysates, which could function as a proteomic library. Crucially, a proteomic library (unlike a random library) is amenable to sequencing by tandem mass spectrometry. This approach was first demonstrated by Huang and colleagues (Huang et al., 2007), wherein rat uterus homogenate was fractionated by Strong Anion Exchange (SAX), treated with phosphatases (to remove endogenous phosphorylation), and heated to inactivate endogenous kinases and all phosphatases. The resulting fractions were incubated with recombinant kinase and *cold* ATP to allow the kinase of interest to phosphorylate the depleted cellular protein fractions (in the absence of ^{32}P -ATP). The reaction mixture was digested with trypsin and phosphoenriched by IMAC, followed by identification by tandem mass spectrometry. Other versions of this approach inactivated endogenous kinases by tryptic digestion prior to SCX fractionation and dephosphorylation (Kettenbach et al., 2012) or by the addition of the irreversible ATP-competitive analog 5'-4-fluorosulphonylbenzoyl adenosine (FSBA) (Knight et al., 2012). While the use of a proteomic library (and therefore the ability to identify sites by tandem mass spectrometry) is a significant advantage, these methods still require large amounts of recombinant kinase, run the risk of contamination in the event of incomplete dephosphorylation or residual endogenous kinase activity, and in some cases require that specificity preferences be queried by peptide (rather than protein) substrates.

The ProPeL approach described in this article has several important benefits and advantages over existing strategies. As the phosphorylation reactions occur *in vivo* in the *E. coli* cytoplasm, the living host produces both the kinase and substrate proteins, and regulates the environment (pH, ionic and co-factor concentrations, etc.). This obviates the need to purify catalytically active kinase, and means that the target kinase interacts with substrates under conditions that are more physiologically relevant than an *in vitro* reaction. The substrates themselves are full-length *E. coli* proteins, which offer two distinct advantages. First, the kinase reaction occurs with phosphoacceptors within protein substrates that are able to adopt a physiologically appropriate folded structure, rather than peptides that may not fully recapitulate the environment surrounding the phosphoacceptor. Second, a proteomic background allows unambiguous sequence and site identification using tandem mass spectrometry. This not only allows for high-throughput sequencing, but also puts each phosphorylation site in sequence context, which allows for intra-motif correlation between positions. As direct phosphorylation of bacterial substrates by the kinase of interest is measured, there is no need for radioactive material (typically ^{32}P -ATP), making ProPeL safer than traditional approaches. The actual reaction and sample preparation are also significantly cheaper than combinatorial peptide library approaches, although access to a mass spectrometer is required.

Critical Parameters

The success of a ProPeL experiment crucially depends on the ability to express soluble, active kinase to facilitate the *in vivo* phosphorylation of bacterial substrates. While the majority of eukaryotic proteins expressed in *E. coli* were easily purified in their correctly folded state (Braun et al., 2002), there are several factors that can affect heterologous protein expression (Dyson et al., 2004). It is reasonable to expect that expressing different foreign kinases within *E. coli* will lead to differential exogenous phosphorylation, necessitating different optimal expression conditions to maximize *in vivo* phosphorylation of bacterial substrates by each kinase. There are several variables that can be adjusted to improve *in vivo* expression and activity.

Choosing the Correct *E. coli* Cell Strain—In our hands, the single variable that has impacted target kinase expression the most has been selecting the correct cell strain. *E. coli* expression of the 61 tRNAs is different from eukaryotic species, which can hinder heterologous protein expression for transcripts that are rich in codons that are under-utilized in *E. coli* (Kane, 1995). This can be ameliorated by exhaustive codon optimization of the target kinase coding sequence, or by the usage of so-called codon-optimized strains. For example, the Rosetta2 strains supply tRNAs for the codons AUA, AGG, AGA, CUA, CCC, GGA, and CGG. Over-expression of these tRNAs can improve expression of the target kinase.

Alternatively, expression of the target kinase may be cytotoxic, preventing any cells that are competent to express the protein from surviving. A commonly used strategy is to switch to a resistant host strain, such as the C41(DE3) and C43(DE3) “Walker” strains, which exhibit elevated ability to express membrane-bound and toxic proteins (Miroux and Walker, 1996). These strains were later characterized as having mutations in the *lacUV5* promoter (which controls expression of T7 RNA polymerase) and therefore exhibited more gradual target protein expression (Wagner et al., 2008).

Typically, we start by expressing a new plasmid in both C41(DE3) and Rosetta2 cells. If the target kinase expresses in the C41(DE3) strain, it implies that codon usage is not a concern, and typically we observe more robust expression in C41(DE3) strains than in Rosetta2. However, when codon utilization appears to be an issue, expression is often *only* successful in Rosetta2, with no expression in C41(DE3). We have also begun to use C43(DE3) cells with some success for kinases that failed to express in C41(DE3).

Protein Induction and Growth Conditions—Protein expression is commonly achieved using the inducible T7 promoter pET system, whereby induction is initiated by addition of an allolactose analog, usually isopropyl β -D-1-thiogalactopyranoside (IPTG) (Donovan et al., 1996). Controlling the concentration of IPTG used and the temperature and duration of induction is critical for optimizing target kinase expression. Typical IPTG concentrations range from 0.1 – 1 mM, and while increasing concentrations can lead to faster protein production, this has the danger of overwhelming the cells and causing cytotoxicity. Similarly, while a longer induction may allow more time for kinase expression, this can be detrimental if the kinase hinders growth.

While IPTG induction is very successful in many instances, induction may overwhelm the cell, inhibiting further protein expression and in some cases killing plasmid-bearing cells. As an alternative to IPTG induction, protein expression can be accomplished using the autoinduction system, which uses a complex media containing both glucose and lactose. The *E. coli* preferentially metabolizes the glucose, growing to high density while target protein expression is suppressed by glucose. As the glucose is depleted the cells switch to lactose, producing allolactose, and inducing target protein expression in robust, high density cultures (Studier, 2005).

Enhancing Protein Solubility—ProPeL relies upon the ability of the target kinase to phosphorylate bacterial proteins *in vivo*. Therefore, it is critical that the kinase is not only expressed, but is also soluble during expression. The major challenge is to prevent the over-expressed kinase from being sequestered in inclusion bodies – insoluble aggregates of mis-folded proteins (Schein, 1989). Inclusion bodies form when exposed, hydrophobic stretches of insoluble, mis-folded, or partially folded proteins stick together via intermolecular β -sheet structures (Fink, 1998). The dynamics of inclusion body formation are such that there are typically a small number of inclusion bodies seeded by an incorrectly folded protein intermediate. Although various studies have suggested refolding strategies both *in vivo* (Zhao et al., 2012) and *in vitro* (Santos et al., 2012), the consensus is that it is critical to prevent inclusion body formation from ever occurring to maximize yield (Schein, 1989; Fink, 1998).

Within the context of ProPeL, there are several strategies that can be attempted for increasing solubility. At the start of a new ProPeL project, it is important to design the correct insert. Reducing the size of the target kinase by only expressing the catalytic domain (if appropriate) will help increase solubility, as lower molecular weight proteins tend to exhibit superior soluble expression (Dyson et al., 2004). The addition of fusion-protein tags such as 6XHis, GST, and MBP may also increase solubility, although there are no universal rules for how a tag may help or hinder solubility, and the inclusion of a tag may interfere with kinase function (Rosano and Ceccarelli, 2014; Guerrero et al., 2015). Lowering the temperature during induction has also been demonstrated to improve protein solubility (Baldwin, 1986). Finally, co-expressing the target protein with molecular chaperones such as the GroELS or DnaK system is a common strategy to aide protein folding (Nishihara et al., 2000; Marco et al., 2007). Unfortunately, to date this step has provided the most significant challenge for ProPeL, with limited success in the cases of protein kinases that appear insoluble.

Troubleshooting

In troubleshooting *in vivo* activity, it is advantageous to first identify whether the issue is protein expression, protein solubility, or kinase activity. The first question that must be answered is a relatively simple one: Is the target kinase expressed? Kinase expression is easily evaluated by lysing an aliquot in the standard denaturing lysis buffer and evaluating expression by SDS-PAGE and Coomassie staining. An example of the desired level of kinase expression is provided in Figure 2. It is also possible to evaluate expression by western blotting, but this may only be necessary if the expected kinase molecular weight is

the same as highly expressed endogenous *E. coli* proteins. If the kinase is not sufficiently expressed as to be easily detected by Coomassie staining, then it is likely not expressed sufficiently to provide adequate *in vivo* activity. It is also worth noting that expression patterns for endogenous *E. coli* proteins change with expression conditions, so it is important to run negative controls (i.e. a kinase dead mutant, see *Strategic Planning*) for each expression condition to generate an accurate background for comparison. If there is no detectable expression, this may indicate one of several problems. Strategies for overcoming codon bias, cytotoxicity, and solubility have been discussed above in the Critical Parameters section. Considerations for kinase activation requirements have been discussed in the Strategic Planning section, and should be used to inform the choice between the Basic Protocol and Alternate Protocol.

Anticipated Results

In our experience, the number of unique phosphorylation sites discovered for a particular kinase of interest most strongly correlates with the level of expression and *in vivo* activity. With the instrumentation setup described in this protocol, we have observed that the total number of unique phosphorylation sites identified in a single run can range from 200–500 sites for one kinase to over 1,500 sites for a different kinase. Technical replicates significantly increase detected phosphopeptides (Ham et al., 2008), although in our experience 2–3 technical replicates are typically sufficient.

Kinase motifs can at times be highly specific, where precise residue positions are favored, presumably due to the unique steric considerations, angles, and electrostatic properties associated with the individual residues. These motifs are frequently represented as consensus sequences, which offer the idealized version of the motif. A classic example is the PKA motif, which today we can summarize as the consensus sequence [R/K][R/K]x[S/T] ϕ , where [R/K] indicates a preference for either arginine or lysine, ϕ represents a hydrophobic amino acid, and “x” indicates no preference. While the PKA motif is well-defined, specificity motifs can also be more general, such as simple recognition of charge or hydrophobicity, and with positional flexibility. It is important to consider motif results holistically, allowing for both specific and general preferences when interpreting the pLogo for a given kinase. It is also informative to evaluate phosphoacceptor motifs separately – often there can be subtle differences between preferences surrounding serine compared with threonine phosphoacceptors, even for the same kinase.

Time Considerations

The Basic Protocol takes approximately 1 week to complete on average. Starting with an overnight bacterial starter culture, expression typically takes up to a day. Cell harvesting, lysis, quantification and SDS-PAGE analysis occurs on day 2, and typically takes a full day. The most efficient workflow is to perform a methanol/chloroform extraction at the end of the day, and allow the protein disc to resolubilize overnight. The tryptic digestion should not proceed longer than 16 hours, so we typically begin reduction and alkylation steps around 2 pm on day 3. Tryptic digestion occurs overnight. SEP-Pak peptide desalting, lyophilization, and TiO₂ enrichment can be accomplished in a single day, but we often store lyophilized peptides (pre-TiO₂) at –20°C overnight. Analysis by LC-MS/MS takes approximately 4

hours for each run including the BSA QC standard analysis. MaxQuant and Andromeda processing of the raw spectra typically takes approximately 0.5–3 hours, depending on the number of simultaneous searches and number of threads available on the server, while data filtering and pLogo analysis can be completed in under an hour. There are several pause points that are noted throughout the protocol. We have successfully stored cell pellets at -80°C for several months, and have stored lyophilized peptides (pre- TiO_2) at -20°C for several months. Cell lysate can be stored at 4°C for a short period (no more than a few days).

Acknowledgments

This work was supported in whole or in part by grants awarded to G.M.C. from the Department of Energy (DE-FG02-02ER63445), and to D.S. from the University of Connecticut Research Foundation, the University of Connecticut Office of the Vice President for Research, and the National Institute of Neurological Disorders and Stroke (1R21NS096516).

Literature Cited

- Baldwin RL. Temperature dependence of the hydrophobic interaction in protein folding. *Proceedings of the National Academy of Sciences*. 1986; 83:8069–8072.
- Braun P, Hu Y, Shen B, Halleck A, Koundinya M, Harlow E, LaBaer J. Proteome-Scale Purification of Human Proteins from Bacteria. *Proceedings of the National Academy of Sciences of the United States of America*. 2002; 99:2654–2659. [PubMed: 11880620]
- Chou MF, Prisis S, Lubner JM, Church GM, Husson RN, Schwartz D. Using Bacteria to Determine Protein Kinase Specificity and Predict Target Substrates. *PLoS ONE*. 2012; 7:e52747. [PubMed: 23300758]
- Chou MF, Schwartz D. Biological sequence motif discovery using motif-x. *Current Protocols in Bioinformatics/Editorial Board, Andreas D. Baxevanis ... [et Al.]*. 2011; Chapter 13(Unit 13):15–24.
- Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nature Biotechnology*. 2008; 26:1367–1372.
- Donovan RS, Robinson CW, Glick BR. Review: optimizing inducer and culture conditions for expression of foreign proteins under the control of the lac promoter. *Journal of Industrial Microbiology*. 1996; 16:145–154. [PubMed: 8652113]
- Dyson MR, Shadbolt SP, Vincent KJ, Perera RL, McCafferty J. Production of soluble mammalian proteins in *Escherichia coli*: identification of protein features that correlate with successful expression. *BMC Biotechnology*. 2004; 4:32. [PubMed: 15598350]
- Fink AL. Protein aggregation: folding aggregates, inclusion bodies and amyloid. *Folding and Design*. 1998; 3:R9–R23. [PubMed: 9502314]
- Gibbs CS, Zoller MJ. Rational scanning mutagenesis of a protein kinase identifies functional regions involved in catalysis and substrate interactions. *Journal of Biological Chemistry*. 1991; 266:8923–8931. [PubMed: 2026604]
- Guerrero F, Ciragan A, Iwai H. Tandem SUMO fusion vectors for improving soluble protein expression and purification. *Protein Expression and Purification*. 2015; 116:42–49. [PubMed: 26297996]
- Ham BM, Yang F, Jayachandran H, Jaitly N, Monroe ME, Gritsenko MA, Livesay EA, Zhao R, Purvine SO, Orton D, et al. The Influence of Sample Preparation and Replicate Analyses on HeLa Cell Phosphoproteome Coverage. *Journal of Proteome Research*. 2008; 7:2215–2221. [PubMed: 18412383]
- Hanks SK, Quinn AM, Hunter T. The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science*. 1988; 241:42–52. [PubMed: 3291115]
- Hansen AM, Chaerkady R, Sharma J, Díaz-Mejía JJ, Tyagi N, Renuse S, Jacob HKC, Pinto SM, Sahasrabudhe NA, Kim MS, et al. The *Escherichia coli* Phosphotyrosine Proteome Relates to Core Pathways and Virulence. *PLOS Pathogens*. 2013; 9:e1003403. [PubMed: 23785281]

- Huang SY, Tsai ML, Chen GY, Wu CJ, Chen SH. A Systematic MS-Based Approach for Identifying in vitro Substrates of PKA and PKG in Rat Uteri. *Journal of Proteome Research*. 2007; 6:2674–2684. [PubMed: 17564427]
- Hutti JE, Jarrell ET, Chang JD, Abbott DW, Storz P, Toker A, Cantley LC, Turk BE. A rapid method for determining protein kinase phosphorylation specificity. *Nature Methods*. 2004; 1:27–29. [PubMed: 15782149]
- Kane JF. Effects of rare codon clusters on high-level expression of heterologous proteins in *Escherichia coli*. *Current Opinion in Biotechnology*. 1995; 6:494–500. [PubMed: 7579660]
- Kettenbach AN, Wang T, Faherty BK, Madden DR, Knapp S, Bailey-Kellogg C, Gerber SA. Rapid Determination of Multiple Linear Kinase Substrate Motifs by Mass Spectrometry. *Chemistry & Biology*. 2012; 19:608–618. [PubMed: 22633412]
- Khokhlatchev A, Xu S, English J, Wu P, Schaefer E, Cobb MH. Reconstitution of Mitogen-activated Protein Kinase Phosphorylation Cascades in Bacteria EFFICIENT SYNTHESIS OF ACTIVE PROTEIN KINASES. *Journal of Biological Chemistry*. 1997; 272:11057–11062. [PubMed: 9110999]
- Knight JD, Tian R, Lee RE, Wang F, Beauvais A, Zou H, Megeney LA, Gingras AC, Pawson T, Figeys D, et al. A novel whole-cell lysate kinase assay identifies substrates of the p38 MAPK in differentiating myoblasts. *Skeletal Muscle*. 2012; 2:5. [PubMed: 22394512]
- Lubner JM, Dodge-Kafka KL, Carlson CR, Church GM, Chou MF, Schwartz D. Cushing's syndrome mutant PKAL205R exhibits altered substrate specificity. *FEBS Letters*. 2017; 591:459–467. [PubMed: 28100013]
- Macek B, Gnad F, Soufi B, Kumar C, Olsen JV, Mijakovic I, Mann M. Phosphoproteome Analysis of *E. coli* Reveals Evolutionary Conservation of Bacterial Ser/Thr/Tyr Phosphorylation. *Molecular & Cellular Proteomics*. 2008; 7:299–307. [PubMed: 17938405]
- de Marco A, Deuerling E, Mogk A, Tomoyasu T, Bukau B. Chaperone-based procedure to increase yields of soluble recombinant proteins produced in *E. coli*. *BMC Biotechnology*. 2007; 7:32. [PubMed: 17565681]
- Miller ML, Jensen LJ, Diella F, Jørgensen C, Tinti M, Li L, Hsiung M, Parker SA, Bordeaux J, Sicheritz-Ponten T, et al. Linear Motif Atlas for Phosphorylation-Dependent Signaling. *Science Signaling*. 2008; 1:ra2–ra2. [PubMed: 18765831]
- Miroux B, Walker JE. Over-production of Proteins in *Escherichia coli*: Mutant Hosts that Allow Synthesis of some Membrane Proteins and Globular Proteins at High Levels. *Journal of Molecular Biology*. 1996; 260:289–298. [PubMed: 8757792]
- Nishihara K, Kanemori M, Yanagi H, Yura T. Overexpression of Trigger Factor Prevents Aggregation of Recombinant Proteins in *Escherichia coli*. *Applied and Environmental Microbiology*. 2000; 66:884–889. [PubMed: 10698746]
- Obenauer JC, Cantley LC, Yaffe MB. Scansite 2.0: proteome-wide prediction of cell signaling interactions using short sequence motifs. *Nucleic Acids Research*. 2003; 31:3635–3641. [PubMed: 12824383]
- O'Shea JP, Chou MF, Quader SA, Ryan JK, Church GM, Schwartz D. pLogo: a probabilistic approach to visualizing sequence motifs. *Nature Methods*. 2013; 10:1211–1212. [PubMed: 24097270]
- Oza JP, Aerni HR, Pirman NL, Barber KW, ter Haar CM, Rogulina S, Amroffell MB, Isaacs FJ, Rinehart J, Jewett MC. Robust production of recombinant phosphoproteins using cell-free protein synthesis. *Nature Communications*. 2015; 6:8168.
- Pinna LA, Ruzzene M. How do protein kinases recognize their substrates? *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*. 1996; 1314:191–225. [PubMed: 8982275]
- Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nature Protocols*. 2007; 2:1896–1906. [PubMed: 17703201]
- Rosano, GL., Ceccarelli, EA. [Accessed June 13, 2016] Recombinant protein expression in *Escherichia coli*: advances and challenges; *Frontiers in Microbiology*. 2014. p. 5 Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4029002/>
- Santos CA, Beloti LL, Toledo MAS, Crucello A, Favaro MTP, Mendes JS, Santiago AS, Azzoni AR, Souza AP. A novel protein refolding protocol for the solubilization and purification of recombinant

- peptidoglycan-associated lipoprotein from *Xylella fastidiosa* overexpressed in *Escherichia coli*. *Protein Expression and Purification*. 2012; 82:284–289. [PubMed: 22306742]
- Schein CH. Production of Soluble Recombinant Proteins in Bacteria. *Nature Biotechnology*. 1989; 7:1141–1149.
- Schwartz D, Gygi SP. An iterative statistical approach to the identification of protein phosphorylation motifs from large-scale data sets. *Nature Biotechnology*. 2005; 23:1391–1398.
- Soares NC, Spät P, Krug K, Macek B. Global Dynamics of the *Escherichia coli* Proteome and Phosphoproteome During Growth in Minimal Medium. *Journal of Proteome Research*. 2013; 12:2611–2621. [PubMed: 23590516]
- Songyang Z, Blechner S, Hoagland N, Hoekstra MF, Piwnicka-Worms H, Cantley LC. Use of an oriented peptide library to determine the optimal substrates of protein kinases. *Current Biology*. 1994; 4:973–982. [PubMed: 7874496]
- Studier FW. Protein production by auto-induction in high-density shaking cultures. *Protein Expression and Purification*. 2005; 41:207–234. [PubMed: 15915565]
- Ubersax JA, Ferrell JE Jr. Mechanisms of specificity in protein phosphorylation. *Nature Reviews Molecular Cell Biology*. 2007; 8:530–541. [PubMed: 17585314]
- Wagner S, Klepsch MM, Schlegel S, Appel A, Draheim R, Tarry M, Högbom M, van Wijk KJ, Slotboom DJ, Persson JO, et al. Tuning *Escherichia coli* for membrane protein overexpression. *Proceedings of the National Academy of Sciences*. 2008; 105:14371–14376.
- Zhao Y, He W, Liu WF, Liu CC, Feng LK, Sun L, Yan YB, Hang HY. Two Distinct States of *Escherichia coli* Cells That Overexpress Recombinant Heterogeneous β -Galactosidase. *Journal of Biological Chemistry*. 2012; 287:9259–9268. [PubMed: 22303013]

Figure 1. Schematic overview of the experimental ProPeL workflow

A kinase of interest is cloned and expressed in *E. coli*. Resulting bacterial phosphorylation is evaluated by SDS-PAGE with Pro-Q Diamond and Coomassie staining. Lysate is digested, phosphoenriched and identified by tandem mass spectrometry. Data sets are visualized with pLogo (O'Shea et al., 2013).

Figure 2. Example of Desired Kinase Expression and Activity

These gels illustrate the desired level of kinase expression and *in vivo* activity with the DYRK1A^{WT} kinase. **(A)** SDS-PAGE with Coomassie staining with robust expression of both (2) DYRK1A^{KD} and (3) DYRK1A^{WT} at the expected molecular weight of 43 kDa, relative to (1) empty vector pET45b. **(B)** Pro-Q Diamond staining reveals robust autophosphorylation and efficient phosphorylation of bacterial substrates over a wide molecular weight range for (3) DYRK1A^{WT} relative to (1) empty vector pET45b or (2) DYRK1A^{KD} negative controls.