Published in final edited form as:

Curr Opin Anaesthesiol. 2013 April ; 26(2): 152–156. doi:10.1097/ACO.0b013e32835e8b34.

Perioperative physiotherapy

Bhakti K. Patel and **Jesse B. Hall**

Department of Pulmonary/Critical Care, University of Chicago Medical Center, Chicago, Illinois, USA

Abstract

Purpose of review—Physiotherapy in the perioperative period is emerging as an important component of postoperative recovery. This review highlights recent advances in the implementation of physiotherapy in the perioperative period and its enhancement of postsurgical outcomes.

Recent findings—Physical therapy in the preoperative period can improve physical deconditioning and potentially affect subsequent postsurgical outcomes. Fast-track surgical programs have highlighted the importance of early ambulation in the postoperative period. Incorporation of this multimodal, evidenced-based approach has been shown to reduce postoperative pulmonary complications and shorten hospital length of stay. Physiotherapy is feasible and well tolerated in patients who remain intubated and mechanically ventilated in the postoperative period. This approach also improves duration of mechanical ventilation and return to functional independence at hospital discharge.

Summary—Timely and early physiotherapy in the perioperative period improves surgical recovery and reduces postoperative complications.

Keywords

early ambulation; early mobilization; fast-track surgery

INTRODUCTION

Traditionally bed rest had been prescribed in the convalescence from surgery as well as medical illness. However by the mid 20th century, there was growing recognition of the deleterious effects of immobilization that are ‘potentiat[ed] by anesthesia, narcotics, or the original illness’ [1]. Initial reports of the benefits and safety of early postoperative ambulation began the paradigm shift toward ‘prompt restoration of surgical patients to normal life’ [2] as an essential feature of postoperative recovery. Recently, a systematic review of 39 trials of bed rest in a variety of conditions demonstrated no therapeutic benefit and instead suggested potential harm [3]. The consequences of immobilization include increased risk of thromboembolism, pulmonary complications, orthostatic intolerance [4],

Correspondence to Jesse B. Hall, MD, Chief, Section of Pulmonary/Critical Care, University of Chicago Medical Center, 5841 S Maryland Ave, MC 6076, Chicago, IL 60637, USA. Tel: +1 773 702 1454; fax: +1 773 702 4754; jhall@medicine.bsd.uchicago.edu.

Conflicts of interest

There are no conflicts of interest.

ileus, and muscle atrophy/weakness. Further, ICU acquired neuromuscular weakness can have residual effects even years after survival from critical illness [5].

Perioperative physiotherapy has emerged as an important intervention in preventing these complications. Prior retrospective studies have suggested that early physical therapy was protective against postoperative complications [6] and associated with improved outcomes in patients with acute respiratory failure [7]. Recent prospective studies have similarly suggested the benefits of mobilization in the preoperative care, with use of multidisciplinary fast-track surgery programs, and in the management of critically ill patients (Table 1). This review will summarize the role of perioperative physical therapy in improving patient outcomes in each of these contexts.

PREOPERATIVE PHYSIOTHERAPY

Pre-existing physical deconditioning can adversely affect surgical outcomes [26]. As such, preoperative rehabilitation has become an attractive option to potentially improve postoperative recovery, although definitive data supporting this notion are somewhat elusive. Herdy *et al.* [8] performed a randomized controlled trial of hospitalized patients awaiting coronary bypass graft surgery (CABG) to receive standard care or preoperative cardiopulmonary rehabilitation for at least 5 days prior to surgery. Patients who received physical therapy had shortened time to extubation, less postoperative pulmonary complications, and shorter hospital length of stay. The role of pulmonary rehabilitation in the preoperative period for patients undergoing lobectomy may improve physiologic parameters and exercise capacity, but its effect on postoperative pulmonary complications remains uncertain [9]. The disparate outcomes in the CABG versus lobectomy patient populations may be more related to the prevention of the complications of in-hospital immobilization rather than benefits of preoperative physical therapy, given that these populations differ significantly in their ability to continue with mobilization strategies in the postoperative period.

Mitigating the effects of preoperative immobilization with physical therapy in extremes of illness has been described in pediatric patients awaiting lung transplant on extracorporeal membrane oxygenation (ECMO). In a small case series of three patients Turner *et al.* [27] demonstrated the feasibility of preoperative physiotherapy while on ECMO and a reduction of post-transplant hospital length of stay by about 50 days. Further study of the role of preoperative physiotherapy in pretransplant patients is warranted.

FAST-TRACK PROGRAMS

The essence of surgical fast-track programs is the employment of evidence-based practice to enhance postoperative recovery and reduce morbidity. This multimodal approach incorporates surgeons, anesthesiologists, physical therapists and nursing into the care team [28]. Fast-track programs employ preoperative assessment and education, evidence-based practice in anesthesia, minimally invasive procedures, analgesia, early feeding and ambulation to enhance and accelerate postoperative recovery. Early physiotherapy has not

been standardized among all programs but generally involves early extubation, sitting in a chair for more than 1 h on the postoperative day, and ambulation by postoperative day one.

The evidence supporting the success of fast-track programs is robust in the colorectal surgical literature [29,30]. Benefits of this multimodal approach include reduced need for ICU stay, faster recovery of normal gastrointestinal function, reduction of postoperative pulmonary complications and shortened hospital stay [10]. A recent meta-analysis of four randomized controlled trials and seven case-controlled studies comparing fast-track surgery to standard care demonstrated a reduced hospital stay by 2.35 days and less morbidity without increase in readmission rates or mortality [11]. Further, Lee *et al.* [31] investigated the contribution of early feeding and mobilization to the benefits of fast-track surgical programs in a randomized control trial. Patients who received early ambulation and feeding had shorter times to recovery, more analgesic-free states, and had a trend toward shorter hospital stay when compared with the patients with late nutrition and bed rest. Fast-track programs with intensive mobilization have been implemented in other surgical procedures including gastric [12,13], pancreatic [14,15,32], hepatic [16,17], abdominal aortic [18–20], esophageal [21,33,34], and total knee replacement surgery [22] with similar enhancement of postoperative outcomes.

The timeliness of mobilization appears important in achieving these beneficial effects on recovery. A case-control study compared 36 patients who underwent lobectomy by axillofateral thoracotomy approach with ambulation at 4 h after surgery versus controls who ambulated on postoperative day one. Patients with late ambulation were twice more likely to require oxygen for more than 2 days. Also, no patients in the early ambulation group had P/F (partial pressure of oxygen/fraction of inspired oxygen) ratios of less than 300 on postoperative day 3, whereas control patients did [35■].

There often are barriers to immediate mobilization in the postoperative period. Concerns over adequate analgesia for postoperative pain especially may thwart early ambulation. However, narcotics may also impede immediate ambulation by causing orthostatic instability. In fact, one retrospective study of patients undergoing laparoscopic gynecologic surgery, described that delayed ambulation [adjusted odds ratio (95% confidence interval), 8.37 (1.23–72.15)] and orthostatic intolerance [adjusted odds ratio (95% confidence interval), 34.78 (11.12–131.72)] was strongly associated with continuous fentanyl infusion [36]. Therefore, a careful balance of adequate analgesia without the associated side-effects is necessary for patients to benefit from enhanced recovery seen with early physiotherapy implemented within fast-track surgery programs. One attractive possibility is to extend the use of postoperative regional anesthetic techniques whenever possible to avoid the undesirable systemic effects of opioids.

EARLY MOBILIZATION OF MECHANICALLY VENTILATED CRITICALLY ILL PATIENTS

In fast-track surgery programs, timely extubation is considered the rate-limiting step for initiation of early mobility. However, this rationale is being questioned by the growing recognition of ICU-acquired weakness, characterized by symmetric limb and axial muscle

weakness that afflicts many patients surviving critical illness [37]. A quarter of patients mechanically ventilated for at least 7 days suffer ICU-acquired weakness [38], but the incidence can reach nearly 100% in patients with sepsis and multiorgan failure [39]. Time on mechanical ventilation is a critical factor as diaphragm muscle atrophy can be seen as early as 18 h after the institution of mechanical ventilation in patients who are passive on the ventilator [40]. Once established, this weakness may persist years after recovery from the original process precipitating critical illness and ICU admission. Accordingly, this process appears to be common, to develop early in the course for many patients, and to confound their recovery for a protracted period of time.

Early mobilization of intubated mechanically ventilated patients has been recently demonstrated as well tolerated, feasible, and beneficial for patients and can prevent or ameliorate ICU-acquired weakness in many patients [23,24,25,41,42]. Bailey *et al.* [41] were the first to describe early mobilization of mechanically ventilated patients who required more than 4 days of mechanical ventilation. Physical therapy was started after physiologic stabilization and almost 70% of patients were able to walk 100 ft by ICU discharge. The main criticisms of this prospective cohort study were that patients were less ill at the time of mobilization and that physiotherapy typically occurred after more than 10 days of admission to the ICU. Schweickert *et al.* [23] published a prospective randomized controlled trial of very early mobility in patients intubated for less than 72 h. Therapy began on average within 36 h of intubation with 89% of sessions occurring during acute lung injury, delirium, renal replacement therapy or vasoactive administration [43]. Patients who underwent early mobilization had more ventilator free days, more delirium free days, and were functionally independent to a greater degree on hospital discharge.

The studies cited above had a predominance of patients admitted to the ICU with medical disorders. Implementation of early mobilization in the surgical ICU may be hindered by unique considerations, such as wound healing, postoperative pain, weight-bearing limitations, and surgical drains [44]. However, there are limited data on the safety and feasibility of early mobilization in the surgical population. Burtin *et al.* [24] performed a randomized controlled trial of early exercise using a bedside bicycle ergometer in patients with expected prolonged stays in the medical and surgical ICUs. A majority of the patients were surgical patients with 84% being intubated during physical therapy. Intervention patients had a higher 6min walk distance, improved quadriceps strength, and a trend to hospital discharge to home. Garzon-Serrano *et al.* [25] performed a prospective observational study on the implementation of physical therapy in a surgical ICU. Patients underwent stepwise mobilization with a passive range of motion in bed, followed by sitting on the edge of the bed, transferring to a chair, standing, and ambulating. Interestingly, physical therapists mobilized patients to higher levels than nurses. This highlights the need for a multidisciplinary team of physical/occupational therapists, nursing, respiratory staff, and physicians to champion early mobilization.

CONCLUSION

Bed rest in the postoperative period can be detrimental to patient recovery. Preoperative physiotherapy in some cases can prime patients for enhanced recovery in the postoperative

period. Fast-track surgery methodology that incorporates intensive mobilization has been shown to improve outcomes. Timeliness of mobilization occurring immediately after surgery or early after institution of mechanical ventilation in cases of respiratory failure can be beneficial in preventing weakness and returning patients to functional independence at time of discharge.

Acknowledgments

None.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

■ of special interest

■■ of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (p. 246).

1. Dock W. The evil sequelae of complete bed rest. *JAMA*. 1944; 125:1083–1085.
2. Powers J. The abuse of rest as a therapeutic measure in surgery. *JAMA*. 1944; 125:1079–1083.
3. Allen C, Glasziou P, Del Mar C. Bed rest: a potentially harmful treatment needing more careful evaluation. *Lancet*. 1999; 354:1229–1233. [PubMed: 10520630]
4. Harper CM, Lyles UM. Physiology and complications after bed rest. *J Am Geriatr Soc*. 1988; 36:1047–1054. [PubMed: 3049751]
5. Herridge MS, Tansey CM, Matte A, et al. Canadian Critical Care Trials Group: functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med*. 2011; 364:1293–1304. [PubMed: 21470008]
6. Das-Neves-Pereira J, Bagan P, Coimbra-Israel A, et al. Fast-track rehabilitation for lung cancer lobectomy: a five year experience. *Eur J Card Thorac Surg*. 2009; 36:383–392.
7. Morris PE, Griffin L, Berry M, et al. Receiving early mobility during an intensive care unit admission is a predictor of improved outcomes in acute respiratory failure. *Am J Med Sci*. 2011; 341:373–377. [PubMed: 21358312]
8. Herdy AH, Marcelli PLB, Vila A, et al. Pre and postoperative cardiopulmonary rehabilitation in hospitalized patients undergoing coronary artery bypass surgery: a randomized controlled trial. *Am J Phys Med Rehabil*. 2008; 87:714–719. [PubMed: 18716482]
9. Nagarajan K, Bennett A, Agostini P, Naidu B. Is preoperative physiotherapy/pulmonary rehabilitation beneficial in lung resection patients? *Interactive Cardiovasc Thorac Surg*. 2011; 13:300–302.
10. Olsen MF, Wennberg E. Fast-track concepts in major open upper abdominal and thoracoabdominal surgery: a review. *World J Surg*. 2011; 35:2586–2593. [PubMed: 21882022]
11. Gouvas N, Tan E, Windsor A, et al. Fast-track vs standard care in colorectal surgery: a meta-analysis update. *Int J Colorectal Dis*. 2009; 24:1119–1131. [PubMed: 19415308]
12. Liu XX, Jiang ZW, Wang ZM, Li JS. Multimodal optimization of surgical care shows beneficial outcome in gastrectomy surgery. *JPEN*. 2010; 34:313–321.
13. Wang D, Kong Y, Zhong B, et al. Fast-track surgery improves postoperative recovery in patients with gastric cancer: a randomized comparison with conventional postoperative care. *Gastrointest Surg*. 2010; 14:620–627.

14. Balzano G, Zerbi A, Braga M, et al. Fast-track recovery programme after pancreaticoduodenectomy reduces delayed gastric emptying. *Br J Surg.* 2008; 95:1387–1393. [PubMed: 18844251]
15. Kennedy EP, Rosato EL, Sauter PK, et al. Initiation of a critical pathway for pancreaticoduodenectomy at an academic institution-the first step in multidisciplinary team building. *J Am Coll Surg.* 2007; 204:917–923. [PubMed: 17481510]
16. van Dam RM, Hendry PO, Coolen MM, et al. Enhanced recovery after surgery (ERAS) group. Initial experience with a multimodal enhanced recovery programme in patients undergoing liver resection. *Br J Surg.* 2008; 95:969–975. [PubMed: 18618897]
17. MacKay G, O'Dwyer PJ. Early discharge following liver resection for colorectal metastases. *Scott Med J.* 2008; 53:22–24.
18. Muehling B, Schelzig H, Steffen P, et al. A prospective randomized trial comparing traditional and fast-track patient care in elective open infrarenal aneurysm repair. *World J Surg.* 2009; 33:577–585. [PubMed: 19137363]
19. Murphy MA, Richards T, Atkinson C, et al. Fast track open aortic surgery: reduced post operative stay with a goal directed pathway. *Eur J Vasc Endovasc Surg.* 2007; 34:274–278. [PubMed: 17587613]
20. Brustia P, Renghi A, Fassiola A, et al. Fast-track approach in abdominal aortic surgery: left subcostal incision with blended anesthesia. *Interact Cardiovasc Thorac Surg.* 2007; 6:60–64. [PubMed: 17669770]
21. Brodner G, Pogatzki E, Van Aken H, et al. A multimodal approach to control postoperative pathophysiology and rehabilitation in patients undergoing abdomin thoracic esophagectomy. *Anesth Analg.* 1998; 86:228–234. [PubMed: 9459224]
22. denHertog A, Gliesche K, Timm J, et al. Pathway-controlled fast-track rehabilitation after total knee arthroplasty: a randomized prospective clinical study evaluating the recovery pattern, drug consumption, and length of stay. *Arch Orthop Trauma Surg.* 2012; 132:1153–1163. [PubMed: 22643801]
23. Schweickert WD, Pohlman MC, Pohlman AS, et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomised controlled trial. *Lancet.* 2009; 373:1874–1882. [PubMed: 19446324]
24. Burtin C, Clerckx B, Robbeets C, et al. Early exercise in critically ill patients enhances short-term functional recovery. *Crit Care Med.* 2009; 37:2499–2505. [PubMed: 19623052]
25. Garzon-Serrano J, Ryan C, Waak K, et al. Early mobilization in critically ill patients: patients' mobilization level depends on healthcare provider's profession. *PMR.* 2011; 3:307–313. Prospective observational study of implementation of early physical therapy in a surgical population. This study demonstrated that physical therapy is feasible and well tolerated in the surgical population, but requires a multidisciplinary approach.
26. Pieper B, Lepczyk M, Caldwell M. Perceptions of the waiting period before coronary artery bypass grafting. *Heart Lung.* 1985; 14:40–44. [PubMed: 3871431]
27. Turner DA, Cheifetz IM, Rehder KJ, et al. Active rehabilitation and physical therapy during extracorporeal membrane oxygenation while awaiting lung transplantation: a practical approach. *Crit Care Med.* 2011; 39:2593–2598. This observational study highlights that physiotherapy can be performed in extremes of illness. [PubMed: 21765353]
28. Kehlet H, Wilmore DW. Evidence-based surgical care and the evolution of fast-track surgery. *Ann Surg.* 2008; 248:189–198. [PubMed: 18650627]
29. Khoo CK, Vickery CJ, Forsyth N, et al. A Prospective randomized controlled trial of multimodal perioperative management protocol in patients undergoing elective colorectal resection for cancer. *Ann Surg.* 2007; 245:867–872. [PubMed: 17522511]
30. Wang Q, Suo J, Jiang J, et al. Effectiveness of fast track rehabilitation vs conventional care in laparoscopic colorectal resection for elderly patients: a randomized trial. *Colorectal Dis.* 2011; 14:1009–1014.
31. Lee TG, Kang SB, Kim DW, et al. Comparison of early mobilization and diet rehabilitation program with conventional care after laparoscopic colon surgery: a prospective randomized controlled trial. *Dis Colon Rectum.* 2011; 54:21–28. [PubMed: 21160309]

32. di Sebastiano P, Festa L, De Bonis A, et al. A modified fast-track program for pancreatic surgery: a prospective single-center experience. *Langenbecks Arch Surg*. 2011; 396:345–351. [PubMed: 20703500]
33. Cerfolio RJ, Bryant AS, Bass CS, et al. Fast tracking after Ivor Lewis esophagogastrrectomy. *Chest*. 2004; 126:1187–1194. [PubMed: 15486381]
34. Neal JM, Wilcox RT, Allen HW, Low DE. Near-total esophagectomy: the influence of standardized multimodal management and intraoperative fluid restriction. *Reg Anesth Pain Med*. 2003; 28:328–334. [PubMed: 12945027]
35. Kaneda H, Saito Y, Okamoto M, et al. Early postoperative mobilization with walking at 4 h after lobectomy in lung cancer patients. *Gen Thorac Cardiovasc Surg*. 2007; 55:493–498. This study describes how the timeliness mobilization occurring early is necessary for improved postsurgical outcomes. [PubMed: 18066640]
36. Iwata Y, Mizota Y, Mizota T, et al. Postoperative continuous infusion of fentanyl is associated with the development of orthostatic intolerance and delayed ambulation in patients after gynecologic laparoscopic surgery. *J Anesth*. 2012; 26:503–508. [PubMed: 22526434]
37. De Jonghe B, Sharshar T, Hopkinson N, et al. Paresis following mechanical ventilation. *Curr Opin Crit Care*. 2004; 10:47–52. [PubMed: 15166849]
38. De Jonghe B, Sharshar T, Lefaucheur JP, et al. Paresis acquired in the intensive care unit: a prospective multicenter study. *JAMA*. 2002; 288:2859–2867. [PubMed: 12472328]
39. Stevens RD, Dowdy DW, Michaels RK, Needham DM. Neuromuscular dysfunction acquired in critical illness: a systematic review. *Intensive Care Med*. 2007; 33:1876–1891. [PubMed: 17639340]
40. Levine S, Nguyen T, Taylor N, et al. Rapid disuse atrophy of diaphragm fibers in mechanically ventilated humans. *N Engl J Med*. 2008; 358:1327–1335. [PubMed: 18367735]
41. Bailey P, Thomsen GE, Spuhler VJ, et al. Early activity is feasible and safe in respiratory failure patients. *Crit Care Med*. 2007; 35:139–145. [PubMed: 17133183]
42. Morris PE, Goad A, Thompson C, et al. Early intensive care unit mobility therapy in the treatment of acute respiratory failure. *Crit Care Med*. 2008; 36:2238–2243. [PubMed: 18596631]
43. Pohlman MC, Schweickert WD, Pohlman AS, et al. Feasibility of physical and occupational therapy beginning from initiation of mechanical ventilation. *Crit Care Med*. 2010; 38:2089–2094. [PubMed: 20711065]
44. Lipshutz AK, Gropper MA. Acquired neuromuscular weakness and early mobilization in the intensive care unit. *Anesthesiology*. 2012; XXX:1–14.

KEY POINTS

- Immobilization in the perioperative period is associated with poor outcomes.
- Timely mobilization early in the postoperative period is key to improving patient recovery.
- Incorporation of early ambulation in fast-track surgery programs has improved postoperative outcomes including shortened hospital stay, less pulmonary complications, and shorter time to analgesic-free states.
- Early physical and occupational therapy is well tolerated and feasible in mechanically ventilated patients and improves return to functional independence at hospital discharge.

Table 1**Effects of perioperative physical therapy**

Preoperative physiotherapy	
Cardiopulmonary rehabilitation prior to coronary bypass graft surgery [8]	Less ventilator days; Less PPC; Shorter hospital length of stay
Pulmonary rehabilitation prior to lobectomy [9]	Improved exercise capacity; No difference in PPC
Fast-track surgery programs	
Colorectal surgery	Reduced need for ICU admission [10]
Gastric surgery [12,13]	Faster recovery of gastrointestinal function [10] Less postoperative pulmonary complications [10]
Pancreatic surgery	Reduced hospital length of stay [10,11] Faster recovery of gastrointestinal function
Hepatic surgery [16,17]	Reduced hospital length of stay Trend for less postsurgical complications Less postsurgical complications [14] Reduced hospital length of stay [14,15] Reduced hospital costs [15] Reduced hospital length of stay
Abdominal aortic surgery	Reduced time to extubation [18] Reduced hospital length of stay [18–20]
Esophageal surgery	Earlier extubation and discharge from ICU [21]
Total knee arthroplasty [22]	Reduced need for analgesic drugs Reduced hospital length of stay
Mechanically ventilated patients	
Critically ill medicine patients [23]	Improved return to functional independence at hospital discharge; shorter duration of delirium; reduced duration of mechanical ventilation
Surgical ICU patients	
Bedside bicycle ergometer [24]	Improved walk distance, quadriceps strength; trend toward discharge to home
Stepwise mobilization to ambulation[25■■■]	Feasible and safe in surgical ICU patients; more effective if implemented by physical/occupational therapists

PPC..