Published in final edited form as:

J Acquir Immune Defic Syndr. 2018 August 01; 78(4): e30–e34. doi:10.1097/QAI.0000000000001698.

Barriers to HIV testing among men who have sex with men in India reached online: implications for interventions

Viraj V. Patel¹, Alpana Dange, MS², Shruta Rawat, MSc², Julia H. Arnsten, MD, MPH¹, Christopher Piña, MD¹, Sarit A. Golub, PhD³, Rohan Pujari, HSc², Ankit Trivedi, HSc², Jack Harrison-Quintana, MPH⁴, and Ken H. Mayer, MD^{5,6}

¹Montefiore Health System/Albert Einstein College of Medicine, Division of General Internal Medicine, Bronx, New York, USA

²The Humsafar Trust, Mumbai, India

³Hunter College, City University of New York, Psychology, New York, United States

⁴Grindr, Grindr for Equality, Washington D.C., United States

⁵The Fenway Institute, Fenway Health, Boston, United States

⁶Beth Israel Deaconess Medical Center, Harvard Medical School, Infectious Diseases, Boston, United States

Keywords

men who have sex with men (MSM); India; HIV testing; social media; Internet; stigma

INTRODUCTION

Indian men who have sex with men (MSM) have a higher HIV prevalence than the general Indian population (4.3% v. 0.3%),¹ and are designated a “High Risk Group” by the National AIDS Control Organization of India and a priority population for targeted HIV prevention interventions.¹ However, the effectiveness of HIV outreach interventions for MSM has been limited, in part due to stigma and criminalization of homosexuality.^{2–4}

Globally and in India, MSM are increasingly using online social networking sites (SNS), such as mobile dating applications or Facebook, to find community and seek partners. Though SNS disrupt traditional face-to-face outreach, SNS also provide access to hard-to-reach populations. Studies among SNS-using Indian MSM have found that up to 50% are

Corresponding Author: Viraj V. Patel, MD, MPH, Assistant Professor of Medicine, Montefiore Health System/Albert Einstein College of Medicine, 3300 Kossuth Avenue; Bronx, NY, USA 10467, Office phone: 718-920-7102; Fax: 718-561-5165; vpatel@montefiore.org.

Findings from this study were presented in part at the 9th International AIDS Society Conference on HIV Science, Scientific Meeting, 23–26 July 2017, Paris, France.

Conflicts of Interest and Sources of Funding: Ken Mayer has unrestricted research grants from Gilead Sciences and ViiV healthcare. The other authors have no conflicts of interest. This study was supported in part by NIH K23MH102118 (PI Viraj V. Patel), Albert Einstein’s Office of Medical Student Research, Albert Einstein College of Medicine’s Global Health Center, the Emory University Center for AIDS Research (NIH P30AI050409), and the Einstein-Rockefeller-CUNY Center for AIDS Research (NIH P30AI051519).

unaware of their HIV status,^{5,6} but were limited to small localities or did not examine correlates of HIV testing.

Utilizing SNS for health interventions could improve HIV prevention and treatment efforts for Indian MSM. Because India is a heterogeneous country with over 1 billion people, large socioeconomic and health disparities, and widely varying HIV prevalence across states, country-wide inferences cannot be made based on a few localities. We therefore conducted a national survey of SNS-using MSM to elucidate factors associated with HIV testing and serostatus awareness.

METHODS

Study design, setting, and participants

Zero-Metres-Away (ZMA) was a cross-sectional, self-administered, anonymous survey conducted from 6 January 2017 to 5 February 2017. Indian MSM were recruited by advertisements on three MSM-specific mobile SNS and LGBTQ Facebook and Instagram groups. The survey was hosted by Surveygizmo (Boulder, Colorado) and began with study information and informed consent. We limited duplicate respondents by using web browser cookies and restricting to single IP addresses. Upon completion, respondents were provided with HIV prevention and testing resources and given the option to enter contact information (unlinked to responses) for a chance to win a 1,000 Indian rupee (approximately \$15 USD) Amazon.co.in gift-card. Inclusion criteria were: (1) 18 years of age, (2) identifying as male (cis- or trans-), (3) anal sex with male or transgender partners in past two years, and (4) born in India and living there at the time of the study. Human subjects research review boards at The Humsafar Trust (an LGBTQ community-based organization in Mumbai, India) and Albert Einstein College of Medicine (Bronx, United States) approved the study.

Survey development and measures

We partnered with The Humsafar Trust to adapt an online survey of MSM and HIV,⁷ translate the survey into Hindi, and refine it to ensure conceptual accuracy.⁸ We assessed sexual identity with categories used in India (panthi, kothi, double-decker, gay/homosexual, bisexual, and straight/heterosexual),^{9–11} but because very few respondents selected panthi, kothi, or double-decker, we reclassified these as gay/homosexual. We asked about past six-month anal sex and condomless anal sex (CAS), and about past 12-month number and types of sexual partners, drug/alcohol use during sex, and sexually transmitted infections (STIs). We measured perceived HIV stigma by calculating mean responses to the following items,¹² which ask agreement on a five-point scale: “Most people in my area would: (1)... discriminate against someone with HIV, (2)...think that people who got HIV through sex or drug use deserve what they have gotten, and (3)...support the rights of a person with HIV to live and work wherever they wanted.” We asked about monthly household income and classified 10,000 Rupees/month as living in poverty and between 10,001–20,000 as low-income. To assess our primary outcome, we dichotomized responses to “Have you ever been tested for HIV?” as yes versus no/don't know. Among those responding yes, we assessed how long ago and setting of last HIV test. For those never tested, we assessed reasons for not HIV testing.^{13, 14}

Analysis

We excluded responses with missing data and characterized the sample using summary statistics. We performed bivariable and multivariable analysis to determine factors associated with ever HIV testing, using generalizing estimating equations for logistic regression to account for clustering by state. We included all variables significant ($p < 0.1$) in bivariable analysis in multivariable models, and report adjusted odds ratios (aOR) and 95% confidence intervals (CI). We conducted sensitivity analyses to explore how missing data ($n = 2,458$ with incomplete surveys) affected estimates, using multiple-imputation with chained equations.¹⁵ Because models run with multiple-imputation yielded similar results, only results from complete case analyses are presented. We used Stata SE, version 15 (StataCorp, College Station, Texas).

RESULTS

Participant characteristics

Of 6,637 MSM meeting inclusion criteria, 4,179 (63%) completed all items used in the present analysis. Median age was 26 years (interquartile range 23–30), most (83%) completed college, many lived in poverty (15%) or were low-income (21%), 18% responded in Hindi, and 0.4% ($n = 17$) identified as transgender MSM. Respondents came from all Indian states (range, $n = 5$ to $n = 1,124$), including 17% rural/semi-rural.

Most respondents identified as either gay/homosexual (49%) or bisexual (46%). Most (75%) had past six-month anal sex and 42% had CAS, with 41% reporting more than six partners in the past 12 months and 67% reporting casual male partners. Even though most (60%) had disclosed sex with/attraction to men to someone, only 23% had disclosed having sex with men to a doctor. Regarding access to HIV testing, only 37% indicated that it was easy/very easy to access free HIV testing, though 47% indicated being aware of a laboratory where they would feel comfortable obtaining testing. Mean score on the three questions assessing perceived HIV stigma was 2.5 ± 0.81 (range 1–5; higher scores indicate greater stigma).

Factors associated with HIV testing

Almost half (47%) had never been HIV tested, with those reporting CAS less likely to have HIV tested than those reporting no CAS (44% vs. 50%, $p < 0.001$). Among those with prior HIV testing ($n = 2,215$), 25% had last tested more than 12 months prior and 5.4% were HIV positive. Multivariable analysis (Table) revealed that younger age, lower income, less education, CAS, no drugs/alcohol during sex, no access to comfortable testing site, paid female sex partner, non-disclosure of sexual identity or behavior, and more perceived stigma were associated with never HIV testing ($p < 0.01$ for all variables).

Reasons for not testing

Most common reasons for not testing were low perceived risk (42%), feeling scared (20%), and not knowing where to test (15%). Reasons did not significantly differ by age, geography (rural versus urban), income, education, or language of survey completion.

DISCUSSION

Among a large, diverse sample of sexually active Indian MSM recruited online, we found that nearly half (47%) of more than 4,000 respondents had never been tested for HIV. This estimate was higher among those reporting CAS (50%) than among those reporting no CAS (44%). We further found that modifiable factors, including access to free testing and availability of comfortable testing sites, were associated with HIV testing. Finally, we found that respondents who had disclosed their sexuality to doctors or others were more likely to have HIV tested.

This is the first study to characterize SNS-using MSM across India and describe their HIV testing behaviors. Prior national or multi-city samples of Indian MSM were recruited through physical venue-based approaches, and demonstrated similar or higher rates of HIV testing, ranging from 51–82%.^{1,16} While the HIV testing prevalence we observed is comparable to smaller studies of Indian MSM recruited online,^{5,6} prior studies have not examined correlates of HIV testing. Additionally, prior studies have not provided country-wide data needed to guide testing interventions for Indian MSM. Our finding that individuals having CAS were less likely to have been HIV-tested is consistent with prior studies and suggests that greater education about HIV is needed.¹⁹

There have been significant efforts towards scaling-up HIV testing across India, and the National AIDS Control Organization guidelines now recommend that sexually active MSM get HIV tested every six months.¹ Our results reveal significant gaps between these recommendations and current HIV testing behaviors among Indian MSM. Scalable interventions that circumvent barriers to HIV testing might include online outreach to MSM to link them to confidential primary care and sexual health services, dissemination of information about MSM-sensitive HIV testing and prevention services, and provision of HIV self-testing.¹⁸ Online outreach could also be used to link individuals to offline community-vetted HIV testing sites in addition to public integrated counseling-testing centers,¹⁷ as we found that most respondents sought testing in the private sector.

The generalizability of our findings is limited partially by high levels of education and income among respondents (e.g., 69.3% completed college compared to 11.5% of Indian men overall.)²⁰ In addition, our survey was cross-sectional, and causality cannot be inferred. Questions may have been misunderstood despite extensive piloting, and survey respondents had to be fluent in English or Hindi. Finally, the survey may have been inaccessible to some rural MSM due to limited or unstable Internet connectivity. Despite these limitations, our results underscore the need for rapid development and implementation of robust online HIV prevention interventions for Indian MSM.

In conclusion, we found that nearly half of online MSM in India had not had an HIV test, despite high education and income levels. These results highlight the need for increased availability and ease of HIV testing. Because low risk perception was a common reason for not testing, more HIV education is also needed. Finally, because social stigma may prevent access to HIV services, both online and structural interventions are needed to address

stigma, foster social acceptance, and increase availability of culturally competent and non-judgmental health services for MSM.

Acknowledgments

We would like to thank all the participants for their time, and to thank Nataly Rios and Drs. Marcus Bachhuber and Maria Zlotorzynska for their support.

References

1. National AIDS Control Organization. [Accessed 10 November 2017] National AIDS Control Organization 2016–2017 Annual Report. Ministry of Health and Family Welfare, Government of India. 2017. Available at: <http://naco.gov.in/sites/default/files/NACO%20ANNUAL%20REPORT%202016-17.pdf>
2. Safren SA, Martin C, Menon S, et al. A survey of MSM HIV prevention outreach workers in Chennai, India. *AIDS Educ Prev*. 2006; 18:323–32. [PubMed: 16961449]
3. Thomas B, Mimiaga MJ, Kumar S, et al. HIV in Indian MSM: Reasons for a concentrated epidemic & strategies for prevention. *Indian J Med Res*. 2011; 134:920. [PubMed: 22310824]
4. Aggarwal P, Bhattar S, Sahani SK, et al. Sexually transmitted infections and HIV in self reporting men who have sex with men: A two-year study from India. *Journal of Infection and Public Health*. 2016; 9:564–70. [PubMed: 26776704]
5. Ekstrand ML, Rawat S, Patankar P, et al. Sexual identity and behavior in an online sample of Indian men who have sex with men. *AIDS Care*. 2017; 29:905–13. [PubMed: 28027656]
6. Welles S, Ross M, Banik S, et al. Demographic and sexual behavior comparisons of Indian and US Internet samples of men who have sex with men. *International Journal of Sexual Health*. 2011; 23:90–101.
7. Zlotorzynska M, Sullivan P, Sanchez T. The Annual American Men's Internet Survey of Behaviors of Men Who Have Sex With Men in the United States: 2015 Key Indicators Report. *JMIR Public Health Surveill*. 2017; 3:e13. [PubMed: 28356240]
8. Hagell P, Hedin P, Meads DM, et al. Effects of method of translation of patient-reported health outcome questionnaires: a randomized study of the translation of the Rheumatoid Arthritis Quality of Life (RAQoL) Instrument for Sweden. *Value in Health*. 2010; 13:424–30. [PubMed: 20070642]
9. Solomon SS, Mehta SH, Srikrishnan AK, et al. High HIV prevalence and incidence among MSM across 12 cities in India. *AIDS*. 2015; 29:723–31. [PubMed: 25849835]
10. Saha MK, Mahapatra T, Biswas S, et al. Burden and correlates of HIV risk among men who have sex with men in Nagaland, India: analysis of sentinel surveillance data. *PloS one*. 2015; 10:e0117385. [PubMed: 25689469]
11. Salve HR, Rai SK, Kant S, et al. Demographic and Sexual Behavior Characteristics of Men Who Have Sex with Men (MSM) Registered in a Targeted Intervention (TI) Program in India. *World Journal of AIDS*. 2015; 5:256.
12. Sanchez T, Smith A, Denson D, et al. Suppl 1: Developing a Web-Based HIV Behavioral Surveillance Pilot Project Among Men Who Have Sex with Men. *The Open AIDS Journal*. 2012; 6:224. [PubMed: 23091580]
13. Woodford MR, Chakrapani V, Newman PA, et al. Barriers and facilitators to voluntary HIV testing uptake among communities at high risk of HIV exposure in Chennai, India. *Global Public Health*. 2016; 11:363–79.
14. Mehta SH, Lucas GM, Solomon S, et al. HIV care continuum among men who have sex with men and persons who inject drugs in India: barriers to successful engagement. *Clinical Infectious Diseases*. 2015; 61:1732–41. [PubMed: 26251048]
15. White IR, Royston P, Wood AM. Multiple imputation using chained equations: issues and guidance for practice. *Stat Med*. 2011; 30:377–99. [PubMed: 21225900]
16. Ramakrishnan L, Ramanathan S, Chakrapani V, et al. Comparison of sexual risk, HIV/STI prevalence and intervention exposure among men who have sex with men and women (MSMW)

- and men who have sex with men only (MSMO) in India: Implications for HIV prevention. *AIDS and Behavior*. 2015; 19:2255–69. [PubMed: 25893657]
17. Anand T, Nitpolprasert C, Trachunthong D, et al. A novel Online-to-Offline (O2O) model for pre-exposure prophylaxis and HIV testing scale up. *J Int AIDS Soc*. 2017; 20:21326. [PubMed: 28362062]
 18. Young SD, Cumberland WG, Nianogo R, et al. The HOPE social media intervention for global HIV prevention in Peru: a cluster randomised controlled trial. *The Lancet HIV*. 2015; 2:e27–32. [PubMed: 26236767]
 19. Vargas SK, Konda KA, Leon SR, et al. The Relationship Between Risk Perception and Frequency of HIV Testing Among Men Who Have Sex with Men and Transgender Women, Lima, Peru. *AIDS and Behavior*. 2018 Jan.8:1–9.
 20. C-8 Appendix Educational Level Graduate And Above By Sex For Population Age 15 And Above, 2011 Census. Office of the Registrar General & Census Commissioner; India: Available at: <http://www.censusindia.gov.in/2011census/C-series/C08A.html> [Accessed: February 17, 2018]

Sociodemographic, behavioral, and structural characteristics associated with HIV testing among online MSM in India

Table

Characteristic	Total N	%	Ever tested n (%)	OR	95% CI	aOR ^a	95% CI
Total	4179	100	2215 (53)				
<u>Sociodemographic</u>							
Age							
18–23	1116	26.7	415 (37)	ref		ref	
24–29	1780	42.6	966 (54)	2.03 ^{***}	1.72 – 2.41	1.65 ^{***}	1.44 – 1.88
30–39	1035	24.8	665 (64)	2.99 ^{***}	2.40 – 3.73	2.25 ^{***}	1.81 – 2.79
40–49	206	4.9	145 (70)	3.53 ^{***}	2.60 – 4.78	3.15 ^{***}	2.29 – 4.33
50+	42	1.0	24 (57)	2.40 ^{***}	1.17 – 4.90	2.11	0.76 – 5.89
Language Survey Taken							
Hindi	790	18.3	362 (48)	0.83 [*]	0.71 – 0.96	0.89	0.79–1.72
English	3522	81.7	1853 (54)	ref		ref	
Highest Education							
Less than High School	110	2.6	43 (39)	ref		ref	
High School	270	6.5	103 (38)	0.96	0.63 – 1.47	1.06	0.57 – 1.97
Some College	322	7.7	144 (45)	1.27	0.81 – 1.98	1.62	0.84 – 3.14
College	2896	69.3	1546 (53)	1.78 ^{***}	1.37 – 2.33	1.62	0.97 – 2.68
Graduate school	580	13.9	378 (65)	2.92 ^{***}	1.99 – 4.28	2.07 [*]	1.17 – 3.64
Household Income (Indian Rs/Month)							
10,000	629	15.1	253 (40)	ref		ref	
10,001–15,000	413	9.9	185 (45)	1.19 [*]	1.00 – 1.42	1.07	0.82 – 1.41
15,001–20,000	468	11.2	267 (57)	1.96 ^{***}	1.50 – 2.56	1.59 ^{**}	1.20 – 2.12
20,001 – 40,000	903	21.6	471 (52)	1.62 ^{***}	1.32 – 1.98	1.15	0.85 – 1.55
> 40,000	1766	42.3	1039 (59)	2.11 ^{***}	1.82 – 2.46	1.31 [*]	1.04 – 1.66
City Size							
Rural	415	9.9	145 (44)	ref		ref	

Characteristic	Total N	%	Ever tested n (%)	OR	Odds of Association with Ever Testing for HIV 95% CI	aOR ^a	95% CI
Semi-rural	385	9.2	200 (51)	1.42 [*]	1.08 – 1.87	1.29	0.92 – 1.81
Urban	1194	28.6	614 (51)	1.41 ^{**}	1.10 – 1.80	1.08	0.81 – 1.45
Metropolis	2179	52.2	1256 (56)	1.77 ^{***}	1.38 – 2.26	1.12	0.83 – 1.52
Behavioral							
Sexual Identity							
Gay/Homosexual	2034	48.7	1168 (57)	ref		ref	
Bisexual	1910	45.8	970 (51)	0.77 ^{**}	0.64 – 0.92	0.96	0.84 – 1.09
Straight/Heterosexual	230	5.5	72 (31)	0.34 ^{***}	0.25 – 0.45	0.65^{**}	0.49 – 0.85
Anal sex in past 6 months	3180	75.4	1742 (55)	1.4 ^{***}	1.26 – 1.55	1.53^{***}	1.36 – 1.73
Condomless anal sex (CAS) in past 6 months	1771	42.4	882 (50)	0.80 ^{***}	0.71 – 0.90	0.68^{***}	0.58 – 0.80
>6 sex partners in past 6 months	1690	40.6	973 (58)	1.48 ^{***}	1.33 – 1.65	1.07	0.86 – 1.33
Sex Partner: Casual Male	3223	66.8	1749 (67)	1.05	0.95 – 1.17		
Sex Partner: Transgender Woman	70	1.7	35 (50)	0.89	0.63 – 1.25		
Sex Partner: Wife or Girlfriend	954	22.8	515 (54)	1.05	0.87 – 1.27		
Sex Partner: Paid Male	323	7.7	199 (62)	1.46 ^{**}	1.14 – 1.88	1.17	0.97 – 1.42
Sex Partner: Paid Female	85	2	36 (42)	0.65 ^{**}	0.46 – 0.90	0.59^{**}	0.38 – 0.84
Drug or alcohol use with sex in past 12 months	1246	29.8	786 (63)	1.80 ^{***}	1.49 – 2.16	1.43^{**}	1.17 – 1.76
Diagnosed with STI in past 12 months	286	6.8	203 (71)	2.29 ^{***}	1.68 – 3.10	1.78^{**}	1.25 – 2.53
Structural							
"Out" to anyone about sex with or attraction to men	2508	60	1483 (59)	1.86 ^{***}	1.59 – 2.18	1.35^{***}	1.18 – 1.54
Disclosed having sex with men to a doctor	938	22.4	740 (79)	4.64 ^{***}	3.74 – 5.76	3.55^{***}	2.76 – 4.58
Easy to access free HIV testing	1559	37.3	953 (61)	1.69 ^{***}	1.51 – 1.88	1.11	0.95 – 1.30
Aware of a comfortable HIV testing site	1965	47	1363 (69)	3.60 ^{***}	3.13 – 4.14	3.01^{***}	2.63 – 3.45
Perceived HIV stigma ^b (Mean ± SD)	2.5	±0.81	2.45 (0.80)	0.84 ^{***}	0.79 – 0.90	0.85^{***}	0.76 – 0.94

* p<.05,

** p<0.01,

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

p < 0.001

g Multivariable logistic regression model adjusted for all other covariates in the column for which data is presented.

g Increasing values indicate higher levels of perceived stigma on a scale of 1–5.