Published in final edited form as:

Arthritis Rheumatol. 2018 July ; 70(7): 977–980. doi:10.1002/art.40454.

Functional Connectivity: Dissecting the Relationship between the Brain and “Pain Centralization” in Rheumatoid Arthritis

Yvonne C. Lee, MD, MMSc¹, Vitaly Napadow, PhD², and Marco L. Loggia, PhD²

¹Division of Rheumatology, Northwestern University Feinberg School of Medicine, Chicago, IL

²Athinoula A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA

Pain is a frequent and disabling symptom experienced by individuals with inflammatory arthritis. Over 50% of patients with inflammatory arthritis report pain visual analog scale (VAS) scores 30/100 mm, despite treatment with disease-modifying antirheumatic drugs (DMARDs) (1). Patients often equate pain with peripheral inflammation, as evidenced by studies demonstrating that pain is the primary factor influencing patient global assessment of disease activity (2). However, recent clinical studies suggest that rheumatoid arthritis (RA) patients may exhibit signs of central sensitization, which may be why treating peripheral inflammation does not always translate into effective pain relief (3). The only study to directly examine the predictive effect of temporal summation, a measure of central sensitization, however, did not observe a significant association between temporal summation and DMARD response (4). Thus, the extent to which RA patients demonstrate neural mechanisms consistent with central sensitization is still unclear.

Advances in neuroimaging have enabled the assessment of functional connectivity between brain regions, allowing researchers to better understand the neural mechanisms underlying spontaneous clinical pain states. Functional brain connectivity assesses synchronization in activity displayed by two or more brain regions when they are “communicating” (e.g., when one region is exchanging information with the other). The connectivity between the insula cortex and a network of brain regions collectively known as the “Default-Mode network” (DMN) (a group of interconnected brain regions including the medial prefrontal cortex (MPFC), posterior cingulate cortex, precuneus, inferior parietal lobule, hippocampal formation and lateral temporal cortex (5)) has attracted particular attention in recent years. In healthy subjects, anterior and middle insula activity typically shows no correlation (or, sometimes, weak negative correlation) with DMN regions. In patients with chronic pain disorders, however, insula subregions can become functionally connected with the DMN. Following the original observation by Napadow et al. in fibromyalgia (FM) patients (6), an elevation of connectivity between the insula and the DMN (or to a specific core DMN

Corresponding author: Yvonne C. Lee, MD, MMSc, Division of Rheumatology, Northwestern University Feinberg School of Medicine, 633 North St. Clair Street, 18th Floor, Chicago, IL 60611. yvonne.lee@northwestern.edu. Telephone: (312) 503-1960. Fax: (312) 503-0994.

DR. YVONNE CLAIRE LEE (Orcid ID : 0000-0002-2105-3393)

Conflicts of interest: YCL reports receiving grant funding from Pfizer. She also served as an unpaid consultant on an advisory committee for Lilly.

region, such as the MPFC) has been documented in several pain conditions, including non-inflammatory and inflammatory chronic low back pain (LBP) (7), (8) (9), osteoarthritis (OA) (8) and migraine (10). Intriguingly, the strength of DMN-insula connectivity was positively correlated with clinical pain severity in many reports (6–8), although at least one study reported negative correlations in the context of an acute migraine attack (10). In addition, DMN-insula connectivity was found to be reduced after successful pharmacological (11) and nonpharmacological treatment (12), therefore raising the possibility that this feature may one day be considered an imaging biomarker of pain perception.

The study by Basu et al. tested the hypothesis that RA patients demonstrate neuronal hallmarks of pain centralization similar to those observed in FM (13). To this end, they enrolled 54 RA patients who met the 2010 American College of Rheumatology (ACR)/European League Against Rheumatism (EULAR) classification criteria and had experienced clinically significant fatigue during the past three months. The subjects underwent functional magnetic resonance imaging (fMRI) and an independent component analysis assessed functional connectivity between four networks of interest, including the DMN, and the rest of the brain. Subjects also completed the 2011 ACR FM survey, a measure of “fibromyalginess” (FMness), which is thought to represent the clinical manifestations of pain centralization (14). In this study, the average score on the 2011 ACR FM survey was 13.2 (the cut-off for the ACR modified preliminary criterion for FM diagnosis is 13), which is compatible with previous studies suggesting that a significant number of RA patients demonstrate comorbid FM. Of note, the ACR FM survey score was significantly positively associated with DMN-insula functional connectivity.

With these results in a population of RA patients, Basu et al. reinforce the generalizability of DMN-insula connectivity as a potential marker across etiologically heterogeneous pain conditions. However, the distinction between this study and studies of other conditions is that, unlike previous reports, there was no correlation between DMN-insula connectivity and spontaneous clinical pain severity, assessed at the time of the scan. By showing that DMN-insula was more associated with FMness than current pain severity, this study suggests another interpretation about the potential functional significance of this marker: not as a marker of “pain intensity” per se, but rather of “pain centralization”. It may be that previous studies demonstrated an association with clinical pain severity because pain intensity is more closely associated with measures of pain centralization (e.g., FMness) in functional pain disorders than in RA, which is known to have a significant peripheral component (e.g., peripheral joint inflammation). Of note, however, an fMRI study of patients with ankylosing spondylitis, a systemic rheumatic disease characterized by inflammatory back pain, did report associations between back pain severity and DMN connectivity with the salience network (of which the insula is a prominent component) (9). Thus, future studies are needed to better understand the relationship between DMN-insula activity and pain severity in systemic inflammatory conditions. In particular, we recommend that studies include different measures of pain (e.g., pain severity, FMness, pain interference pain catastrophizing, etc.) to better define the nature of observed associations.

Strengths of the Basu et al. study include the assessment of multiple relevant clinical characteristics, including age, sex, amitriptyline use, inflammatory disease activity measures (e.g., CRP and the Disease Activity Score in 28 joints), and levels of pain, fatigue, sleep disturbance and depression. By including age, sex, CRP and amitriptyline use in the linear regression models, Basu et al. was able to take into account the roles of these variables as possible confounders of the association between network-whole brain connectivity and FMness. Ultimately, adjusting for these variables did not change the magnitude of the association, which is reassuring that the observed associations are not an artifact of confounding due to relationships between the covariates, network-whole brain connectivity, and FMness. The assessment of pain, fatigue, sleep disturbance, and depression also enabled analyses examining correlations between these variables and DMN-insula connectivity. These analyses were post-hoc and, therefore, should be considered exploratory. Nevertheless, these results provide clues to the underlying cause of the association between DMN-insula connectivity and FMness, suggesting that this association may reflect a distinct phenotype associated with pain centralization.

Another strength of the study is the large sample size. With a study sample of 54 RA patients, this is the largest fMRI study to assess functional connectivity in individuals with RA. A sufficient sample size is required for stable and reproducible results and to enable correction for multiple testing, which is inherent in neuroimaging studies. Each fMRI image volume can be comprised of hundreds of thousands of voxels, or volume elements, each of which is used for separate statistical testing (15). Fortunately, neuroimagers take advantage of the fact that “genuine brain activity” tends to cluster in regions usually spanning many adjacent voxels (as opposed to random noise, which can lead to spurious small clusters in a “salt and pepper” pattern) (16). In this study, per commonly adopted procedure, the statistical significance of each cluster was determined based on the size of the cluster and a family wise error cluster corrected p-value of < 0.05 , thereby accounting for multiple comparisons.

Despite these strengths, there are some limitations. Perhaps the most important is the absence of a control group, which limits whether we can truly conclude that DMN-insula connectivity is significantly altered in RA patients. Another potential limitation is that the data used to perform functional connectivity analyses were collected while the participants were engaged in a cognitive task (the Paced Auditory Serial Attention Test) rather than at rest. While computing connectivity metrics during a task is not uncommon (17), this feature may limit our ability to directly compare the results from the Basu et al.’s study with those from previous studies evaluating the role of DMN in chronic pain, as most or all of them performed connectivity analyses using unconstrained resting-state data.

As the authors note, another primary limitation is generalizability. Inclusion criteria required that participants have a score > 3 on the Chalder Fatigue Binary Scale, restricting the study sample to individuals with clinically meaningful levels of fatigue. Few studies have used the Chalder Fatigue Scale to assess fatigue in RA, and, even fewer have reported on the binary scoring system (18). Thus, it is difficult to ascertain what a score > 3 means in the context of RA patients, and it would be informative to know the proportion of interested participants who were screened out because of this criterion. Future studies are needed to determine if

the relationship between DMN-insula connectivity and FMness holds among individuals with lower levels of fatigue.

Finally, as with all fMRI studies, this study is limited in that it cannot provide direct information on the neurophysiologic processes underlying the observed neuroimaging signals. This study employed a standard fMRI technique using an endogenous contrast mechanism called blood oxygenation level dependent (BOLD) imaging. BOLD fMRI takes advantage of neurovascular coupling in response to brain activity and serves as a proxy for neural activity. However, within each neuroimaging voxel, hundreds of thousands to millions of neurons exist, and the dynamic inter-relationships between these neurons is complex. Thus, the interpretation of fMRI signal to neural activity, or “connectivity” as was used by Basu et al., requires several assumptions; the validity of which may differ from study to study (19).

Despite these limitations, fMRI can provide novel insights into the central nervous system (CNS) pathways involved in the expression of difficult to quantify symptoms, such as pain and fatigue, in RA. This study extends findings found in many previous studies of non-inflammatory pain conditions to RA, a systemic inflammatory condition. Specifically, this study provides evidence highlighting DMN-insula connectivity as one of the most reliable imaging markers of chronic pain reported in the literature thus far.

For rheumatologists, the implications of this observation are broad. First, by identifying associations between aberrant brain functional connectivity and phenotypic characteristics (primarily FMness and secondarily fatigue and sleep disturbance), this study underscores the importance of considering CNS factors, in addition to peripheral joint inflammation, when evaluating symptoms experienced by our RA patients. Second, by showing that despite associations between DMN-insula connectivity and FMness, DMN-insula connectivity was not associated with pain severity, this study highlights the importance of carefully considering the types of pain measures included in studies. FMness includes assessments of widespread pain (e.g., pain distribution) and somatic symptoms, while the pain VAS only assesses pain intensity. Though related, these concepts are inherently different. Third, by providing evidence for a neurobiological underpinning for FMness in RA, this study points toward the potential role of treatment strategies previously shown to modulate DMN-insula connectivity (e.g., acupuncture, GABA analogs) in individuals with RA and high levels of FM symptoms (20).

In summary, this study by Basu et al. is an important step towards understanding the role of the brain in modulating pain in patients with systemic inflammatory conditions. Future studies examining the generalizability (e.g., in studies including RA patients with lower levels of fatigue, studies of newly diagnosed and hence drug-naïve RA patients, etc.), reproducibility (e.g., in observational longitudinal studies), and responsiveness to interventions (e.g., in clinical trials) of these findings are needed. As more neuroimaging studies are performed and published, rheumatologists should become familiar with the strengths and limitations of these types of studies. Characterizing the complex interrelationships between the brain, peripheral nervous system and immune pathways (both peripheral and central) holds promise for the development of safe and effective pain

management strategies, which are sorely needed to improve quality of life for our patients with systemic inflammatory conditions.

Acknowledgments

Funding:

Dr. Lee's work was supported by NIAMS R01 AR064850. Dr. Napadow's work was supported by NCCIH R61 AT009306, NCCIH R01 AT007550, and NIAMS R01 AT007550.

References

1. Rifbjerg-Madsen S, Christensen AW, Christensen R, Hetland ML, Bliddal H, Kristensen LE, et al. Pain and pain mechanisms in patients with inflammatory arthritis: A Danish nationwide cross-sectional DANBIO registry survey. *PLoS One*. 2017; 12(7):e0180014. [PubMed: 28686639]
2. Ferreira RJO, Dougados M, Kirwan JR, Duarte C, de Wit M, Soubrier M, et al. Drivers of patient global assessment in patients with rheumatoid arthritis who are close to remission: an analysis of 1588 patients. *Rheumatology (Oxford)*. 2017; 56(9):1573–8. [PubMed: 28859325]
3. Lee YC, Lu B, Edwards RR, Wasan AD, Nassikas NJ, Clauw DJ, et al. The role of sleep problems in central pain processing in rheumatoid arthritis. *Arthritis Rheum*. 2013; 65(1):59–68. [PubMed: 23124650]
4. Christensen AW, Rifbjerg-Madsen S, Christensen R, Dreyer L, Boesen M, Ellegaard K, et al. Ultrasound Doppler but not temporal summation of pain predicts DAS28 response in rheumatoid arthritis: a prospective cohort study. *Rheumatology (Oxford)*. 2016; 55(6):1091–8. [PubMed: 26983452]
5. Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci*. 2008; 1124:1–38. [PubMed: 18400922]
6. Napadow V, LaCount L, Park K, As-Sanie S, Clauw DJ, Harris RE. Intrinsic brain connectivity in fibromyalgia is associated with chronic pain intensity. *Arthritis Rheum*. 2010; 62(8):2545–55. [PubMed: 20506181]
7. Loggia ML, Kim J, Gollub RL, Vangel MG, Kirsch I, Kong J, et al. Default mode network connectivity encodes clinical pain: an arterial spin labeling study. *Pain*. 2013; 154(1):24–33. [PubMed: 23111164]
8. Baliki MN, Mansour AR, Baria AT, Apkarian AV. Functional reorganization of the default mode network across chronic pain conditions. *PLoS One*. 2014; 9(9):e106133. [PubMed: 25180885]
9. Hemington KS, Wu Q, Kucyi A, Inman RD, Davis KD. Abnormal cross-network functional connectivity in chronic pain and its association with clinical symptoms. *Brain Struct Funct*. 2016; 221(8):4203–19. [PubMed: 26669874]
10. Coppola G, Di Renzo A, Tinelli E, Di Lorenzo C, Scapeccia M, Parisi V, et al. Resting state connectivity between default mode network and insula encodes acute migraine headache. *Cephalalgia*. 2017 333102417715230.
11. Harris RE, Napadow V, Huggins JP, Pauer L, Kim J, Hampson J, et al. Pregabalin rectifies aberrant brain chemistry, connectivity, and functional response in chronic pain patients. *Anesthesiology*. 2013; 119(6):1453–64. [PubMed: 24343290]
12. Napadow V, Kim J, Clauw DJ, Harris RE. Decreased intrinsic brain connectivity is associated with reduced clinical pain in fibromyalgia. *Arthritis Rheum*. 2012; 64(7):2398–403. [PubMed: 22294427]
13. Basu N, Kaplan C, E I, T L, Harris RE, Murray A, et al. Neurobiological features of fibromyalgia are also present among rheumatoid arthritis patients. *Arthritis Rheumatol*. 2018
14. Wolfe F, Clauw DJ, Fitzcharles MA, Goldenberg DL, Hauser W, Katz RS, et al. Fibromyalgia criteria and severity scales for clinical and epidemiological studies: a modification of the ACR Preliminary Diagnostic Criteria for Fibromyalgia. *J Rheumatol*. 2011; 38(6):1113–22. [PubMed: 21285161]
15. Lindquist MA. The statistical analysis of fMRI data. *Statistical Science*. 2008; 23:439–64.

16. Woo CW, Krishnan A, Wager TD. Cluster-extent based thresholding in fMRI analyses: pitfalls and recommendations. *Neuroimage*. 2014; 91:412–9. [PubMed: 24412399]
17. Beaty RE, Kenett YN, Christensen AP, Rosenberg MD, Benedek M, Chen Q, et al. Robust prediction of individual creative ability from brain functional connectivity. *Proc Natl Acad Sci U S A*. 2018
18. Hewlett S, Dures E, Almeida C. Measures of fatigue: Bristol Rheumatoid Arthritis Fatigue Multi-Dimensional Questionnaire (BRAFMQ), Bristol Rheumatoid Arthritis Fatigue Numerical Rating Scales (BRAFNRS) for severity, effect, and coping, Chalder Fatigue Questionnaire (CFQ), Checklist Individual Strength (CIS20R and CIS8R), Fatigue Severity Scale (FSS), Functional Assessment Chronic Illness Therapy (Fatigue) (FACIT-F), Multi-Dimensional Assessment of Fatigue (MAF), Multi-Dimensional Fatigue Inventory (MFI), Pediatric Quality Of Life (PedsQL) Multi-Dimensional Fatigue Scale, Profile of Fatigue (ProF), Short Form 36 Vitality Subscale (SF-36 VT), and Visual Analog Scales (VAS). *Arthritis Care Res (Hoboken)*. 2011; 63(Suppl 11):S263–86. [PubMed: 22588750]
19. Logothetis NK. What we can do and what we cannot do with fMRI. *Nature*. 2008; 453(7197):869–78. [PubMed: 18548064]
20. Napadow V, Harris RE. What has functional connectivity and chemical neuroimaging in fibromyalgia taught us about the mechanisms and management of ‘centralized’ pain? *Arthritis Res Ther*. 2014; 16(5):425. [PubMed: 25606591]