Characterization of a 16SrII subgroup D phytoplasma strain associated with *Calendula officinalis* phyllody in Iran

Seyyed Alireza Esmailzadeh Hosseini¹ · Mohammad Salehi² · Ghobad Babaie³ · Assunta Bertaccini⁴

Received: 7 January 2018 / Accepted: 18 June 2018 / Published online: 26 June 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Calendula officinalis plants with phyllody symptoms (CaoP) were observed in Yazd and Ashkezar (Yazd province, Iran) during 2013–2016. Twenty-one symptomatic and four asymptomatic plants were transferred individually to the greenhouse and potted for the biological and molecular characterization of associated phytoplasma. The dodder transmission from symptomatic potted marigold plants, induced virescence, phyllody and witches' broom symptoms in periwinkle. Total DNAs extracted from symptomatic and symptomless plants and dodder-inoculated periwinkles were subjected to nested PCR assay using primer pairs amplifying phytoplasma ribosomal DNA. Expected PCR amplification was detected in all CaoP plant and dodder-inoculated periwinkle samples. RFLP analysis of the amplicons obtained in direct PCR with P1/P7 primers using *RsaI*, *AluI*, *MseI*, *HinfI* and *HaeIII* restriction enzymes showed profiles identical to each other and related to phytoplasmas in all the 21 positive samples. R16mF2/R16mR2 amplicons from six CaoP plant samples were sequenced where consensus sequences had 100% of identity among each other. R16F2n/R16R2-trimmed sequences (1248 bp) of representative samples from Yazd and Ashkezar were deposited in GenBank under accession numbers KU297202 and MH065715, respectively. BLAST search and phylogenetic analysis showed that the CaoP phytoplasma had 99% homology and clusters with phytoplasmas in group 16SrII. Computer-simulated analysis using *iPhyClassifier* suggests that the CaoP RFLP 16S rRNA gene pattern was identical to 16SrII-D phytoplasmas subgroup. Phytoplasma strains (16SrII-D) were reported as alfalfa witches' broom disease agent previously in the same geographic areas, and it is possible that alfalfa plays a role in the epidemiology of CaoP disease or vice-versa.

Keywords Pot marigold · Alfalfa witches' broom · RFLP · Iran

Introduction

Phytoplasmas are cell wall-less prokaryotes associated to plant diseases worldwide and transmitted mainly by leafhoppers and psyllids and very often causing quite devastating economic losses (Bertaccini et al. 2014). Phytoplasmas mainly produce characteristic disease symptoms including yellowing, discoloration, witches' broom, stunting, virescence and flower phyllody (Lee et al. 2000; Bertaccini and Duduk 2009). Ornamental plants have been reported as hosts of different phytoplasmas worldwide (Chaturvedi et al. 2010; Gera et al. 2006; Marcone et al. 1997; Omar and Alsohim 2016; Rani et al. 2014; Wang and Hiruki 2001) and in particular in Iran, phytoplasmas related to different group from various plants such as *Limonium sinuatum* with phyllody and mild stunting symptoms (16SrI-C), *Gomphocarpus physocarpus* with witches' broom, dwarfing, yellowing and purpling, leaf rolling virescence, decline and

✉ Seyyed Alireza Esmailzadeh Hosseini
saesmailzadeh@iripp.ir

- ¹ Plant Protection Research Department, Yazd Agricultural and Natural Resources Research and Education Center, AREEO, Yazd, Iran
- ² Plant Protection Research Department, Fars Agricultural and Natural Resources Research and Education Center, AREEO, Zarghan, Iran
- ³ Plant Protection Research Department, Chaharmahal and Bakhtiari Agricultural and Natural Resources Research and Education Center, AREEO, Shahrekord, Iran
- ⁴ Department of Agricultural and Food Sciences (DISTAL), Alma Mater Studiorum, University of Bologna, Bologna, Italy

seed sterility (16SrI-B), *Tanacetum parthenium* with stunting and phyllody symptoms (16SrI-B), *Tagetes patula* with witches' broom, virescence, early decline and purpling of leaves (16SrI-B), *Coreopsis lanceolata* with witches' broom, dwarfing and phyllody symptoms (16SrI-B) (Babaie et al. 2007), *Catharanthus roseus* with yellowing, dwarf, witches' broom and phyllody (16SrI-A and 16SrVI-A) (Babaie et al. 2007; Fattahi et al. 2016), *Rudbeckia hirta* with dwarfing, phyllody and virescence (16SrI-A) (Babaie et al. 2007) and *Erysimum cheiri* with phyllody and witches' broom symptoms (16SrII) (Asghari Tazehkand et al. 2010) have been described.

Calendula officinalis L. (family Asteraceae), commonly known as marigold or pot marigold, is cultivated for ornamental purposes and also for many medicinal, culinary and cosmetic uses. Phyllody symptoms named CaoP were observed in a collection of ornamental plants located at the Yazd Agricultural and Natural Resources Research and Education Center in Iran (Esmailzadeh Hosseini et al. 2008) adjacent to alfalfa fields showing epidemic incidence of alfalfa witches' broom (AWB) disease associated with phytoplasma presence (Esmailzadeh Hosseini et al. 2011, 2016d). Molecular and biological characterization of the phytoplasma strain associated with CaoP disease in pot marigold was, therefore, carried out.

Materials and methods

Plant sampling

C. officinalis plants with phyllody symptoms were collected during 2013–2016 at Yazd (31°54'58"N, 54°16'41"E) and Ashkezar (32°03'42"N, 54°10'33"E) (Iran) and transferred to an insect-proof greenhouse. In each area, five pot marigold fields were selected randomly and sampled at five points within 1 m². The number of symptomatic plants and the total number of plants in each sampled square meter were counted for the disease incidence determination. Collectively, 21 symptomatic and 4 asymptomatic plants were used for biological and molecular studies.

Propagation and maintenance of CaoP agent

For propagation and maintenance of the CaoP agent, transmission trials via dodder (*Cuscuta campestris* Yank.) inoculation were carried out in 21 periwinkle [*Catharanthus roseus* (L.) G. Don] and 21 pot marigold plants maintained in an insect-proof greenhouse. Dodder seeds were germinated on moist filter paper and seedlings were transferred to healthy seed-grown sugar beet (*Beta vulgaris* subsp. *vulgaris*) plants. After the colonization of the sugar beet plants, a dodder strand was connected to one symptomatic

C. officinalis plant for 3-week colonization. This dodder was connected with 3–4 leaves of seed-grown periwinkle and healthy pot marigold plants. After 4 weeks, the periwinkle and pot marigold plants were freed from dodder and observed for symptoms of expression. Phytoplasma presence in the dodder-inoculated plants was verified by nested PCR assay.

DNA extraction and polymerase chain reaction (PCR)

The total DNA was extracted from 0.2 g of midrib tissue of fresh leaves from 21 symptomatic and 4 symptomless *C. officinalis* and from the tissues of dodder-inoculated plants using Zhang et al. (1998) procedure. Direct and nested PCR were done using P1/P7 and R16mF2/R16mR2 and R16F2n/R16R2 primer pairs, respectively (Deng and Hiruki 1991; Schneider et al. 1995; Gundersen and Lee 1996). PCR products were examined in 1.2% agarose gels in 1X TBE buffer [108 g Tris-HCl, 55 g boric acid, 40 ml EDTA (0.5M), pH 8.0] and DNA bands, stained with ethidium bromide, were visualized with a UV transilluminator.

RFLP analysis

A preliminary identification of the phytoplasma associated with CaoP was obtained by restriction fragment length polymorphism (RFLP) analysis of P1/P7 PCR products (1.8 kbp fragment). The amplicons were digested separately with each of the *RsaI*, *AluI*, *MseI*, *HinfI* and *HaeIII* restriction enzymes according to the instructions of the manufacturer (Fermentas, Vilnius, Lithuania). The restriction products were then separated by electrophoresis through a 2.5% agarose gel, stained by ethidium bromide and visualized with a UV transilluminator.

Virtual RFLP analysis using *iPhyClassifier* (Zhao et al. 2009) allowed to determine the CaoP phytoplasma ribosomal subgroup affiliation. The RFLP profile of the 1248-bp fragment (R16F2n/R16R2 region of 16S rRNA gene) of the CaoP phytoplasma strain was compared to those of phytoplasmas strains classified in 16SrII subgroups-A to -M with 17 restriction enzymes: *AluI*, *BamHI*, *BfaI*, *BstUI* (*ThaI*), *DraI*, *EcoRI*, *HaeIII*, *HhaI*, *HinfI*, *HpaI*, *HpaII*, *KpnI*, *MboI* (*Sau3AI*), *MseI*, *RsaI*, *SspI* and *TaqI* (Lee et al. 1998).

Sequencing and phylogenetic analysis

Selected R16mF2/R16mR2 primed nested PCR products (1400 bp) of CaoP phytoplasma were directly sequenced using the same primers used for amplification and internal primers designed by the sequencing company. The assembling and trimming of these sequences allow obtaining fragments corresponding to the R16F2n/R2 amplicons.

The homologous sequences search performed by Blast analyses at the National Center for Biotechnology Information (NCBI) (<http://www.ncbi.nlm.nih.gov>) was used to identify the phytoplasmas closest to the strain under study. The R16F2n/R2 sequences of 16S rRNA gene of 29 phytoplasmas including CaoP were aligned, and phylogenetic trees and sequence homologies were generated using MEGA7 software (Kumar et al. 2016). *Acholeplasma laidlawii* was used as an out-group to root the tree after a bootstrapping analysis was performed 1000 times to estimate the stability and support for the branches.

Results

Disease incidence and symptomatology

The characteristic symptoms of CaoP disease were leaf size reduction, yellowing, flower phyllody, virescence, proliferation and sterility, proliferation of axillary buds, witches' broom and stunting (Fig. 1a, b). During the survey, the average disease incidence was of 6.8, 6.4, 8 and 7.8% in Ashkezar fields and 10.2, 11, 10.4 and 12% in Yazd areas, respectively.

Dodder transmission

The success of dodder transmission was different between periwinkle and pot marigold plants: 17 and 14 out of the 21 periwinkles and pot marigold plants, respectively, developed virescence, phyllody and witches' broom symptoms. Differences were observed in latent period dodder-inoculated periwinkle and pot marigold plants with CaoP while symptoms were observed 6–12 and 8–12 weeks, respectively, after inoculation. The CaoP phytoplasma was graft propagated in periwinkle plants with 100% of efficiency, and virescence and phyllody symptoms were observed 4–7 weeks after grafting.

Polymerase chain reaction

After polymerase chain reaction using P1/P7 primers and nested reactions by R16mF2/R16mR2 and R16F2n/R16R2 primer pairs, DNA fragments of ~ 1.8, ~ 1.4 and ~ 1.25 kbp were obtained, respectively, from symptomatic *C. officinalis* and dodder- and graft-inoculated periwinkles and pot marigold, but not from symptomless pot marigold or periwinkle plants (data not shown).

RFLP analysis

Restriction fragment length polymorphism analysis of P1/P7 amplicons using *Rsa*I, *Alu*I, *Mse*I, *Hinf*I and *Hae*III

Fig. 1 Leaf size reduction, yellowing, phyllody, virescence, witches' broom and stunting exhibited by *Calendula officinalis* plants affected by the phyllody disease (a) compared to the healthy plants (b)

Fig. 2 Restriction fragment length polymorphism (RFLP) analysis of PCR products amplified with primers P1/P7 (1800 bp) of *C. officinalis* phyllody (CaoP) phytoplasma and alfalfa witches' broom phytoplasma (AWB) (16SrII-D) using *RsaI*, *AluI*, *MseI*, *HinfI* and *HaeIII* restriction enzymes. M, 100 bp DNA ladder

Fig. 3 Phylogenetic trees constructed using MEGA7 software (Kumar et al. 2016) by the neighbor-joining method (Saitou and Nei 1987). The R16F2n/R2 sequences of 16S rRNA gene from selected phytoplasmas and *A. laidlawii* as the out-group were used. CWBP cactus witches' broom phytoplasma. Bootstrapping was performed 1000 times to estimate the stability and support for the branches and its values > 95% are shown at the branch points. NCBI GenBank accession numbers for sequences are given after the phytoplasma name together with Country of detection in parenthesis. The phytoplasma ribosomal subgroups are listed before the phytoplasma names. The strains under study are in bold. The scale bar represents two nucleotide exchanges per 100 nucleotides

restriction enzymes showed identical patterns referable to those of phytoplasmas in all the positive samples (Fig. 2). The comparison of these patterns with those previously published for P1/P7 phytoplasma amplicons (Khan et al. 2002) clearly indicate identity with the restriction profiles of phytoplasmas enclosed in group 16SrII.

Nucleotide sequence and virtual RFLP analysis

The consensus sequences of the six R16mF2/R16mR2 amplicons obtained from the selected samples collected in Ashkezar and Yazd locations provided 1.2-kbp R16F2n/R2 amplicons and resulted in 100% identical to each other independently from the collecting location or the sampling period. Two sequences of phytoplasma strains from Yazd and Ashkezar were deposited in GenBank under accession numbers KU297202 and MH065715, respectively. BLAST search of these sequences showed a 99% identity with phytoplasmas enclosed in subgroup 16SrII and the phylogenetic analysis confirmed their clustering with strains enclosed in group 16SrII in the '*Candidatus* Phytoplasma australasia'-related strains clade (NCBI GenBank accession numbers Y10097 and JQ868448) (Fig. 3). Virtual RFLP patterns derived from in silico digestions of CaoP phytoplasma

Fig. 4 Comparison of computer-simulated virtual RFLP patterns derived from in silico digestions of phytoplasma 16S rRNA gene (1.2 kb fragment) between *C. officinalis* phyllody phytoplasma (a) and a reference strain of 16SrII subgroup D (b) ‘*Candidatus* Phy-

toplasma australasia’ (GenBank accession number Y10097) using the online *iPhyClassifier* program. MW, molecular weight marker phiX174 *Hind*III digested. Fragment sizes in base pair from top to bottom: 1353; 1078; 872; 603; 310; 281; 271; 234; 194; 118 and 72

sequences exhibited profiles referable to those of members of the 16SrII phytoplasma group and resulted identical to the pattern of phytoplasmas enclosed in the 16SrII-D subgroup (Fig. 4).

Discussion

The phytoplasma associated with *C. officinalis* phyllody in Iran was identified as a member of the 16SrII-D subgroup. The phytoplasma presence was reported in India in the same species showing virescence symptoms (Rani et al. 2014), moreover phytoplasmas belonging to peanut witches’ broom group (16SrII) were detected in Saudi Arabia (Omar and Alsohim 2016) and phytoplasmas belonging to aster yellows (16SrI) group were identified in Italy and Canada, respectively (Marcone et al. 1997; Wang and Hiruki 2001).

A wide range of plant species was reported as infected by peanut witches’ broom phytoplasma group (16SrII) in Iran. This phytoplasma group, according to 16S rRNA gene RFLP analyses, encloses a large number of subgroups: 16SrII-A, 16SrII-B (‘*Ca. P. aurantifolia*’), 16SrII-C, 16SrII-D (‘*Ca. P. australasia*’), 16SrII-E, 16SrII-F, 16SrII-G, 16SrII-H, 16SrII-I, 16SrII-J, 16SrII-K, 16SrII-L (Bertaccini et al. 2014; Lee et al. 1998) and 16SrII-M (Salehi et al. 2015b). Among these groups, the 16SrII-D is associated in Iran with destructive diseases such as alfalfa witches’ broom (Esmailzadeh Hosseini et al. 2015a, b, c, 2016a, b, c; Salehi et al. 2011), tomato witches’ broom (Salehi et al. 2014), parsley phyllody (Salehi et al. 2016d), squash phyllody (Salehi et al. 2015b), garden beet witches’ broom (Mirzaie et al. 2007),

sesame phyllody (Salehi et al. 2016c), sunflower phyllody (Salehi et al. 2015a), carrot witches’ broom (Salehi et al. 2016b) and pomegranate little leaf (Salehi et al. 2016a). The CaoP phytoplasma strains showed RFLP patterns identical to each other in all the positive samples and also identical with alfalfa witches’ broom phytoplasma strain (16srII-D) (Esmailzadeh Hosseini et al. 2016a) identified as the major disease agent in alfalfa growing areas adjacent to *C. officinalis*. It is, therefore, possible to hypothesize that alfalfa plays a role in the epidemiology of *C. officinalis* phyllody disease or vice versa.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Asghari Tazehkand S, Hosseini Pour A, Heydarnejad J, Massumi H, Azadvar M (2010) Identification of phytoplasmas associated with cultivated and ornamental plants in Kerman province. Iran J Phytopathol 158(11–12):713–720
- Babaie G, Khatabi B, Bayat H, Rastgou M, Hosseini A, Salekdeh GH (2007) Detection and characterization of phytoplasma infecting ornamental and weed plants in Iran. J Phytopathol 155(6):368–372
- Bertaccini A, Duduk B (2009) Phytoplasma and phytoplasma diseases: a review of recent research. Phytopathol Mediterr 48:355–378
- Bertaccini A, Duduk B, Paltrinieri S, Contaldo N (2014) Phytoplasmas and phytoplasma diseases: a severe threat to agriculture. Am J Plant Sci 5:1763–1788

- Chaturvedi Y, Rao G, Tiwari A, Duduk B, Bertaccini A (2010) Phytoplasma on ornamentals: detection, diversity and management. *Acta Phytopathol Entomol* 45(1):31–69
- Deng S, Hiruki C (1991) Amplification of 16S rRNA genes from culturable and non-culturable mollicutes. *J Microbiol Methods* 14:53–61
- Esmailzadeh Hosseini SA, Salehi M, Firooz R, Shamszadeh M (2008) Occurrence of marigold phyllody in Yazd province. In: 18th Iranian plant protection congress. Faculty of Agriculture, University of Bu-Ali Sina, Hamedan, Iran, p 418
- Esmailzadeh Hosseini SA, Salehi M, Khanchezar A, Shamszadeh M (2011) The first report of a phytoplasma associated with pot marigold phyllody in Iran. *Bull Insectol* 64(Suppl):109–110
- Esmailzadeh Hosseini SA, Khodakaramian G, Salehi M, Fani SR, Bolok Yazdi HR, Raoufi D, Jadidi O, Bertaccini A (2015a) Status of alfalfa witches' broom phytoplasma disease in Iran. *Phytopathogenic Mollicutes* 5(1-Suppl):65–66
- Esmailzadeh Hosseini SA, Khodakaramian G, Salehi M, Fani SR, Mirchenari SM, Salehi E, Bertaccini A (2015b) Incidence, distribution and economic importance of alfalfa witches' broom disease in Sistan-Baluchestan (Iran) and characterization of associated phytoplasma. *Phytopathogenic Mollicutes* 5(2):84–90
- Esmailzadeh Hosseini SA, Salehi M, Khodakaramian G, Mirchenari SM, Bertaccini A (2015c) An up to date status of alfalfa witches' broom disease in Iran. *Phytopathogenic Mollicutes* 5(1):9–18
- Esmailzadeh Hosseini SA, Khodakaramian G, Salehi M, Bertaccini A (2016a) Characterization of 16SrII group phytoplasmas associated with alfalfa (*Medicago sativa*) witches' broom disease in diverse areas of Iran. *J Crop Prot* 5(4):581–590
- Esmailzadeh Hosseini SA, Khodakaramian G, Salehi M, Bertaccini A (2016b) First report of 16SrVI-A and 16SrXII-A phytoplasmas associated with alfalfa witches' broom diseases in Iran. *J Plant Pathol* 98(2):369
- Esmailzadeh Hosseini SA, Khodakaramian G, Salehi M, Bertaccini A (2016c) Molecular identification and phylogenetic analysis of phytoplasmas associated with alfalfa witches' broom diseases in the western areas of Iran. *Phytopathogenic Mollicutes* 6(1):16–22
- Esmailzadeh Hosseini SA, Salehi M, Mirchenari SM, Bertaccini A (2016d) First report of a 16SrII-D phytoplasma associated with *Calendula officinalis* phyllody in Iran. *New Dis Rep* 34:22
- Fattahi M, Salehi M, Sharzei A, Esmailzadeh Hosseini SA (2016) Partial biological and molecular characteristics of a phytoplasma associated with Behshahr (Mazandaran) periwinkle phyllody. *Iran J Plant Pathol* 52(1):135–141
- Gera A, Maslenin L, Rosner A, Zeidan M, Weintraub PG (2006) Phytoplasma disease in ornamental crops in Israel. *Acta Hort* 722:155–162
- Gundersen DE, Lee I-M (1996) Ultrasensitive detection of phytoplasmas by nested-PCR assays using two universal primer sets. *Phytopathol Mediterr* 35:144–151
- Khan AJ, Botti S, Al-Subhi AM, Gundersen-Rindal DE, Bertaccini AF (2002) Molecular identification of a new phytoplasma associated with alfalfa witches' broom in Oman. *Phytopathology* 92:1038–1047
- Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33(7):1870–1874
- Lee I-M, Gundersen-Rindal DE, Davis RE, Bartoszyk IM (1998) Revised classification scheme of phytoplasmas based on RFLP analyses of 16S rRNA and ribosomal protein gene sequences. *Int J Syst Evol Microbiol* 48:1153–1169
- Lee I-M, Davis RE, Gundersen-Rindal DE (2000) Phytoplasma: phytopathogenic mollicutes. *Ann Rev Microbiol* 54:221–255
- Marcone C, Ragozzino A, Seemüller E (1997) Detection and identification of phytoplasmas in yellows-diseased weeds in Italy. *Plant Pathol* 46:530–537
- Mirzaie A, Esmailzadeh Hosseini SA, Jafari-Nodooshan A, Rahimian H (2007) Molecular characterization and potential insect vector of a phytoplasma associated with garden beet witches broom in Yazd, Iran. *J Phytopathol* 155(4):198–203
- Omar AF, Alsohim AS (2016) Identification of new plant hosts of 16SrII group phytoplasmas in Saudi Arabia. *Phytopathogenic Mollicutes* 6(2):71–76
- Rani A, Misra P, Singh J, Kumar P, Rani R, Shukla P (2014) PCR based detection of phytoplasma association in pot marigold (*Calendula officinalis* L.) and guldawari (*Dendranthema grandiflora* L.). *Asian J Biol Sci* 9(2):238–241
- Saitou N, Nei M (1987) The neighbour-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Salehi M, Izadpanah K, Siampour M, Esmailzadeh Hosseini SA (2011) Polyclonal antibodies for the detection and identification of Fars alfalfa witches' broom phytoplasma. *Bull Insectol* 64(Suppl):59–60
- Salehi E, Salehi M, Taghavi SM, Izadpanah K (2014) 16SrII-D phytoplasma strain associated with tomato witches' broom in Bushehr province. *Iran J Crop Prot* 3(3):377–388
- Salehi M, Esmailzadeh Hosseini SA, Salehi E (2015a) Characterisation of a phytoplasma associated with sunflower phyllody in Fars, Isfahan and Yazd provinces of Iran. *New Dis Rep* 3:6
- Salehi M, Siampour M, Esmailzadeh Hosseini SA, Bertaccini A (2015b) Characterization and vector identification of phytoplasmas associated with cucumber and squash phyllody in Iran. *Bull Insectol* 68(2):311–319
- Salehi M, Esmailzadeh Hosseini SA, Rasoulpour R, Salehi E, Bertaccini A (2016a) Identification of a phytoplasma associated with pomegranate little leaf disease in Iran. *Crop Prot* 87:50–54
- Salehi M, Esmailzadeh Hosseini SA, Salehi E, Bertaccini A (2016b) Molecular and biological characterization of a 16SrII phytoplasma associated with carrot witches' broom in Iran. *J Plant Pathol* 98(1):83–90
- Salehi M, Esmailzadeh Hosseini SA, Salehi E, Bertaccini A (2016c) Genetic diversity and vector transmission of phytoplasmas associated with sesame phyllody in Iran. *Folia Microbiol* 62(2):99–109
- Salehi M, Esmailzadeh Hosseini SA, Salehi E, Bertaccini A (2016d) Occurrence and characterization of a 16SrII-D subgroup phytoplasma associated with parsley witches' broom disease in Iran. *J Phytopathol* 164(11–12):996–1002
- Schneider B, Seemüller E, Smart CD, Kirkpatrick BC (1995) Phylogenetic classification of plant pathogenic mycoplasma-like organisms or phytoplasmas. In: S Razin, Tully JG (eds) *Molecular and diagnostic procedures in mycoplasmaology*. Academic Press, San Diego, pp 369–380
- Wang K, Hiruki C (2001) Use of heteroduplex mobility assay for identification and differentiation of phytoplasmas in the aster yellows group and the clover proliferation group. *Phytopathology* 91:546–552
- Zhang YP, Uyemoto JK, Kirkpatrick BC (1998) A small-scale procedure for extracting nucleic acids from woody plants infected with various phytoplasmas for PCR assay. *J Virol Methods* 71:45–50
- Zhao Y, Wei W, Lee I-M, Shao J, Suo X, Davis RE (2009) Construction of an interactive online phytoplasma classification tool, iPhyClassifier, and its application in analysis of the peach X-disease phytoplasma group (16SrIII). *Int J Syst Evol Microbiol* 59:2582–2593