Production of a highly concentrated probiotic culture of *Lactococcus lactis* CECT 539 containing high amounts of nisin

Mónica Costas Malvido¹ · Elisa Alonso González¹ · David Outeiriño² · Nelson Pérez Guerra¹

Received: 23 November 2017 / Accepted: 18 June 2018 / Published online: 25 June 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

In this study, probiotic biomass and nisin productions by *Lactococcus lactis* CECT 539 were followed in two realcalized fed-batch cultures in diluted whey (DW) supplemented with KH_2PO_4 up to a total phosphorus concentration of 0.459 g/L. Increased biomass (5.12 g/L, 2.18×10^{10} CFU/mL) and nisin (235.23 BU/mL) concentrations were obtained in the culture fed concentrated whey and concentrated mussel processing waste (CMPW) medium supplemented with glucose up to a concentration of 400 g/L (CMPW + G medium) compared with similar fed-batch fermentations in DW medium. In the second fed-batch fermentation, the feeding medium CMPW + G was supplemented with KH_2PO_4 up to a TP concentration of 3.21 g/L. With this approach, increased production of biomass (5.49 g/L, 2.33×10^{10} CFU/mL) and nisin (257.59 BU/mL) was obtained. Considering the substantial availability of these wastes at very low prices from local dairy and mussel processing plants in Galicia, their use as culture media could offer an attractive alternative for a low-cost production of probiotic biomass and nisin at a high scale.

Keywords Fed-batch fermentation · Mussel processing waste · Nisin · Probiotic culture · Whey

Abbreviations

DW	Diluted whey medium
CW	Concentrated whey medium
MPW	Mussel processing waste medium
CMPW	Concentrated mussel processing waste medium
CG	400 g/L concentrated glucose
CMPW + G	Concentrated mussel processing waste medium supplemented with glucose up to a concentration of 400 g/L
TS	Total sugars concentration
TP	Total phosphorus concentration
CMPWCW-I	Realcalized fed-batch fermentation in DW medium containing 0.459 g TP/L with feeding with a mixture of CW and CMPW + G medium

CMPWCW-II	Realcalized fed-batch fermentation in DW medium containing 0.459 g TP/L with feeding with a mixture of CW and CMPW + G medium supplemented with KH_2PO_4 to obtain the same C/P relationship noted in the MRS medium
CDW	Cell dry weight concentration (g/L)
LA	Lactic acid concentration (g/L)
AA	Acetic acid concentration (g/L)
B	Butane-2,3-diol concentration (g/L)
G	Glucose concentration (g/L)
L	Lactose concentration (g/L)
Pr	Protein concentration (g/L)
TN	Total nitrogen concentration (g/L)
TP	Total phosphorus concentration (g/L)
Nis	Nisin concentration (BU/mL or IU/mL)
VCC	Viable cell count (CFU/mL)
P_{CDW}	Cell dry weight productivity (g/L/h)
P_{Nis}	Nisin productivity (BU/mL/h)
P_{LA}	Lactic acid productivity (g/L/h)
P_{AA}	Acetic acid productivity (g/L/h)
P_{B}	Butane-2,3-diol productivity (g/L/h)
$Y_{\text{Nis/CDW}}$	Yield of nisin on cell dry weight (BU/mg)
$Y_{\text{CDW/TS}}$	Yield of cell dry weight on total sugars consumed (g/g)

✉ Nelson Pérez Guerra
nelsonpg@uvigo.es

¹ Department of Analytical and Food Chemistry, Food Science and Technology Faculty, University of Vigo, Ourense Campus, As Lagoas s/n, 32004 Ourense, Spain

² Department of Chemical Engineering, Faculty of Sciences, University of Vigo, Ourense Campus, As Lagoas s/n, 32004 Ourense, Spain

$Y_{\text{Nis/TS}_c}$	Yield of nisin on total sugars consumed (BU/mg)
$Y_{\text{CDW/TN}_c}$	Yield of cell dry weight on total nitrogen consumed (g/g)
$Y_{\text{Nis/TN}_c}$	Yield of nisin on total nitrogen consumed (BU/mg)
E_L	Efficiency in lactose consumption (g lactose consumed/g lactose supplied)
E_G	Efficiency in glucose consumption (g glucose consumed/g glucose supplied)
E_{TN}	Efficiency in total nitrogen consumption (g TN consumed/g TN supplied)
E_{TP}	Efficiency in total phosphorus consumption (g TP consumed/g TP supplied)

Introduction

In recent years, there has been a considerable interest in using strains of viable lactic acid bacteria (genus *Lactobacillus*, *Pediococcus*, *Streptococcus* and *Enterococcus*) as probiotic microorganisms to replace the use of antibiotics in feeds in the European Union (Anadón et al. 2006). This interest is mainly attributed to the notion that the consumption of these probiotic bacteria can produce beneficial physiological effects on farm animals (Matijašić et al. 2004; Guerra et al. 2007a) and reduce the side effects associated with the excessive use of antibiotics, including allergic reactions, presence of antibiotic residues in edible animal products and the development of pathogenic and antibiotic-resistant bacteria (Pascual et al. 1999; Thorlacius and Jeppsson 2003; Fajardo et al. 2008).

In addition, given that these bacteria can produce bacteriocins and organic acids (Guerra and Pastrana 2003), their use as additives in animal feeds could allow control of the growth of undesirable harmful bacteria in the feed and the gut of the host, as observed in a previous study with weaned piglets (Guerra et al. 2007a). Consequently, the animals could expend lower amounts of energy fighting harmful bacteria, so they can use the energy obtained from the nutrients more efficiently for their own growth, as occurs when the animals consume a feed with antibiotics (Landers et al. 2012).

Thus, an efficient fermentation procedure should be designed to produce a highly concentrated culture of the probiotic bacterium to ensure that a higher number of cells can reach the host's gut and colonize it. With this approach, higher concentrations of bacteriocins and organic acids (mainly lactic acid) could also be produced, given that these products are commonly synthesized as growth-associated metabolites (Guerra et al. 2007b).

A previous study demonstrated that *Lactococcus lactis* CECT 539, a nisin-producing strain, exhibited many of the

probiotic criteria when a highly concentrated culture of this strain was used as an additive in a feed for weaned piglets (Guerra et al. 2007a). To produce this culture, realkalized fed-batch fermentation (CWCL culture) in diluted whey (DW, ~20 g of lactose/L) was performed using two lactose-containing substrates, concentrated whey (CW, ~50 g of lactose/L) and 400 g/L concentrated lactose (CL), as feeding media (Guerra and Pastrana 2003).

Given that the feeding with 400 g/L concentrated glucose (CG) increased stimulation of both probiotic biomass and nisin production compared with feeding with CL (Costas et al. 2016), mixtures of lactose- and glucose-containing substrates or only glucose-containing substrates were used to feed the cultures in subsequent studies (Costas et al. 2017). In these cases, the cultures were fed CW medium and CG (CWCG culture) or a concentrated mussel processing waste medium (CMPW, ~100 g of glucose/L) and CG (CMPWCG culture). The biomass and nisin levels obtained in the latter fermentation were considerably increased compared with control cultures (fed CL or CG) as well as CWCL and CWCG cultures (Costas et al. 2016, 2017). In these fermentations, both the CW and CMPW media were used as sources of sugars, protein, nitrogen, phosphorus, vitamins and mineral salts, whereas CG and CL were used as carbon sources.

According to these results, the joint use of CW and CMPW media as feeding substrates could represent a good alternative to continue improving the biomass and nisin synthesis by *L. lactis* CECT 539 in DW medium in realkalized fed-batch cultures. However, to our knowledge, no information is available on the use of CW and CMPW media as feeding substrates for the fed-batch productions of probiotic biomass and nisin.

Therefore, the aim of the present study was to design a realkalized fed-batch culture in DW medium using a mixture of these two feeding media to produce a highly concentrated probiotic culture of *L. lactis* CECT 539 containing high amounts of nisin. As a previous step, the batch production of biomass and nisin in DW medium was studied as a function of the initial total sugars and phosphorus concentrations to select the more appropriate fermentation substrate for the fed-batch fermentation. For this purpose, CMPW medium and KH_2PO_4 salt were used as carbon and phosphorus sources, respectively. This study was performed by combining response surface methodology with kinetic analysis of different cultures.

Materials and methods

Bacterial strains, media and culture conditions

The two bacteria used in this study (*L. lactis* CECT 539 and *Carnobacterium piscicola* CECT 4020) were provided

by the Spanish Type Culture Collection (CECT, Valencia, Spain) as freeze-dried cultures in glass ampoules. The two cultures were rehydrated with De Man, Rogosa and Sharpe (MRS) broth. After mixing, the suspensions were transferred into culture tubes containing MRS broth or agar slant, incubated at 30 °C for 24 h and subcultured weekly to obtain stock cultures. Prior to the batch and fed-batch fermentations, working cultures of *L. lactis* CECT 539 (the bacteriocin producer) and *C. piscicola* CECT 4020 (the target microorganism in the nisin activity assay) were prepared by inoculating 1 mL of the corresponding stock culture in 50 mL of MRS medium, followed by incubation at 30 °C for 12 h.

The culture media used in this study were prepared with whey [both concentrated (CW: the liquid remaining after the first cheese pressing) and diluted (DW: CW mixed with washing waters)] and mussel processing waste (MPW), which were obtained from local dairy and mussel processing coastal plants of the Galician region (Spain), respectively.

The MPW was first concentrated by ultrafiltration at 100 kDa to obtain an average glycogen level of approximately 100 g/L (CMPW medium), according to the methods described in a previous work (Murado et al. 1993).

Before being used as fermentation substrates, DW, CW and CMPW media were acidified to pH 4.5 with 5 M HCl and thermally treated (121 °C for 15 min) to denature the thermolabile protein fraction that precipitates at acidic pH values and hinders biomass measurements (Guerra et al. 2007b). The precipitates were then removed by centrifugation (12,000×g for 15 min). Given that *L. lactis* CECT 539 is not an amylolytic bacterium, the glycogen present in the CMPW medium (Murado et al. 1993) was saccharified to glucose by enzymatic hydrolysis at 40 °C for 1 h by mixing a volume of San Super 240 L commercial preparation of α -amylase (Novo Nordisk, Denmark) with 1000 volumes of CMPW medium (Guerra et al. 2007b). The mean composition of the resulting DW, CW and CMPW media is presented in Table 1.

Given that glucose was a better carbon source than lactose for nisin production by *L. lactis* CECT 539 (Costas et al. 2017), the batch production of biomass and nisin was studied in media prepared by mixing 100 volumes of DW medium with different volumes of CMPW medium before performing the realkalized fed-batch cultures. However, due to the low total phosphorus concentration present in the CMPW medium (Table 1), the concentration of this nutrient in the resulting mixed media decreased considerably. Thus, batch productions of biomass and nisin were studied as a function of the concentrations of total sugars (TS) and phosphorus (TP) using an orthogonal central composite design to select the most appropriate

Table 1 Mean composition (g/L) of the substrates used as culture media

Nutrient	DW	CW	CMPW
Lactose	22.61 ± 0.05	51.35 ± 0.46	–
Glucose	–	–	101.72 ± 0.17
Total nitrogen	0.455 ± 0.01	0.994 ± 0.02	0.540 ± 0.01
Total phosphorus	0.240 ± 0.02	0.472 ± 0.01	0.060 ± 0.00
Protein	2.07 ± 0.01	4.57 ± 0.04	3.47 ± 0.03

fermentation medium for the realkalized fed-batch cultures. In addition, the time courses of the growth, product formation, pH and nutrient consumption corresponding to each point of the experimental matrix were followed for 21 h. The TS levels in the mixed media ranged from the initial TS concentration of the DW medium (22.61 g/L) to that (51.35 g/L) of the CW medium (Tables 1, 2). The TP levels ranged from the initial TP concentration of the DW medium (0.240 g/L) to that (0.631 g/L) of the MRS medium (Table 2).

After mixing the corresponding volumes of DW and CMPW substrates to obtain the TS concentrations (actual value) defined in Table 2, the resulting mixed media were sterilized (121 °C/15 min). Then, each culture medium was supplemented with the corresponding sterile solution of KH_2PO_4 previously adjusted to pH 7.0 with 4 M NaOH (Guerra et al. 2001) to obtain the different TP levels defined in Table 2. The mean compositions of the different media obtained are presented in Table 2.

Batch fermentations in the different mixed media were performed under the same conditions [2% (v/v) inoculation rate, 30 °C, 200 rpm] as those described in a previous work (Costas et al. 2016). The culture samples, which comprise two experimental units (two flasks), were withdrawn at 3 h and used for analytical determinations (pH, biomass, nisin, total sugars, nitrogen and phosphorus, glucose, lactose and lactic acid).

Duplicate fixed-volume realkalized fed-batch cultures were performed, as previously described (Guerra and Pas-trana 2003; Costas et al. 2016, 2017) in two 13-L fermenters using the optimum fermentation medium obtained from the previous batch experiments. The first fermentation (named as CMPWCW-I) was fed a mixture of CW (without supplementation) and CMPW medium supplemented with glucose up to a concentration of 400 g/L (CMPW + G medium). The second fed-batch culture (named as CMPWCW-II) was fed a mixture of CW (without supplementation) and CMPW + G medium supplemented with KH_2PO_4 up to a TP concentration of 3.21 g/L.

Table 2 Experimental domain and codification of the independent variables (TS: total sugars and TP: total phosphorus) used in the experimental design, mean composition of the resulting culture media

and values of biomass (CDW) and nisin (Nis) calculated with models (1) and (2), respectively

No.	Independent variables		Actual values		Concentration of the other nutrients				Response variables	
	Coded values									
	TS	TP	TS (g/L)	TP (g/L)	Pr (g/L)	TN (g/L)	G (g/L)	L (g/L)	CDW (g/L)	Nis (BU/mL)
1	1	1	48.32	0.589	2.53 ± 0.02	0.479 ± 0.002	33.06 ± 0.09	15.26 ± 0.08	0.500	15.75
2	1	−1	48.32	0.281	2.53 ± 0.02	0.479 ± 0.002	33.06 ± 0.09	15.26 ± 0.08	0.492	14.97
3	−1	1	25.64	0.589	2.12 ± 0.03	0.453 ± 0.003	3.90 ± 0.07	21.74 ± 0.04	0.715	22.07
4	−1	−1	25.64	0.281	2.12 ± 0.03	0.453 ± 0.003	3.90 ± 0.07	21.74 ± 0.04	0.624	20.45
5	1.267	0	51.35	0.435	2.58 ± 0.05	0.483 ± 0.001	36.95 ± 0.05	14.40 ± 0.03	0.503	16.07
6	−1.267	0	22.61	0.435	2.07 ± 0.01	0.450 ± 0.001	–	22.61 ± 0.05	0.707	23.57
7	0	1.267	36.98	0.631	2.32 ± 0.03	0.466 ± 0.002	18.48 ± 0.03	18.50 ± 0.03	0.588	18.72
8	0	−1.267	36.98	0.240	2.32 ± 0.03	0.466 ± 0.002	18.48 ± 0.03	18.50 ± 0.03	0.550	17.28
9	0	0	36.98	0.435	2.32 ± 0.03	0.466 ± 0.002	18.48 ± 0.03	18.50 ± 0.03	0.631	20.82
10	0	0	36.98	0.435	2.32 ± 0.03	0.466 ± 0.002	18.48 ± 0.03	18.50 ± 0.03	0.627	20.87
11	0	0	36.98	0.435	2.32 ± 0.03	0.466 ± 0.002	18.48 ± 0.03	18.50 ± 0.03	0.633	21.15
12	0	0	36.98	0.435	2.32 ± 0.03	0.466 ± 0.002	18.48 ± 0.03	18.50 ± 0.03	0.625	20.31
13	0	0	36.98	0.435	2.32 ± 0.03	0.466 ± 0.002	18.48 ± 0.03	18.50 ± 0.03	0.622	21.66

Pr, TN, G and L are the mean concentrations of proteins, total nitrogen, glucose and lactose, respectively, in the corresponding culture media
No. number of the experiment

Analytical methods

Concentrations of biomass (as cell dry weight, CDW), lactic acid (LA), acetic acid (AA), butane-2,3-diol (B), glucose (G), lactose (L), total sugars (TS), protein (Pr), total nitrogen (TN) and phosphorus (TP) were determined in triplicate using methods described previously (Guerra and Pastrana 2003; Costas et al. 2016). Total viable cell numbers [as colony forming units (CFU)/mL] were enumerated in triplicate samples on MRS agar at 30 °C for 48 h (Fajardo et al. 2008).

Nisin activity determination

Cell-free supernatants (CFS, adjusted at pH 6.0) obtained after heating (80 °C/3 min) and centrifuging (27 200×g/15 min at 4 °C) acidified culture samples (pH 3.5) of *L. lactis* were used to determine the nisin activity (Nis) against *C. piscicola* CECT 4020 (test microorganism) according to the photometric method described by Cabo et al. (2001). Nisin activity [bacteriocin units (BU)/mL] was quantified as indicated by Murado et al. (2002). One BU/mL was defined as the reciprocal of the dilution of CFS causing 50% growth inhibition compared with control tubes, in which the diluted CFS was replaced by distilled sterile water (Fajardo et al. 2008). For comparative purposes, BU/mL values were converted to IU/

mL from a standard curve (IU vs. BU). Nisin standards, ranging from 1 to 1000 IU/mL, were obtained from a stock nisin solution (1000 IU/mL) prepared by dissolving Nisaplin™ (10⁶ IU/g, 2.5% nisin, Aplin and Barrett Ltd., Dorset, UK) in 0.02 M HCl (pH 2).

Statistical analyses

The different batch cultures were statistically compared in terms of the concentrations of the culture variables (culture pH, CDW, Nis, LA, G, L, TS, TN and TP) at the optimum fermentation time. The realkalized fed-batch cultures CMPWCW-I and -II were statistically compared with two similar fed-batch fermentations (CWCG and CMPWCG) of *L. lactis* CECT 539 in terms of the levels of the culture variables (culture pH, CDW, VCC, Nis, LA, AA, B, G, L, TS, TN and TP) and the fermentation parameters (productivities, yields and efficiencies). All the statistical comparisons were performed using the software package IBM SPSS Statistics for Windows (Version 21.0, Armonk, NY: IBM Corp., 2012) with the level of significance set at 5% (Costas et al. 2017).

An orthogonal central composite design based on two levels and two variables was used to study the effect of addition of the carbon (CMPW medium) and phosphorus (KH₂PO₄) sources on the production of biomass and nisin in DW medium. First, growth kinetics and bacteriocin production were followed at each experimental point (Table 2).

Subsequently, the experimental biomass (CDW, in g/L) and nisin (Nis, in BU/mL) data were smoothed using the logistic models (1) and (2), respectively (Guerra 2014):

$$CDW = \frac{CDW_{\max}}{1 + e^{(c-b \cdot t)}}, \quad \text{being} \quad c = \ln\left(\frac{CDW_{\max}}{CDW_{t=0h}} - 1\right), \quad (1)$$

$$Nis = \frac{Nis_{\max}}{1 + e^{(A_1 + A_2 \cdot t + A_3 \cdot t^2)}} - \frac{Nis_{\max}}{1 + e^{(A_1)}}, \quad (2)$$

where $CDW_{\max}$ and $Nis_{\max}$ are the maximum concentrations of biomass (g/L) and nisin (BU/mL); b (h^{-1}), A_1 (dimensionless), A_2 (h^{-1}) and A_3 (h^{-2}) are constants; and t is the time (h).

Finally, the smoothed biomass and nisin levels obtained at the optimal fermentation time were used in the abovementioned orthogonal central composite design as the response variables. Then, the results were analyzed using the experimental design module of the Statistica software package (Statistica 12.0 for Windows computer program manual; StatSoft Inc., 2013, Tulsa, OK, USA). The statistical significance of the empirical models and their coefficients were analyzed using the Fisher's F test for analysis of variance (ANOVA) and Student's t test, respectively.

Results and discussion

As a previous step to the design of the realkalized fed-batch fermentations, batch growth and nisin production by *L. lactis* CECT 539 were studied as a function of the TS and TP concentrations using response surface methodology and empirical modeling (Table 2). The fermentation media used consisted of mixtures of 100 volumes of DW medium with different volumes of CMPW medium, which were then supplemented with KH_2PO_4 . This salt was recognized as the best phosphorus source for nisin production (De Vuyst and Vandamme 1993). The results obtained (Figs. 1, 2) demonstrate that all the cultures stopped at 12 h of fermentation. After this time, the rates of biomass and nisin production and nutrient consumption decreased considerably.

With regard to the consumption of the carbon source (Fig. 2), glucose was the first sugar consumed by *L. lactis* CECT 539, and lactose consumption only started when the monosaccharide was completely consumed (cultures 3, 4, 6 and in that performed at the optimum conditions). In the other cases (cultures 1, 2, 5, 7, 8, and 9–13), only glucose was consumed, whereas the initial concentration of lactose remained constant throughout the cultures (Fig. 2).

In these batch fermentations, a low availability of nutrients (Pr, TN, TP and TS) in the cultures did not seem to

explain the growth cessation observed from the 12 h of incubation, because complete depletion of these nutrients had not occurred at this time (Figs. 1, 2).

Possible explanations of growth cessation could be the exhaustion of some micronutrient (amino acid, vitamin or salt) essential for the growth of the nisin-producing strain (Guerra et al. 2001) and/or the low pH values in the cultures at 12 h of incubation. These pH values were very close to the pH value of 4.72 (Fig. 1), at which the growth of *L. lactis* CECT 539 stopped in the batch culture in unsupplemented DW medium (Guerra et al. 2001; Costas et al. 2016). In fact, pH values at 12 h of incubation in the different cultures were considerably low compared with the optimum pH range (between 6.0 and 6.5) for efficient nutrient transport in *L. lactis* and *L. cremoris* strains (Poolman and Konings 1988).

Therefore, the concentrations of biomass and nisin predicted by models (1) and (2), respectively, at 12 h of fermentation were used in the experimental design to study the effects of the independent variables TP and TS on growth and bacteriocin synthesis by *L. lactis* CECT 539 (Table 2; Fig. 3).

The empirical models obtained for each dependent variable (CDW and Nis) with their significant coefficients ($P < 0.05$) were as follows:

$$CDW(g/L) = 0.63 - 0.08 \cdot TS + 0.02 \cdot TP - 0.02 \cdot TS \cdot TP - 0.01 \cdot TS^2 - 0.03 \cdot TP^2, \quad (3)$$

$$Nis(BU/mL) = 21.0 - 2.96 \cdot TS + 0.59 \cdot TP - 0.74 \cdot TS^2 - 1.88 \cdot TP^2. \quad (4)$$

The values of P (greater than 0.05) and F (less than 6.390) as well as the high values for the determination coefficients obtained for the two models (Table 3) indicated their high significance at the 95% confidence level.

The response surfaces generated with the empirical models (3) and (4) are presented in Fig. 3. Maximum biomass (0.725 g/L) and nisin (23.57 BU/mL) production at 12 h of incubation were obtained at $TS = 22.61$ g/L (coded value = -1.267) and $TP = 0.480$ g/L (coded value = 0.294) and at $TS = 22.61$ g/L and $TP = 0.459$ g/L (coded value = 0.156), respectively (Fig. 3). Thus, although the addition of CMPW volumes to the DW medium led to increase in both protein and total nitrogen concentrations in the resulting mixed media (Table 2), the increase in the TS concentration inhibited both growth and bacteriocin production by *L. lactis* CECT 539 (Figs. 1, 3).

Given that nisin has been classified as a pH-dependent primary metabolite in DW medium (Guerra et al. 2007b), the effects of the initial TP and TS concentrations on bacteriocin production can be analyzed by determining the

Fig. 1 Kinetics of culture pH, growth, product synthesis and nutrient consumption by *L. lactis* CECT 539 in batch cultures at the initial TS and TP concentrations defined in Table 1 and at the optimum TS and TP levels. Data reported are means $\pm$ standard deviations of two repeated experiments and three replicate measurements. The solid lines drawn through the experimental CDW and Nis data were obtained according to the logistic models (1) and (2), respectively

effects of the two independent variables on the growth of *L. lactis* CECT 539. Thus, the inhibition that the increase in the TS concentration produced on biomass production by *L. lactis* CECT 539 could be related to a reduction in the specific sugar uptake rate and consequently the activities of key glycolytic enzymes (phosphofructokinase, pyruvate kinase and lactate dehydrogenase). This effect results in significant reductions in glycolytic flux and growth rates in *L. lactis* strains (Papagianni et al. 2007a). The increase in TP from 0.240 to 0.480 g/L likely stimulated sugar intake and induced an increase in fructose 1,6-bisphosphate concentrations that allowed the cells to metabolize glucose efficiently, stimulating the growth of the strain CECT 539 (Levering et al. 2012; Goel et al. 2012). On the other hand, the growth inhibition observed at TP levels greater than 0.480 g/L could be explained by the fact that at these concentrations, the inorganic phosphate inhibited pyruvate kinase activity in *L. lactis* CECT 539 as previously observed for different *L. lactis* strains (Collins and Thomas 1974; Coccagn-Bousquet et al. 1996; Goel et al. 2012).

Duplicate batch cultures were performed under the optimized conditions for biomass and nisin synthesis to prove the reliability and accuracy of the empirical models (Experiment 14 in Figs. 1, 2) for predicting both productions. However, given that the optimum TP values for high biomass and nisin productions differed, the experiments were performed in DW medium (TS = 22.61 g/L) supplemented with KH_2PO_4 to obtain a TP concentration of 0.459 g/L (Figs. 1, 2). This TP concentration was selected as the optimum TP value, because the biomass concentration (0.725 g/L) predicted by the empirical model at TP = 0.480 g/L was similar to that (0.721 g/L) obtained at TP = 0.459 g/L when TS was fixed at a value of 22.61 g/L.

Under the optimum conditions, the experimental CDW and Nis concentrations obtained at 12 h of incubation were 0.722 g/L and 23.66 BU/mL, respectively, which are consistent with the values predicted by the empirical models (0.721 g/L and 23.57 BU/mL). These results indicate the high predictive capacity of the two empirical models obtained (Fig. 3).

Taking into account these observations, a realkalized fed-batch culture (CMPWCW-I fermentation) was performed using the DW medium containing 0.459 g TP/L as fermentation medium and the CMPW + G and CW media as feeding substrates. The feeding media were added to the growing

culture to replenish the amounts of total sugars consumed by the nisin-producing strain in each realkalization and feeding cycle (Costas et al. 2017). This approach could offer an attractive possibility to obtain further increase in the concentrations of probiotic biomass and nisin in DW medium while maintaining the TS concentration in the culture medium at non-inhibitory values (≤ 22.61 g/L). The results obtained are presented in Figs. 4, 5 and 6.

From a kinetic point of view, it can be noted that the CMPWCW-I culture evolved in a similar manner as that of previous realkalized fed-batch cultures of *L. lactis* CECT 539 in DW medium (Guerra and Pastrana 2003; Costas et al. 2016, 2017). Thus, the following features were observed in the CMPWCW-I fermentation:

- (i) nisin and biomass production (the left side of Fig. 5) and the pH values at the end of each realkalization and feeding cycle (the left side of Fig. 4) increased, revealing two logistic profiles (0–96 and 96–204 h) and suggesting that nisin was produced as a pH-dependent primary metabolite;
- (ii) proteins, nitrogen and phosphorus were consumed in parallel with biomass production and relatively low concentrations of these nutrients (0.459, 0.164 and 0.062 g/L, respectively) remained in the medium at the end of the incubation (Fig. 6);
- (iii) homolactic fermentation shifted to mixed-acid fermentation at 96 h of incubation (Fig. 5) when the nitrogen concentration in the media reached a value of 0.210 g/L (Fig. 6);
- (iv) the total amount of glucose consumed (147.98 g/L) was considerably increased ($P < 0.05$) compared with lactose (19.83 g/L) probably due to the repression that the presence of the monosaccharide produces on the expression of the *L. lactis* lactose operon (van Rooijen et al. 1992; Garrigues et al. 1997).

On the other hand, the concentration of viable cells was 2.00×10^{10} CFU/mL at 156 h of incubation and increased slightly until obtaining a final level of 2.18×10^{10} CFU/mL (the left side of Fig. 5), which was four orders of magnitude increased ($P < 0.05$) compared with the recommended dose of viable probiotic (10^6 CFU of probiotic per g or mL) necessary to observe beneficial effects in the host (Guerra et al. 2007a).

From comparisons between the CWCG, CMPWCG and CMPWCW-I cultures (Table 4), it can be noted that the highest levels of biomass, viable cells count and nisin were obtained in the CMPWCW-I fermentation ($P < 0.05$). However, the highest lactic acid, acetic acid and butane-2,3-diol concentrations were obtained in the CMPWCG culture ($P < 0.05$). In fact, the CDW, VCC and Nis concentrations obtained in the CMPWCW-I culture increase by 1.03-,

Fig. 2 Consumption of glucose (G) and lactose (L) and lactic acid production by *L. lactis* CECT 539 in the corresponding batch cultures shown in Fig. 1. Data reported are means $\pm$ standard deviations of two repeated experiments and three replicate measurements

1.07- and 1.05-fold compared with those obtained in the CMPWCG fermentation, which contained lactic acid, acetic acid and butane-2,3-diol concentrations that increased by 1.05-, 1.10- and 1.03-fold ($P < 0.05$).

However, given that the incubation times in these three fermentations were different (Table 4), it is more reasonable to compare the productivities obtained in these cultures than the final amounts of products synthesized. With this approach, the final biomass, nisin, lactic acid, acetic acid and butane-2,3-diol productivities in the CMPWCW-I culture were higher ($P < 0.05$) than those obtained in both the CWCG and CMPWCG fermentations (Table 4). This finding was probably due to an increase in the availability of nutrients (proteins, total nitrogen and phosphorus) and micronutrients (amino acids, vitamins or mineral salts) in the fermentation medium due to the feeding with CMPW and CW media. This finding suggests that the nisin-producing strain utilized these nutrients more efficiently for product formation in the CMPWCW-I culture. Thus, with this approach, a more concentrated and viable culture of *L. lactis* CECT 539 was produced compared with those obtained in the CWCG and CMPWCG fermentations.

The CMPWCW-I culture was also characterized with increased $Y_{\text{Nis}/X}$, $Y_{X/\text{TNc}}$ and $Y_{\text{Nis}/\text{TNc}}$ values, indicating that the strain CECT 539 was more productive and utilized the nitrogen source for both the biomass and bacteriocin synthesis in a more effective manner compared with the CMPWCG and CWCG cultures. However, the feeding with CMPW and CW media did not significantly improve ($P > 0.05$) the yields $Y_{X/\text{TSc}}$ and $Y_{\text{Nis}/\text{TSc}}$ with respect to the CMPWCG culture (Table 4). This finding was likely due to a similar efficiency in the utilization of the carbon sources (glucose and lactose) for both biomass and nisin production in both fermentations.

In the CMPWCW-I culture, the efficiencies in lactose (E_L) and glucose (E_G) consumption increased ($P < 0.05$), the efficiency in phosphorus consumption (E_{TP}) was similar, but the efficiency in nitrogen consumption (E_{TN}) decreased compared with the CMPWCG fermentation (Table 4). This observed decrease in the E_{TN} value in the CMPWCW-I fermentation could be related to increased accumulation of a nitrogen fraction that is difficult to metabolize by the strain CECT 539 in the culture medium as a consequence of the joint use of two protein-containing feeding substrates (CMPW and CW). In fact, DW, CW and CMPW media contain a nitrogen fraction that is not

consumed by the nisin-producing strain and remained in the fermentation media at the end of the cultures, as previously observed in both batch (Guerra et al. 2001) and real-kalized fed-batch cultures of *L. lactis* CECT 539 (Guerra et al. 2007b). For example, at 204 h of incubation (when the CMPWCW-I culture was stopped), the strain CECT 539 consumed increased amounts of nitrogen (0.446 g/L) in the CMPWCW-I fermentation compared with the CMPWCG culture (0.390 g/L). However, the cumulative level of nitrogen added in the first culture (0.709 g/L) at this time was considerably increased ($P < 0.05$) compared with the latter fermentation (0.513 g/L).

Considering the low total phosphorus concentration in the CMPW medium compared with DW and CW media (Table 1), it could be reasonable to think that its use as a feeding substrate could provoke a reduction in the TP concentration in the fermentation medium. Thus, new real-kalized fed-batch fermentation (CMPWCW-II) was performed under the same conditions as the previous culture. However, in this case, the CMPW + G medium was supplemented with KH_2PO_4 to obtain the same C/P relationship noted in the MRS medium. To avoid the use of a feeding medium with a relatively high phosphorus concentration, which could lead to the accumulation of this nutrient in the fermentation medium, supplementation with KH_2PO_4 was performed by considering the initial TS concentration (101.72 g/L) of the CMPW medium. Then, the initial TP concentration in this substrate after phosphorous supplementation was 3.21 g/L.

From the comparison between the two real-kalized fed-batch cultures (CMPWCW-I and -II), it could be noted that all the culture variables evolved in a similar manner in both cases (Figs. 4, 5, 6). However, in the CMPWCW-II culture, the acidification rate in each real-kalization cycle (Fig. 4; Table 4) and the productions of CDW, VCC, Nis, LA, AA and B (Fig. 5) were slightly increased (1.07-, 1.07-, 1.10-, 1.08-, 1.09- and 1.12-fold, respectively) compared with the CMPWCW-I culture ($P < 0.05$). In addition, the amounts of TS, G, L, Pr and TP consumed were similar ($P > 0.05$) in both real-kalized fed-batch cultures, but TN consumption was considerably increased ($P < 0.05$) in the latter fermentation (the right side of Fig. 6).

Supplementing the CMPW medium with KH_2PO_4 improved the productivities (P_{CDW} , P_{Nis} , P_{LA} , P_{AA} and P_{B}), the yields ($Y_{\text{Nis}/\text{CDW}}$, $Y_{\text{Nis}/\text{TSc}}$, $Y_{\text{CDW}/\text{TNc}}$ and $Y_{\text{Nis}/\text{TNc}}$), and the efficiencies in nutrient consumption (except E_L , which decreases slightly) in the CMPWCW-II culture. Higher increases were observed for the E_{TP} values (Table 4) and probably due to a high availability of TP in the culture medium as a consequence of the feeding with the phosphorus-supplemented CMPW medium.

Fig. 3 Response surfaces generated according to the predictions of models (3) and (4) and showing the effect of the initial concentrations of total sugars (TS) and phosphorus (TP) on biomass (CDW) production and nisin activity by *L. lactis* CECT 539 in culture media prepared with mixtures of DW and CMPW media supplemented with KH_2PO_4

Table 3 Significance of the coefficients and summarized data of analysis of variance (ANOVA) of models (3) and (4) obtained according to the experimental design defined in Table 2

Significance of factors

Factor	Biomass production model (3)			Nisin production model (4)		
	Coefficient $\pm$ SE	<i>t</i> (7)	<i>P</i> value	Coefficient $\pm$ SE	<i>t</i> (4)	<i>P</i> value
Constant	0.63 ± 0.00	194.7	0.00 ^a	21.0 ± 0.22	95.9	0.00 ^a
TS	-0.08 ± 0.00	-31.0	0.00 ^a	-2.96 ± 0.18	-16.1	0.00 ^a
TP	0.02 ± 0.00	-3.78	0.00 ^a	0.59 ± 0.18	3.19	0.03 ^a
TS·TP	-0.02 ± 0.00	-5.74	0.00 ^a	-0.21 ± 0.24	-0.86	0.44
TS ²	-0.01 ± 0.00	-3.78	0.00 ^a	-0.74 ± 0.22	-3.42	0.03 ^a
TP ²	-0.03 ± 0.00	-10.8	0.00 ^a	-1.88 ± 0.22	-8.65	0.00 ^a
TP ²	-0.03 ± 0.00	-10.8	0.00 ^a	-1.88 ± 0.22	-8.65	0.00 ^a

ANOVA

Source	SS	<i>df</i>	MS	<i>F</i> value	<i>P</i> value	<i>r</i> ²	Adj <i>r</i> ²
Biomass production model (3)							
Lack of fit	2.8×10^{-4}	3	9.4×10^{-5}	4.20	0.10 ^b	0.99	0.99
Pure error	8.9×10^{-5}	4	2.2×10^{-5}				
Total SS	6.3×10^{-2}	12					
Bacteriocin production model (4)							
Lack of fit	1.5×10^{-2}	3	5.0×10^{-3}	0.02	0.99 ^b	0.99	0.98
Pure error	9.7×10^{-1}	4	2.4×10^{-1}				
Total SS	87.7	12					

SE standard error, SS sum of squares, *df* degrees of freedom, MS mean of squares

^aSignificant at $P < 0.05$

^bSignificant at $P > 0.05$

However, the recourse of supplementing the CMPW medium with KH_2PO_4 clearly led to increased accumulation of total phosphorus in the fermentation medium (the lower left side of Fig. 6); however, the final TP concentration in the fermentation medium (0.152 g/L) was reduced compared with the initial TP concentration (0.460 g/L) in the DW medium.

Thus, at the end of the incubations, the initial Pr, TN and TP concentrations of the fermentation substrate (DW medium with 0.460 g TP/L) were reduced by 77.8, 63.9 and 86.6% in the CMPWCW-I culture and 82.0, 70.5 and 67.0% in the CMPWCW-II culture, respectively. This finding was due to the joint effect of the consumption of these nutrients by the strain CECT 539 and the sample extraction.

Fig. 4 Time course of the culture pH in the realkalized fed-batch cultures CMPWCW-I and CMPWCW-II of *L. lactis* CECT 539. Data reported are means $\pm$ standard deviations of two repeated experiments and three replicate measurements

Fig. 5 Cumulative concentrations of biomass (CDW), viable cell counts [log (CFU/mL)], nisin (Nis), lactic acid (LA), acetic acid (AA) and butane-2,3-diol (B) produced by *L. lactis* CECT 539 in the realkalized fed-batch cultures CMPWCW-I and CMPWCW-II. Data reported are means $\pm$ standard deviations of two repeated experiments and three replicate measurements

Table 5 presents comparisons of the maximum levels of biomass and bacteriocin, their volumetric productivities, and the yield $Y_{\text{Nis/CDW}}$ obtained in the CMPWCW-II

culture with other fed-batch fermentation processes with different *L. lactis* strains and culture media. Although the highest biomass production was obtained in the

Fig. 6 Time course of the remaining (left side) and consumed (right side) concentrations of total sugars (TS and TSc), glucose (G and Gc), lactose (L and Lc), proteins (Pr and Prc), total nitrogen (TN and TNc) and phosphorus (TP and TPc) in the realkalized fed-batch cultures CMPWCW-I and CMPWCW-II. Data reported are means $\pm$ standard deviations of two repeated experiments and three replicate measurements

Table 4 Comparison of the maximum mean concentration of products (CDW, VCC, Nis, LA, AA and B) and values of fermentation parameters (productivities, yield factors and efficiencies) obtained at the end of the realkalized fed-batch fermentations of *L. lactis* CECT 539 in DW medium fed with mixtures of CW and CG (CWCG culture), CMPW and CG (CMPWCG culture) (Costas et al. 2017), or CMPW and CW (CMPWCW-I and CMPWCW-II cultures)

Parameters	CWCG	CMPWCG	CMPWCW-I	CMPWCW-II
Fermentation time (h)	168	264	204	204
CDW (g/L)	2.97	4.95	5.12	5.49
VCC (CFU/mL)	1.35×10^{10}	2.03×10^{10}	2.18×10^{10}	2.33×10^{10}
Nis (BU/mL)	108.00	223.98	235.23	257.59
LA (g/L)	13.86	33.37	31.77	34.20
AA (g/L)	1.85	3.83	3.47	3.79
B (g/L)	3.05	6.68	6.47	7.28
P_{CDW} (g/L/h)	1.8×10^{-2}	1.9×10^{-2}	2.5×10^{-2}	2.7×10^{-2}
P_{Nis} (BU/mL/h)	0.64	0.85	1.15	1.26
P_{LA} (g/L/h)	0.08	0.13	0.16	0.17
P_{AA} (g/L/h)	1.1×10^{-2}	1.4×10^{-2}	1.7×10^{-2}	1.9×10^{-2}
P_B (g/L/h)	1.8×10^{-2}	2.5×10^{-2}	3.2×10^{-2}	3.6×10^{-2}
$Y_{Nis/CDW}$ (BU/mg)	36.38	45.27	46.69	46.92
$Y_{CDW/TSc}$	0.03	0.03	0.03	0.03
$Y_{Nis/TSc}$	0.96	1.38	1.40	1.47
$Y_{CDW/TNc}$	6.82	11.16	11.30	11.41
$Y_{Nis/TNc}$	248.19	505.07	527.55	535.65
E_L	0.44	0.66	0.74	0.73
E_G	0.95	0.89	0.90	0.91
E_{TN}	0.72	0.79	0.63	0.68
E_{TP}	0.63	0.57	0.56	0.87

CDW, Nis, LA, AA and B are the final concentrations of cell dry weight (biomass), nisin, lactic acid, acetic acid and butane-2,3-diol, respectively. VCC is the viable cell count. P_{CDW} , P_{Nis} , P_{LA} , P_{AA} , and P_B are the corresponding volumetric productivities. $Y_{Nis/CDW}$, $Y_{Nis/TSc}$ and $Y_{Nis/TNc}$ are the nisin yield coefficients [amounts of nisin (Nis) synthesized (BU) per mg of CDW produced, and total sugars (TSc) or nitrogen (TNc) consumed]. $Y_{CDW/TSc}$ and $Y_{CDW/TNc}$ are the biomass yield coefficients (g of CDW per g of TSc or TNc). E_L , E_G , E_{TN} and E_{TP} are the efficiencies [g of nutrients (lactose (L), glucose (G), total nitrogen (TN) or phosphorus (TP)) consumed per g of nutrients supplied]

CMPWCW-II process, the fed-batch processes with the strains ATCC 11454 (Lv et al. 2005; Papagianni et al. 2007b), NIZO 22186 (Pongtharangkul and Demirci 2006) and W28 (Wu et al. 2009) exhibited increased nisin titers and P_{CDW} , P_{Nis} and $Y_{Nis/CDW}$ levels. This finding is mainly attributed to the notion that the culture media used in the latter fermentation process were supplemented with complex nitrogen sources (e.g., yeast extract and peptones), amino acids, vitamins and minerals, which have growth- and bacteriocin-promoting properties (Guerra and Pastrana 2002; Vázquez et al. 2004). However, the high cost of the complex nitrogen sources makes them unsuitable for large-scale production of probiotic biomass and nisin (Guerra and Pastrana 2002).

As observed in the present work, both the biomass and nisin productions obtained in the fed-batch fermentations with the strains CECT 539 (Vázquez et al. 2004), ATCC 11454 (Lv et al. 2005; Papagianni et al. 2007b), NIZO 22186 (Pongtharangkul and Demirci 2006) and W28 (Wu et al. 2009) were also higher than the corresponding batch cultures.

Conclusions

The main contribution of this study is the design of a realkalized fed-batch culture based on the combination of feeding culture media prepared with effluents (CMPW and CW) of the food industry for cost-effective production of a highly concentrated and viable probiotic culture of *L. lactis* CECT 539 in DW containing a high concentration of nisin. The biomass and bacteriocin levels obtained in this culture were considerably increased compared with those obtained in previous realkalized fed-batch cultures in DW medium fed mixtures of CW and CG (CWCG culture) or CMPW and CG (CMPWCG culture). This approach has the additional advantage of considerably reducing the initial nutrient (Pr, TN and TP) content of these types of wastes (DW, CW and CMPW) and could consequently serve as an appropriate method to reduce their contamination effect.

Table 5 Comparison of the maximum mean concentrations of biomass and nisin obtained in the realkalized fed-batch fermentation CMPWCW-II with *L. lactis* CECT 539 and those obtained by other

researchers in different fed-batch fermentation processes with similar or different *L. lactis* strains

Parameters	<i>L. lactis</i> strains					
	CECT 539 ^a	CECT 539 ^b	ATCC 11454 ^c	ATCC 11454 ^d	NIZO 22186 ^e	W28 ^f
Fermentation time (h)	204	123	10	13	12	14
CDW (g/L)	5.49	1.59	3.55	3.35	1.15	4.42
Nis (IU/mL)	3829	120	4961	6100	4188	4490
P_{CDW} (g/L/h)	2.7×10^{-2}	1.3×10^{-2}	3.6×10^{-1}	2.6×10^{-1}	9.6×10^{-2}	3.2×10^{-1}
P_{Nis} (IU/mL/h)	18.8	1.0	496.1	469.2	349.0	320.7
$Y_{Nis/CDW}$ (IU/mg)	697.4	75.5	1397.4	1820.9	3641.7	1015.8

CDW and Nis are the maximum concentrations of cell dry weight (biomass) and nisin. P_{CDW} and P_{Nis} are the corresponding volumetric productivities. $Y_{Nis/CDW}$ is the nisin yield coefficient (amounts of nisin (Nis) synthesized (IU) per mg of CDW produced)

^aResults obtained in the present study in the CMPWCW-II culture

^bResults obtained by Vázquez et al. (2004) in a realkalized fed-batch culture on a synthetic medium containing glucose, six salts, five vitamins and nine amino acids

^cResults obtained by Lv et al. (2005) in a fed-batch culture in a complex medium containing sucrose, yeast extract, soy peptone and salts

^dResults obtained by Papagianni et al. (2007a, b) in a fed-batch glucostat culture in the synthetic MCD medium containing glucose, 18 amino acids and 14 vitamins among other ingredients

^eResults obtained by Pongtharangkul and Demirci (2006) in fed-batch culture at constant pH (6.8) in a complex medium containing sucrose, peptone, yeast extract and salts

^fResults obtained by Wu et al. (2009) in a variable volume fed-batch fermentation at constant pH (6.8) in a fermentation medium containing corn syrup, sucrose, peptone, yeast extract and salts

Acknowledgements The research presented in this paper was financially supported by the Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, Spain (project CAL01-045-C2-2) and The Xunta de Galicia, Spain (project PGIDT00BIO1E). We thank Coren S.C.L. for their collaboration in the elaboration of this work. Results concerning the evaluation of *L. lactis* CECT 539 probiotic properties were included in a previous publication (Guerra et al. 2007a).

Compliance with ethical standards

Conflict of interest All the authors declare that they have no conflicts of interest.

References

- Anadón A, Martínez MR, Aranzazu M (2006) Probiotics for animal nutrition in the European Union. Regulation and safety assessment. Regul Toxicol Pharmacol 45:91–95
- Cabo ML, Murado MA, González MP, Pastoriza L (2001) Effects of aeration and pH gradient on nisin production. A mathematical model. Enzyme Microb Technol 29:264–273
- Cocaign-Bousquet M, Garrigues C, Loubiere P, Lindley ND (1996) Physiology of pyruvate metabolism in *Lactococcus lactis*. Antonie Van Leeuwenhoek 70:253–267
- Collins LB, Thomas TD (1974) Pyruvate kinase of *Streptococcus lactis*. J Bacteriol 120:52–58
- Costas M, Alonso E, Guerra NP (2016) Nisin production in realkalized fed-batch cultures in whey with feeding with lactose- or glucose-containing substrates. Appl Microbiol Biotechnol 100:7899–7908
- Costas M, Alonso E, Outeiriño D, Fajardo P, Guerra NP (2017) Combination of food wastes for an efficient production of nisin in realkalized fed-batch cultures. Biochem Eng J 123:13–23
- De Vuyst L, Vandamme EJ (1993) Influence of the phosphorus and nitrogen source on nisin production in *Lactococcus lactis* subsp. *lactis* batch fermentations using a complex medium. Appl Microbiol Biotechnol 40:17–22
- Fajardo P, Fuciños C, Méndez J, Pastrana L, Guerra NP (2008) Performance and intestinal coliform counts in weaned piglets fed a probiotic culture (*Lactobacillus casei* subsp. *casei* CECT 4043) or an antibiotic. J Food Prot 71:1797–1805
- Garrigues C, Loubiere P, Lindley ND, Cocaign-Bousquet M (1997) Control of the shift from homolactic acid to mixed-acid fermentation in *Lactococcus lactis*: predominant role of the NADH/NAD⁺ ratio. J Bacteriol 179:5282–5287
- Goel A, Santos F, de Vos WM, Teusink B, Molenaar D (2012) Standardized assay medium to measure *Lactococcus lactis* enzyme activities while mimicking intracellular conditions. Appl Environ Microbiol 78:134–143
- Guerra NP (2014) Modeling the batch bacteriocin production system by lactic acid bacteria by using modified three-dimensional Lotka–Volterra equations. Biochem Eng J 88:115–130
- Guerra NP, Pastrana L (2002) Modelling the influence of pH on the kinetics of both nisin and pediocin production and characterization of their functional properties. Proc Biochem 37:1005–1015
- Guerra NP, Pastrana L (2003) Enhancement of nisin production by *Lactococcus lactis* in periodically re-alkalized cultures. Biotechnol Appl Biochem 38:157–167
- Guerra NP, Rua ML, Pastrana L (2001) Nutritional factors affecting the production of two bacteriocins from lactic acid bacteria on whey. Int J Food Microbiol 70:267–281
- Guerra NP, Fajardo P, Méndez J, Cachaldora P, Pastrana L (2007a) Production of four potentially probiotic lactic acid bacteria and

- their evaluation as feed additives for weaned piglets. *Anim Feed Sci Technol* 134:89–107
- Guerra NP, Torrado A, López C, Fajardo P, Pastrana L (2007b) Dynamic mathematical models to describe the growth and nisin production by *Lactococcus lactis* subsp. *lactis* CECT 539 in both batch and re-alkalized fed-batch cultures. *J Food Eng* 82:103–113
- Landers TF, Cohen B, Wittum TE, Larson EL (2012) A review of antibiotic use in food animals: perspective, policy, and potential. *Public Health Rep* 127:4–22
- Levering J, Musters MWJM, Bekker M, Bellomo D, Fiedler T, de Vos WM, Hugenholtz J, Kreikemeyer B, Kummer U, Teusink B (2012) Role of phosphate in the central metabolism of two lactic acid bacteria—a comparative systems biology approach. *FEBS J* 279:1274–1290
- Lv W, Cong W, Cai Z (2005) Effect of sucrose on nisin production in batch and fed-batch culture by *Lactococcus lactis*. *J Chem Technol Biotechnol* 80:511–514
- Matijašić BB, Stojkovic S, Salobir J, Malovrh S, Rogelj I (2004) Evaluation of the *Lactobacillus gasseri* K7 and LF221 strains in weaned piglets for their possible probiotic use and their detection in the faeces. *Anim Res* 53:35–44
- Murado MA, Siso MIG, González MP, Montemayor MI, Pastrana L, Pintado J (1993) Characterization of microbial biomasses and amylolytic preparations obtained from mussel processing waste treatment. *Bioresour Technol* 43:117–125
- Murado MA, González MP, Vázquez JA (2002) Dose-response relationships: an overview, a generative model and its application to the verification of descriptive models. *Enzyme Microb Technol* 31:439–455
- Papagianni M, Avramidis N, Filiouis G (2007a) Glycolysis and the regulation of glucose transport in *Lactococcus lactis* spp. *lactis* in batch and fed-batch culture. *Microb Cell Fact* 6:16
- Papagianni M, Avramidis N, Filiouis G (2007b) Investigating the relationship between the specific glucose uptake rate and nisin production in aerobic batch and fed-batch glucostat cultures of *Lactococcus lactis*. *Enzyme Microb Technol* 40:1557–1563
- Pascual M, Hugas M, Badiola JJ, Monfort JM, Garriga M (1999) *Lactobacillus salivarius* CTC2197 prevents *Salmonella enteritidis* colonization in chickens. *Appl Environ Microbiol* 65:4981–4986
- Pongtharangkul T, Demirci A (2006) Effects of fed-batch fermentation and pH profiles on nisin production in suspended-cell and biofilm reactors. *Appl Microbiol Biotechnol* 73:73–79
- Poolman B, Konings WN (1988) Relation of growth of *Streptococcus lactis* and *Streptococcus cremoris* to amino acid transport. *J Bacteriol* 170:700–707
- Thorlacius H, Jeppsson B (2003) Anti-infective effects of lactobacilli in the gastrointestinal tract. *Curr Med Chem Antiinfect Agents* 2:263–267
- van Rooijen RJ, Gasson MJ, de Vos WM (1992) Characterization of the *Lactococcus lactis* lactose operon promoter: contribution of flanking sequences and LacR repressor to promoter activity. *J Bacteriol* 174:2273–2280
- Vázquez JA, Cabo ML, González MP, Murado MA (2004) The role of amino acids in nisin and pediocin production by two lactic acid bacteria. A factorial study. *Enzyme Microb Technol* 34:319–325
- Wu Z, Wang L, Jing Y, Li X, Zhao Y (2009) Variable volume fed-batch fermentation for nisin production by *Lactococcus lactis* subsp. *lactis* W28. *Appl Biochem Biotechnol* 152:372–382