Published in final edited form as:

AIDS Behav. 2018 July ; 22(7): 2277–2283. doi:10.1007/s10461-018-2048-y.

Comparison of self-report to biomarkers of recent HIV infection: Findings from the START Trial

Katherine E. Schlusser¹, Shweta Sharma², Pola de la Torre³, Giuseppe Tambussi⁴, Rika Draenert⁵, Angie N. Pinto⁶, Julia A. Metcalf⁷, Danielle German⁸, James D. Neaton², and Oliver Laeyendecker^{1,9} for the INSIGHT START Study Group

¹Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

²Division of Biostatistics, School of Public Health, University of Minnesota, Minneapolis, Minnesota, USA

³Cooper University Hospital, Camden, New Jersey, USA

⁴IRCCS-Ospedale San Raffaele, Milano, Italy

⁵Klinikum der Universität Munich, Medizinische Klinik IV, Section Clinical Infectious Diseases, Munich, Germany

⁶The Kirby Institute, UNSW Australia, Sydney, Australia

⁷Division of Clinical Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA

⁸Department of Health, Behavior and Society, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland, USA

⁹Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA

Abstract

Background—Identifying individuals with recent HIV infection is critical to research related to viral reservoirs, outbreak investigations and intervention applications. A multi-assay algorithm (MAA) for recency of infection was used in conjunction with self-reported date of infection and documented date of diagnosis to estimate the number of participants recently infected in the Strategic Timing of AntiRetroviral Treatment (START) trial.

Methods—We tested samples for three groups of participants from START using a MAA: 1) 167 individuals who reported being infected 6 months before randomization; 2) 771 individuals who did not know their date of infection but were diagnosed within 6 months before randomization; and 3) as controls for the MAA, 199 individuals diagnosed with HIV 2 years before

randomization. Participants with low titer and avidity and a baseline viral load >400 copies/mL were classified as recently infected.

Results—A significantly higher percentage of participants who self-reported being infected 6 months were classified as recently infected compared to participants diagnosed 2 years (65% [109/167] vs. 2.5% [5/199], $p<0.001$). Among the 771 individuals who did not know their duration of infection at randomization, 206 (26.7%) were classified as recently infected. Among those diagnosed with HIV in the 6 months prior to enrollment, the 373 participants who reported recent infection ($n=167$) or who had confirmed recent infection by the MAA ($n=206$) differed significantly on a number of baseline characteristics from those who had an unknown date of infection and were not confirmed by the MAA ($n=565$). Participants recently infected by self-report and/or MAA were younger, more likely to be Asian, less likely to be black, less likely to be heterosexual, more likely to be enrolled at sites in the U.S., Europe or Australia, and have higher HIV RNA levels.

Conclusions—There was good agreement between self-report of recency of infection and the MAA. We estimate that 373 participants enrolled in START were infected within 6 months of randomization. Compared to those not recently infected, these participants were younger, had higher HIV RNA levels and were more likely to come from high income countries and from populations such as MSM with more regular HIV testing.

INTRODUCTION

Self-report of the duration of HIV infection is an important measure for HIV research. Individuals with recent infection can be treated early in their infection, which may limit the size of their viral reservoir¹ and enhance CD4+ recovery^{2,3}. Additionally, newly diagnosed individuals who are recently infected are more likely to be linked to HIV infected individuals who do not know their infection status⁴. Thus, identifying recently infected individuals and then determining their social network and testing those individuals for HIV can decrease the number of HIV infected people unaware of their infection status⁴. Furthermore, if these individuals reduce their risky behaviors as a result of knowing that they are HIV positive, the rate of HIV transmission would be reduced. For these reasons accurate self-report of newly diagnosed individuals is critical for HIV clinical trials⁵ and contact tracing⁶. However, the accuracy of self-reported recent HIV infection is not known.

There are many reasons why accurate self-report of recent infection would be useful both scientifically and for public health. Individuals with recent infection have smaller viral reservoirs¹ and are being enrolled for cure protocol studies (Research In Viral Eradication of HIV Reservoirs, protocol # ISRCTN83717528), and prioritized for public health follow-up and partner notification. When using biomarkers to determine recent infection, there are multiple confounding factors that impact their ability to correctly discriminate the duration of infection at the individual level and correctly estimate population level incidence. These include race⁷, gender⁸, infecting subtype^{9–11}, set point viral load^{12,13}, AIDS^{14,15}, viral suppression^{13,14,16}, and viral breakthrough¹⁷. A multi assay algorithm (MAA) uses a combination of biomarkers with orthogonal properties for misclassification to help maximize the precision in distinguishing recent from non-recent infection¹⁸. This method

has been demonstrated to accurately estimate population level incidence and to accurately identify recent infection at the individual level^{19,20}.

We applied this methodology to a subset of participants in the Strategic Timing of AntiRetroviral Treatment (START) trial²¹ who were: 1) diagnosed with HIV within 6 months before randomization and also reported being infected within the 6 months before randomization, 2) diagnosed with HIV within 6 months before randomization and did not know their date of infection, and 3) were diagnosed 2 years before randomization (controls). Our goal was to validate self-report of recency of HIV infection and to use the MAA in combination with self-reported date of infection and documented date of diagnosis to estimate the number of participants enrolled in START with early infection (< 6 months prior to randomization). By identifying a cohort of recently infected individuals randomized to immediate or deferred treatment who will be in long term follow up until 2021, we will be able to determine the health benefits of treatment early in infection.

METHODS

START is a multinational randomized clinical trial comparing immediate versus deferred antiretroviral treatment (ART) initiation for major clinical outcomes. The study enrolled 4684 HIV-positive ART-naïve individuals from 35 countries, between April 2009 and December 2013. Participants had to have two CD4 counts >500 cells/mm³ at least 2 weeks apart within the 60 days before enrollment and be at least 18 years of age²¹. The study results were unblinded in May 2015 due to clear and significant clinical benefit of immediate ART initiation²¹.

At study entry, in addition to demographics, CD4+ count, HIV RNA level and likely mode of HIV infection, two dates were recorded: 1) the date the participant was first diagnosed with HIV as documented in the participant's record; and 2) if known, the month and year the participant believes he/she was first infected with HIV. Plasma samples were collected at entry for future research purposes and stored centrally. These stored samples were tested with the MAA for three groups of participants. Group 1 included 167 participants who were diagnosed and reported being infected 6 months prior to randomization. Group 2 included 771 participants who were diagnosed 6 months prior to randomization and who reported the date of infection was unknown. Those diagnosed with HIV between 6 and 24 months of randomization were not studied (n=1551), as well as those who were diagnosed within 6 months but reported being infected for > 6 months (n=518). Group 3 consisted of 199 randomly chosen participants who were diagnosed with HIV 2 years prior to randomization (controls). All samples tested were from individuals who consented to and had stored plasma samples.

Baseline plasma samples were tested by the MAA, which was developed for cross-sectional incidence testing¹⁹. All samples were tested with the Limiting Antigen-Avidity assay (LAg-Avidity)²² and those with a LAg-Avidity normalized optical density <3.0 were further tested by the Johns Hopkins avidity modified GS HIV-1/HIV-2 Plus O EIA²³. As all participants had a CD4+ count >500 cells/mm³ at enrollment, those who had a LAg-Avidity normalized optical density <2.9, had an avidity index below 85% and a baseline viral load >400

copies/mL were classified as recently infected. This testing algorithm had been validated in samples from the U.S. where the average duration that an individual appeared recently infected was 146 days¹⁹ and no sample from an individual infected >4 years was classified as recent. For the algorithm used, it was previously determined that for individuals classified as recent by this testing methodology, 95% were infected <1 year¹⁹.

Baseline characteristics of Groups 1–3 were summarized with simple summary statistics. The following baseline characteristics were compared: age, gender, race, geographic location, likely mode of HIV infection, baseline CD4+ and CD8+ cell count, and baseline HIV RNA level. These baseline characteristics were chosen to describe the groups because we hypothesized that those with recent infection would differ from those who were not recently infected on many of the characteristics. Specifically, those with recent infection were hypothesized to be younger, men who have sex with men, and enrolled by sites in high-income countries where they were more likely to be tested. Those with early infection would also be more likely to have elevated HIV RNA levels. Proportions were compared using chi-squared tests and medians compared using Wilcoxon rank sum tests. Logistic regression analysis which considered multiple factors was used for three comparisons performed in the following order: 1) within group 1, participants classified as recently infected by the MAA versus those who were not; 2) participants classified as recently infected by the MAA in Group 2 versus all participants in Group 1; and 3) all participants in Group 1 plus participants in Group 2 who were classified as recently infected by the MAA versus participants in Group 2 who were not classified as recently infected by the MAA. Odds ratios were used to compare groups. The first comparison was made in order to determine whether those who self-reported recent infection differed according to the MAA result. The 2nd comparison was made to determine whether those classified as recently infected by the MAA who did not know the date of their infection differed from those who self-reported recent infection. The 3rd comparison was made to describe the differences in baseline characteristics between those we classified as recently infected by self-report or MAA with those who were not classified as recently infected. Statistical analyses were performed using SAS, version 9.3 (SAS Institute, Cary, NC). A p-value < 0.05 was considered significant. All p-values cited are 2-sided and not adjusted for multiple comparisons.

RESULTS

Sample Description and MAA Results

Among the 180 participants who reported being infected within 6 months of randomization, 167 (93%) consented to storage of specimens for future HIV research and had a baseline specimen collected. Of those diagnosed with HIV in the 6 months prior to randomization with unknown date of infection, 93% (771/830) had baseline samples for analysis. 1605 participants reported a diagnosis of HIV two or more years before randomization, 1539 (96%) consented to store specimens, and 199 of those participants (13%) were randomly chosen for this study as controls. Characteristics of the three groups studied are shown in Supplemental Table I, overall and according to MAA results.

A significantly higher percentage of individuals who reported being infected 6 months were recently infected by the MAA compared to controls, participants who were diagnosed

with HIV for at least 2 years before enrolling in START (65% [109/167] vs. 2.5% [5/199]; $\chi^2(1, N=366)=166.8, p<0.001$) (Table I). Additionally, the group who reported being infected 6 months had lower LAg-Avidity values than the group diagnosed 2 or more years before enrollment (median normalized optical density 1.86 [IQR 0.99, 3.01] vs. 4.53 [3.84, 4.99]; $\chi^2(1, N=366)=162.5, p<0.001$).

Among the 771 participants who were diagnosed with HIV within 6 months prior to enrollment and reported the date of infection as unknown, 206 (26.7%) were recently infected by the MAA. The median LAg-Avidity results for those 206 participants was slightly higher than for the 109 participants in Group 1 who were recently infected by the MAA (OD-n 1.62 [1.05, 2.16] versus 1.46 [0.70, 1.94]; $\chi^2(1, N=314)=4.5, p=0.034$).

Comparison of Baseline Characteristics of Participants Recently Infected Versus Not Recently Infected

Among participants in Group 1 (diagnosed with HIV within the past 6 months and HIV infected within the past 6 months) the only baseline characteristic that differed between those classified as recently infected by the MAA and those who were not was HIV RNA level (26,323 copies/mL versus 19,114 copies/mL; $\chi^2(1, N=167)=2.6, p=0.11$ in univariate analysis and $\chi^2(1, N=165)=4.8, p=0.028$ in multivariate analysis) (Supplemental Tables I and II).

Those in Group 2 (diagnosed with HIV within the past 6 months and with a date of HIV infection unknown) classified as recently infected by the MAA were similar to participants in Group 1 with two exceptions, the percentage of participants who were MSM (69.4% versus 82.6%; $\chi^2(1, N=373)=7.6, p=0.006$ in univariate analysis and $\chi^2(1, N=373)=4.3, p=0.039$ in multivariate analysis) and CD8+ count (945 versus 1,086 cells/mm³; $\chi^2(1, N=371)=7.3, p=0.007$ in univariate analysis and $\chi^2(1, N=373)=10.5, p=0.001$ in multivariate analysis, Table II and Supplemental Table I).

Table III (and Supplemental Table III) compares participants in Group 1 together with those in Group 2 who were classified as recently infected by the MAA (recently infected group) to participants in Group 2 who were not classified as recently infected by the MAA. There were a number of differences between these two groups. In the multivariate analysis, participants recently infected by self-report and/or MAA were younger, more likely to be Asian, less likely to be black, less likely to be heterosexual, more likely to be enrolled at sites in the U.S., Europe or Australia, and have higher HIV RNA levels.

DISCUSSION

In START we demonstrated that nearly two-thirds of participants who were diagnosed with HIV within 6 months before enrollment and reported HIV infection within 6 months before enrollment had MAA findings which confirmed their recent infection, whereas, only 2.5% of participants who were diagnosed with HIV more than 2 years before enrollment (controls) had positive assay results. Those reporting recent infection and confirmed by the assay did not differ in terms of baseline characteristics from those who were not. This observation

taken together with the limitations of the MAA assay used¹⁹, indicate that the classification of participants as recently infected by self-report in START (Group 1) is likely accurate.

There are several limitations to this study. The gold standard used to identify/confirm recent infection was the combination of biomarkers as part of a MAA. The MAA used has a window period of 146 days for HIV-1 clade B infected individuals^{19,26}. This means that the average duration an individual will appear as recently infected by this MAA is 4.8 months (146 days, 95%CI 122 to 170 days) after seroconversion. In START, the median time from diagnosis to enrollment was 2.4 months for those who reported being infected 6 months. MAA was based on a plasma sample taken at the time of randomization. Had a plasma sample been available at the time of HIV diagnosis to study the agreement of the MAA and self-reported recent infection, agreement would likely have been higher. Furthermore, of the five individuals who were known to be infected more than two years, but were classified as recently infected by the MAA, three were subjects where the time between diagnosis and blood draw was < 4 years. Additionally, four of these five individuals were from countries where HIV-1 non-B clades circulate²⁴. Since the incidence assays used have different properties depending on the infecting clade^{10,11,22}, it is not surprising that the majority of the misclassified long term infected samples come from HIV-1 non-B endemic areas.

Among participants with an unknown infection date who were diagnosed with HIV in the 6 months prior to enrollment, 206 (27%) were recently infected according to the MAA. These 206 participants had similar baseline characteristics to those who self-reported being recently infected. Thus, taken together with those self-reporting recent infection, 373 participants in START were infected within 6 months before enrollment. Compared to the 565 participants diagnosed with HIV in the 6 months prior to enrollment who were not recently infected, the 373 participants who were recently infected were younger, more likely to be Asian, less likely to be black, less likely to be heterosexual, more likely to be enrolled at sites in the U.S., Europe or Australia, and have higher HIV RNA levels.

There are several explanations for why particular factors were associated with recent infection in the START trial population. The entry requirements for the trial required high CD4 counts which would select for individuals who were early in their infection or with a less pathogenic strain of HIV. Therefore the finding that higher viral load was associated with recent infection is not surprising. The majority of heterosexual participants in the trial came from Africa, where frequent testing is not the norm. For example a recent study in Zambia, 40% of men had never been tested for HIV in their lifetime.³³ The increased odds of Asians being recently infected is not surprising as most of those subjects came from Thailand which has a long history of HIV prevention and 75% of MSM reporting getting tested for HIV in the past year²⁵.

Although we do not know of any published studies comparing self-report of recency of HIV infection to biomarkers, a few studies have compared self-report of HIV status to biomarkers of HIV infection. For example, Fisher et al. compared enzyme-linked immunosorbent assay test results to self-report of HIV status, finding that the specificity of self-report was above 99% at baseline and follow-up while the sensitivity was 32% at baseline and 61% at follow-up²⁶. Additionally, Ross et al. compared Western blot results to self-reported HIV infection

and found that the sensitivity of self-report was 93% and the specificity was above 99%²⁷. However in our study there were no pre-determined reasons for misinforming the study staff about their date of HIV infection. Our results showed a similarly high level of self-report specificity compared to these two studies while our sensitivity was lower than that observed by Ross et al. and slightly higher than the sensitivity seen by Fisher et al. at follow-up^{26,27}.

In summary, in the START trial 373 participants with recent HIV infection were enrolled. To our knowledge, this subgroup represents the largest study of immediate versus deferred ART among individuals recently infected with HIV.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We wish to thank the participants and clinical staff of the study. This study was supported in part by the Division of Intramural Research, National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH). Support was provided by the NIH Clinical Center, National Cancer Institute, National Heart, Lung, and Blood Institute, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institute of Mental Health, National Institute of Neurological Disorders and Stroke, National Institute of Arthritis and Musculoskeletal and Skin Diseases, Agence Nationale de Recherches sur le SIDA et les Hépatites Virales (France), National Health and Medical Research Council (Australia), National Research Foundation (Denmark), Bundesministerium für Bildung und Forschung (Germany), European AIDS Treatment Network, Medical Research Council (United Kingdom), National Institute for Health Research, National Health Service (United Kingdom), and University of Minnesota. Antiretroviral drugs were donated to the central drug repository by AbbVie, Bristol-Myers Squibb, Gilead Sciences, GlaxoSmithKline/ViiV Healthcare, Janssen Scientific Affairs, and Merck (UM1-AI068641 and UM1-AI120197). Additional support was provided by the HIV Prevention Trials Network sponsored by NIAID, National Institute of Child Health and Human Development, National Institute of Drug Abuse, National Institute of Mental Health, and the Office of AIDS Research, of the NIH DHHS (UM1 AI068613), and NIAID (R01 AI095068).

References

1. Ostrowski M, Benko E, Yue FY, et al. Intensifying Antiretroviral Therapy With Raltegravir and Maraviroc During Early Human Immunodeficiency Virus (HIV) Infection Does Not Accelerate HIV Reservoir Reduction. *Open forum infectious diseases*. Dec.2015 2(4):ofv138. [PubMed: 26512359]
2. Le T, Wright EJ, Smith DM, et al. Enhanced CD4+ T-cell recovery with earlier HIV-1 antiretroviral therapy. *The New England journal of medicine*. Jan 17; 2013 368(3):218–230. [PubMed: 23323898]
3. Okulicz JF, Le TD, Agan BK, et al. Influence of the timing of antiretroviral therapy on the potential for normalization of immune status in human immunodeficiency virus 1-infected individuals. *JAMA internal medicine*. Jan; 2015 175(1):88–99. [PubMed: 25419650]
4. Friedman SR, Downing MJ Jr, Smyrnov P, et al. Socially-integrated transdisciplinary HIV prevention. *AIDS and behavior*. Oct; 2014 18(10):1821–1834. [PubMed: 24165983]
5. Mastro TD, Kim AA, Hallett T, et al. Estimating HIV Incidence in Populations Using Tests for Recent Infection: Issues, Challenges and the Way Forward. *Journal of HIV AIDS surveillance & epidemiology*. Jan 01; 2010 2(1):1–14. [PubMed: 21743821]
6. Dennis AM, Murillo W, de Maria Hernandez F, et al. Social network-based recruitment successfully reveals HIV-1 transmission networks among high-risk individuals in El Salvador. *Journal of acquired immune deficiency syndromes (1999)*. May 01; 2013 63(1):135–141. [PubMed: 23364512]
7. Laeyendecker O, Brookmeyer R, Oliver AE, et al. Factors associated with incorrect identification of recent HIV infection using the BED capture immunoassay. *AIDS Res Hum Retroviruses*. Aug; 2012 28(8):816–822. [PubMed: 22014036]

8. Mullis CE, Munshaw S, Grabowski MK, et al. Differential specificity of HIV incidence assays in HIV subtypes A and D-infected individuals from Rakai, Uganda. *AIDS Res Hum Retroviruses*. Aug; 2013 29(8):1146–1150. [PubMed: 23641870]
9. Parekh BS, Hanson DL, Hargrove J, et al. Determination of mean recency period for estimation of HIV type 1 Incidence with the BED-capture EIA in persons infected with diverse subtypes. *AIDS Res Hum Retroviruses*. Mar; 2011 27(3):265–273. [PubMed: 20954834]
10. Longosz AF, Serwadda D, Nalugoda F, et al. Impact of HIV subtype on performance of the limiting antigen-avidity enzyme immunoassay, the bio-rad avidity assay, and the BED capture immunoassay in Rakai, Uganda. *AIDS Res Hum Retroviruses*. Apr; 2014 30(4):339–344. [PubMed: 24083837]
11. Kassanjee R, Pilcher CD, Keating SM, et al. Independent assessment of candidate HIV incidence assays on specimens in the CEPHIA repository. *AIDS*. Oct 23; 2014 28(16):2439–2449. [PubMed: 25144218]
12. Laeyendecker O, Rothman RE, Henson C, et al. The effect of viral suppression on cross-sectional incidence testing in the Johns Hopkins hospital emergency department. *Journal of acquired immune deficiency syndromes (1999)*. Jun 01; 2008 48(2):211–215. [PubMed: 18520680]
13. Kassanjee R, Pilcher CD, Busch MP, et al. Viral load criteria and threshold optimization to improve HIV incidence assay characteristics. *AIDS*. Sep 24; 2016 30(15):2361–2371. [PubMed: 27454561]
14. Hayashida T, Gatanaga H, Tanuma J, Oka S. Effects of low HIV type 1 load and antiretroviral treatment on IgG-capture BED-enzyme immunoassay. *AIDS Res Hum Retroviruses*. Mar; 2008 24(3):495–498. [PubMed: 18327979]
15. Longosz AF, Mehta SH, Kirk GD, et al. Incorrect identification of recent HIV infection in adults in the United States using a limiting-antigen avidity assay. *AIDS*. May 15; 2014 28(8):1227–1232. [PubMed: 24513567]
16. Marinda ET, Hargrove J, Preiser W, et al. Significantly diminished long-term specificity of the BED capture enzyme immunoassay among patients with HIV-1 with very low CD4 counts and those on antiretroviral therapy. *Journal of acquired immune deficiency syndromes (1999)*. Apr 01; 2010 53(4):496–499. [PubMed: 20306555]
17. Wendel SK, Mullis CE, Eshleman SH, et al. Effect of natural and ARV-induced viral suppression and viral breakthrough on anti-HIV antibody proportion and avidity in patients with HIV-1 subtype B infection. *PloS one*. 2013; 8(2):e55525. [PubMed: 23437058]
18. Laeyendecker O, Kulich M, Donnell D, et al. Development of methods for cross-sectional HIV incidence estimation in a large, community randomized trial. *PloS one*. 2013; 8(11):e78818. [PubMed: 24236054]
19. Konikoff J, Brookmeyer R, Longosz AF, et al. Performance of a limiting-antigen avidity enzyme immunoassay for cross-sectional estimation of HIV incidence in the United States. *PloS one*. 2013; 8(12):e82772. [PubMed: 24386116]
20. Huynh D, Laeyendecker O, Brookmeyer R. A serial risk score approach to disease classification that accounts for accuracy and cost. *Biometrics*. Dec; 2014 70(4):1042–1051. [PubMed: 25156309]
21. Lundgren JD, Babiker AG, Gordin F, et al. Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection. *The New England journal of medicine*. Aug 27; 2015 373(9):795–807. [PubMed: 26192873]
22. Duong YT, Kassanjee R, Welte A, et al. Recalibration of the Limiting Antigen Avidity EIA to Determine Mean Duration of Recent Infection in Divergent HIV-1 Subtypes. *PloS one*. 2015; 10(2):e0114947. [PubMed: 25710171]
23. Longosz AF, Morrison CS, Chen PL, et al. Immune responses in Ugandan women infected with subtypes A and D HIV using the BED capture immunoassay and an antibody avidity assay. *Journal of acquired immune deficiency syndromes (1999)*. Apr 01; 2014 65(4):390–396. [PubMed: 24583613]
24. Hemelaar J, Gouws E, Ghys PD, Osmanov S. Global trends in molecular epidemiology of HIV-1 during 2000–2007. *Aids*. Mar 13; 2011 25(5):679–689. [PubMed: 21297424]

25. Hensen B, Lewis JJ, Schaap A, et al. Frequency of HIV-testing and factors associated with multiple lifetime HIV-testing among a rural population of Zambian men. *BMC public health*. Sep 24.2015 15:960. [PubMed: 26404638]
26. Fisher DG, Reynolds GL, Jaffe A, Johnson ME. Reliability, sensitivity and specificity of self-report of HIV test results. *AIDS care*. May; 2007 19(5):692–696. [PubMed: 17505932]
27. Ross MW, Loxley W, Wodak A, Buzolic A, Monheit B, Stowe A. Drug users' self-reported false-positive HIV status. *American journal of public health*. Sep; 1993 83(9):1349–1351. [PubMed: 8363021]

Table I

Results from the MAA for the three HIV infection groups.

	HIV Diagnosis Date 6 Months		HIV Diagnosis Date 2 Years (Group 3)
	HIV Infection Date 6 Months (Group 1)	HIV Infection Date Unknown (Group 2)	
No. tested	167	771	199
Months between diagnosis and enrollment, median [IQR ^a]	2.4 [1.8, 3.1]	2.9 [2.1, 4.2]	55.1 [36.7, 85.2]
MAA ^b positive, n(%)	109 (65.3)	206 (26.7)	5 (2.5)
LA _g -Avidity OD-n ^c , median [IQR]	1.86 [0.99, 3.01]	3.46 [2.14, 4.42]	4.53 [3.84, 4.99]

^aIQR: interquartile range^bMAA: multi algorithm assay^cOD-n: normalized optical density

Table II

Comparison of participants in Group 2 confirmed as recently infected by MAA (unknown HIV infection date, n=206) with those in Group 1 (self-reported HIV infection 6 months, n=167); all participants diagnosed with HIV 6 months before randomization.

	Univariate Models ^a			Multivariate Model ^b		
	OR [95 % CI]	χ ^{2c}	P-value	OR [95 % CI]	χ ^{2c}	P-value
Age, per 5 years	1.11 [1.00, 1.23]	3.55	0.06	1.06 [0.94, 1.19]	0.86	0.35
Female (ref=male)	1.86 [0.93, 3.72]	3.10	0.08	0.97 [0.36, 2.61]	0.003	0.96
Race/ethnicity (ref=white)		2.23	0.69		0.94	0.92
Asian	1.37 [0.73, 2.56]	0.96	0.33	1.22 [0.59, 2.54]	0.29	0.59
Black	1.24 [0.67, 2.29]	0.48	0.49	0.96 [0.47, 1.95]	0.01	0.91
Latino/Hispanic	0.98 [0.54, 1.77]	0.004	0.95	1.01 [0.53, 1.91]	0.0005	0.98
Other	0.68 [0.26, 1.81]	0.59	0.44	0.72 [0.26, 1.97]	0.42	0.52
Likely mode of HIV infection (ref=heterosexual)		8.59	0.014		6.15	0.046
MSM	0.45[0.26, 0.80]	7.56	0.006	0.43 [0.19, 0.96]	4.25	0.039
Other	0.82 [0.30, 2.23]	0.15	0.70	1.16 [0.37, 3.65]	0.07	0.80
High income region ^d (ref = low-middle income)	0.91 [0.60, 1.37]	0.22	0.64	1.01 [0.62, 1.65]	0.003	0.96
CD4 cell count (per 100 cells/mm ³)	0.99 [0.86, 1.13]	0.05	0.83	1.03 [0.88, 1.20]	0.11	0.74
CD8 cell count (per 100 cells/mm ³)	0.94 [0.91, 0.98]	7.32	0.007	0.93 [0.88, 0.97]	10.47	0.001
HIV RNA (log ₁₀ copies/mL)	1.11 [0.83, 1.49]	0.54	0.46	1.34 [0.96, 1.87]	3.03	0.08
No. MAA confirmed	206			206		
Total pts	373			371		

CI: confidence interval; dF=degree(s) of freedom; MAA: multi algorithm assay, MSM: men having sex with men; OR: Odds ratio

^aOdds ratios from individual logistic regression models.

^bOdds ratios from one multivariate logistic regression model with all listed variables. Model limited to participants with complete data.

^c χ^2 tests for univariate models are 1dF, N=373 except for the omnibus race (4dF, N=373) and likely mode of HIV infection (2dF, N=373) tests. Multivariate model significance tests are based on the same dF with N=371.

^dHigh income region included the United States, Europe, and Australia. Low-middle includes Latin America, Asia, and Africa

Table III

Comparison of participants in Group 1 (self-reported HIV infection 6 months, n=167) combined with those in Group 2 confirmed as recently infected by the MAA (unknown HIV infection date, n=206) versus those in Group 2 not confirmed as recently infected (n=565); all participants diagnosed with HIV 6 months before randomization.

	Univariate Models ^a			Multivariate Model ^b		
	OR [95 % CI]	χ^2	P-value	OR [95 % CI]	χ^2	P-value
Age, per 5 years	0.88 [0.82, 0.95]	12.65	<0.001	0.90 [0.83, 0.97]	7.36	0.007
Female (ref=male)	0.26 [0.18, 0.37]	51.79	<0.001	0.75 [0.44, 1.27]	1.15	0.28
Race/ethnicity (ref=white)		68.40	<0.001		31.94	<0.001
Asian	1.72 [1.07, 2.77]	5.04	0.024	3.14 [1.78, 5.53]	15.69	<0.001
Black	0.28 [0.20, 0.40]	49.44	<0.001	0.63 [0.41, 0.98]	4.29	0.038
Latino/Hispanic	1.04 [0.69, 1.56]	0.03	0.86	1.26 [0.81, 1.99]	1.04	0.31
Other	0.58 [0.32, 1.06]	3.18	0.08	0.78 [0.41, 1.50]	0.55	0.46
Likely mode of HIV infection (ref=heterosexual)		76.69	<0.001		13.12	0.001
MSM	4.06 [2.97, 5.55]	76.67	<0.001	2.31 [1.44, 3.70]	11.99	<0.001
Other	2.60 [1.44, 4.70]	10.00	0.002	2.32 [1.17, 4.59]	5.81	0.016
High income region ^d (ref = low-middle income)	2.14 [1.63, 2.82]	29.49	<0.001	1.62 [1.15, 2.28]	7.58	0.006
CD4 cell count (per 100 cells/mm ³)	0.92 [0.85, 0.99]	4.81	0.028	1.08 [0.99, 1.19]	3.00	0.08
CD8 cell count (per 100 cells/mm ³)	1.01 [0.98, 1.04]	0.50	0.48	0.98 [0.95, 1.01]	1.44	0.23
HIV RNA (log ₁₀ copies/mL)	1.99 [1.68, 2.36]	63.30	<0.001	1.83 [1.50, 2.22]	36.38	<0.001
<i>No. recently infected (MAA or self-report)</i>	373				371	
<i>Total pts</i>	938				931	

^aOdds ratios from individual logistic regression models.

^bOdds ratios from one multivariate logistic regression model with all listed variables. Model limited to participants with complete data.

^c χ^2 tests for univariate models are 1dF, N=938 except for the omnibus race (4dF, N=938) and likely mode of HIV infection (2dF, N=938) tests. Multivariate model significance tests are based on the same dF with N=931.

^dHigh income region included the United States, Europe, and Australia. Low-middle includes Latin America, Asia, and Africa