CONFLICT OF INTEREST

The authors have nothing to disclose.

REFERENCES

1. Chang JW, Yeh KY, Wang CH, Yang TS, Chiang HF, Wei FC, et al. Malignant melanoma in Taiwan: a prognostic study of 181 cases. *Melanoma Res* 2004;14:537-541.
2. Bradford PT, Goldstein AM, McMaster ML, Tucker MA. Acral lentiginous melanoma: incidence and survival patterns in the United States, 1986-2005. *Arch Dermatol* 2009;145:427-434.
3. Ito T, Moroi Y, Oba J, Nakahara T, Takeuchi S, Uchi H, et al. The prognostic value of a reverse transcriptase-PCR assay of sentinel lymph node biopsy for patients with cutaneous melanoma: a single-center analysis in Japan. *Melanoma Res* 2012;22:38-44.
4. Phan A, Touzet S, Dalle S, Ronger-Savlé S, Balme B, Thomas L. Acral lentiginous melanoma: a clinicoprognostic study of 126 cases. *Br J Dermatol* 2006;155:561-569.
5. Krementz ET, Feed RJ, Coleman WP 3rd, Sutherland CM, Carter RD, Campbell M. Acral lentiginous melanoma. A clinicopathologic entity. *Ann Surg* 1982;195:632-645.
6. Slingluff CL Jr, Vollmer R, Seigler HF. Acral melanoma: a review of 185 patients with identification of prognostic variables. *J Surg Oncol* 1990;45:91-98.
7. Pryor JG, Bourne PA, Yang Q, Spaulding BO, Scott GA, Xu H. IMP-3 is a novel progression marker in malignant melanoma. *Mod Pathol* 2008;21:431-437.
8. Yu L, Xu H, Wasco MJ, Bourne PA, Ma L. IMP-3 expression in melanocytic lesions. *J Cutan Pathol* 2010;37:316-322.
9. Sheen YS, Liao YH, Lin MH, Chiu HC, Jee SH, Liao JY, et al. Insulin-like growth factor II mRNA-binding protein 3 expression correlates with poor prognosis in acral lentiginous melanoma. *PLoS One* 2016;11:e0147431.
10. Sheen YS, Liao YH, Lin MH, Chu CY, Ho BY, Hsieh MC, et al. IMP-3 promotes migration and invasion of melanoma cells by modulating the expression of HMGA2 and predicts poor prognosis in melanoma. *J Invest Dermatol* 2015;135:1065-1073.

<https://doi.org/10.5021/ad.2018.30.4.499>

Terra Firma-Forme Dermatositis Developed after Thyroidectomy: Clinical and Dermoscopic Features

Jungyoon Moon, Minsoo Kim¹, Jong Seo Park, Jaewon Lee, Je-Ho Mun

Department of Dermatology, Seoul National University College of Medicine, Seoul, Korea, ¹Department of Dermatology, The University of Auckland, Auckland, New Zealand

Dear Editor:

Terra firma-forme dermatosis (TFFD) is an idiopathic benign skin disorder characterized by a dirty appearance of brownish pigmentation resulting from incomplete maturation

of keratinocytes¹. TFFD is thought to be prevalent in clinical practice, but it has been often ignored due to unawareness and indifference to the disease entity. Herein, we introduce a case of TFFD, which occurred on the ante-

Received March 31, 2017, Revised September 8, 2017, Accepted for publication September 17, 2017

Corresponding author: Je-Ho Mun, Department of Dermatology, Seoul National University Hospital, 101 Daehak-ro, Jongno-gu, Seoul 03080, Korea. Tel: 82-2-2072-2417, Fax: 82-2-742-7344, E-mail: jehomun@gmail.com
ORCID: <https://orcid.org/0000-0002-0734-2850>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © The Korean Dermatological Association and The Korean Society for Investigative Dermatology

rior neck after robot-assisted thyroidectomy.

A 36-year-old woman was referred to our clinic with a 3-month history of an asymptomatic dirt-like brownish skin lesion on her anterior neck (Fig. 1A). Widely scattered tiny papules with fine crusts were observed; some of which were fused to form dark, velvety textured plaques. Despite normal daily cleansing to remove the lesion, it gradually expanded. The patient denied any discomfort such as pruritus. There were no previous medical problems apart from papillary thyroid carcinoma, which had been treated with robot-assisted total thyroidectomy 4 months prior. A KOH smear was negative. Dermoscopic examination showed multiple subtle brown polygons arranged in a tile-like pattern and each plat was separated by flesh-colored furrows (Fig. 1B). Based on the suspicion of TFFD, we attempted to rub the lesion firmly with alcohol swabs (Fig. 1C), exposing normal skin (Fig. 1D). There was no recurrence for a month.

TFFD is a bizarre condition which presents as dirt-like patches or plaques that are resistant to the usual cleansing method but resolved with alcohol swabs¹. According to a

report evaluating 11 cases of TFFD, almost two-third of the patients represented with TFFD at the surgical site post-operatively². Surgical procedures related to TFFD included coronary bypass grafting, thyroidectomy, total knee replacement, and laparotomy for perforated duodenal ulcer. This suggests that TFFD can be manifested as a phenomenon associated with surgery. Altered sensation due to nerve injury following surgery can elicit fear and prevent patients from proper skin cleansing. In addition, applying several dressing materials such as povidone iodine may trigger an excessive accumulation of sweat and sebum on the surgical site². Although our patient did not present with inappropriate washing habits, reluctance or hesitation to aggressive scrubbing may have been a contributing factor in the etiology of TFFD.

TFFD is easily misdiagnosed as it may clinically resemble other dermatological conditions. Dermatitis neglecta is the most challenging disease to be distinguished from TFFD³. It is more prevalent among patients with poor hygiene and is treated with standard cleansing, while TFFD patients usually have good hygiene and require vigorous alcohol rubs for restoration. Other differential diagnoses include confluent and reticulated papillomatosis, acanthosis nigricans and pityriasis versicolor³. Dermoscopy may aid the diagnosis of challenging cases and typically shows large pigmented polygonal areas with flat morphology arranged in a mosaic pattern in TFFD, which is consistent with our case⁴. On the other hand, poorly defined bordered, brownish pigmentation called a sulci and gyri pattern is visible in confluent and reticulated papillomatosis, focal dark brown spots on a yellowish background with aberrant crista cutis in acanthosis nigricans, and a pigmented scaly network consisting of brown bands in pityriasis versicolor⁵.

This is a case of TFFD which arose following thyroidectomy and successfully diagnosed based on the patient's history and dermoscopic findings. Awareness of its key dermoscopic features and its association with surgery is fundamental in the diagnosis of TFFD.

CONFLICT OF INTEREST

The authors have nothing to disclose.

REFERENCES

1. Moon J, Kim MW, Yoon HS, Cho S, Park HS. A case of terra firma-forme dermatosis: differentiation from other dirty-appearing diseases. *Ann Dermatol* 2016;28:413-415.
2. Ashique KT, Kaliyadan F, Goyal T. Terra firma-forme dermatosis: report of a series of 11 cases and a brief review

Fig. 1. (A) A brownish pigmented lesion on the patient's anterior neck. (B) Dermoscopy showing brown colored polygonal structures arranged in a tile-like pattern. (C) The lesion being cleared by hard alcohol swabs. (D) Normal skin revealed after the procedure.

- of the literature. *Int J Dermatol* 2016;55:769-774.
3. Chun SW, Lee SY, Kim JB, Choi HM, Ro BI, Cho HK. A case of terra firma-forme dermatosis treated with salicylic acid alcohol peeling. *Ann Dermatol* 2017;29:83-85.
 4. Abdel-Razek MM, Fathy H. Terra firma-forme dermatosis:

case series and dermoscopic features. *Dermatol Online J* 2015;21(10):20.

5. Zhou H, Tang XH, De Han J, Chen MK. Dermoscopy as an ancillary tool for the diagnosis of pityriasis versicolor. *J Am Acad Dermatol* 2015;73:e205-e206.

<https://doi.org/10.5021/ad.2018.30.4.501>

A Case of Low-Grade Fibromyxoid Sarcoma Arising in the Finger

Byeol Han, Sook-Ja Son, June Hyunkyung Lee, Tae Young Han

Department of Dermatology, Nowon Eulji Medical Center, Eulji University, Seoul, Korea

Dear Editor:

A 46-year-old man presented with a 5-year history of a nodule on his finger. Physical examination revealed a tender, solitary skin-colored irregular-shaped nodule measuring 1.4×1.4 cm, located on the left fourth finger (Fig. 1). Histological examination revealed a spindle-cell tumor in an admixture of fibromyxoid matrix in the dermis (Fig. 2A). The tumor consisted of alternating hypercellular myxoid and hypocellular fibrous stroma with an abrupt transition from one to the other. The tumor cells were bland spindle-shaped cells and stellate cells, arranged in a swirling or fascicular growth pattern (Fig. 2B). A few curvilinear vessels were observed. Mitotic figures were detected in 5/50 high power fields. Immunohistochemical studies showed reactivity for vimentin, CD99, CD68, and epithelial membrane antigen (EMA), whereas the tumor cells

were negative for CD34, S100, smooth muscle actin, and desmin. These findings were consistent with low-grade fibromyxoid sarcoma (LGFMS).

LGFMS is a rare soft tissue tumor, which typically occurs in the deep soft tissues of the proximal extremities or trunk in young to middle-aged adults^{1,2}.

Histologically, LGFMS consists of bland fibroblasts with a whorled or linear arrangement, alternating between paucicellular fibrous zones and myxoid zones¹⁻³. There is a

Fig. 1. A solitary skin-colored firm nodule on the middle phalanx of the left fourth finger.

Received September 5, 2017, Revised September 12, 2017, Accepted for publication September 17, 2017

Corresponding author: Tae Young Han, Department of Dermatology, Nowon Eulji Medical Center, Eulji University, 68 Hangeulbiseok-ro, Nowon-gu, Seoul 01830, Korea. Tel: 82-2-970-8580, Fax: 82-2-974-1577, E-mail: dermahan@gmail.com

ORCID: <https://orcid.org/0000-0003-0317-9635>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © The Korean Dermatological Association and The Korean Society for Investigative Dermatology