CASE REPORT

A novel cause of postmenopausal bleeding in an immunosuppressed patient

Neill Paul Kiely,¹ Lyndal Anderson,² Harpreet Arora,³ Christopher Benness³

¹Obstetrics and Gynaecology, The Royal Hospital for Women, Randwick, New South Wales, Australia

²Pathology, Royal Prince Alfred Hospital, Camperdown, New South Wales, Australia

³Obstetrics and Gynaecology, Royal Prince Alfred Hospital, Camperdown, New South Wales, Australia

Correspondence to
Dr Neill Paul Kiely,
neilldiver@gmail.com

Accepted 18 June 2018

SUMMARY

Malakoplakia is a rare histiocytic disease first described in 1902 by Michaelis and Gutmann. It is associated with host immunocompromise including chronic inflammatory conditions, infectious conditions or malnutrition. Here, we report the case of uterine malakoplakia as a rare cause of postmenopausal bleeding in an immunocompromised patient.

BACKGROUND

Malakoplakia is a rare histiocytic disease first described in 1902 by Michaelis and Gutmann.¹ Macroscopically, it can appear as yellow plaques and derived its name from the Greek words *malakos* (soft) and *plakos* (plaques). Microscopically, it is characterised by the presence of distinctive von Hanseman cells (histiocytic infiltrates) and Michaelis-Gutmann bodies (mineralised laminated cytoplasmic inclusions). It is associated with host immunocompromise including chronic inflammatory conditions, infectious conditions or malnutrition. More specifically, patients with organ transplantation, allergic conditions, chemotherapy, AIDS and malignancy are at increased risk. The urinary tract is the most frequent site of disease, in particular the urinary bladder.² The first description of malakoplakia outside of the urinary tract was in 1958, occurring in multiple other organs, most commonly the gastrointestinal tract, with only 18 cases reported in the literature of malakoplakia of the uterus. Malakoplakia is often asymptomatic, diagnosed incidentally or presenting with problematic vaginal bleeding.³ We present a unique case of concomitant bladder and uterine malakoplakia in an immunocompromised patient who presented with recurrent postmenopausal bleeding.

CASE PRESENTATION

A 65-year-old woman was referred by her urologist to the gynaecology outpatient clinic for review of her postmenopausal bleeding, on a background of known urinary bladder malakoplakia diagnosed in 2012 and vaginal malakoplakia diagnosed in 2014. She had a normal pelvic ultrasound (US) and MRI with ongoing vaginal bleeding. She was admitted for examination under anaesthesia, hysteroscopy and cystoscopy. She had a significant medical history including liver transplantation in 2008 for treatment of complications of autoimmune hepatitis which had been complicated by hyperacute rejection and managed with methylprednisone. She had

remained stable on a combination of tacrolimus, mycophenolate and prednisone. Additional medical history included steroid-induced type 2 diabetes mellitus, osteopenia, chronic renal impairment and recurrent urinary tract infections. She underwent an uncomplicated cystoscopy and hysteroscopy with findings of an atrophic vagina, 5 cm uterine cavity and abnormal soft yellow nodules on the anterior and posterior uterine walls which were suspicious for malakoplakia, and biopsies were taken. On cystoscopy, similar yellow nodules were identified within the urinary bladder.

INVESTIGATIONS

Pathological findings

Microscopy of the uterine curettings showed diffuse inflammatory infiltration. A diagnosis of malakoplakia was confirmed by the presence of von Hanseman histiocytes (figure 1) characterised by cells with distinctive cell borders, oval to round nuclei and abundant eosinophilic and granular cytoplasm. Several contained calcified Michaelis-Gutmann bodies. These were also present in the extracellular spaces. The Michaelis-Gutmann bodies were positive in the periodic acid-Schiff reaction (figure 2), and calcium was confirmed on the von Kossa stain (figure 3).

OUTCOME AND FOLLOW-UP

Within 2 hours postoperatively, she had a septic shower requiring admission to the intensive care unit with inotropic support. She was diagnosed with *Escherichia coli* (*E. coli*) bacteraemia and developed acute-on-chronic renal failure requiring haemodialysis. She recovered and was discharged on postoperative day 10.

DISCUSSION

Postmenopausal bleeding in all women requires prompt investigation for malignancy, especially in the context of chronic immunosuppression. This case highlights an additional differential diagnosis in the chronically immunosuppressed population. Malakoplakia is most commonly encountered in the mucosal surface of urinary and gastrointestinal tracts; rarely involving the genital tract.⁴

It is postulated that an acquired defect in intracellular degradation of bacteria is responsible for the development of this disease as it occurs in chronic immunosuppression⁵. Abnormal macrophage activity has been attributed to inadequate intracellular eradication of phagocytosed bacteria, leading to

To cite: Kiely NP, Anderson L, Arora H, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-225642

Figure 1 H&E stain showing von Hanseman histiocytes.

accumulation of calcium and iron on bacterial glycolipid within cellular phagolysosomes.⁵ The formation of Michaelis-Gutmann bodies has been thought to be the result of fusion and then calcification of the phagolysosomes within macrophages. The mechanism of failed exocytosis from the macrophage is still unclear; however, it is thought that it may be a function of decreased intracellular cyclic guanosine monophosphate and increased cyclic AMP resulting in reduced microtubular and lysosomal functioning.⁵

Coliform bacteria, in particular *E. coli*, are the most common organisms implicated in malakoplakia. Other organisms including *Rhodococcus equinovirus*, *Proteus* and *Klebsiella* have been associated with malakoplakia. Our patient developed an *E. coli* bacteraemia following biopsy of the bladder and uterine plaques and suffered from chronic *Klebsiella* and *E. coli* urinary tract infections.

Despite the rarity of malakoplakia in the female genital tract, it has been reported in all of the genital organs, with the vagina being the most common location.² Malakoplakia of the female genital tract is most commonly described in older patients with a mean age of 70 at diagnosis.³ Clinical investigations and examinations are often non-diagnostic, and this in combination with the rarity of the disease poses challenges for the diagnosis of

Figure 2 Michaelis-Gutmann bodies seen in the periodic acid-Schiff reaction.

Figure 3 von Kossa stain showing Michaelis-Gutmann bodies.

malakoplakia in the female genital tract.⁵ The clinical significance of this case being significant disease burden without reliable detection on MRI, pelvic US and CT.

The treatment of malakoplakia involves long-term antibiotic treatment, surgical excision or a combination of both.⁵ Appropriate antibiotic selection and duration are important to ensure eradication of malakoplakia. However, many studies have found that although antibiotic treatment may assist in reduction of lesion size, surgical intervention is necessary when the involved areas are extensive and multifocal.⁴ Of concern in this case is recurrence and progression of disease despite long-term antibiotic treatment. In conclusion, this case highlights the importance of further investigation of postmenopausal bleeding in the absence of disease on non-invasive imaging.

Learning points

- ▶ Postmenopausal bleeding in all women requires prompt investigation for malignancy, especially in the context of chronic immunosuppression.
- ▶ As our population ages and medical care improves, larger numbers of chronically immunosuppressed patients are reaching menopause and beyond.
- ▶ This case highlights an additional differential diagnosis when investigating postmenopausal bleeding in the chronically immunosuppressed population.
- ▶ Close monitoring and early identification of clinical deterioration in postoperative patients is important.

Contributors NPK is the primary author, the article was reviewed and edited by LA. HA and CB were primary clinicians involved in this case.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

REFERENCES

- 1 Michealis L, Gutmann C. Ueber Einschlüsse in Blasentumoren. *Ztschr Kin Med* 1902;47:208–15.

- 2 Yousef GM, Naghibi B, Hamodat MM, *et al.* Malakoplakia outside the urinary tract. *Arch Pathol Lab Med* 2007;131:297–300.
- 3 Kawai K, Fukuda K, Tsuchiyama H, *et al.* Malacoplakia of the endometrium. An unusual case studied by electron microscopy and a review of the literature. *Acta Pathol Jpn* 1988;38:531–40.
- 4 Chou SC, Wang JS, Tseng HH, *et al.* Malacoplakia of the ovary, fallopian tube and uterus: a case associated with diabetes mellitus. *Pathol Int* 2002;52:789–93.
- 5 Ramdial PK, Sing Y, Chotey NA, *et al.* Concomitant malacoplakia and granuloma inguinale of the cervix in acquired immune deficiency syndrome. *Int J Gynecol Pathol* 2008;27:282–7.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow