Published in final edited form as:

Clin Cancer Res. 2018 July 15; 24(14): 3456–3464. doi:10.1158/1078-0432.CCR-17-3615.

Identification and validation of a 3-gene methylation classifier for HPV-based cervical screening on self-samples

Wina Verlaet¹, Barbara C. Snoek¹, Daniëlle A.M. Heideman¹, Saskia M. Wilting^{1,3}, Peter J.F. Snijders¹, Putri W. Novianti^{1,2}, Annina P. van Splunter¹, Carel F.W. Peeters², Nienke E. van Trommel⁴, Leon F.A.G. Massuger⁵, Ruud L.M. Bekkers⁵, Willem J.G. Melchers⁶, Folkert J. van Kemenade⁷, Johannes Berkhof², Mark A. van de Wiel², Chris J.L.M. Meijer¹, and Renske D.M. Steenbergen^{1,*}

¹Cancer Center Amsterdam, Department of Pathology, VU University Medical Center, Amsterdam, The Netherlands ²Department of Epidemiology and Biostatistics, Amsterdam Public Health research institute, VU University Medical Center, Amsterdam, The Netherlands

⁴Department of Gynecology, Antoni van Leeuwenhoek Hospital/Netherlands Cancer Institute, Amsterdam, The Netherlands ⁵Department of Obstetrics and Gynaecology, Radboud University Medical Center, Nijmegen, The Netherlands ⁶Department of Medical Microbiology, Radboud University Medical Center, Nijmegen, The Netherlands ⁷Department of Pathology, Erasmus MC Cancer Institute, Rotterdam, The Netherlands

Abstract

Purpose—Offering self-sampling of cervico-vaginal material for high-risk human papillomavirus (hrHPV) testing is an effective method to increase the coverage in cervical screening programs. Molecular triage directly on hrHPV-positive self-samples for colposcopy referral opens the way to full molecular cervical screening. Here, we set out to identify a DNA methylation classifier for detection of cervical precancer (CIN3) and cancer, applicable to lavage and brush self-samples.

Experimental Design—We determined genome-wide DNA methylation profiles of 72 hrHPV-positive self-samples, using the Infinium Methylation 450K Array. The selected DNA methylation markers were evaluated by multiplex quantitative methylation-specific PCR (qMSP) in both

*Corresponding author: Renske D.M. Steenbergen, PhD, Department of Pathology, VU University Medical Center, PO Box 7057, 1007 MB Amsterdam, The Netherlands, Telephone number: +31-20-4442331, r.steenbergen@vumc.nl.

³Current address: Department of Medical Oncology, Erasmus MC Cancer Institute, Erasmus University Medical Center, Rotterdam, The Netherlands

Conflict of interest:

P. Snijders, C. Meijer, R. Steenbergen and D. Heideman have minority stake in Self-Screen B.V., a spin-off company of VU University Medical Center Amsterdam, which owns patents related to this work. P. Snijders has been on the speaker's bureau of Roche, Abbott, Gen-Probe, Qiagen and Seegene. He is consultant for Crucell Holland B.V. C. Meijer has participated in the sponsored speaker's bureau of Merck and Qiagen, and served occasionally on the scientific advisory board of Qiagen and Merck. C. Meijer has occasionally been a consultant for Qiagen and was a minority shareholder of Diassay B.V. till March 2016. He owns a small number shares of Qiagen and is part-time director of Self-Screen. D. Heideman has been on the speaker's bureau of Qiagen, and serves occasionally on the scientific advisory boards of Pfizer and Bristol-Meyer Squibb. J. Berkhof has received consultancy fees from Roche, GlaxoSmithKline and Merck, and received travel support from DDL. All fees were collected by his employer. R. Bekkers received grants from Roche and SP-MSD and has participated in the speakers' bureau of Roche. All other authors have no conflict of interest to declare.

hrHPV-positive lavage (n=245) and brush (n=246) self-samples from screening cohorts. Subsequently, logistic regression analysis was performed to build a DNA methylation classifier for CIN3 detection applicable to self-samples of both devices. For validation, an independent set of hrHPV-positive lavage (n=199) and brush (n=287) self-samples was analyzed.

Results—Genome-wide DNA methylation profiling revealed 12 DNA methylation markers for CIN3 detection. Multiplex qMSP analysis of these markers in large series of lavage and brush self-samples yielded a 3-gene methylation classifier (*ASCL1*, *LHX8* and *ST6GALNAC5*). This classifier showed a very good clinical performance for CIN3 detection in both lavage (AUC=0.88; sensitivity=74%; specificity=79%) and brush (AUC=0.90; sensitivity=88%; specificity=81%) self-samples in the validation set. Importantly, all self-samples from women with cervical cancer scored DNA methylation-positive.

Conclusion—By genome-wide DNA methylation profiling on self-samples, we identified a highly effective 3-gene methylation classifier for direct triage on hrHPV-positive self-samples, which is superior to currently available methods.

Introduction

Organized cytology-based cervical screening programs using physician-collected cervical scrapes have led to a substantial decrease in cervical cancer incidence and mortality in high-income countries (1). However, a considerable subset of women does not attend cervical screening (non-attendees), which compromises the effectiveness of the screening program (2). Previous studies have shown that offering self-sampling of cervico-vaginal specimens (self-samples) for high-risk human papillomavirus (hrHPV) testing (hrHPV self-sampling) to non-attendees, increases the attendance to cervical screening. Up to 30% of the invited non-attendees returned their self-sample to the laboratory for hrHPV testing (3–6). Importantly, the diagnostic accuracy of hrHPV testing on self-samples for cervical intraepithelial neoplasia grade 3 and cervical cancer (CIN3+) is similar to hrHPV-screening of physician-collected cervical scrapes (7,8). Therefore, offering hrHPV self-sampling as an alternative to conventional scrapes has just been implemented in the new hrHPV-based cervical screening program in The Netherlands. Partial substitution of hrHPV testing on physician-collected scrapes in cervical screening programs by hrHPV self-sampling can be envisioned in the near future.

Although hrHPV testing has a higher sensitivity for CIN3+ compared to cytology, its 3–5% lower specificity for CIN3+ necessitates the use of a triage test to distinguish women with clinically relevant disease from those with irrelevant, transient hrHPV infections to prevent over-referral and overtreatment. Currently, cytology is the most widely accepted triage tool. Since cytology cannot be reliably performed on self-sampled material (9–11), women with hrHPV-positive self-samples need to visit a physician for an additional cervical scrape for cytology. This may lead to loss to follow-up, delay the diagnostic track, and is less feasible in low income countries given the lack of adequate infrastructure and limited number of trained practitioners (8,12,13). Therefore, molecular triage testing directly applicable to self-sampled material from hrHPV-positive women is preferred.

We and others have shown that DNA methylation analysis of tumor suppressor genes on self-samples is well feasible and effective to detect CIN3+ using quantitative methylation-specific PCR (qMSP) (12–16). DNA methylation analysis has already shown competitive clinical performance versus other triage options in cervical scrapes, whereas improvements in performance on self-samples are conceivable. Previous findings have shown that DNA methylation markers originally discovered in tissue specimens and tested on hrHPV-positive cervical scrapes are not necessarily of clinical value when applied to hrHPV-positive self-samples (17). This is likely due to the cellular composition of self-samples, which contain fewer disease-related cells. Therefore, self-samples may display distinct epigenetic signatures compared to physician-collected cervical specimens. Hence, DNA methylation marker discovery screens directly performed on self-samples are more likely to yield the most informative DNA methylation markers for hrHPV-positive self-samples.

In this study, we describe the identification and validation of a DNA methylation classifier for the detection of CIN3 and cervical cancer in hrHPV-positive self-samples. A genome-wide DNA methylation marker discovery for CIN3 detection was performed using the Infinium 450K BeadChip array to 72 hrHPV-positive self-samples from a screening cohort of non-attendees. The identified candidate DNA methylation markers were evaluated by multiplex qMSP in unique, large series of lavage-based (n=245; further referred to as “lavage self-samples”) and brush-based (n=246; further referred to as “brush self-samples”) self-samples from screening cohorts of non-attendees to build an optimal DNA methylation classifier for detection of CIN3 that is applicable to self-samples of both devices. The clinical performance of the obtained DNA methylation classifier was subsequently validated by multiplex qMSP on an independent series of lavage (n=199) and brush (n=287) self-samples.

Materials and Methods

Clinical specimens

Discovery set: Case-control series for DNA methylation marker discovery screen—For genome-wide DNA methylation marker discovery for CIN3 detection, hrHPV-positive lavage self-samples collected using the Delphi screener (Delphi Bioscience, Scherpenzeel, The Netherlands) were obtained from a screening cohort of non-attendees (PROTECT-1 trial (3); NTR792) (n=72) (Figure 1; Discovery screen). Detailed characteristics of study design and clinical specimens, inclusion criteria and follow-up procedures have been described previously (3). Array data from a pilot experiment of 12 self-samples for power calculations revealed a ratio of 3 (hrHPV-positive controls) to 4 (CIN3) for proper marker discovery. Therefore, the discovery series comprised hrHPV-positive lavage self-samples from 29 control women, who either had histological evidence of absence of CIN2+ (CIN1) or displayed hrHPV clearance combined with normal cytology in follow-up (further referred to as hrHPV-positive controls; median age 36; range: 31–56), and 39 cases histologically diagnosed with CIN3 (median age 36; range: 31–62). Controls and cases were matched according to age and hrHPV type to the extent of sample availability. The hrHPV types in controls were 8 HPV16, 4 HPV51, 4 HPV52, 4 HPV56, 3 HPV45, 2 HPV35, 2 HPV58, 2 HPV66, 1 HPV33 and 1 HPV39; the hrHPV types in CIN3

were 21 HPV16, 6 HPV31, 4 HPV52, 3 HPV33, 3 HPV56, 2 HPV51, 2 HPV68, 1 HPV18, 1 HPV35, 1 HPV39, 1 HPV45 and 1 HPV66. Additionally, hrHPV-positive lavage self-samples from women histologically diagnosed with cervical squamous cell carcinoma (SCC; n=4) were included (median age 49; range: 42-61). The hrHPV types in SCC were 2 HPV16, 1 HPV31 and 1 HPV45.

Building set: Case-control series to build a DNA methylation classifier—To build a DNA methylation classifier for CIN3 detection, both hrHPV-positive lavage self-samples (n=245; PROTECT-1 trial (3); excluding samples used for the discovery screen) and brush self-samples collected using a VibaBrush (Rovers, Oss, The Netherlands) (n=246; PROTECT-2 trial (4); NTR1851) were obtained from screening cohorts of non-attendees who reached a study-endpoint and all of which were not pre-selected (Figure 1; Building a DNA methylation classifier; Supplementary Fig. S1). Detailed characteristics of study design and clinical specimens, inclusion criteria and follow-up procedures have been described previously (4). Available lavage self-samples of 214 hrHPV-positive controls (controls; median age 41; range 31-62) and 31 women histologically diagnosed with CIN3 (cases; median age 36; range 31-62) were included. Brush self-samples included 174 hrHPV-positive controls (controls; median age 37; range 30-62) and 72 women histologically diagnosed with CIN3 (cases; median age 36; range 31-61).

Validation set: Independent series to validate the DNA methylation classifier—To validate the clinical performance of the DNA methylation classifier, independent series of both hrHPV-positive lavage (n=199) and brush (n=287) self-samples, all of which were not pre-selected, were used (Figure 1; Validation of DNA methylation classifier; Supplementary Fig. S1). For lavage self-samples, hrHPV-positive samples collected using the Delphi Screener (Delphi Bioscience) were obtained from a screening cohort of non-attendees who reached a study-endpoint in the PROTECT-3 trial (methylation-arm; NTR2606) (12). Detailed characteristics of study design and clinical specimens, inclusion criteria and follow-up procedures have been described previously (12). Half of the available samples in this trial were randomly chosen for evaluation in the current study. These were supplemented with an independent series of 4 lavage self-samples from women with SCC who participated in the PROTECT-1 trial (3). The total lavage series comprised 134 hrHPV-positive controls (median age 38; range 33-63), 22 women with CIN2 (median age 38; range 33-58), 35 women with CIN3 (median age 38; range 33-48), 7 women with SCC (median age 48; range 38-61) and 1 woman with adenocarcinoma (AdCA; age 33). For brush self-samples, hrHPV-positive samples collected using the Evalyn brush (Rovers) were obtained from a screening cohort of non-attendees who reached a study-endpoint in the PROTECT-3B trial (NTR3350) (18). Detailed characteristics of study design and clinical specimens, inclusion criteria and follow-up procedures have been described previously (18). These were supplemented with an independent series of 4 brush self-samples from women with SCC and 1 brush self-samples from a woman with adenocarcinoma in situ (ACIS) who participated in the PROTECT-2 trial (4) and 7 brush self-samples from women with SCC and 5 brush self-samples from women with AdCA who visited the gynecology clinic (METC15.1468/X15MET study). The total brush series comprised 178 hrHPV-positive controls (median age 39; range 33-63), 28 women with CIN2 (median age 38; range 33-53),

56 women with CIN3 (median age 38; range 33-59), 16 women with SCC (median age 44; range 29-75), 1 woman with ACIS (age 41) and 8 women with AdCA (median age 44; range 27-62).

This study followed the ethical guidelines of the Institutional Review Board of VU University Medical Center and Antoni van Leeuwenhoek Hospital/Netherlands Cancer Institute. All participants in the PROHTECT and X15MET trials gave informed consent.

Infinium HumanMethylation450 BeadChip and data pre-processing

Before application, quality of the DNA was assessed by Qubit BR dsDNA measurement and visual evaluation of DNA integrity on an agarose gel. Genome-wide DNA methylation profiling was performed by Infinium HumanMethylation450 BeadChip (Illumina, San Diego, CA, USA). Data are available from the NCBI Gene Expression Omnibus (GEO) through series accession number GSE99511 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE99511>). Data were pre-processed and checked for sample and probe quality. Four samples (1 hrHPV-positive control and 3 CIN3) were excluded based on the correlation heatmap results (Supplementary Fig. S2). For further data pre-processing, see Supplementary Methods.

hrHPV and DNA methylation testing

For sample processing, hrHPV testing and DNA methylation analysis, see Supplementary Methods and Supplementary Table S1. HrHPV-positivity was determined for all samples. HrHPV genotypes were defined in a subset of the classifier building set only. In each multiplex qMSP assay, 3 targets and the housekeeping gene β -actin (*ACTB*) were combined as described before (19). Target DNA methylation values were normalized to reference gene *ACTB* and the calibrator using the comparative Ct method ($2^{-Ct} \times 100$) to obtain Ct ratios (20). The Ct ratios were square-root transformed. Only samples for which sufficient DNA material was available and which achieved a *ACTB* Ct value <30 were included.

Statistical analyses

Discovery screen: Genome-wide DNA methylation array data—After pre-processing of Infinium data, we applied adaptive group-regularized logistic ridge regression (GRridge) (21). We incorporated auxiliary information (referred to as co-data) in building the GRridge classification model, namely *p*-values from a similar study in cervical tissue specimens using the same array platform (Farkas *et al.* (22)), and standard deviation of each probe in the current dataset. Using informative co-data has been shown to enhance the identification of valuable markers in rather impure samples, such as self-samples (Supplementary Fig. S3). More details regarding the GRridge model by incorporating such information is provided in Supplementary Methods and elsewhere (21). Posthoc forward selection was applied to the GRridge model to render a model of DNA methylation markers. The performance of the GRridge model was visualized by a receiver operating characteristics (ROC) curve, obtained by leave-one-out-cross-validation, and quantified by area under the curve (AUC). Predicted probabilities, representing the risk for an underlying CIN3, were calculated using the GRridge model. Hierarchical clustering of the 28 DNA

methylation markers was performed to further select the genes that were most discriminative between CIN3 and hrHPV-positive controls.

Building and validation of DNA methylation classifier: qMSP data—To compare DNA methylation levels between 2 groups (hrHPV-positive controls and CIN3), the Wilcoxon rank sum test (two-sided) was applied on the square-root transformed Ct ratios. Statistical significance was set at $p < .05$.

To build a DNA methylation classifier, classical logistic regression analysis was performed on qMSP data to select relevant DNA methylation markers for CIN3 detection in both lavage and brush self-samples (detailed description in Supplementary Methods and Supplementary Fig. S4). In brief, logistic regression analysis followed by stepwise selection and backward elimination was performed on the combination of lavage and brush self-sample datasets (to encourage overlap) to obtain an initial marker panel of 2 DNA methylation markers for both self-sample types. Forward selection on the separate lavage and brush datasets suggested the addition of a 3rd DNA methylation marker, which was particularly relevant for the brush dataset, without harming the performance in the lavage dataset. Since DNA methylation in CpG islands has been shown to increase with age (23), we included age as a factor in the DNA methylation classifier. Supplementary Table S2 shows the p -value and contribution (coefficient/sd) of age and the 3rd DNA methylation marker *ST6GALNAC5* in the 3-gene methylation classifier. These two factors were included in the classifier since exclusion of age and *ST6GALNAC5* resulted in a lower performance in particularly the brush self-samples. Predicted probabilities and 95% confidence intervals (CI) were calculated for all analyzed samples using the logistic regression models of the DNA methylation classifier for lavage and brush self-samples. The clinical performance of the logistic regression models in both classifier building and validation sets was visualized by a ROC curve and evaluated by AUC calculation. The ROC curves show the sensitivity and specificity for the complete spectrum of different thresholds in predicted probabilities using the logistic regression models. A threshold was fixed for predicted probabilities corresponding to 80% specificity (lavage self-samples: 0.053; brush self-samples: 0.240) based on the classifier building set and subsequently evaluated in the independent validation set for CIN3 sensitivity and specificity. Additionally, the DNA methylation classifier at a fixed threshold was applied on self-samples from women with CIN2, SCC and ACIS/AdCA to evaluate the positivity rates in these disease categories. A classification and regression tree (CART) algorithm, which renders a DNA methylation classifier using marker-based cut-offs, was built for comparison to the continuous values obtained by regression. For the details of the CART method, see Supplementary Methods.

Results

An overview of the study design is given in Figure 1

Discovery of DNA Methylation markers in hrHPV-positive self-samples

In total, we obtained 68 genome-wide DNA methylation profiles of hrHPV-positive lavage self-samples from a screening cohort of non-attendees, of which 64 (28 controls and 36 women with CIN3) were suitable to identify DNA methylation markers for CIN3 detection

(Figure 1; Discovery screen). Adaptive group-regularized ridge regression, GRridge (21), and variable selection on the DNA methylation profiles from women with and without CIN3 yielded a panel of 28 DNA methylation markers with discriminatory power for CIN3 (AUC of 0.77). Hierarchical clustering of all 28 DNA methylation markers showed that 12 methylated genes, i.e., *ACAN*, *ASCL1*, *LHX8*, *MYADM*, *NRG3*, *RGS7*, *ST6GALNAC3*, *ST6GALNAC5*, *WDR17*, *ZNF582*, *ZNF583* and *ZNF781*, were mostly contributing to the discrimination of women with and without CIN3 (Figure 1; Hierarchical clustering, Figure 2A, Table 1 and Supplementary Fig. S5). Evaluation of the DNA methylation profiling data from 4 hrHPV-positive lavage self-samples from women with SCC confirmed high DNA methylation levels for all these 12 DNA methylation markers (Figure 2B and Supplementary Fig. S5).

Building a DNA methylation classifier using hrHPV-positive lavage and brush self-samples

Next, the 12 most discriminative DNA methylation markers from the discovery screen were further analyzed using multiplex qMSP in large series of hrHPV-positive lavage self-samples (n=245) and brush self-samples (n=246) from women with and without CIN3 from two screening cohorts (Figure 1; Building a DNA methylation classifier). In both lavage and brush self-samples, all genes except *ACAN* (in lavage only; $p < .05$), showed significantly increased methylation levels ($p < .001$) in self-samples from women with CIN3 compared to hrHPV-positive controls (Figure 3).

To build an optimal DNA methylation classifier for detection of CIN3, which is applicable to different self-sample types, logistic regression analysis followed by stepwise selection and backward elimination was performed on the combined dataset of lavage and brush self-sample qMSP results (see Materials and Methods, Supplementary Methods and Supplementary Fig. S4). This revealed a 3-gene methylation classifier for CIN3 detection in both self-sample types, consisting of *ASCL1*, *LHX8* and *ST6GALNAC5* (Supplementary Fig. S4 and Supplementary Table S2). This 3-gene methylation classifier showed a very good clinical performance for CIN3 detection in both hrHPV-positive lavage (AUC=0.90) and brush (AUC=0.86) self-samples (Figure 4A and B, black lines). At the threshold corresponding to a specificity of 80% in hrHPV-positive controls, 83% (25 of 30) of lavage self-samples and 76% (52 of 68) of brush self-samples from women with CIN3 were DNA methylation-positive (Supplementary Fig. S6).

Validation of DNA methylation classifier

To validate the clinical performance of the 3-gene methylation classifier, an independent, large series of hrHPV-positive lavage self-samples (n=199) and brush self-samples (n=287) was analyzed by multiplex qMSP (Figure 1; Validation of DNA methylation classifier). Solely hrHPV-positive controls and CIN3 from independent screening cohorts were used for validation of the 3-gene methylation classifier. This showed a comparable clinical performance for CIN3 detection as observed in the above described classifier building set, in both hrHPV-positive lavage (AUC=0.88) and brush (AUC=0.90) self-samples (Figure 4A and B, grey lines). The predefined threshold corresponding to an 80% specificity in the classifier building set (see above) was applied to this validation set. This resulted in a CIN3 sensitivity of 74% (26 of 35) in lavage self-samples and 88% (49 of 56) in brush self-

samples, at 79% and 81% specificity in hrHPV-positive controls, respectively (Supplementary Fig. S6). To confirm these findings, we applied an alternative method (CART) on both lavage and brush self-samples, which rendered similar results to those shown here (Supplementary Methods, Table S3, Fig. S7 and S8).

Furthermore, this validation series also comprised self-samples from women with CIN2 from a screening cohort. Fifty percent of these lavage self-samples (11 of 22) and brush self-samples (14 of 28) were DNA methylation-positive (Supplementary Fig. S6). Importantly, all 23 SCC (7 lavage self-samples and 16 brush self-samples; Supplementary Fig. S6), and all ACIS (1 brush self-sample) and AdCA (1 lavage self-sample and 8 brush self-samples) scored DNA methylation-positive (Supplementary Fig. S9).

Discussion

Here, we identified a DNA methylation classifier consisting of 3 methylated gene promoters, *ASCL1*, *LHX8* and *ST6GALNAC5*, for the detection of CIN3 and cervical cancer in hrHPV-positive self-samples and validated the clinical performance in large series of both cervical lavage and brush self-samples from independent screening cohorts of non-attendees.

Previous publications showed that CIN lesions detected by DNA methylation analysis do not completely overlap with those detected by cytology (24). In fact, DNA methylation analysis tends to preferably detect cervical cancer and advanced high-grade precursor lesions, defined as CIN2/3 associated with a persistent hrHPV infection of >5 years. Women with advanced CIN2/3 are presumed to have a high short-term progression risk to cancer and are therefore in need of immediate referral and treatment (24,25). Cytology on the other hand, detects both early and advanced CIN lesions with a moderate sensitivity of 65%-80%, and cannot be reliably applied to self-samples, requiring a visit to the physician (9–11). DNA methylation markers are applicable on self-samples and have the potential to reduce the risk for undetected cervical cancers and advanced CIN2/3. Contrarily, women with a negative DNA methylation marker test would have a low short-term cancer progression risk, indicating that immediate colposcopy referral is unnecessary. To prevent over-referral and overtreatment in hrHPV-based self-sampling, direct triage testing by DNA methylation markers in self-sampled material enables the identification of only those hrHPV-positive women with clinically relevant disease who are in need of treatment and it allows for full molecular cervical self-screening.

This is the first study performing a discovery screen directly on self-samples, which allowed us to define the most optimal DNA methylation classifier for direct molecular triage testing on hrHPV-positive self-sampled material. Our 3-gene methylation classifier showed a very good and reproducible clinical performance for detection of CIN3 in both hrHPV-positive lavage (classifier building set AUC=0.90; classifier validation set AUC=0.88) and brush (classifier building set AUC=0.86; classifier validation set AUC=0.90) self-samples. This indicates that it represents a universal triage test for both self-sample devices. Furthermore, the combined analysis of the 3-gene methylation classifier and a reference gene in a single multiplex assay saves material, costs and time and allows for (semi)high-throughput screening.

To select the most discriminatory DNA methylation markers for CIN3 from our discovery screen on hrHPV-positive self-samples, which are rather impure due to an overrepresentation of non-disease related cells, we applied our recently proposed GRridge model (21). This method enables objective use of co-data and was shown to potentially outperform other prediction methods (Supplementary Fig. S3) (26). In particular publicly available DNA methylation data from relatively pure cervical tissue specimens, obtained by the same array platform, proved to be useful co-data (22). The validity of this approach is supported by the identification of the 3 DNA methylation classifier genes, that have all been previously described in DNA methylation studies on cervical cancer (22,27,28). The combination of GRridge (on array data) and classical logistic regression analysis (on qMSP data) enabled us to build a highly discriminative methylation classifier for CIN3 detection consisting of *ASCL1*, *LHX8* and *ST6GALNAC5*. The narrow range of the 95% CI of the predicted probabilities (i.e., the methylation classifier value; range 0 to 1) in both lavage and brush self-samples supports a good representation of the disease state (case vs control) in the population by the 3-gene methylation classifier (Supplementary Fig. S10). Comparison of the 3 markers in HPV16-positive self-samples to self-samples positive for other hrHPV types (non-HPV16), in the subset of samples with HPV typing information, revealed no significant difference in DNA methylation levels in both lavage and brush self-samples, except for *LHX8* in HPV16 versus non-HPV16 controls of lavage self-samples (p -value = 0.03; Supplementary Fig. S11).

ASCL1, achaete-scute family bHLH transcription factor 1, is a proneural transcription factor and functions as a main regulator of differentiation in neurogenesis (29). *LHX8*, LIM homeobox 8, is a highly conserved transcription factor regulating cell fate in neurogenesis, tooth morphogenesis and oogenesis (30). *ST6GALNAC5*, ST6 N-Acetylgalactosaminide Alpha-2,6-Sialyltransferase 5, is a transmembrane sialyltransferase involved in the biosynthesis of gangliosides on the cell surface (31). Next to cervical cancer, *LHX8* methylation has been detected in breast cancer (32), *ST6GALNAC5* methylation has been described in colorectal cancer studies (33) and *ASCL1* methylation has been detected in oral and colorectal cancer (34,35).

Of the previously described DNA methylation markers tested in self-samples (12–14,17,24), the DNA methylation panel *FAM19A4/miR124-2* showed the best clinical performance in a large screening cohort. Analysis of the same study cohorts as used in present study showed a CIN3+ sensitivity of 70.5% in lavage and 69.4% in brush self-samples at a specificity of 67.8% and 76.4%, respectively (14). Within the CIN3+ group, 68% of CIN3 and all cancers were detected in both self-sample types. Other DNA methylation marker panels, such as *JAM3/EPB41L3/TERT/C13ORF18*, have only been analyzed in small selected series of self-samples (15,16). A combination of DNA methylation markers with HPV16/18 genotyping results in higher sensitivities compared with solely DNA methylation, however at the cost of severe lower specificities due to detection of early CIN2/3 (14,36). Our 3-gene methylation classifier shows a better sensitivity for CIN3 than other assays in both lavage (74%) and brush (88%) self-samples in a similar screening population, at a higher specificity of 79% and 81%, respectively. These findings emphasize the validity and importance of our approach to perform the DNA methylation marker discovery directly on self-sampled material. Furthermore, the 3-gene methylation classifier detected all self-samples from

women with SCC. Importantly, all self-samples from women with SCC showed very high predicted probabilities (median: 1.00; range: 0.54-1.00), which accentuates the value of our 3-gene methylation classifier for detection of cervical cancer. Additionally, all self-samples from women with ACIS and AdCA scored DNA methylation-positive, indicating that glandular lesions are also detected by our 3-gene methylation classifier. Nevertheless, further evaluation of cervical glandular lesions and other rare cervical cancer types is warranted.

A limitation of our study is that we used cohorts of non-attending women. Therefore, further confirmation in a regular population-based population is warranted. In addition, the Infinium 450K array is limited to 485,577 CpG measurements. A new version of this platform, the Infinium MethylationEPIC Beadchip array, covers over 850,000 CpG sites and would yield more discovery data, especially in the enhancer regions (37). Although the Infinium 450K array is not fully genome-wide, and may yield partly different results than other methylome analysis methods, all 485,577 probes cover 99% of Refseq genes and 96% of all CpG islands with multiple probes per gene and CpG island (38). Furthermore, this array is one of the most widely accepted methods for genome-wide DNA methylation profiling and it is cost-effective (39).

In conclusion, by genome-wide DNA methylation profiling on self-samples obtained from a screening trial, we identified and validated an effective 3-gene methylation classifier for detection of CIN3 and cervical cancer in both lavage and brush self-samples from hrHPV-positive women. Moreover, this 3-gene methylation classifier showed an improved clinical performance compared to current (complex) triage strategies for the management of hrHPV-positive self-samples (13). Our findings indicate that a transition towards full molecular self-screening in hrHPV-based cervical screening programs is feasible.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Lise De Strooper, Bart Hesselink, Maarten van der Salm, Saskia Doorn, Martijn Bogaarts and Dénira Agard for excellent technical assistance. In addition, we thank dr. S. Farkas for providing the raw data of her study (22). This work was supported by the European Research Council (ERC advanced 2012-AdG; 322986; Mass-Care) to C.J.L.M. Meijer and by ZonMw (Netherlands Organisation for Health Research and Development; 91216012) to M.A. van de Wiel.

References

1. Peto PJ, Gilham PC, Fletcher O, Matthews FE. The cervical cancer epidemic that screening has prevented in the UK. *Lancet*. 2004; 364:249–56. [PubMed: 15262102]
2. Bos AB, Rebolj M, Habbema JDF, Van Ballegooijen M. Nonattendance is still the main limitation for the effectiveness of screening for cervical cancer in the Netherlands. *Int J Cancer*. 2006; 119:2372–5. [PubMed: 16858676]
3. Gok M, Heideman DAM, Van Kemenade FJ, Berkhof J, Rozendaal L, Spruyt JWM, et al. HPV testing on self collected cervicovaginal lavage specimens as screening method for women who do not attend cervical screening: cohort study. *Br Med J*. 2010; 340:c1040. [PubMed: 20223872]
4. Gok M, Van Kemenade FJ, Heideman DAM, Berkhof J, Rozendaal L, Spruyt JWM, et al. Experience with high-risk human papillomavirus testing on vaginal brush-based self-samples of

- non-attendees of the cervical screening program. *Int J cancer*. 2012; 130:1128–35. [PubMed: 21484793]
5. Bais AG, Van Kemenade FJ, Berkhof J, Verheijen RHM, Snijders PJF, Voorhorst F, et al. Human papillomavirus testing on self-sampled cervicovaginal brushes: An effective alternative to protect nonresponders in cervical screening programs. *Int J Cancer*. 2007; 120:1505–10. [PubMed: 17205514]
 6. Racey CS, Withrow DR, Gesink D. Self-collected HPV Testing Improves Participation in Cervical Cancer Screening: A Systematic Review and Meta-analysis. *Can J Public Heal*. 2013; 104:159–66.
 7. Snijders PJF, Verhoef VMJ, Arbyn M, Ogilvie G, Minozzi S, Banzi R, et al. High-risk HPV testing on self-sampled versus clinician-collected specimens: A review on the clinical accuracy and impact on population attendance in cervical cancer screening. *Int J Cancer*. 2013; 132:2223–36. [PubMed: 22907569]
 8. Arbyn M, Verdoodt F, Snijders PJF, Verhoef VMJ, Suonio E, Dillner L, et al. Accuracy of human papillomavirus testing on self-collected versus clinician-collected samples: A meta-analysis. *Lancet Oncol*. 2014; 15:172–83. [PubMed: 24433684]
 9. Rijkaart DC, Berkhof J, Van Kemenade FJ, Coupé VMH, Hesselink AT, Rozendaal L, et al. Evaluation of 14 triage strategies for HPV DNA-positive women in population-based cervical screening. *Int J Cancer*. 2012; 130:602–10. [PubMed: 21400507]
 10. Dijkstra MG, Van Niekerk D, Rijkaart DC, Van Kemenade FJ, Heideman DAM, Snijders PJF, et al. Primary hrHPV DNA Testing in Cervical Cancer Screening: How to Manage Screen-Positive Women? A POBASCAM Trial Substudy. *Cancer Epidemiol Biomarkers Prev*. 2014; 23:55–63. [PubMed: 23733907]
 11. Garcia F, Barker B, Santos C, Brown EM, Muñoz T, Giuliano A, et al. Cross-sectional Study of Patient- and Physician- Collected Cervical Cytology and Human Papillomavirus. *Obstet Gynecol*. 2003; 102:266–72. [PubMed: 12907098]
 12. Verhoef VMJ, Bosgraaf RP, Van Kemenade FJ, Rozendaal L, Heideman DAM, Hesselink AT, et al. Triage by methylation-marker testing versus cytology in women who test HPV-positive on self-collected cervicovaginal specimens (PROTECT-3): A randomised controlled non-inferiority trial. *Lancet Oncol*. 2014; 15:315–22. [PubMed: 24529697]
 13. Luttmer R, De Strooper LMA, Steenbergen RDM, Berkhof J, Snijders PJF, Heideman DAM, et al. Management of high-risk HPV-positive women for detection of cervical (pre)cancer. *Expert Rev Mol Diagn*. 2016; 16:961–74. [PubMed: 27459506]
 14. De Strooper LMA, Verhoef VMJ, Berkhof J, Hesselink AT, De Bruin HME, Van Kemenade FJ, et al. Validation of the FAM19A4 / mir124-2 DNA methylation test for both lavage- and brush-based self-samples to detect cervical (pre) cancer in HPV-positive women. *Gynecol Oncol*. 2016; 141:341–7. [PubMed: 26921784]
 15. Boers A, Bosgraaf RP, van Leeuwen RW, Schuurin E, Heideman DAM, Massuger LFAG, et al. DNA methylation analysis in self-sampled brush material as a triage test in hrHPV-positive women. *Br J Cancer*. 2014; 111:1095–101. [PubMed: 25032730]
 16. Eijssink JJH, Yang N, Lendvai A, Klip HG, Volders HH, Buikema HJ, et al. Detection of cervical neoplasia by DNA methylation analysis in cervico-vaginal lavages, a feasibility study. *Gynecol Oncol*. 2011; 120:280–3. [PubMed: 21093897]
 17. Hesselink AT, Heideman DAM, Steenbergen RDM, Gök M, Van Kemenade FJ, Wilting SM, et al. Methylation marker analysis of self-sampled cervico-vaginal lavage specimens to triage high-risk HPV-positive women for colposcopy. *Int J Cancer*. 2014; 135:880–6. [PubMed: 24474183]
 18. Bosgraaf RP, Verhoef VMJ, Massuger LFAG, Siebers AG, Bulten J, de Kuyper-de Ridder GMD, et al. Comparative performance of novel self-sampling methods in detecting high-risk human papillomavirus in 30,130 women not attending cervical screening. *Int J cancer*. 2015; 136:646–55. [PubMed: 24923998]
 19. Snellenberg S, De Strooper LMA, Hesselink AT, Meijer CJLM, Snijders PJF, Heideman DAM, et al. Development of a multiplex methylation-specific PCR as candidate triage test for women with an HPV-positive cervical scrape. *BMC Cancer*. 2012; 12:551. [PubMed: 23176198]
 20. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. *Nat Protoc*. 2008; 3:1101–8. [PubMed: 18546601]

21. van de Wiel MA, Lien TG, Verlaet W, van Wieringen WN, Wilting SM. Better prediction by use of co-data: adaptive group-regularized ridge regression. *Stat Med*. 2016; 35:368–81. [PubMed: 26365903]
22. Farkas SA, Milutin-Gašperov N, Grce M, Nilsson TK. Genome-wide DNA methylation assay reveals novel candidate biomarker genes in cervical cancer. *Epigenetics*. 2013; 8:1213–25. [PubMed: 24030264]
23. Christensen BC, Houseman EA, Marsit CJ, Zheng S, Wrensch MR, Wiemels JL, et al. Aging and environmental exposures alter tissue-specific DNA methylation dependent upon CPG island context. *PLoS Genet*. 2009; 5:e1000602. [PubMed: 19680444]
24. Steenbergen RDM, Snijders PJF, Heideman DAM, Meijer CJLM. Clinical implications of (epi)genetic changes in HPV-induced cervical precancerous lesions. *Nat Rev Cancer*. 2014; 14:395–405. [PubMed: 24854082]
25. De Strooper LMA, Meijer CJLM, Berkhof J, Hesselink AT, Snijders PJF, Steenbergen RDM, et al. Methylation analysis of the FAM19A4 gene in cervical scrapes is highly efficient in detecting cervical carcinomas and advanced CIN2/3 lesions. *Cancer Prev Res*. 2014; 7:1251–7.
26. Novianti PW, Snoek BC, Wilting SM, van de Wiel MA. Better diagnostic signatures from RNAseq data through use of auxiliary co-data. *Bioinformatics*. 2017; 33:1572–4. [PubMed: 28073760]
27. Boers A, Wang R, van Leeuwen RW, Klip HG, De Bock GH, Hollema H, et al. Discovery of new methylation markers to improve screening for cervical intraepithelial neoplasia grade 2/3. *Clin Epigenetics*. 2016; 8:29. [PubMed: 26962367]
28. Clarke MA, Luhn P, Gage JC, Bodelon C, Dunn ST, Walker J, et al. Discovery and validation of candidate host DNA methylation markers for detection of cervical precancer and cancer. *Int J cancer*. 2017; 141:701–10. [PubMed: 28500655]
29. Vasconcelos FF, Castro DS. Transcriptional control of vertebrate neurogenesis by the proneural factor *Ascl1*. *Front Cell Neurosci*. 2014; 8:412. [PubMed: 25520623]
30. Zhou C, Yang G, Chen M, He L, Xiang L, Ricupero C, et al. *Lhx6* and *Lhx8*: Cell fate regulators and beyond. *FASEB J*. 2015; 29:4083–91. [PubMed: 26148970]
31. Drolez A, Vandenhaute E, Delannoy CP, Dewald JH, Gosselet F, Cecchelli R, et al. *ST6GALNAC5* expression decreases the interactions between breast cancer cells and the human blood-brain barrier. *Int J Mol Sci*. 2016; 17:1309.
32. Tommasi S, Karm DL, Wu X, Yen Y, Pfeifer GP. Methylation of homeobox genes is a frequent and early epigenetic event in breast cancer. *Breast cancer Res*. 2009; 11:R14. [PubMed: 19250546]
33. Øster B, Thorsen K, Lamy P, Wojdacz TK, Hansen LL, Birkenkamp-Demtröder K, et al. Identification and validation of highly frequent CpG island hypermethylation in colorectal adenomas and carcinomas. *Int J Cancer*. 2011; 129:2855–66. [PubMed: 21400501]
34. Jin B, Yao B, Li J-L, Fields CR, Delmas AL, Liu C, et al. *DNMT1* and *DNMT3B* modulate distinct polycomb-mediated histone modifications in colon cancer. *Cancer Res*. 2009; 69:7412–21. [PubMed: 19723660]
35. Li Y-F, Hsiao Y-H, Lai Y-H, Chen Y-C, Chen Y-J, Chou J-L, et al. DNA methylation profiles and biomarkers of oral squamous cell carcinoma. *Epigenetics*. 2015; 10:229–36. [PubMed: 25612142]
36. Verhoef VMJ, Heideman DAM, Van Kemenade FJ, Rozendaal L, Bosgraaf RP, Hesselink AT, et al. Methylation marker analysis and HPV16/18 genotyping in high-risk HPV positive self-sampled specimens to identify women with high grade CIN or cervical cancer. *Gynecol Oncol*. 2014; 135:58–63. [PubMed: 25111387]
37. Moran S, Arribas C, Esteller M. Validation of a DNA methylation microarray for 850,000 CpG sites of the human genome enriched in enhancer sequences. *Epigenomics*. 2016; 8:389–99. [PubMed: 26673039]
38. Bibikova M, Barnes B, Tsan C, Ho V, Klotzle B, Le JM, et al. High density DNA methylation array with single CpG site resolution. *Genomics*. 2011; 98:288–95. [PubMed: 21839163]
39. Yong W-S, Hsu F-M, Chen P-Y. Profiling genome-wide DNA methylation. *Epigenetics Chromatin*. 2016; 9:26. [PubMed: 27358654]

Translational Relevance

Offering self-sampling of cervico-vaginal specimens for hrHPV testing to non-attendees increases the attendance rate in cervical screening. However, an additional triage test directly applicable on self-sampled material is necessary to identify hrHPV-positive women at risk for progression to cervical cancer. Since cytology, the wide-accepted triage method, cannot be reliably performed on self-sampled material, there is an urgent need for molecular triage markers. This is the first study performing a genome-wide DNA methylation discovery directly on self-samples, which allowed us to define the most optimal DNA methylation markers. We identified and validated a highly effective 3-gene methylation classifier (*ASCL1*, *LHX8* and *ST6GALNAC5*) for detection of cervical precancer and cancer in both lavage and brush self-samples from hrHPV-positive women, which outperforms currently available methods. These findings could greatly improve the clinical management of women with hrHPV-positive self-samples and indicate that a transition to a full molecular self-screening approach in cervical screening programs is feasible.

Figure 1. Experimental setup of the study.

All self-samples were obtained from screening cohorts of non-attendees, except 7 SCC and 5 AdCA brush self-samples in the validation set.

Figure 2. Heatmap of the 28 DNA methylation markers in the discovery screen.

Hierarchical clustering of the 28 Infinium 450K BeadChip probes, each probe corresponds to a DNA methylation marker. Low (blue) to high (purple) DNA methylation levels (arcsine square-root transformed beta values) are displayed for each DNA methylation marker (cg-numbers of the probes). **(A)** DNA methylation data of self-samples from hrHPV-positive controls (green; n=28) and from women with CIN3 (orange; n=36). The samples are ordered by predicted probability. The 12 DNA methylation markers above the black line showed the most discriminative DNA methylation profile between women with and without CIN3. **(B)** DNA methylation data of self-samples from women with SCC (red; n=4).

Figure 3. Differential DNA methylation levels of the 12 candidate methylation markers in hrHPV-positive self-samples.

DNA methylation levels represented by the square-root transformed Ct ratios (y-axis) in (A) lavage self-samples from hrHPV-positive controls (n=214) and women with CIN3 (n=31; x-axis), and (B) brush self-samples from hrHPV-positive controls (n=174) and women with CIN3 (n=72; x-axis). The 3 genes left of the black line are included in the 3-gene methylation classifier. * $p < .05$; *** $p < .001$; NS: not significant.

Figure 4. Clinical performance of the 3-gene methylation classifier for CIN3 detection in hrHPV-positive lavage and brush self-samples.

ROC curve and AUC of the 3-gene methylation classifier for CIN3 detection in (A) lavage and (B) brush self-samples in classifier building set (grey) and validation set (black).

Table 1

The 12 candidate DNA methylation markers from the discovery screen.

Infinium BeadChip probe	Chr.	Chr. Location	Gene name
cg08272731	1	75602167	<i>LHX8</i>
cg14156405	1	241520286	<i>RGS7</i>
cg20707222	1	76540222	<i>ST6GALNAC3</i>
cg23243867	1	77334045	<i>ST6GALNAC5</i>
cg27486637	4	176987174	<i>WDR17</i>
cg10401879	10	83634276	<i>NRG3</i>
cg20718350	12	103352294	<i>ASCL1</i>
cg06675190	15	89346205	<i>ACAN</i>
cg13499300	19	54369556	<i>MYADM</i>
cg02763101	19	56904945	<i>ZNF582</i>
cg00796360	19	56915650	<i>ZNF583</i>
cg14587524	19	38183262	<i>ZNF781</i>

Chr.: Chromosome