Published in final edited form as:

Psychooncology. 2018 March ; 27(3): 879–885. doi:10.1002/pon.4584.

Costs of an Ostomy Self-Management Training Program for Cancer Survivors

Mark C. Hornbrook, PhD¹, Martha D. Cobb, MS, MEd, CWOCN², Nancy J. Tallman, BSN, WOCN³, Janice Colwell, RN, MS, CWOCN, APN⁴, Ruth McCorkle, RN, PhD, FAAN⁵, Elizabeth Ercolano, RN, MSN, DNSc⁵, Marcia Grant, RN, PhD, FAAN⁶, Virginia Sun, RN, PhD⁶, Christopher S. Wendel, MS^{7,8}, Judith H. Hibbard, DrPH, MPH⁹, and Robert S. Krouse, MD^{10,11}

¹Kaiser Permanente Center for Health Research, Portland, Oregon

²University of Arizona College of Nursing, Tucson, AZ

³Wound Ostomy Continence Nurse, Unaffiliated, Tucson, AZ

⁴University of Chicago School of Medicine, Chicago, IL

⁵Yale University School of Nursing, New Haven, CT

⁶City of Hope National Medical Center/Beckman Research Institute, Duarte, CA

⁷University of Arizona College of Medicine, Tucson, Arizona

⁸Southern Arizona Veterans Affairs Health Care System, Tucson, AZ

⁹University of Oregon Health Policy Research Group, Eugene, OR

¹⁰University of Pennsylvania School of Medicine, Philadelphia, PA

¹¹CMC Veterans Affairs Medical Center, Philadelphia, PA

Abstract

Objective—To measure incremental expenses to an oncologic surgical practice for delivering a community-based, ostomy nurse-led, small-group, behavior skills-training intervention to help bladder and colorectal cancer survivors understand and adjust to their ostomies and improve their health-related quality of life, as well as assist family caregivers to understand survivors' needs and provide appropriate supportive care.

Methods—The intervention was a five-session group behavior skills training in ostomy self-management following the principles of the Chronic Care Model. Faculty included Wound, Ostomy, Continence Nurses (WOCNs) utilizing an ostomy care curriculum. A gender-matched

Address Correspondence and Reprint Requests to: Mark C. Hornbrook PhD, Senior Investigator Emeritus, Kaiser Permanente Center for Health Research, 3800 North Interstate Avenue, Portland, OR 97227-1110, (503) 780-9613, mark.c.hornbrook@gmail.com.

NOTE

This analysis is a supplement to an article that previously appeared in *Psycho-Oncology*: Krouse RS, Grant M, McCorkle R, Wendel CS, Cobb MD, Tallman NJ, Ercolano E, Sun V, Hibbard JH, Hornbrook MC. A chronic care ostomy self-management program for cancer survivors. *Psycho-Oncology* 2016;25: 574-581. doi:10.1002/pon.4078; PMID:26804708 NIHMS 749541.

The authors do not have any conflicts of interest.

peer-in-time buddy was assigned to each ostomy survivor. The four-session Survivor curriculum included: Self-management practice and solving immediate ostomy concerns; Social well-being; Healthy lifestyle; and a Booster session. The single Family Caregiver session was co-led by a WOCN and an ostomy Peer staff member and covered relevant caregiver and Ostomate support issues. Each cohort required eight weeks to complete the intervention. Non-labor inputs included ostomy supplies, teaching materials, automobile mileage for WOCNs, mailing, and meeting space rental. Intervention personnel were employed by the University of Arizona. Labor expenses included salaries and fringe benefits.

Results—The total incremental expense per intervention cohort of four Survivors was \$7,246, or \$1,812 per patient.

Conclusions—A WOCN-led group self-help ostomy survivorship intervention provided affordable, effective, care to cancer survivors with ostomies.

Keywords

Oncology; Colorectal Cancer; Bladder Cancer; Ostomy; Self-Care; Chronic Care Model; Cancer Survivorship; Costs; Social Support; Wound and Ostomy Care Nursing

BACKGROUND

Emulating the development of consensus care for colorectal and urostomy surgery,¹ the WOCN Society Consensus Conference on discharge criteria from home care for persons with new fecal or urinary diversions recommended that "... every person with an ostomy should have access to an ostomy nurse specialist in the ambulatory care setting who can provide ongoing education, counseling, and assistance with physical problems associated with the individual's ostomy."² The WOCN research literature emphasizes the importance of post-discharge care for stoma patients for physical, psychological, and social problems.^{3, 4} From 20% to 71% of persons with ostomies experience complications such as pouch leakage and peristomal skin problems.⁴ Peristomal skin problems affect one-third of colostomy patients and two-thirds of urostomy and ileostomy patients.⁴ Post-discharge support and education are needed to prevent, diagnose, and treat peristomal skin problems.⁵

As one step towards this goal, the authors pilot-tested the Ostomy Self-Management Training program (OSMT) to improve patients' health-related quality of life (HRQOL) through a systematic program driven by Ostomates' needs.⁶ Formal teaching by hospital-based Wound, Ostomy, Continence Nurses (WOCNs) may begin preoperatively and continue postoperatively, but usually does not continue after discharge.⁷ WOCNs are typically hospital or Skilled Nursing Facility (SNF) employees, and most hospitals and SNFs have not developed outpatient clinics staffed by WOCNs. As hospital stays have shortened, so has the time available for ostomy teaching and practice in inpatient settings. Current peri-operative and long-term ostomy care is also constrained by lack of formal post-discharge follow-up with WOCNs. Time constraints faced by WOCNs, absence of reimbursement for such activities, and lack of research are factors contributing to this care gap. Ostomates and their families are often left to trial-and-error methods to improve self-management.⁶ Only when severe complications arise do surgeons and/or WOCNs become

involved in a patient's recovery after discharge. In hospital settings, other forms of patient information, such as booklets and websites, may be biased from ostomy product companies, anecdotal without rigorous analysis from patient advocacy groups, or variable in content.⁹

OSMT is a novel outpatient self-management educational and coaching program to assist Ostomates who have completed their cancer treatments to understand and adjust to their ostomies.⁶ The goals of OSMT are to reduce the post-discharge care deficit and assist with adaptation and long-term needs for cancer survivors with an ostomy.⁶ Based on the Chronic Care Model using planned, proven strategies, management, and patient activation,^{10, 11} self-management skills and strategies are essential components of OSMT, as are partnerships formed between patients and providers to empower patients so they can increasingly manage their own care. We drew upon extensive research on survivors with ostomies^{6, 9, 12–23} and the skills of our WOCNs^{7, 18, 19} to design the OSMT program to be participative with peer support from other ostomy survivors as well as spouses, partners, and family caregivers. Previous articles have described OSMT and demonstrated impacts on patient activation and self-efficacy to manage their ostomies, as well as improved HRQOL for Ostomates.^{6, 19}

SPECIFIC AIM

The specific aim of this analysis was to conduct an economic analysis of a community-based, ostomy nurse-led, peer-Ostomate assisted, small-group, behavior skills-training intervention designed to help Ostomates (e.g., urostomies, ileostomies, colostomies) to understand and adjust to their ostomies and to improve their health-related quality of life, as well as to assist their designated informal caregivers, where a “Peer-Ostomate” is an ostomy survivor of the same gender and similar age as the study subject with considerable experience in self-care.

METHODS

Human Subjects

All study procedures were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000, and were approved by the University of Arizona Human Subjects Protection Program (FWA00004218). The University of Arizona IRB approval number for this project is 0800000679R002. Written informed consent was obtained from each participant.

Setting

Our demonstration was based in an academic medical center—The University of Arizona College of Medicine. We assumed that the OSMT program is sponsored by a surgical practice that performs ostomy surgeries on cancer patients and that the surgeons provide continuing ostomy care to patients after they are discharged from their surgical inpatient stays.

Program Scale

Our implementation of OSMT was four cancer patients per intervention cohort, one cohort at a time. Our program formally covered both ostomy and cancer care. WOCNs' scopes of practice include care of many other types of surgical procedures, traumatic injuries, pressure sores, and causes of bladder and bowel incontinence. The study WOCNs addressed all concerns raised by ostomy patients and their family caregivers. They provided referral information for questions outside their knowledge scope.

Participants

Cancer survivors with ostomies were recruited from multiple sources and methods in the Tucson, AZ metropolitan area, including distribution of brochures, direct surgeon or ostomy nurse referral, email solicitations, and, most commonly, from the Tucson Ostomy Support Group peer group sessions. Our pilot demonstration included Ostomates (urostomies, colostomies) and their spouses/significant others/family caregivers. OSMT enrolled 35 patients. Time between ostomy surgery and baseline survey averaged 201 days, with a range of 22 to 1,626 days. Nine participants dropped out of the intervention before completing the full curriculum: one before Session 1, three before Session 2, and five before Session 4. Reasons for drop-out included death (2), illness (2), early ostomy reversal (1), travel distance (1), and no longer wishing to participate (3).²³

Program Faculty and Curriculum

Program faculty were two experienced ostomy nurses (MDC, NJT) who were trained in the program curriculum by a board-certified cancer surgeon (RSK). Following the principles of the Chronic Care Model as a guide^{10, 11}, a five-session program was designed and implemented. The WOCNs focused on training Ostomates to become problem solvers, rather than simply giving them a health professional's solution. For example, potential problem areas (skin problems, pouch selections) were identified by the WOCNs, and the survivor group was asked how they would approach solving these problems. Those who experienced the problem shared what actions they took with others in the group, demonstrating problem-solving strategies. Two Peer Ostomates, one male and one female, were trained in the intervention, attended each session, and worked with participants between formal sessions. The program curriculum is described in Table 1 and in more detail elsewhere.¹ One intervention session was devoted to family caregivers only to share their challenges and coping strategies among themselves, and the WOCNs could reinforce key coping skills and fill ostomy care knowledge gaps.

Perspective of this Economic Analysis

Our perspective was incremental resources required by an oncologic surgical practice to implement this WOCN/family-based Ostomate support program. We did not measure the direct and indirect economic costs incurred by Ostomates and their family caregivers to participate in this intervention other than Ostomates' health-related quality of life before, during, and after the intervention.²³

RESULTS

Labor Inputs

Three classes of labor inputs included cancer surgeon, WOCNs, and an administrative assistant (Table 2). The cancer surgeon was the medical provider who designed the OSMT rehabilitation program, trained two WOCNs to serve as the primary interventionists, trained the administrative assistant to support the intervention program, and facilitated the regular project team meetings. The WOCNs traded-off leading all five intervention sessions, but both served as co-leads for the session with spouses/significant others/family caregivers of the ostomy survivors. A part-time administrative assistant served as the communications hub for the intervention team and program participants, arranged for meeting space, ordered ostomy supplies and brochures, prepared meeting materials for each session, recruited patients for each cohort, scheduled patients into intervention groups, scheduled ostomy buddies for each session, oriented participants to the intervention by telephone prior to the first session, consented intervention participants, conducted follow-back telephone calls with participants who missed any intervention sessions to problem solve the reason(s) for absence and encourage them to attend the next session, entered data from surveys, and produced minutes of intervention team meetings.

The time input of the cancer surgeon included eight one-hour intervention team meetings per cohort of four ostomy patients—two meetings prior to the first group session to manage recruitment for the new series, five intervention team meetings to review each group session, and one meeting after the last group session to review the final status of each participant—for a total of 8 hours direct time per four-patient cohort.

Time input from the lead WOCN included: 0.5 hour pre-session preparation time; 1 hour/session face-to-face time with participants and ostomy buddies; 0.5 hour post-session check-in, and 0.5 hour round-trip travel time to/from meeting for a total of 2.5 hours per face-to-face intervention session. With five sessions, this equaled 12.5 hours per cohort plus eight one-hour weekly team meetings for a total of 20.5 hours per intervention cohort. The time input of a second WOCN was eight hours of intervention team meetings, and 2.5 hours (work + travel time) for the session with family members/informal caregivers, for a total of 10.5 hours per cohort.

Intervention team meetings reviewed the previous week's intervention group session and the status of each participant and family caregiver, and prepared for the next group session as well as individual feedback to patients and caregivers. Since these activities are participant- and session-specific, they must be repeated for each session and participant. These labor costs vary with the number of participants and number of sessions. We found that experienced WOCNs required only modest fixed training time before the start of the intervention sessions. Note that Table 1 does not contain any labor costs for pre-program training. Labor costs start with a pre-session staff meeting to discuss each patient cohort.

The administrative assistant worked 25% time for each four-patient cohort for eight work weeks performing the tasks identified above.

Non-Labor Inputs

Non-labor inputs and costs are described in Table 2. WOCNs were reimbursed for round-trip mileage to intervention sessions. Patients and gender-matched ostomy buddies arranged for their own transportation to and from group sessions. We produced a color descriptive brochure for the intervention program and printed copies to mail to participants. Postage costs were incurred for recruitment and orientation mailings. Ostomy supplies from various manufacturers were purchased and demonstrated during intervention sessions. Patients were instructed on differences among product brands with respect to price and quality and on how to cut the ostomy wafers to fit their own stomas. Some patients were accustomed to pre-cut wafers. We assumed that the intervention could be replicated by cancer surgeons and WOCNs, and, therefore, assumed that incremental costs were more appropriate for this analysis than fully-allocated direct and indirect costs of a surgical practice. We assumed that the surgical practice would already have telephone and office support, including a conference room where group sessions could be conducted. We also assumed that free public-use conference rooms were available in the community to reduce lengthy travel times for participants. All intervention staff had university e-mail accounts and personal smart phones.

Intervention Health-Related Quality of Life Outputs

Previously we showed that our self-management training program improved and sustained scores on multiple HRQOL outcome measures, including Patient Activation ($p=0.0004$), Self-Efficacy ($p=0.006$), Total HRQOL ($p=0.01$), Physical Well-being ($p=0.005$), and Social Well-being ($p=0.002$). Survivor Anxiety was significantly reduced by end of follow-up ($p=0.047$).²³

Total Costs

The total incremental cost of a complete four-session program with four ostomy patients was \$7,246, with a per-patient cost of \$1,812 (2014 dollars).

DISCUSSION

Program costs did not vary appreciably by attendance rates because of the largely fixed labor resources required to provide the intervention. Increasing the number of participants per session over four would likely weaken the power of the intervention; the WOCNs felt that four was the maximum number of dyads that could be effectively and equitably handled in a one-hour group session. In large cities, larger class sizes could be tested—up to five or six participants— but at the risk of reducing OSMT effect sizes. Medley conducted an incremental budget analysis under local reimbursement rules and demonstrated the cost-effectiveness of adding a WOCN to a joint inpatient-outpatient service in one hospital.²⁴

Clinical Implications

Cancers and treatment aftereffects may cause functional and emotional losses that could benefit from outpatient rehabilitation, distinct from routine physician-delivered primary care.² OSMT should be prescribed by cancer surgeons as a component of post-ostomy rehabilitation.²⁵ Participation should be based on patient needs, not recency of the ostomy

procedure. Local OSMT programs should be linked to national and local cancer support programs, such as the American Cancer Society (ACS) national program office and local ACS chapters.

Research Implications

Program costs were approached from a short-term provider perspective. Future research is needed on induced decreases and/or increases in medical care utilization and expenses caused by the intervention over time, such as fewer post-operative complications, lower incidence and severity of post-operative depression for patients and their family caregivers, higher levels of post-operative physical and social activities for both patients and family caregivers, fewer post-operative skin infections, reduced emergency room and outpatient visits for post-surgical complications, better management of post-operative diet and bowel function, and lower rates of hospital readmissions for post-surgical complications. Data needs include health insurance claims and payments, coinsurance and deductible payments by families, and uncovered out-of-pocket health care and long-term care expenses for intervention and observation-only comparison groups. Research is also needed to evaluate extending health insurance coverage for this WOCN-based intervention and the effects of out-of-pocket medical care outlays on families' economic assets.

A telehealth version of OSMT may improve inclusiveness and cost-effectiveness. A newly-funded project by the Patient-Centered Outcomes Research Institute, entitled "Ostomy Telehealth for Cancer Survivors" (PCOR grant #1507-31690, PI: Robert S. Krouse MD), is a multi-site randomized trial of a telehealth version of OSMT to test the feasibility and acceptability of a web-based health intervention for ostomy survivors. Future research should examine innovative models of care and support for Ostomates and their family caregivers, including an integrated face-to-face small-group plus telehealth model. Variations on intervention team composition and intervention intensity are also needed to better understand program cost and outcome trade-offs.

Study Limitations

Our participant numbers were small; we could not explore economies of scale or learnings from experience with greater numbers and variety of patients. Travel distance to meeting facility, lack of familiarity with our program, and concerns about coping with unfamiliar environments were oft-cited reasons for not accepting our recruitment invitations. The oncologic surgeon leading this study strongly valued this WOCN-based intervention program, which limits replicability to similarly-motivated surgeons. Cost-effectiveness or cost-utility analyses are required to determine OSMT's actual value for money.

CONCLUSIONS

Our demonstration sets the stage for larger studies in different types of communities utilizing innovative implementation methods responsive to wishes and needs of cancer survivors with ostomies. Because the OSMT intervention has been shown to benefit cancer survivors with ostomies²³, had reasonable program costs (\$1,812 per patient), we recommend consideration for coverage by health insurers. Based on qualitative input from

informal caregivers of our study participants, future studies should focus on improving our understanding of caregivers' responsibilities, stressors, and unmet resource needs for their OSMT support role.

Acknowledgments

The authors graciously acknowledge Octavio Bojorquez from Arizona Cancer Center for collecting the data on OSMT activities, resources, and costs.

FUNDING

This work was supported by grants from the National Cancer Institute at the National Institutes of Health: Grant number R21 CA133337 (University of Arizona, PI: Robert S. Krouse MD), and Arizona Cancer Center Support grant number 5P30CA023074-35 (University of Arizona, PI: Anne Cress MD). An unrestricted donation was received from the Sun Capital Partners Foundation. The contents of this work are solely the responsibility of the authors and do not necessarily represent the official views of the National Cancer Institute or the National Institutes of Health, or any other funders. The funders had no role in the design and conduct of the study, collection, management, analysis and interpretation of the data, preparation, review or approval of the manuscript.

References

- 1 Johnston D, Miller D, Frecea M, McKenzie M. Enhanced recovery after surgery and fecal diversions: development of a best practice guideline. *J Wound Ostomy Continence Nurs.* 2016; 43(3):227–228. [PubMed: 27163680]
- 2 Colwell JC, Kupsick PT, McNichol LL. Outcome criteria for discharging the patient with a new ostomy from home health care: a WOCN Society consensus conference. *J Wound Ostomy Continence Nurs.* 42(3):269–273.
- 3 Johnson T. Follow-up care of stoma patients: a systematic literature review. *Gastrointest Nurs.* 2012; 10(9):30–36.
- 4 Williams J. Healthy peristomal skin is essential. *Br J Nurs.* 2011; 20(22):1408. [PubMed: 22241485]
- 5 Beitz JM, Colwell JC. Ostomy care management approaches to stomal and peristomal complications: a narrative descriptive study. *J Wound Ostomy Continence Nurs.* 2016; 43(3):263–268. [PubMed: 26938334]
- 6 Grant M, McCorkle R, Hornbrook MC, et al. Development of a chronic care ostomy self-management program. *Journal of Cancer Education: The Official Journal of the American Association for Cancer Education.* 2013; 28:70–78. [PubMed: 23104143]
- 7 Tallman NJ, Cobb MD, Grant M, et al. Colostomy Irrigation: Issues Most Important to Wound, Ostomy and Continence Nurses. *J Wound Ostomy Continence Nurs.* 2015; 42:487–493. [PubMed: 26336046]
- 8 Krouse RS, Grant M, Wendel CS, et al. A mixed-methods evaluation of health-related quality of life for male veterans with and without intestinal stomas. *Dis Colon Rectum.* 2007; 50:2054–2066. [PubMed: 17701071]
- 9 Wasserman M, Baxter NN, Rosen B, et al. Systematic review of internet patient information on colorectal cancer surgery. *Dis Colon Rectum.* 2014; 57:64–69. [PubMed: 24316947]
- 10 Wagner EH, Austin BT, Davis C, et al. Improving chronic illness care: translating evidence into action. *Health Aff (Millwood).* 2001; 20:64–78. [PubMed: 11816692]
- 11 McCorkle R, Ercolano E, Lazenby M, Schulman-Green D, Schilling LS, Lorig K, Wagner EH. Self-management: Enabling and empowering patients living with cancer as a chronic illness. *CA: A Cancer Journal for Clinicians.* 2011; 61:50–62. DOI: 10.3322/caac.20093 [PubMed: 21205833]
- 12 Grant M, Ferrell B, Dean G, et al. Revision and psychometric testing of the City of Hope Quality of Life-Ostomy Questionnaire. *Qual Life Res.* 2004; 13:1445–1457. [PubMed: 15503840]
- 13 Krouse R, Grant M, Ferrell B, et al. Quality of life outcomes in 599 cancer and non-cancer patients with colostomies. *J Surg Res.* 2007; 138:79–87. [PubMed: 17196990]

- 14Krouse RS, Herrinton LJ, Grant M, et al. Health-related quality of life among long-term rectal cancer survivors with an ostomy: Manifestations by gender. *J Clin Oncol*. 2009; 27:4664–4670. [PubMed: 19720920]
- 15Sun V, Grant M, McMullen CK, et al. Surviving colorectal cancer: long-term, persistent ostomy-specific concerns and adaptations. *J Wound Ostomy Continence Nurs*. 2013; 40:61–72. [PubMed: 23222968]
- 16Krouse RS, Grant M, Rawl SM, et al. Coping and acceptance: the greatest challenge for veterans with intestinal stomas. *Journal of Psychosomatic Research*. 2009; 66:227–233. [PubMed: 19232235]
- 17McMullen CK, Hornbrook MC, Grant M, et al. The greatest challenges reported by long-term colorectal cancer survivors with stomas. *J Support Oncol*. 2008; 6:175–182. [PubMed: 18491686]
- 18Altschuler A, Ramirez M, Grant M, et al. The influence of husbands' or male partners' support on women's psychosocial adjustment to having an ostomy resulting from colorectal cancer. *J Wound Ostomy Continence Nurs*. 2009; 36:299–305. [PubMed: 19448512]
- 19McMullen CK, Hornbrook MC, Herrinton LJ, et al. Improving survivorship care for long-term colorectal cancer survivors: Key findings of a 5-year study. *Clinical Medicine and Research*. 2010; 1:32–33.
- 20Grant M, McMullen CK, Altschuler A, et al. Irrigation practices in long-term survivors of colorectal cancer with colostomies. *Clin J Oncol Nurs*. 2012; 16:514–519. [PubMed: 23022935]
- 21Grant M, McMullen CK, Altschuler A, et al. Dietary adjustments reported by colorectal cancer (CRC) survivors with permanent ostomies. *J Cancer Educ Supplement*. 2007:22.
- 22Cobb MD, Grant M, Tallman NJ, et al. Colostomy irrigation: current knowledge and practice of WOC nurses. *J Wound Ostomy Continence Nurs*. 2015; 42:65–70. [PubMed: 25325284]
- 23Krouse RS, Grant M, McCorkle R, Wendel CS, Cobb MD, Tallman NJ, Ercolano E, Sun V, Hibbard JH, Hornbrook MC. A chronic care ostomy self-management program for cancer survivors. *Psycho-Oncology*. 2016; 25(5):574–581. DOI: 10.1002/pon.4078 [PubMed: 26804708]
- 24Medley JA. Cost-effectiveness of a WOC Advanced Practice Nurse in the acute care and outpatient setting. *J Wound Ostomy Continence Nurs*. 2014; 41(4):307–310. [PubMed: 24918767]
- 25Pergolotti M, Williams GR, Campbell C, et al. Occupational therapy for adults with cancer: why it matters. *Oncologist*. 2016; 21:314–319. [PubMed: 26865588]

Appendix. Annual Salary Rates for OSMT Intervention Personnel

Annual Salary Rates for Intervention Personnel based on the University of Arizona salary and wage structure	Annual Salary ^a	Fringe Benefits @ 55% ^a	Total Annual Labor Cost ^a
Specialist MD annual salary + fringe benefits	\$260,000	\$130,000	\$390,000
WOCN annual salary + fringe benefits	\$83,200	\$41,600	\$124,800
Administrative Assistant annual salary + fringe benefits	\$52,000	\$26,000	\$78,000

^a2014 dollars

AUTHOR INFORMATION

Mark C. Hornbrook, PhD, Senior Investigator Emeritus, Kaiser Permanente Center for Health Research, 3800 North Interstate Avenue, Portland, OR 97227-1110, Tel: 503-780-9613, Fax: 503-335-2428, mark.c.hornbrook@gmail.com

Martha Cobb, MS, MEd, CWOCN, Wound, Ostomy and Continence Nurse (retired), University of Arizona College of Nursing, Tucson, AZ 85721, Tel: 520-290-6167, Fax: none, TwinHillsCobbs@aol.com

Nancy J. Tallman, BSN, WOCN, Wound, Ostomy, and Continence Nurse, Unaffiliated, Tucson, AZ 85737, Tel: 520-797-6903, Fax: none, nancyjtallman@comcast.net

Janice Colwell, RN, MS, CWOCN, FAAN, Advanced Practice Nurse, University of Chicago Medicine, 5841 S. Maryland Ave, MC 9081, Chicago, IL 60637, Tel: 773-702-9371, Fax: 773-834-1779, janice.colwell@uchospitals.edu

Ruth McCorkle, PhD, FAAN, Florence Wald Professor of Nursing and Professor of Chronic Diseases, Yale University School of Nursing and Yale Cancer Center, 100 Church Street South, Room 268, New Haven, CT 06535, Tel: 203-737-5501, Fax: 203-737-2414, ruth.mccorkle@yale.edu

Elizabeth Ercolano, DNSc, RN, Associate Research Scientist, Yale University School of Nursing, Yale West Campus, PO Box 27399, West Haven, CT 06516-0972, Ph: 203-737-2193, Elizabeth.ercolano@yale.edu

Marcia Grant, RN, PhD, FAAN, Distinguished Professor, Department of Nursing Research & Education, City of Hope National Medical Center, 1500 East Duarte Road, Duarte, CA 91010, Tel: (626) 256-4673 ext. 63483, Fax: (626) 301-8941, marciagrants13@gmail.com

Virginia Sun, RN, PhD, Assistant Professor, Division of Nursing Research & Education, Department of Population Sciences, Cancer Control and Population Sciences Program, City of Hope National Medical Center, 1500 East Duarte Road, Duarte, CA 91010, Tel: 626-256-4673 x63122, Fax: 626-301-8941, vsun@coh.org

Christopher S. Wendel, MS, Biostatistician, University of Arizona College of Medicine, Arizona Center on Aging, 1807 E. Elm St, Tucson, AZ 85719, Tel: 520-237-4624, Fax: None, cwendel@email.arizona.edu, tucvares@theriver.com

Judith H. Hibbard, DrPH, Professor Emerita, Department of Planning, Public Policy, & Management, Research Professor, Health Policy Research Group, University of Oregon, 119 Hendricks Hall, University of Oregon, Eugene, OR 97403-1209, Tel: 541 346-3364, Fax: 541 346-2040, jhibbard@uoregon.edu

Robert S. Krouse, MD, FACS, Chief, Surgical Service, CMC Veterans Affairs Medical Center, Instructor of Surgery, University of Pennsylvania School of Medicine, CMC VAMC Surgical Service (112), 3900 Woodland Ave., Philadelphia, PA 19104, Tel: (215) 823-5880, Fax: (215) 823-4309, robert.krouse@va.gov (CMC VAMC), robert.krouse@uphs.upenn.edu (University of Pennsylvania)

Table 1OSMT Intervention Session Syllabus^{a,b}

Session	Content
1	Self-management; Ostomates' immediate concerns(e.g., definitions, associated disease states); daily care; nutritional needs; impact on feelings, skin care and clothing changes. Interactive teaching methods with hands-on practice with equipment, pouches, and belts. Discussion of potential ostomy complications. Assignment: Monitor ostomy output (frequency and consistency) and problems with pouching system.
2	Discuss completed assignment of personal OSMT goals. Address social well-being, problems of social/interpersonal relationships, public appearances, being prepared for emergencies, intimacy and sexuality, and communication skills within the cultural framework of the individual participant and their family in home and social contexts. Fatigue as a model of long-term effects of cancer.
3	Family caregivers attend a separate session akin to Ostomates' Session 2, covering the same topics, but specifically tailored to support adjustment of CGs. Additional content as needed to ensure CG has achieved a comfort level with ostomy care. Focus for the CG customized to their cultural framework and home environment. Includes HRQOL assessment of CG and self-evaluation of their participation.
4	Discuss completed assignment. Promoting healthy lifestyles via nutritional management, physical activity, overcoming barriers, improving psychological health, and improving attitudes. Participants encouraged to set new priorities, evaluate friends, and work on changing negative attitudes. Traveling tips included. Discuss potential ostomy complications. Assignment: Review personal goals in regards to ostomy care.
5	Booster Intervention content driven by group demands and needs. CGs are invited to this session to help ensure a well-rounded understanding of issues, comfort with ostomy care, and anticipated future dilemmas. Discuss attainment of personal goals for participation in OSMT.

^aGrant M, McCorkle R, Hornbrook MC, Wendel CS, Krouse R. Development of a chronic care ostomy self-management program. *J Cancer Educ* 2013; **28**: 70–78.

^bKrouse RS, Grant M, McCorkle R, Wendel CS, Cobb MD, Tallman NJ, Ercolano E, Sun V, Hibbard JH, Hornbrook MC. A chronic care ostomy self-management program for cancer survivors.. *Psycho-Oncology*. 2016; **25**: 574–581. PMID:26804708, NIHMS 749541, DOI: 10.1002/pon.4078

Table 2

Incremental Resource Costs per OSMT Intervention Cohort

Personnel ^a	Tasks	Quantity and Cost of Resources per OSMT Intervention Cohort ^b	Total Cost per Cohort ^b	Per Patient Cost ^b
Medical Director/Cancer Surgeon	Provide medical supervision of WOCNs; provide supervision of Administrative Assistant; conduct 8 one-hour intervention team meetings with WOCN and Administrative Assistant; telephone consultation with WOCN as needed.	8 weekly intervention team meetings (1 hr./meeting × 8 meetings × \$125/hr × 155% fringe benefits) ^b	\$1,550	
WOCNs	Prepare and set-up for Ostomate intervention group meetings; lead intervention group sessions; conduct in-person individual counseling of patients immediately before and immediately after group sessions; provide PRN telephone counseling of patients and ostomy buddies between group sessions; maintain record of session attendance; attend regular meetings of intervention team; participate in Spouse/Significant Other/Family Caregiver Session	WOCN #1: 5 sessions × 2.5 hours/session (0.5 hr. pre-session preparation time + 1 hr. /session face-to-face time + 0.5 hrs. post-session check-in + 0.5 hr. round-trip travel time to/from meeting = 2.5 hrs.) = 12.5 hours, plus 8 hours of team meetings = 20.5 hours × \$48/hr. × 1.55 fringe benefits and overhead costs WOCN #2: 1 session × 2.5 hours = 2.5 hrs., plus 8 hours of team meetings = 10.5 hrs. × \$48/hr. × 1.55 fringe benefits and overhead costs	\$1,525 \$781	
Ostomate Peer Buddies	2 ostomy peer buddies for 4 sessions	0.5 hr. pre-meeting preparation time + 1 hr./session face-to-face time + 0.5 hr. post-session check-ins + 0.5 hr. round-trip travel time to meeting place = 2.5 hrs./session × 4 sessions × 2 buddies = 20 hrs./cohort × \$24/hr. × 1.55 fringe benefits ^b	\$744	
Administrative Assistant	Recruit patients into intervention; set up session times and places; inform participants and staff of meeting places and times; send out session reminders; order ostomy supplies and brochures; consent patients, contact no-shows to ascertain reason(s) for not attending; attend regular meetings of intervention team.	8 hrs./week × \$25/hr. × 8 work weeks × 1.55 fringe benefits ^b	\$2,480	
Non-Labor Inputs	Description	Quantity and Cost of Resources per Intervention Cohort	Total Cost per Cohort ^b	Per Patient Cost
Transportation	Auto mileage reimbursement for WOCN (2 WOCNs × 8 intervention staff meetings = 16 trips + 5 trips to group sessions = 21 trips)	WOC nurse mileage to/from meetings = 10 mi RT × 21 trips × \$0.55/mi	\$116	
Photocopying	Purchase & photocopy intervention materials for WOCNs and participants	Intervention brochures = 4 @ \$1 ea. = \$4 + \$30 for stoma equipment brochures = \$34	\$34	
Supplies	Ostomy supplies for teaching and demonstration	Stoma wafers, ostomy bags, skin protectant, water bottles, small duffle bag, disposable wipes, etc. —donated by supply companies	Donated	

Personnel ^a	Tasks	Quantity and Cost of Resources per OSMT Intervention Cohort ^b	Total Cost per Cohort ^b	Per Patient Cost ^b
Postage	Mailings to participants	Postage for mailings	\$16	
Telephone	Telephone service to participants and among staff members	Use surgeon's medical office telephone line + WOCN/Buddy personal cell phones	\$0	
Internet	Assume all intervention staff have an internet service provider as part of their household utility expenses	Use household Internet Service Provider	\$0	
E-mail	Assume all intervention staff have a personal internet appliance, e.g., smartphone, tablet, laptop, personal computer, etc.	Use university/hospital/household email service	\$0	
Meeting Room	Surgeon's medical office facility &/or free community meeting room space	Donated meeting space	\$0	
Total Cost/Cohort and Average Cost/Patient:			\$7,246	\$1,812

^a Intervention personnel were employed by the University of Arizona; labor costs included salaries and fringe benefits

^b 2014 dollars