Published in final edited form as:

Addict Behav. 2018 October ; 85: 107–112. doi:10.1016/j.addbeh.2018.05.024.

Little cigars and cigarillos: Affect and perceived relative harm among U.S. adults, 2015

Ban A. Majeed^{a,*}, Amy Nyman^b, Kymberle L. Sterling^c, and Paul Slovic^{d,e}

^aDivision of Epidemiology, Department of Population Health Sciences, Medical College of Georgia, Augusta University, Augusta, GA, USA

^bTobacco Center of Regulatory Science (TCORS), School of Public Health, Georgia State University, Atlanta, GA, USA

^cBattelle Memorial Institute, Atlanta, GA, USA

^dDecision Research, Eugene, OR, USA

^eDepartment of Psychology, University of Oregon, Eugene, OR, USA

Abstract

Introduction: Similar to cigarette smoking, consumption of cigars delivers nicotine and byproducts of tobacco combustion and elevates the risk of addiction, illness, and premature death. This study examined the relationship of affect, perceived relative harm, and LCC smoking behavior among U.S. adults.

Methods: Data were from Tobacco Products and Risk Perceptions Survey conducted in 2015. The study included a probability based sample of 6051 adults (18+) drawn from an online research panel. A current LCC smoker was defined as having ever smoked LCCs and was currently smoking LCCs every day, some days, or rarely. Participants were asked whether smoking LCCs was less harmful, had about the same level of harm, or was more harmful than smoking regular cigarettes. Feelings about LCCs were collected using word association technique. Descriptive and multinomial logistic regression analyses were conducted.

Results: About 7% of the study participants were current LCC smokers. Adults with positive feelings had four-fold the adjusted odds to be current LCC smokers. Perceiving LCCs to be less harmful had 2.7 higher adjusted odds of being current LCC smokers.

Conclusions: Compared to cigarettes, LCCs evoked more positive feelings among adults and these positive feelings were strongly associated with both perceiving LCCs as less harmful than cigarettes and with current LCC smoking. Cessation and prevention interventions would benefit

*Corresponding author at: 1120 15th street, AE-1036. Augusta, GA 3012-4900, USA., bmajeed@augusta.edu (B.A. Majeed).
Contributors

All authors were involved in the design of the study. Majeed conducted the statistical analysis and wrote the first draft of the manuscript and all authors contributed to and have approved the final manuscript.

Conflict of interest
None.

from applying the principles of social marketing in which information is provided not only to inform consumers but also to evoke negative feelings and associations with LCC smoking.

Keywords

Affect; Harm perception; Little cigars; Cigarillos; Feelings

1. Introduction

Similar to cigarette smoking, consumption of cigars—including large, premium cigars, cigarillos, as well as little filtered (cigarette-like) cigars—delivers nicotine and byproducts of tobacco combustion (e.g. carbon monoxide, nitrosamines, nitrogen oxide, and ammonia) and thus elevate the risk of addiction, premature death, and illness (Blank, Nasim, Hart Jr, & Eissenberg, 2011; Boffetta, Pershagen, Jockel, et al., 1999; Chang, Corey, Rostron, & Apelberg, 2015; Rickert, Trivedi, Momin, Wagstaff, & Lauterbach, 2011). Cigars vary in size, filter, tip, and characterizing flavors (Corey, King, Coleman, et al., 2014). In 2009, the Family Smoking Prevention and Tobacco Control Act (TCA) gave the Food and Drug Administration (FDA) the power to regulate cigarettes (Administration FaD, 2017). During the same year, the FDA prohibited the sales of cigarettes, *but not cigars*, that contain any artificial or natural flavors other than tobacco and menthol because flavors were strategically used to target youth and increase prevalence of smoking initiation (Administration FaD, 2017). Cigar characterizing flavors, particularly attractive to young people, are shown to increase the appeal of cigar smoking by masking the harshness and smell of tobacco (Delnevo, Giovenco, Ambrose, Corey, & Conway, 2015).

Little cigars and cigarillos (LCCs), often sold in singles or small packs, have been marketed as the less expensive and less harmful substitute for cigarettes (Malone, Yerger, & Pearson, 2001; Messer, White, Strong, et al., 2015; Sterling, Fryer, & Fagan, 2016). Although LCC smoking exposes smokers to nicotine and other toxicants, known to impair vascular endothelial function (Liu et al., 2016) and cause oral, esophageal, and lung cancer (Chang et al., 2015), LCC smokers tend to perceive them as less harmful and less addictive than cigarettes (Cohn, Cobb, Niaura, & Richardson, 2015). Perceptions about the potential harm and addictiveness of LCCs have been linked to intention of future behavior among adult cigarette smokers (Sterling, Majeed, Nyman, & Eriksen, 2017). In cigarette smoking research, the role of feelings on smoking initiation is well-documented (Slovic, 2012). Cigarette smoking often begins in adolescence, with beginning smokers acting in response to emotion-based media appeals and social pressures that supply immediate positive associations with smoking, but no information on the risks of illness or addiction caused by smoking (Slovic, 2012). The immediate feelings an individual uses to judge the level of risk is termed the *Affect Heuristic* (FM, Ali, Paul, & JS, 2000). The risk perception theory based on the *Affect Heuristic* posits that information on risk could reduce the emotional (affective) favorability of a given behavior and that information on benefit could increase the favorability, thus influencing feelings and perceptions of harm, and in turn may discourage or promote the behavior (FM et al., 2000). Cigarette ads are designed to exploit feelings and generate positive imagery associated with smoking which in turn reduces the perceived harm and promote smoking behavior (Slovic, 2001; Slovic, Peters, Finucane, & Macgregor, 2005).

In contrast with data on the relationship between affect, risk perception, and cigarette smoking, data on factors influencing perceptions of harm and the role of affect in shaping the beliefs about LCCs are limited. Therefore, based on the research on affect, risk, and decision making, we conducted this study on the perceptions of harm of LCCs relative to cigarettes and the impact of feelings on LCC smoking. The objectives of the current study were to explore perceptions of harm associated with LCC smoking relative to cigarette smoking, and examine the relationship of affect, perceived relative harm, and LCC smoking behavior among U.S. adults.

2. Methods

Data were from the Tobacco Products and Risk Perceptions Survey, an online cross-sectional survey conducted in August–September 2015. This annual survey was administered by the Tobacco Center of Regulatory Science at Georgia State University. The overall goal of the survey was to investigate the perception of multiple tobacco products (i.e. cigarettes, electronic cigarettes, hookah, little cigars and cigarillos) how the risk perception relates to the individual's decision to use tobacco products. The study used a probability sample drawn from an online research panel designed to be representative of the U.S. population, known as *KnowledgePanel*. Since 2009, address-based sampling (ABS) has been employed to recruit panelists. This sampling methodology covers about 97% of U.S. households including those with unlisted telephone numbers, with no landline telephones, and has no access to the internet or no device to access the internet. Currently, 55,000 adults aged 18 and older have joined *KnowledgePanel*.

Final stage survey completion rate was 76.0% ($N = 6051$ adults aged 18 years and older). After exclusion of respondents who were unaware of LCCs or had missing values on the awareness variable, the final analytical sample used in the current study was 5105 adults. The study was approved by the Institutional Review Board of Georgia State University.

2.1. Measures

2.1.1. Affect—Consistent with previous research on affect and decision making (Slovic, 2012), word association technique was used to elicit feelings toward LCCs. After word association, participants are typically asked to assign a degree of negativity or positivity to the associated word/image. In prior studies these associations have been shown to be predictive of both preferences and behavior (Slovic, 2012). Using an open-ended (text) question, the study participants who were aware of LCCs were asked to report the first thought or image that comes to mind when hearing the phrase “little cigars, cigarillos, or filtered cigars.” Data were collected on affect associated with the reported images or thoughts associated with LCCs using this question, “how do you feel about this thought or image?” Response options included very bad, somewhat bad, both good and bad, somewhat good, and very good, measured on five-point scale ranging from -2 to $+2$.

First image or thought and its associated affect were also elicited in response to the term “cigarette.” The response categories “very good” and “somewhat good” were combined into “good,” and “very bad” and “somewhat bad” were grouped and labeled “bad,” creating a

three-response category (good, neutral, bad) variable to represent affect (feelings) about the first thought or image associated with LCC smoking.

2.1.2. Relative harm perceptions of LCC smoking compared to cigarettes—

All those who were aware of LCCs were asked whether smoking LCCs was less harmful, had about the same level of harm, or was more harmful than smoking regular cigarettes. Participants could also select “I don’t know” in response to this question.

2.1.3. Perceived addictiveness of LCCs—One direct measure of perceived addictiveness of LCCs was used. Participants were asked whether people can become addicted to LCCs; response options were yes, no, and “I don’t know.”

2.1.4. LCC smoking status—Participants were grouped into three mutually exclusive groups: current, former, and never LCC smokers. A current LCC smoker was defined as having ever smoked LCCs and was currently smoking LCCs every day, some days, or rarely. Participants who reported ever smoking LCCs and responded *not at all* to the LCC use now question were categorized as former smokers. Never LCC smokers were participants who responded *no* when asked whether they have ever smoked LCCs, even one or two puffs (Agaku, King, Husten, et al., 2014). Study participants who had reported they were not aware of LCCs prior to this study were classified as never users.

2.1.5. Cigarette smoking status—To assess cigarette smoking status, we used the commonly used measure for defining current, former, and never smokers (Messer et al., 2015). Adults who reported smoking at least 100 cigarettes in their lifetime and were currently smoking every day or some days were categorized as current cigarette smokers. Adults who have smoked 100 cigarettes in their lifetime and responded *not at all* to the “smoke now” question were classified as former cigarette smokers. Never cigarette smokers were adults who reported not having smoked at least 100 cigarettes in their lifetime.

Demographic characteristics included in this study were sex, age, race/ethnicity, educational attainment, annual household income.

2.2. Statistical analysis

Analyses were conducted in January 2017 using Stata/MP 13.1 (StataCorp, College Station, TX) to assess the relationship of images and feelings to LCC smoking and perception of harm, and to compute weighted estimates representative of the U.S. adults. Survey specific weighting variables were used to account for the complex survey design and survey non-response. Weighted percentages overall and by LCC smoking status were estimated. To determine the characteristics associated with perceptions of relative harm of LCCs and with LCC smoking (current and former), we conducted multivariable multinomial logistic regression analyses. Adjusted odds ratio (AOR) and 95% confidence intervals (CI) were estimated. Significance level was set at $p < .05$.

3. Results

The sample included 66.1% white, non-Hispanic, 11.6% Black, non-Hispanic, and 51.4% females. More details on the demographic characteristics of the study sample is included in a previous publication (Majeed, Weaver, Gregory, et al., 2017). In 2015, among all adults, 7.1% (95% CI: 6.3, 8.0) were current LCC smokers, 25.2% (95% CI: 23.9, 26.6) were former LCC smokers, and 67.7% (95% CI: 66.2, 69.1) were never LCC smokers. Whereas 66.2% of the study participants rated the images associated with cigarettes as 'bad', 55.1% rated the images associated with LCCs as 'bad' (Fig. 1). Compared to proportions of participants who felt 'neutral' or 'good' about images associated with *cigarettes*, higher proportions assigned 'neutral' or 'good' feelings to images associated with LCCs.

The majority of the study participants (65.0%; 95% CI: 63.3, 66.7) thought LCCs were as harmful as cigarettes; 7.6% (95% CI: 6.8, 8.6) less harmful; 9.8% (95% CI: 8.7, 10.9) more harmful than cigarettes; and 17.6% (95% CI: 16.3, 19.0) said 'I don't know'. Table 1 shows that, compared with perceiving LCCs to be equally harmful as cigarettes, expressing *good or neutral* feelings about LCC images was associated with higher odds of the perception that LCC smoking was less harmful than cigarettes (Table 1). Adults who thought LCCs were *not addictive* had a 9.6 fold higher adjusted odds to think LCC were less harmful. Males, those who were 18–24 years, and who were current LCC smoker were more likely to perceive LCCs as less harmful than cigarettes. Compared to White adults, Black adults were less likely to perceive LCCs as less harmful than cigarettes. Compared to never cigarette smokers, current cigarette smokers had 60% reduction in the adjusted odds of perceiving LCCs as less harmful than cigarettes.

Adults who expressed *good or neutral* feelings about images associated with LCCs had four-folds the adjusted odds to be current LCC smokers than never LCC smokers (Table 2). Perceiving LCCs to be less harmful than cigarettes (compared to be equally harmful) had 2.7 higher adjusted odds of being current LCC smokers. Men were more likely than women to be current or former LCC smokers than to be never LCC smokers (Table 2). Self-identification as Non-Hispanic Black compared to Non-Hispanic White was associated with a 2.4-fold increase in the adjusted odds of being a current LCC smoker. Current cigarette smokers were more likely than never smokers to be current LCC smokers. No significant differences were observed in LCC smoking status by levels of education or household income.

4. Discussion

This study showed that compared to cigarettes, LCCs evoked more positive feelings (positive affect) among adults and that these positive feelings were strongly associated with both perceiving LCCs as less harmful than cigarettes and with current LCC smoking. Consistent with prior research (Sterling, Berg, Thomas, Glantz, & Ahluwalia, 2013), the current study revealed that current LCC smoking is more prevalent among African Americans, males, younger adults, and current cigarette smokers. Efforts to reduce the burden of premature death and disease caused by combustible tobacco products should target minority populations and young adults. In order to be effective, health communication

campaigns must be tailored to the characteristics of the population at risk (Schmid, Rivers, Latimer, & Salovey, 2008).

The finding that the majority of the study participants correctly perceived the harm of LCCs to be equal to that of cigarettes is encouraging. Similar to cigarette smoking, use of cigars (large, cigarillos, and little filtered cigars) may lead to premature mortality and morbidity. A systematic review of 22 studies on the effect of current exclusive cigar smoking among adults, documented elevated risk of developing oral, esophageal, laryngeal, pancreatic, and lung cancer, as well as coronary heart disease (CHD) and aortic aneurysm (Chang et al., 2015). Yet, some individuals are still misinformed or uninformed about the harm of LCCs. This study revealed that perceiving LCCs as not addictive and favorable feelings may contribute perceiving LCCs as less harmful than cigarettes.

Young adults and current LCC smokers are especially at risk of misperceiving the harm of LCC compared to cigarettes. Possible reasons for this relationship may be related to LCC product design (flavor and packaging), altering cigars (freaking), use as blunts and pattern of use. First) a recent review of 20 qualitative studies on the role of flavor in tobacco products on harm perception showed that flavored tobacco products, including electronic cigarettes, hookah, and LCCs, are believed to be less harmful than cigarettes (Kowitz et al., 2017). A study among young adult LCC smokers revealed that flavored LCCs were perceived as less harmful than cigarettes because of attractive flavors (Sterling, Fryer, & Fagan, 2016) and package design, particularly the use of images of fruit on the package (Sterling, Fryer, Pagano, & Fagan, 2016). Second) some LCC users, specifically smokers of Black and Mild, modify their cigars before smoking by removing the filter paper (inner binder), a.k.a. 'cancer paper.' This modification, known as 'freaking' (Nasim et al., 2014), 'hying' was believed to reduce the harm of cigar smoking (Jolly, 2008). Third) another reason for the perception of reduced harm could be due to the use of cigars as blunts to smoke marijuana, substance perceived as less harmful than tobacco (Sterling, Fryer, & Fagan, 2016). Lastly, amount smoked and inhalation style (perceived 'no inhalation') of cigar smoking is different than that of cigarette smoking, thus, some users may perceive cigars as less harmful. Health education campaigns should provide accurate information regarding the addictive nature of nicotine in LCCs, as well as correct misperceptions surrounding the use of unaltered and altered LCCs.

In May 2016, the Food and Drug Administration (FDA) extended its regulatory authority over products that meet the definition of tobacco products, including large cigars, cigarillos, and little cigars (Food and Drug Administration H, 2016). Under this rule, the FDA requires the display of health warnings on LCC packages and cigar advertisements (Food and Drug Administration H, 2016). The required warning statements for use on LCC packages include: addiction or nicotine statement, risk of mouth and throat cancer, risk of lung cancer, not being a safe alternative to cigarettes, and reproductive harm (Food and Drug Administration H, 2016). Our study provides baseline information on the perceived relative harm and perceived addictiveness of LCCs among U.S. adults that may change over time in response to the implementation of new regulations. Continuous monitoring of LCC smoking as well as the perceptions of harm and addictiveness among U.S. adults, not only among

current smokers, is required to assess the potential impact of future regulatory and communication efforts.

Consistent with research on cigarette smoking, the current study provides evidence that implicit attitudes (affect) toward LCCs influence both perception of harm and LCC smoking behavior (FM et al., 2000; Marks Anthony, O'Neill, & Hine Donald, 2011; Slovic, 2001). The role of both cognitive and affective (feeling) components of harm perception in cigarette smoking has been documented (Marks Anthony et al., 2011). For example, repeated exposure to cigarette ads was shown to induce positive feelings for smoking and reduce the individual's perceived risk (Slovic, 2001). Given that feelings are subject to change, employing a similar strategy could prove effective to curb the rise in LCC smoking. LCC smoking cessation and prevention interventions would benefit from applying the principles of social marketing in which information is provided not only to inform consumers but also to evoke negative feelings and associations with LCC smoking.

In the current study, the estimated prevalence of current LCC smoking among U.S. adults (7.1%) was higher than that documented in previous studies, suggesting a rise in the prevalence of LCC smoking. Data from the 2012–2013 National Adult Tobacco Survey revealed that 5.8% of U.S. adults reported smoking at least 50 cigars, cigarillos, or filtered little cigars during their lifetime and now smoked “every day” or “some days” or “rarely (Agaku et al., 2014)”. To guide and evaluate tobacco control interventions, continuous monitoring of cigarette and non-cigarette tobacco product use (Corey et al., 2014), is critical, using consistent measures to allow for comparability across multiple studies. Consistent with previous research (Corral, Landrine, Simms, & Bess, 2013; Ehlke & Cohn, 2016; Loukas, Batanova, Fernandez, & Agarwal, 2015; Milam, Bone, Byron, et al., 2013; Nyman, Sterling, Weaver, Majeed, & Eriksen, 2016), the current study showed that those who are Black, non-Hispanic, young adults, males, and current cigarette smokers are at higher risk of LCC smoking and in turn higher risk for tobacco related diseases. Flavors (Kostygina, Glantz, & Ling, 2016; Sterling, Fryer, Nix, & Fagan, 2015), affordable prices (Nyman et al., 2016), and targeted advertisement may explain the high uptake of LCCs among these subpopulations. Cantrell, et al. documented the high availability of LCCs, aggressive exterior advertisement, and lower prices of LCCs in African American and young adult predominant neighborhoods (Cantrell, Kreslake, Ganz, et al., 2013; Ribisl, D'Angelo, Feld, et al., 2017). Furthermore, this study confirms that perception of reduced harm compared to cigarettes is positively related to LCC smoking (Sterling et al., 2017; Sterling, Fryer, Pagano, Jones, & Fagan, 2016). Harm perceptions and personal characteristics can inform and guide future LCC-related research, policy, and interventions. For example, the impact of health communications and tobacco counter-marketing efforts tailored to the audience characteristics is maximized (Schmid et al., 2008).

5. Limitations

Use of an online research panel (*KnowledgePanel*) to draw the study sample may raise concerns about generalizability of the results to the U.S. adults. However, *KnowledgePanel* is a probability-based and designed to be representative of non-institutionalized USA adults. Given the cross-sectional nature of the survey, it cannot be determined whether the

perception of reduced relative harm and associating LCC thoughts and images with positive feelings preceded or followed LCC smoking behavior. Thus, causal inferences are not warranted. Longitudinal or experimental study designs would be useful to remove concerns about causality. The survey focused on eliciting perceptions in relation to LCCs used with tobacco and did not include blunts, thus, the findings may not reflect LCCs use with other products than tobacco. Finally, the study relied on self-reporting of LCC smoking and thus may raise issues of response bias.

6. Conclusions

The study suggests that favorable affect plays an important role in LCC smoking behavior and in shaping harm perceptions. The results document that current LCC smoking is more prevalent among African Americans, males, younger adults, and current cigarette smokers. The findings have implications for policy, research, and public health interventions. Enhanced surveillance efforts tracking the prevalence of use and perceptions of LCCs could benefit evaluation of new implemented LCC-related policy. Experimental research on strategies to evoke negative feelings about LCCs is necessary to inform health communication efforts. Cessation services should be provided to populations at higher risk of LCC smoking, namely, African Americans and young adults. Targeted health communication efforts to correct information and create negative feelings toward LCCs could prove effective in reducing the burden of LCC smoking among vulnerable populations.

Acknowledgment

Funding for this study was provided by Grant Number P50DA036128 from the National Institutes of Health, National Institute of Drug Abuse (NIH/NIDA) and Food and Drug Administration (P50DA036128), Center for Tobacco Products (FDA CTP). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or the Food and Drug Administration. FDA and NIH had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

References

- Administration FaD (2017). *Tobacco Control Act*. <https://www.fda.gov/TobaccoProducts/Labeling/RulesRegulationsGuidance/ucm246129.htm>, Accessed date: July 2017.
- Agaku IT, King BA, Husten CG. Tobacco product use among adults—United States, 2012–2013 MMWR. Morbidity and Mortality Weekly Report. 2014; 63(25):542–547. [PubMed: 24964880]
- Blank MD, Nasim A, Hart A Jr., Eissenberg T. Acute effects of cigarillo smoking Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco. 2011; 13(9):874–879. [PubMed: 21504888]
- Boffetta P, Pershagen G, Jockel KH. Cigar and pipe smoking and lung cancer risk: A multicenter study from Europe Journal of the National Cancer Institute. 1999; 91(8):697–701. [PubMed: 10218507]
- Cantrell J, Kreslake JM, Ganz O. Marketing little cigars and cigarillos: Advertising, price, and associations with neighborhood demographics American Journal of Public Health. 2013; 103(10): 1902–1909. [PubMed: 23948008]
- Chang CM, Corey CG, Rostron BL, Apelberg BJ. Systematic review of cigar smoking and all cause and smoking related mortality BMC Public Health. 2015; 15:390. [PubMed: 25907101]
- Cohn A, Cobb CO, Niaura RS, Richardson A. The other combustible products: Prevalence and correlates of little cigar/cigarillo use among cigarette smokers Nicotine & Tobacco Research:

- Official Journal of the Society for Research on Nicotine and Tobacco. 2015; 17(12):1473–1481. [PubMed: 25634932]
- Corey CG, King BA, Coleman BN. Little filtered cigar, cigarillo, and premium cigar smoking among adults—United States, 2012–2013 MMWR. Morbidity and Mortality Weekly Report. 2014; 63(30): 650–654. [PubMed: 25078654]
- Corral I, Landrine H, Simms DA, Bess JJ. Polytabacco use and multiple-product smoking among a random community sample of African-American adults BMJ Open. 2013; 3(12):e003606.
- Delnevo CD, Giovenco DP, Ambrose BK, Corey CG, Conway KP. Preference for flavoured cigar brands among youth, young adults and adults in the USA Tobacco Control. 2015; 24(4):389–394. [PubMed: 24721967]
- Ehlke SJ, Cohn AM. Was it the drink? The conditioned association of alcohol and desire to quit smoking on the dual use of little cigars/cigarillos and cigarettes among men and women Addictive Behaviors. 2016; 59:48–51. [PubMed: 27070096]
- FM L, Ali A, Paul S, JS M. The affect heuristic in judgments of risks and benefits Journal of Behavioral Decision Making. 2000; 13(1):1–17.
- Food and Drug Administration H. Deeming tobacco products to be subject to the Federal Food, drug, and cosmetic act, as amended by the family smoking prevention and tobacco control act; restrictions on the sale and distribution of tobacco products and required warning statements for tobacco products. Final rule Federal Register. 2016; 81:28973–29106. [PubMed: 27192730]
- Jolly DH. Exploring the use of little cigars by students at a historically black university Preventing Chronic Disease. 2008; 5(3):A82. [PubMed: 18558032]
- Kostygina G, Glantz SA, Ling PM. Tobacco industry use of flavours to recruit new users of little cigars and cigarillos Tobacco Control. 2016 Jan; 25(1):66–74. DOI: 10.1136/tobaccocontrol-2014-051830 [PubMed: 25354674]
- Kowitt SD, Meernik C, Baker HM, Osman A, Huang LL, Goldstein AO. Perceptions and experiences with flavored non-menthol tobacco products: A systematic review of qualitative studies International Journal of Environmental Research and Public Health. 2017; 14(4)
- Liu J, Wang X, Narayan S, Glantz SA, Schick SF, Springer ML. Impairment of endothelial function by little cigar secondhand smoke Tobacco Regulatory Science. 2016; 2(1):56–63. [PubMed: 26753171]
- Loukas A, Batanova M, Fernandez A, Agarwal D. Changes in use of cigarettes and non-cigarette alternative products among college students Addictive Behaviors. 2015; 49:46–51. [PubMed: 26046401]
- Majeed BA, Weaver SR, Gregory KR. Changing perceptions of harm of e-cigarettes among U.S. adults, 2012–2015 American Journal of Preventive Medicine. 2017; 52(3):331–338. [PubMed: 28341303]
- Malone RE, Yerger V, Pearson C. Cigar risk perceptions in focus groups of urban African American youth Journal of Substance Abuse. 2001; 13(4):549–561. [PubMed: 11775082]
- Marks Anthony DG, O'Neill G, Hine Donald W. Role of affect, expectancies and dual processes of cognition in predicting adult cigarette smoking Australian Journal of Psychology. 2011; 60(3): 160–167.
- Messer K, White MM, Strong DR. Trends in use of little cigars or cigarillos and cigarettes among U.S. smokers, 2002–2011 Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco. 2015; 17(5):515–523. [PubMed: 25239955]
- Milam AJ, Bone LR, Byron MJ. Cigarillo use among high-risk urban young adults Journal of Health Care for the Poor and Underserved. 2013; 24(4):1657–1665. [PubMed: 24185161]
- Nasim A, Blank MD, Cobb CO, Berry BM, Kennedy MG, Eissenberg T. How to freak a Black & Mild: A multi-study analysis of YouTube videos illustrating cigar product modification Health Education Research. 2014; 29(1):41–57. [PubMed: 24162926]
- Nyman AL, Sterling KL, Weaver SR, Majeed BA, Eriksen MP. Little cigars and cigarillos: Users, perceptions, and reasons for use Tobacco Regulatory Science. 2016; 2(3):239–251. [PubMed: 27413772]

- Ribisl KM, D'Angelo H, Feld AL. Disparities in tobacco marketing and product availability at the point of sale: Results of a national study *Preventive Medicine*. 2017; 105:381–388. [PubMed: 28392252]
- Rickert WS, Trivedi AH, Momin RA, Wagstaff WG, Lauterbach JH. Mutagenic, cytotoxic, and genotoxic properties of tobacco smoke produced by cigarillos available on the Canadian market *Regulatory Toxicology and Pharmacology: RTP*. 2011; 61(2):199–209. [PubMed: 21821091]
- Schmid KL, Rivers SE, Latimer AE, Salovey P. Targeting or tailoring? Maximizing resources to create effective health communications *Marketing Health Services*. 2008; 28(1):32–37.
- Slovic P, (2001). Cigarette smokers: Rational actors or rational fools? In Slovic P. (Ed.). *Smoking: Risk, perception, & policy* (pp. 97–124). Thousand Oaks, CA, US: Sage Publication .
- Slovic P. The 'value' of smoking: An editorial *Health, Risk & Society*. 2012; 14(5):409–414.
- Slovic P, Peters E, Finucane ML, Macgregor DG. Affect, risk, and decision making *Health Psychology: Official Journal of the Division of Health Psychology American Psychological Association*. 2005; 24(4s):S35–S40.
- Sterling K, Berg CJ, Thomas AN, Glantz SA, Ahluwalia JS. Factors associated with small cigar use among college students *American Journal of Health Behavior*. 2013; 37(3):325–333. [PubMed: 23985179]
- Sterling K, Fryer C, Pagano I, Jones D, Fagan P. Association between menthol-flavoured cigarette smoking and flavoured little cigar and cigarillo use among African-American, Hispanic, and white young and middle-aged adult smokers *Tobacco Control*. 2016; 25(Suppl. 2):ii21–ii31. [PubMed: 27856997]
- Sterling KL, Fryer CS, Fagan P. The most natural tobacco used: A qualitative investigation of young adult Smokers' risk perceptions of flavored little cigars and cigarillos *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*. 2016; 18(5):827–833. [PubMed: 26175458]
- Sterling KL, Fryer CS, Nix M, Fagan P. Appeal and impact of characterizing flavors on young adult small cigar use *Tobacco Regulatory Science*. 2015; 1:42–53. [PubMed: 27917396]
- Sterling KL, Fryer CS, Pagano I, Fagan P. Little cigars and cigarillos use among young adult cigarette smokers in the United States: Understanding risk of concomitant use subtypes *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*. 2016; 18(12): 2234–2242. [PubMed: 27613889]
- Sterling KL, Majeed BA, Nyman A, Eriksen M. Risk perceptions of little cigar and cigarillo smoking among adult current cigarette smokers *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*. 2017; 19(11):1351–1358. [PubMed: 27659275]

HIGHLIGHTS

- Participants were more likely to have favorable feeling about LCCs than cigarettes.
- Those who thought LCCs were not addictive also thought they were less harmful.
- Adults who had favorable feelings about LCCs were more likely to smoke them.
- Favorable affect has important role in LCC smoking and in harm perceptions.

Fig. 1.

Feelings about images associated with “little cigars, cigarillos, or filtered cigars” and “cigarette” among U.S. adults, 2015.

Table 1

Characteristics associated with perceptions of relative harm of little cigars and cigarillos among U.S. adults, 2015.

Participant characteristics	Perceptions of relative harm ^a		
	Less harmful	More harmful	"I don't know"
	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
Feelings about images associated with LCCs			
Good or neutral	1.8 ** (1.3, 2.4)	0.6 ** (0.5, 0.9)	1.0 (0.8, 1.2)
Bad	1.0	1.0	1.0
Perceived addictiveness of LCCs			
Not addictive	9.6 ** (5.5, 17.0)	2.6 * (1.2, 5.6)	2.9 ** (1.4, 6.3)
Yes addictive	1.0	1.0	1.0
"I don't know"	3.9 ** (2.6, 5.9)	1.8 * (1.1, 3.1)	7.9 ** (6.1, 10.4)
Sex			
Male	1.5 * (1.1, 2.0)	1.1 (0.8, 1.5)	1.0 (0.8, 1.2)
Female	1.0	1.0	1.0
Age (years)			
18–24	2.4 ** (1.3, 4.3)	1.7 (0.8, 3.4)	0.7 (0.4, 1.2)
25–34	1.1 (0.7, 1.7)	1.7 * (1.1, 2.5)	0.7 (0.5, 1.0)
35–44	0.9 (0.5, 1.4)	1.2 (0.7, 2.0)	0.7 (0.5, 1.0)
45–54	1.2 (0.8, 1.9)	1.1 (0.7, 1.7)	0.7 (0.5, 1.0)
55–64	1.3 (0.9, 2.1)	1.2 (0.8, 1.8)	0.9 (0.7, 1.3)
65 +	1.0	1.0	1.0
Race/ethnicity			
White, NH	1.0	1.0	1.0
Black, NH	0.4 ** (0.2, 0.7)	1.1 (0.7, 1.7)	1.6 * (1.1, 2.2)
Other, NH	0.8 (0.3, 1.8)	1.9 * (1.1, 3.4)	2.2 ** (1.2, 3.8)
Hispanic	0.8 (0.5, 1.2)	0.8 (0.6, 1.3)	1.2 (0.9, 1.7)
Education			
< High school	1.0	1.0	1.0
High school	0.6 (0.3, 1.2)	0.8 (0.4, 1.6)	0.7 (0.5, 1.1)
Some college	0.8 (0.4, 1.5)	1.3 (0.7, 2.6)	0.6 * (0.4, 0.9)
College degree +	0.8 (0.4, 1.7)	1.1 (0.5, 2.2)	0.7 (0.4, 1.1)
Household income			
\$24,999 or less	1.0	1.0	1.0
\$25,000–\$84,999	1.2 (0.7, 1.9)	0.7 (0.5, 1.0)	1.1 (0.8, 1.5)
\$85,000 or more	1.1 (0.7, 1.9)	0.6 ** (0.4, 0.9)	0.9 (0.6, 1.3)
Cigarette smoking status			
Current smoker	0.4 ** (0.3, 0.7)	1.7 ** (1.1, 2.5)	1.2 (0.9, 1.7)
Former smoker	0.5 ** (0.4, 0.8)	1.4 (1.0, 2.1)	0.9 (0.7, 1.1)

Participant characteristics	Perceptions of relative harm ^a		
	Less harmful	More harmful	"I don't know"
	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
Never smoker	1.0	1.0	1.0
LCC smoking status			
Current smoker	1.8[*] (1.1, 3.0)	1.4 (0.9, 2.3)	0.5[*] (0.3, 0.9)
Former smoker	1.3 (1.0, 1.9)	1.1 (0.8, 1.4)	0.7^{**} (0.5, 0.9)
Never smoker	1.0	1.0	1.0

Bold face indicates statistical significance. All variables appear in the table were included in the model. NH: Non-Hispanic.

^{**}
p < .01.

^{*}
p < .05.

^a About the same harm was used as the reference group in the multinomial logistic regression mode.

Table 2

Characteristics of smokers of little cigars and cigarillos among U.S. adults, 2015.

Participant characteristics	LCC smoking status	
	Current LCC smoker AOR (95% CI)	Former LCC smoker AOR (95% CI)
Feelings about images associated with LCCs		
Good or neutral	4.0 ** (2.8, 5.7)	1.6 ** (1.4, 2.0)
Bad	1.0	1.0
Perceived relative harm of LCCs		
Less harmful	2.7 ** (1.7, 4.4)	1.4 (1.0, 2.0)
About the same	1.0	1.0
More harmful	1.6 (1.0, 2.6)	1.1 (0.8, 1.5)
"I don't know"	0.6 * (0.4, 0.9)	0.7 ** (0.6, 0.9)
Sex		
Male	5.6 ** (3.9, 7.9)	3.3 ** (2.8, 4.0)
Female	1.0	1.0
Age (years)		
18–24	8.9 ** (4.6, 17.3)	1.3 (0.8, 2.1)
25–34	8.3 ** (4.8, 14.3)	2.4 ** (1.8, 3.2)
35–44	5.6 ** (3.3, 9.6)	1.1 (0.8, 1.5)
45–54	3.7 ** (2.1, 6.3)	1.1 (0.9, 1.5)
55–64	2.2 ** (1.3, 3.8)	1.1 (0.9, 1.5)
65 +	1.0	1.0
Race/ethnicity		
White, NH	1.0	1.0
Black, NH	2.4 ** (1.5, 3.9)	0.8 (0.6, 1.2)
Other, NH	1.2 (0.6, 2.3)	0.8 (0.5, 1.3)
Hispanic	1.4 (0.9, 2.2)	1.0 (0.7, 1.3)
Education		
< High school	1.0	1.0
High school	0.9 (0.5, 1.7)	1.1 (0.7, 1.8)
Some college	1.3 (0.6, 2.5)	1.6 (1.0, 2.6)
College degree +	1.0 (0.5, 2.0)	1.7 (1.0, 2.7)
Household income		
\$24,999 or less		
\$25,000–\$84,999	0.6 (0.4, 1.0)	1.0 (0.8, 1.4)
\$85,000 or more	0.6 (0.4, 1.0)	1.0 (0.8, 1.4)
Cigarette smoking status		
Current smoker	14.5 ** (9.4, 22.3)	6.3 ** (4.8, 8.3)
Former smoker	3.1 ** (2.0, 4.9)	5.7 ** (4.6, 7.0)

Participant characteristics	LCC smoking status	
	Current LCC smoker	Former LCC smoker
	AOR (95% CI)	AOR (95% CI)
Never smoker	1.0	1.0

Bold face indicates statistical significance. Note, *never smoker of LCCs* was the reference group in the multinomial logistic regression. NH: Non-Hispanic.

p < .01.

p < .05.