Published in final edited form as:

J Nucl Cardiol. 2018 August ; 25(4): 1148–1155. doi:10.1007/s12350-017-1158-7.

Molecular Imaging of Calcific Aortic Valve Disease

Jae-Joon Jung, PhD, Farid Jadbabaie, MD, and Mehran M. Sadeghi, MD

Cardiovascular Molecular Imaging Laboratory, Section of Cardiovascular Medicine and Yale Cardiovascular Research Center, Yale School of Medicine, New Haven, CT; and Veterans Affairs Connecticut Healthcare System, West Haven, CT

Summary

Calcific aortic valve disease (CAVD) can progress to symptomatic aortic stenosis in a subset of patients. The severity of aortic stenosis and the extent of valvular calcification can be evaluated readily by echocardiography, CT, and MRI using well-established imaging protocols. However, these techniques fail to address optimally other important aspects of CAVD, including the propensity for disease progression, risk of complications in asymptomatic patients, and the effect of therapeutic interventions on valvular biology. These gaps may be addressed by molecular imaging targeted at key biological processes such as inflammation, remodeling, and calcification that mediate the development and progression of CAVD. In this review, recent advances in valvular molecular imaging, including ^{18}F -fluorodeoxyglucose (FDG) and ^{18}F -sodium fluoride (NaF) PET, and matrix metalloproteinase-targeted SPECT imaging in the preclinical and clinical settings are presented and discussed.

Keywords

Aortic stenosis; Aortic valve; Inflammation; Calcification; Atherosclerosis; PET; SPECT; Matrix metalloproteinases; FDG; Sodium Fluoride

Calcific aortic valve disease (CAVD) is the most common cause of aortic stenosis (AS), which according to recent US statistics affects ~0.4% of the general population¹. The prevalence of AS increases with age with moderate to severe AS affecting nearly 3% of patients 75 years old¹. In addition, 17,000 deaths and 55,000 hospitalizations are annually attributable to aortic valve disease (not just AS). The Natural history of CAVD involves a long asymptomatic period during which the initial fibrotic thickening of aortic valve leaflets with limited calcification (aortic sclerosis) progresses to more extensive valvular calcification and ultimately, hemodynamically significant aortic stenosis. The resulting pressure overload promotes left ventricular hypertrophy, an adaptive response that underlies the typically long period of asymptomatic disease present despite gradual worsening of aortic stenosis. The rise in wall stress may lead to impairment of coronary perfusion and development of sub-endocardial ischemia, which in turn triggers cardiomyocyte apoptosis and myocardial fibrosis. It is postulated that eventually, ventricular decompensation triggers angina, heart failure, syncope, or sudden death. While symptomatic AS is indeed the

hallmark of severe CAVD, not all patients with moderate AS progress to symptomatic disease. Even in patients with severe AS (peak systolic velocity ≥ 4 m/s by Doppler echocardiography), ~30% do not develop symptoms over a 5 year period². On the other hand, a non-negligible minority of such asymptomatic patients die of sudden cardiac death², highlighting the complex nature of the relation between aortic valve structure and physiology (including the severity of AS), ventricular response, and symptoms.

In addition to age, the number of leaflets is another major risk factor for the development of CAVD. Indeed, while bicuspid aortic valve (BAV) is found in ~1.4% of live births, BAV accounts for a large fraction of aortic valve surgeries³⁻⁵. Reflecting the similarity between CAVD and atherosclerosis, male sex, smoking, hypertension, diabetes, hypercholesterolemia, and elevated Lp(a) are additional risk factors for the development of CAVD⁴. Many of the factors that increase the risk of CAVD development, are also risk factors for CAVD progression. In particular, disease progression is accelerated with the severity of stenosis, higher extent of calcification, older age and presence of BAV⁴. Moreover, aortic valve area and left ventricular hypertrophy are independent predictors of the development of symptoms². Importantly, the presence of CAVD, even in the absence of severe disease or symptoms, portends a higher risk of all-cause mortality^{2, 6, 7}. The predictors of all-cause mortality in aortic stenosis include age, chronic renal failure, inactivity, aortic valve velocity², left ventricle mass⁸, and left ventricle wall stress⁹.

Currently, there is no known treatment to slow down the progression of CAVD, and surgical or transcatheter aortic valve replacement remain the only effective therapeutic options for advanced, symptomatic disease. Despite considerable overlap between the risk factors for CAVD and atherosclerosis and promising results of observational studies¹⁰, randomized clinical trials of statins have failed to demonstrate benefit in slowing down the progression of aortic valve calcification or stenosis^{11, 12}. Inadequate duration of statin therapy and late stage of the disease, as well as the pro-calcific effects of statins may have contributed to this failure. In addition to statins, several other approaches are currently under investigation to prevent CAVD progression¹³. Examples include Niacin [NCT02109614, to reduce Lp(a) levels], denosumab or alendronic acid (NCT02132026, to target bone metabolism), tadalafil (NCT01275339, to inhibit phosphodiesterase Type 5), and ataciguat (NCT02481258, to promote guanylate cyclase activity). Irrespective of the drug tested, considerable heterogeneity of CAVD progression makes it difficult to demonstrate therapeutic effectiveness in clinical trials. For instance in the Simvastatin and Ezetimibe in Aortic Stenosis (SEAS) trial, the mean ($\pm$ SD) change in peak aortic-jet velocity during a median follow-up of 52.2 months was 0.62 ± 0.61 meters per second in the placebo group¹². It is reasonable to assume that by targeting therapeutic interventions to those patients who are at the highest risk for progression, it is easier to establish any potential therapeutic efficacy.

The initial diagnosis and evaluation of CAVD and AS is typically based on clinical findings and echocardiography. In addition, a subset of patients (e.g., those with inconclusive echocardiography results) may require invasive evaluation of the valve. However, classical non-invasive imaging techniques, echocardiography, CT and MRI, which focus on aortic valve anatomy and physiology (e.g., calcification, area, flow rates and transvalvular gradient) and associated aortic and myocardial abnormalities (focal and diffuse myocardial

fibrosis, global and basilar longitudinal strain) inadequately inform of the patient progression risk and prognosis. This risk is directly related to valvular (and myocardial) pathobiology, which can be appraised by molecular imaging. As such, in vivo assessment of valvular biology by molecular imaging can potentially help clinical decision making regarding the need for aortic valve replacement in patients with moderate AS who are to undergo coronary artery bypass grafting, and possibly in patients with severe asymptomatic AS, who might be at increased risk for sudden cardiac death. Selection of patients who might benefit from emerging medical therapies and assessment of the response to therapeutic interventions within a relatively short period of time are additional applications of such techniques for drug development. Last, but certainly not least, molecular imaging can help address important gaps in CAVD pathophysiology through in vivo assessment of aortic valve biology and serial imaging in the same subject.

Pathobiology

CAVD was once presumed to be a degenerative disease associated with passive calcium deposition in leaflets. Over the past two decades, advances in our understanding of aortic valve pathobiology have led to a paradigm shift of our view of the disease to an active process of valvular thickening and calcification starting with asymptomatic, hemodynamically insignificant aortic sclerosis that progresses to aortic stenosis¹⁴.

Aortic valve leaflets are covered by valvular endothelial cells (VECs) and are composed of three layers, namely the collagen-rich fibrosa on the aortic side, the proteoglycan-rich spongiosa, and the elastin-rich ventricularis,¹⁵. The valve-specific, fibroblast-like valvular interstitial cells (VICs) are interspersed within these layers and play a central role in fibrocalcific remodeling of the leaflets¹⁶. Environmental factors, biomechanical perturbations and genetic predisposition contribute to the leaflet remodeling process, which culminates in the development of hemodynamically significant aortic stenosis.

VEC activation induced by changes in shear stress or other stimuli, e.g., atherosclerotic risk factors, initiates the development of CAVD by promoting subendothelial lipid retention and oxidation, and recruitment of inflammatory cells to the leaflet^{17, 18}. Inflammation, VIC transformation, angiogenesis, extracellular matrix remodeling and calcification are pathologic hallmarks of CAVD. Infiltrated macrophages and T cells may trigger tissue remodeling by promoting oxidative stress, release of proteases, e.g., matrix metalloproteinases and cathepsins, and differentiation of quiescent VICs into myofibroblasts. This promotes fibrotic remodeling of the valve. In parallel, these cells may undergo osteogenic differentiation, resulting in the formation of calcific nodules, a process which starts at the base of the fibrosa layer^{19–21 22}. Ultimately, the thickening, fibrosis and calcification of the leaflets leads to hemodynamically significant aortic stenosis in a subset of patients. Of note, while the immune-inflammatory (oxidative stress-driven) pathway underlies most cases of CAVD, an alternative hyperphosphatemia-driven pathway of calcification appears to dominate in patients with chronic kidney disease. These issues are discussed in detail in elsewhere^{14, 23}.

Molecular Imaging

a) Preclinical studies

The small size of aortic valve and motion are challenges to in vivo imaging. Accordingly, a number of studies have relied on ex vivo imaging modalities [near-infrared fluorescence (NIRF) imaging, and magnetic resonance imaging and spectroscopy (MRI/MRS)] to image various aspects of valvular pathobiology in rodent models of CAVD, including endothelial cell activation, proteolytic activity, phagocytic activity, osteogenic activity and angiogenesis^{24–26}. While valuable as complement to conventional molecular and valvular biology techniques, the inherent limitations of these techniques preclude their application in longitudinal studies and ultimately, clinical implementation. This is in sharp contrast with emerging nuclear imaging-based approaches, which can be readily applied to non-invasive imaging in humans.

MMPs play a key role in extracellular matrix remodeling. Several members of MMP family, MMP-1, -9, and -12, are amongst the most highly up-regulated genes in CAVD²⁷. Inflammatory cells are major sources of MMP production and activity. Accordingly, MMP activation and inflammation are closely intertwined in CAVD. The feasibility and correlates of MMP-targeted imaging was recently shown in apolipoprotein E^{-/-} mice fed a Western diet for up to 9 months to induce CAVD. In this model, a subset of animals develops hemodynamically significant aortic stenosis. In vivo microSPECT-CT imaging using RP805, a ^{99m}Tc-labeled tracer that binds to several activated MMPs, demonstrated specific in vivo aortic valve uptake of the tracer (Figure 1)²⁸. Interestingly, the MMP signal peaked at 6 months of Western diet, while key features of CAVD (leaflet thickening, valvular calcification, and aortic stenosis) were most pronounced after 9 months. This suggests that MMPs are implicated in CAVD progression and MMP imaging may serve as a predictive tool in CAVD. A significant correlation between valvular macrophage staining (CD68) and MMP signal in vivo suggested that this technique may be used for detection of valvular inflammation and remodeling. Whether this technique can be used for imaging CAVD in the absence of atherosclerotic lesions (which co-exist in the aforementioned model) or tracking the response to therapeutic interventions remains to be determined.

b) Clinical studies

FDG PET—Despite some concern regarding its specificity, ¹⁸F-FDG PET has been extensively used for imaging vascular inflammation and has served as a non-invasive tool for early assessment of the effect of novel therapeutic interventions in clinical trials. Indeed, several studies have shown a close relationship between the ¹⁸F-FDG signal and macrophage content of atherosclerotic plaques^{29–33}. The potential role of ¹⁸F-FDG PET in aortic valve disease was first investigated in a retrospective analysis of whole body ¹⁸F-FDG PET/CT images acquired for oncological staging³⁴. In this study, FDG uptake [expressed as target-to-blood ratio (TBR)] was significantly higher in patients with aortic stenosis (n=42) than in the matched control group (Figure 2). Reminiscent of MMP imaging data in murine CAVD, when categorized based on severity of stenosis and calcification, ¹⁸F-FDG uptake was significantly increased in patients with mild and moderate disease compared to control patients, but not in those with severe disease.

Another retrospective study evaluated the relation between aortic valve ^{18}F -FDG signal in 111 patients without active cancer or aortic stenosis who underwent at least 2 ^{18}F -FDG PET/CT studies within a period of 1 to 5 years³⁵. When categorized as non-progressors or progressors based on aortic valve calcium score determined at baseline and follow-up time points, the ^{18}F -FDG signal (expressed as SUVmax) in aortic valve was found to be higher in the progressor group compared to non-progressor group. A similar difference was seen in the subset of subjects without aortic valve calcification at baseline. Interestingly, in this study the ^{18}F -FDG signal could independently predict subsequent calcification after adjusting for cardiovascular risk factors.

In a prospective study of ^{18}F -FDG PET (and ^{18}F -NaF) imaging involving 121 subjects (including 20 controls), the aortic valve ^{18}F -FDG signal (maximum TBR) was significantly higher in patients with aortic stenosis than in controls and the uptake modestly increased with the severity of stenosis³⁶. A subset of the subjects of the original cohort (n=30) was followed for one year, during which 12 subjects underwent aortic valve replacement (AVR) for symptomatic AS. In the remaining subjects without severe AS, there was no correlation between the baseline ^{18}F -FDG signal and 1 year-change in calcium score³⁷. In those subjects who underwent valve replacement, there was no correlation between the ^{18}F -FDG signal in vivo and CD68 staining of the surgical specimens. A follow up evaluation of the full cohort of the subjects (including the controls) at two years showed weak correlations between the baseline ^{18}F -FDG signal and aortic calcification (CT) and stenosis (echocardiography). In addition, the baseline ^{18}F -FDG signal was able to predict clinical outcomes (composite of cardiovascular death and aortic valve replacement) independently of age and sex³⁸.

While the non-linear relation between ^{18}F -FDG signal in aortic valve and the severity of aortic stenosis is intriguing, existing data raise many questions about the biological and clinical significance of the ^{18}F -FDG signal in the valve. In addition, uptake in the myocardium is a practical barrier to broader implementation of aortic valve ^{18}F -FDG PET imaging, indicating that alternative, more robust techniques are needed for evaluation of CAVD in humans.

NaF PET— ^{18}F -NaF is retained at the sites of calcification through binding to hydroxyapatite via an exchange mechanism with hydroxyl groups³⁹. However, the biological basis of ^{18}F -NaF uptake in aortic valve is not well defined. Studies in human atherosclerotic lesions have linked the NaF signal in atherosclerosis to the sites of calcification. As such, in carotid endarterectomy specimens NaF uptake ex vivo detected by autoradiography corresponded to the areas of calcification (Alizarin red staining), but not CD68 (macrophages), CD31 (endothelial cells) or alpha-smooth muscle actin (smooth muscle cells) expression⁴⁰. The higher ^{18}F -NaF retention at the sites of microcalcification as opposed to larger masses of calcified material has been attributed to the relatively larger surface area of the former available for ^{18}F -NaF binding.

The potential of ^{18}F -NaF PET for imaging valvular biology in CAVD was brought up by retrospective analysis of aortic valve signal in ^{18}F -NaF PET-CT studies of a small group of subjects (n=5) with cancer or rheumatological disease and concomitant aortic stenosis (and

control subjects without aortic valve calcification), which showed significantly higher ^{18}F -NaF (both maximal SUVs and TBRs) in those with aortic stenosis⁴¹. The aforementioned prospective study of 121 subjects with aortic sclerosis or stenosis of different severity or control subjects without apparent aortic valve disease who underwent both ^{18}F -FDG and ^{18}F -NaF PET/CT shortly followed this initial observation and demonstrated higher ^{18}F -NaF signal in aortic valve of patients with CAVD than controls with the intensity of the signal increasing with the severity of the aortic stenosis³⁶. Of note, while 91% of patients with aortic stenosis had increased ^{18}F -NaF uptake, only 35% showed increased ^{18}F -FDG uptake. The poor correlation between aortic valve signal from these tracers indicated that they target different aspects of aortic valve biology. The one-year follow-up study of a subset of the original cohort (n=30, including 12 subjects who had undergone surgical AVR and 18 with asymptomatic disease) reported a statistically significant correlation between the baseline ^{18}F -NaF signal and changes in calcium score. A similar correlation was found between the baseline and subsequent changes in calcium score³⁷. Histological analysis of the aortic valve from those subjects who underwent AVR during this one-year period showed an association (albeit modest) between the baseline ^{18}F -NaF signal and markers of calcification (alkaline phosphatase and osteocalcin), but not inflammation (CD68). Importantly, areas of osteocalcin expression in the leaflet extended beyond the area of Von Kossa staining (established calcium) and corresponded to the area with maximal ex vivo ^{18}F -NaF uptake on autoradiography. The two-year follow up evaluation of this cohort reported the development of new sites of macrocalcification corresponding to the sites of baseline ^{18}F -NaF signal, and a good correlation ($r^2=0.64$) between the baseline ^{18}F -NaF signal and subsequent changes in calcium score (Figure 3). However, the relationship with echocardiographic measures of hemodynamic progression was much weaker, highlighting the role of other factors (fibrosis, distribution of calcification) in aortic stenosis³⁸. Consistent with the observations at 1-year follow up, ^{18}F -NaF PET-CT was not an independent predictor of clinical outcomes at 2 years after correction for baseline CT-based calcium scores.

Perspective and conclusions

The utility of ^{18}F -NaF PET as a marker of disease activity in CAVD is currently under evaluation in several clinical trials [NCT02132026 (SALTIRE2), NCT02740088, NCT03095313]. While potentially useful as a tool to track the effect of therapeutic interventions on valvular calcification, ^{18}F -NaF PET imaging probably misses the fibrotic component of the disease and additional molecular imaging tools are necessary to fully evaluate different aspects of this pathology. The size of aortic valve and its continuous motion are major hurdles to imaging quantitative assessment of valvular biology by molecular imaging. In the case of ^{18}F -FDG, despite the emergence of several protocols aimed at suppressing myocardial uptake of the tracer, consistent, complete suppression remains an additional challenge. Recent advances in image acquisition and quantification methodology have improved the image quality and reproducibility of aortic valve ^{18}F -NaF PET images, and more progress is expected to lead to more reliable data for clinical studies^{42, 43}. While PET is a powerful tool for clinical imaging, the spatial resolution of microPET is a challenge to imaging in the mouse and microSPECT is more suitable in this setting. Preclinical studies of CAVD are also hampered by the paucity of relevant animal

models, and in this regard, emergence of non-atherosclerotic models of the disease with leaflet calcification (similar to human disease) would be of great value. Furthermore, despite many similarities between valvular and vascular calcification, there are also key differences between the two processes that should not be overlooked⁴⁴. Nevertheless, a number of new tracers that are under evaluation for imaging atherosclerosis may be of value in CAVD⁴⁵. The limitations of ¹⁸F-FDG and ¹⁸F-NaF PET imaging highlight the need for such new approaches to detect key aspects of CAVD pathophysiology.

In conclusion, molecular imaging may address some of the existing gaps in CAVD management. Accordingly, there is a great need for non-invasive, highly specific molecular imaging tools with real potential for clinical translation. As only a subset of patients with aortic sclerosis progress to advanced CAVD, detection of early molecular and functional abnormalities in aortic valve disease that predict the future risk for disease progression can help identify subjects who would most benefit from novel therapeutic approaches to prevent aortic valve stenosis. In addition to its clinical relevance, the development of an imaging approach for tracking valvular biology in CAVD will help advance our understanding of pathophysiology and potentially lead to novel medical therapies for this prevalent disease.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Sources

This work was supported by grants from NIH (R01-HL114703, R01-HL138567), and Department of Veterans Affairs (10-BX001750).

References

1. Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J, Fornage M, Gillespie C, Isasi CR, Jimenez MC, Jordan LC, Judd SE, Lackland D, Lichtman JH, Lisabeth L, Liu S, Longenecker CT, Mackey RH, Matsushita K, Mozaffarian D, Mussolino ME, Nasir K, Neumar RW, Palaniappan L, Pandey DK, Thiagarajan RR, Reeves MJ, Ritchey M, Rodriguez CJ, Roth GA, Rosamond WD, Sasson C, Towfighi A, Tsao CW, Turner MB, Virani SS, Voeks JH, Willey JZ, Wilkins JT, Wu JH, Alger HM, Wong SS, Muntner P. Heart Disease and Stroke Statistics-2017 Update: A Report From the American Heart Association. *Circulation*. 2017
2. Pellikka PA, Sarano ME, Nishimura RA, Malouf JF, Bailey KR, Scott CG, Barnes ME, Tajik AJ. Outcome of 622 adults with asymptomatic, hemodynamically significant aortic stenosis during prolonged follow-up. *Circulation*. 2005; 111:3290–5. [PubMed: 15956131]
3. Hoffman JI, Kaplan S. The incidence of congenital heart disease. *J Am Coll Cardiol*. 2002; 39:1890–900. [PubMed: 12084585]
4. Lindman BR, Clavel MA, Mathieu P, Iung B, Lancellotti P, Otto CM, Pibarot P. Calcific aortic stenosis. *Nat Rev Dis Primers*. 2016; 2:16006. [PubMed: 27188578]
5. Masri A, Svensson LG, Griffin BP, Desai MY. Contemporary natural history of bicuspid aortic valve disease: a systematic review. *Heart*. 2017; 103:1323–1330. [PubMed: 28490615]
6. Otto CM, Lind BK, Kitzman DW, Gersh BJ, Siscovick DS. Association of aortic-valve sclerosis with cardiovascular mortality and morbidity in the elderly. *N Engl J Med*. 1999; 341:142–7. [PubMed: 10403851]

7. Rosenhek R, Kloor U, Schemper M, Scholten C, Heger M, Gabriel H, Binder T, Maurer G, Baumgartner H. Mild and moderate aortic stenosis. Natural history and risk stratification by echocardiography. *Eur Heart J*. 2004; 25:199–205. [PubMed: 14972419]
8. Gerds E, Rossebø AB, Pedersen TR, Cioffi G, Lønnebakken MT, Cramariuc D, Rogge BP, Devereux RB. Relation of Left Ventricular Mass to Prognosis in Initially Asymptomatic Mild to Moderate Aortic Valve Stenosis. *Circulation: Cardiovascular Imaging*. 2015; 8:e003644. [PubMed: 26489804]
9. Ng ACT, Prihadi EA, Antoni ML, Bertini M, Ewe SH, Ajmone Marsan N, Leung DY, Delgado V, Bax JJ. Left ventricular global longitudinal strain is predictive of all-cause mortality independent of aortic stenosis severity and ejection fraction. *Eur Heart J Cardiovasc Imaging*. 2017
10. Bellamy MF, Pellikka PA, Klarich KW, Tajik AJ, Enriquez-Sarano M. Association of cholesterol levels, hydroxymethylglutaryl coenzyme-A reductase inhibitor treatment, and progression of aortic stenosis in the community. *J Am Coll Cardiol*. 2002; 40:1723–30. [PubMed: 12446053]
11. Cowell SJ, Newby DE, Prescott RJ, Bloomfield P, Reid J, Northridge DB, Boon NA. A randomized trial of intensive lipid-lowering therapy in calcific aortic stenosis. *N Engl J Med*. 2005; 352:2389–97. [PubMed: 15944423]
12. Rossebø AB, Pedersen TR, Boman K, Brudi P, Chambers JB, Egstrup K, Gerds E, Gohlke-Barwolf C, Holme I, Kesaniemi YA, Malbecq W, Nienaber CA, Ray S, Skjaerpe T, Wachtell K, Willenheimer R, Investigators S. Intensive lipid lowering with simvastatin and ezetimibe in aortic stenosis. *N Engl J Med*. 2008; 359:1343–56. [PubMed: 18765433]
13. Marquis-Gravel G, Redfors B, Leon MB, Gnreux P. Medical Treatment of Aortic Stenosis. *Circulation*. 2016; 134:1766. [PubMed: 27895025]
14. Yutzy KE, Demer LL, Body SC, Huggins GS, Towler DA, Giachelli CM, Hofmann-Bowman MA, Mortlock DP, Rogers MB, Sadeghi MM, Aikawa E. Calcific aortic valve disease: a consensus summary from the Alliance of Investigators on Calcific Aortic Valve Disease. *Arterioscler Thromb Vasc Biol*. 2014; 34:2387–93. [PubMed: 25189570]
15. Hinton RB Jr, Lincoln J, Deutsch GH, Osinska H, Manning PB, Benson DW, Yutzy KE. Extracellular matrix remodeling and organization in developing and diseased aortic valves. *Circ Res*. 2006; 98:1431–8. [PubMed: 16645142]
16. Rajamannan NM, Evans FJ, Aikawa E, Grande-Allen KJ, Demer LL, Heistad DD, Simmons CA, Masters KS, Mathieu P, O'Brien KD, Schoen FJ, Towler DA, Yoganathan AP, Otto CM. Calcific aortic valve disease: not simply a degenerative process: A review and agenda for research from the National Heart and Lung and Blood Institute Aortic Stenosis Working Group. Executive summary: Calcific aortic valve disease-2011 update. *Circulation*. 2011; 124:1783–91. [PubMed: 22007101]
17. Butcher JT, Nerem RM. Valvular endothelial cells and the mechanoregulation of valvular pathology. *Philosophical transactions of the Royal Society of London Series B, Biological sciences*. 2007; 362:1445–57. [PubMed: 17569641]
18. Suckosky P, Balachandran K, Elhammali A, Jo H, Yoganathan AP. Altered shear stress stimulates upregulation of endothelial VCAM-1 and ICAM-1 in a BMP-4- and TGF-beta1-dependent pathway. *Arteriosclerosis, thrombosis, and vascular biology*. 2009; 29:254–60.
19. New SE, Aikawa E. Molecular imaging insights into early inflammatory stages of arterial and aortic valve calcification. *Circulation research*. 2011; 108:1381–91. [PubMed: 21617135]
20. Chester AH, Taylor PM. Molecular and functional characteristics of heart-valve interstitial cells. *Philosophical transactions of the Royal Society of London Series B, Biological sciences*. 2007; 362:1437–43. [PubMed: 17569642]
21. Hinz B, Phan SH, Thannickal VJ, Galli A, Bochaton-Piallat ML, Gabbiani G. The myofibroblast: one function, multiple origins. *The American journal of pathology*. 2007; 170:1807–16. [PubMed: 17525249]
22. Mohler ER 3rd, Gannon F, Reynolds C, Zimmerman R, Keane MG, Kaplan FS. Bone formation and inflammation in cardiac valves. *Circulation*. 2001; 103:1522–8. [PubMed: 11257079]
23. Aikawa E, Libby P. A Rock and a Hard Place. *Circulation*. 2017; 135:1951. [PubMed: 28507250]
24. Aikawa E, Nahrendorf M, Sosnovik D, Lok VM, Jaffer FA, Aikawa M, Weissleder R. Multimodality molecular imaging identifies proteolytic and osteogenic activities in early aortic valve disease. *Circulation*. 2007; 115:377–86. [PubMed: 17224478]

25. Waters EA, Chen J, Allen JS, Zhang H, Lanza GM, Wickline SA. Detection and quantification of angiogenesis in experimental valve disease with integrin-targeted nanoparticles and 19-fluorine MRI/MRS. *J Cardiovasc Magn Reson*. 2008; 10:43. [PubMed: 18817557]
26. Aikawa E, Aikawa M, Libby P, Figueiredo JL, Rusanescu G, Iwamoto Y, Fukuda D, Kohler RH, Shi GP, Jaffer FA, Weissleder R. Arterial and aortic valve calcification abolished by elastolytic cathepsin S deficiency in chronic renal disease. *Circulation*. 2009; 119:1785–94. [PubMed: 19307473]
27. Bosse Y, Miqdad A, Fournier D, Pepin A, Pibarot P, Mathieu P. Refining molecular pathways leading to calcific aortic valve stenosis by studying gene expression profile of normal and calcified stenotic human aortic valves. *Circ Cardiovasc Genet*. 2009; 2:489–98. [PubMed: 20031625]
28. Jung JJ, Razavian M, Challa AA, Nie L, Golestani R, Zhang J, Ye Y, Russell KS, Robinson SP, Heistad DD, Sadeghi MM. Multimodality and molecular imaging of matrix metalloproteinase activation in calcific aortic valve disease. *J Nucl Med*. 2015; 56:933–8. [PubMed: 25908827]
29. Font MA, Fernandez A, Carvajal A, Gamez C, Badimon L, Slevin M, Krupinski J. Imaging of early inflammation in low-to-moderate carotid stenosis by 18-FDG-PET. *Frontiers in bioscience*. 2009; 14:3352–60.
30. Hyafil F, Cornily JC, Rudd JH, Machac J, Feldman LJ, Fayad ZA. Quantification of inflammation within rabbit atherosclerotic plaques using the macrophage-specific CT contrast agent N1177: a comparison with 18F-FDG PET/CT and histology. *J Nucl Med*. 2009; 50:959–65. [PubMed: 19443582]
31. Rudd JH, Narula J, Strauss HW, Virmani R, Machac J, Klimas M, Tahara N, Fuster V, Warburton EA, Fayad ZA, Tawakol AA. Imaging atherosclerotic plaque inflammation by fluorodeoxyglucose with positron emission tomography: ready for prime time? *J Am Coll Cardiol*. 2010; 55:2527–35. [PubMed: 20513592]
32. Tawakol A, Migrino RQ, Hoffmann U, Abbata S, Houser S, Gewirtz H, Muller JE, Brady TJ, Fischman AJ. Noninvasive in vivo measurement of vascular inflammation with F-18 fluorodeoxyglucose positron emission tomography. *J Nucl Cardiol*. 2005; 12:294–301. [PubMed: 15944534]
33. Zhang Z, Machac J, Helft G, Worthley SG, Tang C, Zaman AG, Rodriguez OJ, Buchsbaum MS, Fuster V, Badimon JJ. Non-invasive imaging of atherosclerotic plaque macrophage in a rabbit model with F-18 FDG PET: a histopathological correlation. *BMC nuclear medicine*. 2006; 6:3. [PubMed: 16725052]
34. Marincheva-Savcheva G, Subramanian S, Qadir S, Figueroa A, Truong Q, Vijayakumar J, Brady TJ, Hoffmann U, Tawakol A. Imaging of the aortic valve using fluorodeoxyglucose positron emission tomography increased valvular fluorodeoxyglucose uptake in aortic stenosis. *J Am Coll Cardiol*. 2011; 57:2507–15. [PubMed: 21679852]
35. Abdelbaky A, Corsini E, Figueroa AL, Subramanian S, Fontanez S, Emami H, Hoffmann U, Narula J, Tawakol A. Early aortic valve inflammation precedes calcification: a longitudinal FDG-PET/CT study. *Atherosclerosis*. 2015; 238:165–72. [PubMed: 25525744]
36. Dweck MR, Jones C, Joshi NV, Fletcher AM, Richardson H, White A, Marsden M, Pessotto R, Clark JC, Wallace WA, Salter DM, McKillop G, van Beek EJ, Boon NA, Rudd JH, Newby DE. Assessment of valvular calcification and inflammation by positron emission tomography in patients with aortic stenosis. *Circulation*. 2012; 125:76–86. [PubMed: 22090163]
37. Dweck MR, Jenkins WS, Vesey AT, Pringle MA, Chin CW, Malley TS, Cowie WJ, Tsampasian V, Richardson H, Fletcher A, Wallace WA, Pessotto R, van Beek EJ, Boon NA, Rudd JH, Newby DE. 18F-sodium fluoride uptake is a marker of active calcification and disease progression in patients with aortic stenosis. *Circ Cardiovasc Imaging*. 2014; 7:371–8. [PubMed: 24508669]
38. Jenkins WS, Vesey AT, Shah AS, Pawade TA, Chin CW, White AC, Fletcher A, Carlidge TR, Mitchell AJ, Pringle MA, Brown OS, Pessotto R, McKillop G, Van Beek EJ, Boon NA, Rudd JH, Newby DE, Dweck MR. Valvular (18)F-Fluoride and (18)F-Fluorodeoxyglucose Uptake Predict Disease Progression and Clinical Outcome in Patients With Aortic Stenosis. *J Am Coll Cardiol*. 2015; 66:1200–1. [PubMed: 26338001]
39. Blau M, Ganatra R, Bender MA. 18 F-fluoride for bone imaging. *Seminars in nuclear medicine*. 1972; 2:31–7. [PubMed: 5059349]

40. Irkle A, Vesey AT, Lewis DY, Skepper JN, Bird JLE, Dweck MR, Joshi FR, Gallagher FA, Warburton EA, Bennett MR, Brindle KM, Newby DE, Rudd JH, Davenport AP. Identifying active vascular microcalcification by 18F-sodium fluoride positron emission tomography. *Nature communications*. 2015; 6:7495.
41. Hyafil F, Messika-Zeitoun D, Burg S, Rouzet F, Benali K, Iung B, Vahanian A, Le Guludec D. Detection of 18fluoride sodium accumulation by positron emission tomography in calcified stenotic aortic valves. *Am J Cardiol*. 2012; 109:1194–6. [PubMed: 22264594]
42. Pawade TA, Cartlidge TR, Jenkins WS, Adamson PD, Robson P, Lucatelli C, Van Beek EJ, Prendergast B, Denison AR, Forsyth L, Rudd JH, Fayad ZA, Fletcher A, Tuck S, Newby DE, Dweck MR. Optimization and Reproducibility of Aortic Valve 18F-Fluoride Positron Emission Tomography in Patients With Aortic Stenosis. *Circ Cardiovasc Imaging*. 2016; 9
43. Doris MK, Rubeaux M, Pawade T, Otaki Y, Xie Y, Li D, Tamarappoo BK, Newby D, Berman DS, Dweck MR, Slomka PJ, Dey D. Motion-corrected imaging of the aortic valve with 18F-NaF PET/CT and PET/MR: a feasibility study. *J Nucl Med*. 2017
44. Weiss RM, Miller JD, Heistad DD. Fibrocalcific aortic valve disease: opportunity to understand disease mechanisms using mouse models. *Circ Res*. 2013; 113:209–22. [PubMed: 23833295]
45. Tavakoli S, Vashist A, Sadeghi MM. Molecular imaging of plaque vulnerability. *J Nucl Cardiol*. 2014; 21:1112–28. quiz 1129. [PubMed: 25124827]

Figure 1.

In vivo matrix metalloproteinase-targeted imaging in apolipoprotein E^{-/-} mouse fed on Western diet for up to 9 months to induce CAVD. Top panels: Examples of transverse contrast-enhanced CT, ^{99m}Tc-RP805 (matrix metalloproteinase-targeted) SPECT, and fused images in a control, wild-type mouse, and an apolipoprotein E^{-/-} mouse fed on Western diet for 6 months. Bottom panel: Background-corrected quantification of aortic valve ^{99m}Tc-RP805 uptake in control animals and apolipoprotein E^{-/-} mice fed on Western diet for up to 9 months. Arrows point to aortic valve area. cpv = counts per voxel. **P* < 0.05, ***P* < 0.01, ****P* < 0.0001, *n* = 7–13 in each group. This research was originally published in Journal of Nuclear Medicine (Jung et al.²⁸, © by the Society of Nuclear Medicine and Molecular Imaging, Inc.).

Figure 2. ^{18}F -FDG imaging of CAVD. Examples of CT images (A) and fused ^{18}F -FDG-PET/CT images (B) of aortic valve in patients with different stages of CAVD. (Arrows point to ^{18}F -FDG uptake in aortic valve. AS: aortic stenosis. Adapted with permission from Elsevier and Copyright Clearance Center, Journal of the American College of Cardiology,³⁴ © 2011.

Figure 3. ^{18}F -NaF imaging and progression of valvular calcification in CAVD. Baseline CT images (left) and fused ^{18}F -NaF-PET/CT images (middle), and 2 years follow up CT images (right) in two patients with CAVD. Adapted with permission from Elsevier and Copyright Clearance Center, Journal of the American College of Cardiology,³⁸ © 2015.