Published in final edited form as:

Annu Rev Virol. 2015 November ; 2(1): 525–548. doi:10.1146/annurev-virology-100114-055204.

Viruses in Rodent Colonies: Lessons Learned from Murine Norovirus

Stephanie M. Karst¹ and Christiane E. Wobus^{2,*}

¹Department of Molecular Genetics and Microbiology, University of Florida, Gainesville, FL, 32610, USA

²Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI, 48109, USA

Abstract

Noroviruses (NoVs) are highly prevalent, positive-sense RNA viruses that infect a range of mammals, including humans and mice. Murine noroviruses (MuNoVs) are the most prevalent pathogens in biomedical research colonies, and they have been used extensively as a model system for human noroviruses (HuNoVs). Despite recent successes in culturing HuNoV in the laboratory and a small animal host, studies of human viruses have inherent limitations. Thus, the versatility of the MuNoV system with its native host, reverse genetics and cell culture systems will continue to provide important insights into NoV and enteric virus biology. In the current review, we will summarize recent findings from MuNoV that increase our understanding of enteric virus pathogenesis and highlight similarities between human and murine NoVs that underscore the value of MuNoV to inform studies of HuNoV biology. We also discuss the potential of endemic MuNoV infections to impact other disease models.

Keywords

Enteric virus; Calicivirus; Positive-sense RNA virus; Animal model; Virus-microbiota-host interplay

Introduction

The breeding of specific-pathogen-free (SPF) mice by exclusion of common mouse pathogens has enabled biomedical research to increase consistency of disease models across laboratories and institutions by removing potentially confounding effects of natural infections. However, the definition of SPF is unavoidably restricted to previously identified pathogens, most often discovered by their association with specific disease patterns. Murine noroviruses (MuNoVs) cause asymptomatic infections in many mouse strains including immunocompetent mice and those lacking adaptive immune responses (1, 2). Thus, their existence went unrecognized until 2003 when we serendipitously discovered the first MuNoV, called MNV-1, in our research facility (2). Since then, diagnostic MuNoV testing

*Corresponding author: University of Michigan, Department of Microbiology and Immunology, 5622 MSII, 1150 West Medical Center Dr. Ann Arbor, MI, 48109-5620, Tel.: 734-647-9599, Fax: 734-764-3562, cwobus@umich.edu.

has been developed (3) and routine surveillance in mouse facilities has revealed that MuNoVs are the most prevalent type of pathogen in biomedical research facilities worldwide (4–7). Moreover, many MuNoV strains circulate in mouse research colonies and in the wild, with over 100 different MuNoV sequences currently deposited in GenBank.

With the recent advances in high-throughput sequencing technology, identification of infectious agents no longer depends on their ability to cause overt symptoms. In analogy with commensal bacteria, the awareness of the virome (viruses quiescently inhabiting the mammalian host) and its likely influences on the health of the mammalian host is increasing (8). Thus, the academic research community needs to tackle the difficult question of which virome constituents are tolerable in mouse colonies and how to control for their inevitable variability. Our analysis of MuNoVs twelve years after their initial discovery – in terms of their utility as a norovirus (NoV) model system and their potential to impact other research models as members of the mouse virome – is meant to inform and stimulate this conversation. Thus, this review will focus on (i) how MuNoV research has advanced our understanding of general NoV biology including cell tropism, promotion of viral pathogenesis by commensal bacteria, persistence, and immunity; and (ii) the implications of endemic MuNoV infections on other disease models considering their presence in many barrier facilities.

Noroviruses: The Basics

Norovirus Genomic Organization

NoVs are positive-sense, single-stranded RNA viruses that constitute a genus in the *Caliciviridae* family (9). The genome is a ~7.5 kb polyadenylated RNA with a viral protein called VPg (viral protein, genome-linked) covalently attached to the 5' end, and comprised of three or four open reading frames (ORFs) (Fig. 1A). ORF1 encodes the non-structural proteins, which are expressed as a polyprotein that is cleaved by the virus-encoded protease into six mature proteins (NS1/2 = N-terminal protein, NS3 = p22, NS4 = 3A-like, NS5 = VPg, NS6 = protease, NS7 = RNA-dependent RNA polymerase) (Fig. 1A). ORF2 encodes the major capsid protein VP1, which is divided into shell (S) and protruding (P) domains, and forms non-enveloped, icosahedral particles that are 27–40 nm in size (Fig. 1B). ORF3 encodes the minor capsid protein VP2, of which a few copies in the interior of the virion are thought to play a role in genome packaging (10, 11). MuNoVs also contain a fourth ORF that is expressed in an alternate reading frame from ORF2 and encodes virulence factor 1 (VF1), an antagonist of the innate immune response (12, 13).

Norovirus Phylogeny

NoVs display a high degree of genetic diversity and are classified based on their VP1 sequences into six genogroups (GI–GVI) (14) and a proposed GVII (15). Human NoVs (HuNoVs) segregate in GI, GII, and GIV, with GII also containing porcine strains. GIII contains bovine and ovine strains; GV contains murine and rat strains; and GIV, GVI and the proposed GVII contain canine strains (15). Genogroups are further subdivided into genotypes with GII genotype 4 (GII.4) strains causing the majority of outbreaks and recurrent epidemics in humans (16, 17). The relationship between genetic and serologic

diversity of NoVs remains largely unknown. One exception is a study of seven MuNoV strains, which segregated into a single genotype and serotype (13). In spite of the genetic and serologic similarity between identified MuNoV strains, they display significant phenotypic variability as will be highlighted in this review. Considering the much greater genetic diversity among HuNoVs, it is likely that they are even more phenotypically variable.

Norovirus Epidemiology

HuNoV infections occur in all age groups and are associated with significant morbidity, mortality and economic losses. They cause ~20% of all cases of acute gastroenteritis worldwide (18) and an estimated 200,000 deaths in young children in developing countries annually (19). In the United States alone, these viruses cause approximately 21 million infections, ~70,000 hospitalizations and ~800 deaths (20). After introduction of a new rotavirus vaccine, HuNoVs have become the most common cause of severe childhood diarrhea (21–23). The estimated costs of food-borne outbreaks due to HuNoVs in the United States range between 2.2 – 5.8 billion US\$ per year (24, 25). The high prevalence of NoVs is attributable to their great stability in the environment and their highly infectious nature. NoV particles remain infectious for weeks to months in the environment (26–28), and the estimated 50% human infectious dose is between 18 and 2800 particles depending on the study (29, 30). Other features that make NoVs particularly well-adapted at causing large outbreaks are their multiple routes of transmission (e.g., exposure to contaminated food, water, or fomites; person-to-person spread; exposure to aerosolized vomitus particles) and asymptomatic long-term shedding (31–33).

A detailed understanding of the biology of these prevalent pathogens is critical to develop effective prevention and treatment strategies. Identification of MuNoVs allowed the research community to study NoV infection in a tractable and genetically manipulable small animal model system. Moreover, until recently (34), MuNoVs were the only NoVs that could be cultured in the laboratory (35). Together with multiple reverse genetics systems that permit genetic manipulation of the MuNoV genome (36–38), these tools have greatly advanced our understanding of NoV biology (39, 40). NoV replication strategies were just comprehensively reviewed (41). Thus, we will focus herein on findings in the MuNoV system and their contribution to our understanding of NoV and enteric virus biology.

Norovirus Tropism for Immune Cells

The cell tropism of HuNoVs has long eluded the research community as these viruses appear to infect very few cells *in vivo* (42), complicating detection of viral antigen in tissue sections from infected hosts. Extensive efforts to culture HuNoVs in established intestinal epithelial cell (IEC) lines have been unsuccessful to date (43–46). While infection of IECs *in vivo* cannot be ruled out by these studies, the data outlined below demonstrate immune cells are MuNoV and HuNoV target cells.

Murine noroviruses infect dendritic cells and macrophages

MuNoVs infect macrophages and dendritic cells in culture and *in vivo* (35, 47). Infection in culture induces apoptosis and cell lysis (13, 48). Although a HuNoV did not replicate in blood-derived macrophages or dendritic cells *in vitro* (42), detection of HuNoV antigen in cells resembling or confirmed to be dendritic cells or macrophages in tissue sections of infected hosts suggests that they may be infected *in vivo* (42, 49–51). To reach their target cells, MuNoVs can cross the intestinal epithelium by transcytosis through microfold (M) cells in the absence of viral replication (52, 53). M cells are specialized IECs that transport particulate antigens from the lumen into the host to initiate immune responses but are also used by numerous enteric pathogens to overcome host defenses (54, 55). Thus, it is likely that HuNoVs can also use M cells to breach the intestinal epithelial barrier.

Murine noroviruses infect B cells

MuNoVs also infect B cells in culture (34). However, B cell infection is distinct from that of dendritic cells and macrophages. First, MuNoVs display a low infectivity in mature B cells, with 5–15% of B cells actively infected compared to >90% of macrophages (34). Second, MuNoV infection of mature B cells does not result in a loss in cell viability (34). Further studies are required to confirm whether infection is truly noncytopathic or due to the low infectivity rate. Finally, mature B cells become persistently infected by MuNoVs despite high levels of progeny virus production and release into the culture supernatants (34). Multiple lines of evidence corroborate that B cells are also *in vivo* MuNoV targets – virus titers are reduced in mice lacking functional B cells compared to wild-type mice; viral genomes can be amplified from purified Peyer's patch B cells of infected wild-type mice; and viral nonstructural protein can be detected in Peyer's patch B cells of infected STAT1^{-/-} and IL-10^{-/-} mice (1, 34, 56).

A human norovirus infects cultured B cells

MuNoV infection of B cells (34) combined with HuNoV antigen in intestinal B cells of infected chimpanzees (49) prompted us to test whether HuNoVs similarly infect this cell type. Indeed, a GII.4 HuNoV productively infects the human B cell line BJAB (34). Similar to MuNoV, HuNoV infection of B cells in culture appears noncytopathic, and HuNoV can be transcytosed across a polarized epithelium to infect underlying B cells in a co-culture system (34). Ongoing studies aim to enhance the robustness of this B cell cultivation system since it currently supports only low levels of viral replication, which is in apparent contradiction with the high levels of virus shed in the stool of infected people (31). Whether HuNoVs replicate in other cell types either *in vitro* or *in vivo*, in particular macrophages, dendritic cells, and IECs, needs to be determined in future studies.

Norovirus Interactions with Carbohydrates

Human and murine noroviruses bind glycans in a virus strain-specific manner

In analogy to the related feline calicivirus (FCV) (57, 58), NoV infection of cells starts with viral attachment to cell surface glycans followed by subsequent binding to a protein receptor, although the latter have yet to be identified for NoVs. Most HuNoVs bind histo-blood group

antigens (HBGAs) in a strain-dependent manner (e.g., reviewed in: (59, 60)). Binding to HBGAs also extends to bovine (61) and canine NoVs (62), but not MuNoVs (58) or feline NoVs (63). HBGAs, including the ABO and Lewis families of antigens, are neutral carbohydrates attached to lipids or proteins presented on multiple cell types including IECs and secreted into body secretions like saliva (reviewed in: (58)). Certain HuNoV strains also bind heparan sulfate (64), sialic acid (65), and β -galactosylceramide (GalCer) (66), demonstrating the plasticity of glycan binding. Like HuNoV, MuNoV binding to host cell glycans is virus strain-dependent (67). Certain MuNoV strains bind to terminal α 2,3-linked sialic acid residues on gangliosides and glycoproteins, while others recognize yet to be defined glycans on N-linked proteins (67, 68).

Conserved features of human and murine norovirus glycan interactions

Although binding of human and murine NoVs to different glycans likely reflects an adaptation to the respective host species (67, 69), common features of virus-glycan interactions emerge. First, the glycan binding sites are located in a similar region of the VP1 P domain (67) topologically (Fig. 2A–D) and at the sequence level (Fig. 2E). Second, the outer-most loops of the MuNoV and HuNoV P domains are highly flexible and present in two conformations, open and closed (70) (Fig. 2F). For MuNoVs, the loop flexibility is important for escape from a neutralizing antibody in addition to strict disruption of the antibody-antigen binding interface (71). Future studies are needed to determine whether an anti-HuNoV antibody directed to the corresponding loop (72) is neutralizing and whether conformational changes in the HuNoV capsid can mediate escape.

Carbohydrates may act as norovirus attachment receptors

Although the presence of certain HBGAs is linked to susceptibility to specific HuNoV strains (reviewed in: (73)), the mechanism by which HBGAs facilitate HuNoV infection remains to be determined. HBGAs are not entry receptors since the presence of HBGAs on the cell surface does not permit infection of non-susceptible cells (74). Instead, HBGAs likely play a role as HuNoV attachment or co-receptors via concentrating viral particles on the cell surface (75). Free HBGAs also increase HuNoV attachment to, and promotes infection of, susceptible B cells (34). HuNoV-based clustering of glycosphingolipids causes membrane invaginations (76), thus binding to glycosphingolipids by murine and human NoVs may represent the first step in NoV entry. Although HBGAs linked to lipids and/or proteins are synthesized by IECs, immune cells, a NoV target cell population, lack the biosynthetic enzymes and only absorb a low level of HBGA-linked glycolipids from the plasma (77). Thus, additional mechanistic studies are needed to solidify the role(s) of different forms of HBGAs during HuNoV infection.

Noroviruses and Bacteria

An emerging theme of enteric virus pathogenesis is their exploitation of commensal bacteria for the purpose of enhancing infection (78, 79). Analyzing viral infections in germ-free mice devoid of an intestinal microbiota and antibiotic (Abx)-treated mice with a significantly reduced intestinal microbial load demonstrated that infection, and in some cases

transmission, of different types of enteric viruses – poliovirus, reovirus, mouse mammary tumor virus (MMTV), and rotavirus – is enhanced by the microbiota (80–83).

Commensal bacteria promote murine norovirus infection of mice

Mirroring results for these other enteric viruses, commensal bacteria also promote MuNoV infections (Fig. 3A). Commensal bacteria enhance acute MNV-1 and MNV-3 infections as virus titers in mucosal tissues were reduced in Abx-treated mice (34). Commensal bacteria also facilitate the establishment of persistent MNV-CR6 infection (84) because persistent intestinal infection was not observed in Abx-treated mice but fecal transplantation from microbially colonized (but not Abx-treated) mice restored the ability of MNV-CR6 to establish persistence. A third study demonstrated reduced MNV-CR6 infection in germ-free mice indicating that bacteria enhance, but are not absolutely required for, MuNoV infection (85). This is consistent with findings that MuNoV infection of cultured cells (34, 35), and its transcytosis across the epithelial barrier *in vitro* (52), do not require bacteria.

Commensal bacteria stimulate human norovirus infection of B cells

The importance of commensal bacteria for enteric MuNoV infection led Jones et al. (34) to test whether HuNoV infection of cultured cells was enhanced by the presence of bacteria. Indeed, removal of commensal bacteria from virus-positive stool via filtration drastically reduced viral infectivity of human B cells, although viral titers in the inoculum were unaffected; and a specific commensal bacteria – *Enterobacter cloacae* – was able to restore infectivity of the filtered virus inoculum (34).

In contrast to poliovirus and MMTV (80, 81, 83), NoVs do not appear to bind bacterial LPS: Free LPS was unable to stimulate HuNoV infection of B cells (34), and the LPS sensor TLR4 and its downstream signaling molecules were dispensable for establishment of persistent MuNoV infections (84). Instead, HBGAs expressed on certain commensal bacteria (86) promote HuNoV infection of B cells while a non-HBGA-expressing bacterial strain did not (34). Importantly, synthetic HBGA restored infectivity of filtered virus inoculum comparably to HBGA-expressing bacteria (34). A model emerges whereby a HuNoV binds an HBGA-expressing commensal bacteria in the gut lumen, the virus-bacteria complex is transcytosed across the intestinal epithelium, and the bacterial glycan stimulates viral attachment to underlying intestinal B cells (Fig. 3B) (87). Overall, identification of commensal bacteria as a stimulatory factor for *in vivo* MuNoV infection led directly to the development of the first HuNoV culture system, underscoring the power of animal models in advancing our understanding of human pathogens.

Many open questions remain regarding the physiological significance of NoV interactions with commensal bacteria. For example, what are the immune implications of virus binding to commensal bacteria? Co-recognition of bacterial and viral ligands may result in a tolerogenic environment that suppresses robust antiviral immunity similar to findings with MMTV (81). Consistent with this idea, MuNoV infection of wild-type mice is only weakly inflammatory, while infection of IL-10^{-/-} mice results in intestinal inflammation but only in the presence of commensal bacteria (Fig. 4A) (56). This suggests that the anti-inflammatory cytokine IL-10 suppresses immune responses to a NoV in a bacteria-dependent manner.

Norovirus Persistence and Immunity

The immune mechanisms that control acute NoV infections and protective immunity determinants were reviewed recently (40). In brief, type I interferon (IFN) and components of the adaptive immune system coordinate control and clearance of primary NoV infections; and antibody and CD4⁺ T cells are critical for mediating protective immunity to a secondary NoV challenge, although there are virus strain-specific distinctions in the magnitude of memory immune responses and the overall duration of these responses is fairly short-lived. Below we focus on the ability of NoVs to establish prolonged, or persistent, infections; and on more recent studies pertaining to NoV immunity that illuminate (i) a critical role for type III IFN in controlling persistent enteric infection, and (ii) influences of nutritional status and co-infection on anti-NoV immune responses.

Murine noroviruses can persist in the colon, contributing to prolonged fecal shedding

The classical symptoms of HuNoV infection (e.g., vomiting, diarrhea and nausea) develop and resolve within a few days but viral shedding following symptomatic and asymptomatic infections lasts for weeks to months (31, 88, 89). Moreover, immunocompromised patients can be chronically infected (90), generating highly diverse viral populations which may serve as reservoirs of phenotypically distinct HuNoV strains (91, 92). Similar to HuNoVs, most MuNoVs are shed in the feces for extended periods of time with the exception of MNV-1 (13). Long-term shedding of MuNoVs has been associated with ongoing low-level persistent infection of the colon (93). A single amino acid difference in the NS1/2 protein between acute MNV-1 and persistent MNV-CR6 regulates colonic tropism and establishment of persistence (94). In addition, reduced antiviral CD8⁺ T cell are induced during persistent MNV-CR6 infection compared to T cell responses during acute MNV-1 infection (95), suggesting that immune evasion contributes to the ability of a NoV to establish persistence. The cell type acting as a persistent reservoir of NoVs is undefined, but we speculate that it is B cells considering they are persistently infected by MuNoVs in culture (34).

Type III interferon controls persistent murine norovirus infection of the colon

Although type I interferon (IFN) plays a critical role in preventing severe MuNoV infections and limits replication in peripheral tissues (1, 2, 96), the type I IFN receptor (*Ifnar1*) is dispensable for controlling colonic infection and fecal shedding of persistent MNV-CR6 (96). Instead, the type III IFN, or IFN λ , receptor (*IFNlr1*) controls intestinal infection independent of type I IFN signaling or adaptive immune cells. IFN λ treatment one day after MNV-CR6 infection prevented persistent infection, while IFN λ treatment after establishment of persistent infection resulted in viral clearance from tissues and feces. This suggests type III IFN inhibition of intestinal infection occurs indirectly (Fig. 4B), a conclusion further supported by the following findings. First, IFN λ (but not IFN β) treatment of permissive dendritic cells in culture failed to inhibit MNV-CR6 replication. Second, bone marrow chimera experiments showed increased MNV-CR6 shedding into the feces when *IFNlr1*^{-/-} recipient mice received wild-type donor (immune) cells but not when wild-type recipient mice received *IFNlr1*^{-/-} donor cells. Finally, the ability of a MuNoV

strain to establish persistence correlated with a lack of induction of IFN λ (but not IFN β) in infected tissues.

Overall, findings from MuNoV indicate that NoVs evade both innate (e.g., IFN induction) and adaptive (e.g., robust CD8⁺ T cell response) immunity to establish persistence (12, 95). The influence of commensal bacteria on persistence is complex as underscored by the observation that MNV-CR6 persistence in *IFN λ IR^{-/-}* mice is not only enhanced compared to wild-type control mice but it also occurs in a bacteria-independent manner; whereas persistence establishment in wild-type mice is dependent on bacteria (84). Thus, commensal bacteria may normally suppress IFN λ and thereby facilitate enteric viral infections.

Malnutrition and helminth co-infections suppress anti-norovirus immunity

People in developing countries are susceptible to more severe enteric infections than those in industrialized nations and they fail to respond as robustly to oral vaccines (97–100). The basis for these phenomena is multi-factorial and studying these processes in tractable animal models is critical to the development of strategies to enhance protection in this particularly vulnerable population. Studies using the MuNoV model have made important contributions. First, malnourished mice developed more severe and prolonged MNV-1 infections, and they failed to develop protective immunity to a secondary challenge, which correlated with a severely muted antiviral mucosal IgA response (101). They also supported enhanced viral evolution, raising the possibility that malnourished individuals could represent reservoirs of novel viral variants. Second, the antiviral CD8⁺ T cell response to MNV-1 or MNV-CR6 was reduced in mice co-infected with the helminth *Trichinella spiralis* (102). This reduction was associated with a potent activation of alternatively-activated macrophages, suggesting an immunomodulatory role for these innate effector cells. Because MuNoV infections are regulated by the intestinal microbiota, nutrition, and co-infections with other common enteric pathogens, they represent an excellent model with which to study the complex relationships that ultimately dictate disease outcome.

Development of norovirus antivirals

Although the rapid nature of typical NoV infections may limit the utility of therapeutic antiviral treatment strategies in immunocompetent individuals, NoV drug discovery efforts are needed to treat the high number of immunocompromised patients chronically infected with HuNoVs (90). Moreover, prophylactic drugs that limit viral transmission can greatly reduce the economic costs associated with norovirus infections (25). Due to the difficulties in developing a HuNoV cell culture system (43–46), the inability to orally infect mice with HuNoV (51), and the high cost of HuNoV large animal studies, many investigators have used the MuNoV model for initial drug discovery often complemented by studies in HuNoV replicon-bearing cells (103). Several recent reviews on the topic of NoV antiviral drug development have been published (e.g., (104, 105)). Thus, we will focus on examples that have used the MuNoV model for drug development and efficacy testing both *in vitro* and *in vivo*.

Antivirals targeting viral proteins

While all NoV proteins represent putative targets for antiviral development, most work to date has focused on the viral protease and polymerase (105). The conserved nature of these two NoV proteins (106, 107) increases the likelihood that findings from MuNoVs will translate to HuNoVs in future clinical trials. Furthermore, the NoV RNA-dependent RNA polymerase (RdRp) is similar to other RdRps from positive-strand RNA viruses (107), allowing the use of previously developed RdRp inhibitors such as nucleoside analogs 2'-C-methylcytidine (2CMC) and favipiravir. 2CMC and the closely related analog 2'-F'-2CMC displayed antiviral activity against MuNoV *in vitro* in the low micromolar range and drug-resistant escape mutants were not detected through 30 passages (108, 109). Importantly, mice lacking type I and II IFN receptors treated with 2CMC prior to MNV-1 infection were protected from developing diarrhea and they developed protective immunity to a subsequent lethal challenge (110). 2CMC also has potential as a prophylactic in preventing NoV transmission (111) and was effective in clearing HuNoV from replicon-bearing cells (110). However, the likelihood of widespread use in humans is low given the previously observed side effects of the 2CMC derivative Valopicitabine (also called NM283) in clinical trials of HCV-infected patients (112) (http://www.natap.org/2007/HCV/071607_03.htm).

Favipiravir (T-705) is in late stage clinical development for influenza virus treatment and exhibits antiviral activity against multiple other RNA viruses in animal models (e.g., (113–116)). Favipiravir prevents MuNoV replication in culture early during infection (117). More importantly, it reduced MuNoV titers *in vivo* and virus was eliminated from the feces in the majority of persistently infected animals (118). Favipiravir is a guanosine and adenosine analog (119) that induced mutations in the MuNoV genome *in vivo* (118), providing *in vivo* proof that lethal mutagenesis (i.e., increasing mutations in the viral genome to cause extinction of the viral population) can be used therapeutically. Considering the broad spectrum antiviral activity of favipiravir (120), this drug will likely exhibit antiviral activity against diverse HuNoVs strains.

Antivirals targeting host proteins

Numerous host proteins facilitate NoV infections, and those representing candidate drug targets were recently reviewed (104). However, *in vivo* studies of host-targeted antivirals that reduce NoV infections without significant toxicity remain scarce. One example is WP1130, a small molecule that inhibits a subset of cellular deubiquitinases (DUBs) (121). Ubiquitin modifying enzymes, including DUBs, play critical regulatory functions in many cellular processes and can influence viruses at all stages of the viral life cycle. WP1130 exhibits antiviral activity in cell culture against several RNA viruses, including MuNoV in murine macrophages and Norwalk virus in replicon-containing cells by induction of the unfolded protein response through cellular DUB inhibition (122). Oral administration of WP1130 inhibited MuNoV infection in the proximal small intestine of mice, although it failed to block infection in more distal regions of the intestine because of the poor solubility and/or bioavailability of the compound. These studies suggest that cellular DUBs represent promising targets for host-targeted broad spectrum antivirals, but future efforts are needed to identify more effective derivatives.

Although anti-NoV drug development efforts are still at an early stage, studies with MuNoV have identified many putative targets. The recent development of a HuNoV culture model (34) now provides an important tool to test the translatability of MuNoV findings to HuNoV, thus significantly enhancing future efforts.

Influence of Murine Norovirus on Biomedical Research Models

MuNoVs are highly prevalent in mouse colonies at academic research institutions and in pet stores (4, 123). Most strains of mice naturally infected with a MuNoV do not display noticeable signs of illness, thus their presence easily goes unrecognized. Eradicating MuNoVs from mouse colonies is time-consuming and expensive (124), thus many academic research institutions tolerate the presence of natural MuNoVs. This can significantly compromise experimental interpretability as exemplified by a large microarray analysis of germ-free versus conventionally housed mice in which certain conventional animals were contaminated with MuNoV causing upregulation of antiviral genes (125). Considering the tropism of MuNoVs for innate and adaptive immune cells (34, 35) and their ability to establish persistent infections (2, 13), multiple studies have directly tested the influence of MuNoV infection on various models of inflammatory diseases and other infections. As summarized below, these studies demonstrate that MuNoV infections can have effects on intestinal inflammatory conditions dependent on host genetic background and MuNoV strain but most MuNoV infections minimally impact other disease models.

Effects of murine norovirus infections on extra-intestinal inflammatory and infectious conditions

With the exception of inflammatory bowel disease (IBD) models discussed below, studies examining the effect of MuNoV infection on inflammatory conditions have observed no or modest conditional phenotypes. MNV-4 infection resulted in a modest increase in the size of atherosclerotic lesions in *Ldlr*^{-/-} mice fed a high-fat, high-cholesterol diet, but not in those fed a high-fat, high-sucrose diet (126). MNV-4 infection did not affect the development or severity of obesity or insulin resistance in wild-type C57BL/6 mice fed a high-fat diet or in *Ldlr*^{-/-} mice (126, 127); nor did it affect *Helicobacter*-induced colon cancer in *Smad3*^{-/-} mice (128).

Similarly, MuNoVs have no to modest effects on other infections. Persistent MNV-G infection enhanced a subsequent mouse parvovirus (MPV) infection and extended the duration of MPV shedding (129). However, neither acute MNV-1 nor persistent MNV-CR6 infections altered the severity of, or immunity to, influenza A virus or vaccinia virus infections (130). MNV-1 also failed to affect murine cytomegalovirus (MCMV) infection establishment or reactivation, although MNV-1 infection modestly reduced the T cell response to MCMV (131). Furthermore, MNV-CR6 did not alter the immune response to Friend retrovirus (132).

Murine norovirus effects on inflammatory bowel disease models

Crohn's disease, a common form of IBD in people, is associated with genetic polymorphisms in numerous loci including the autophagy gene *ATG16L1* (133, 134). Mice

with an *ATG16L1* mutation are susceptible to intestinal abnormalities displaying features of Crohn's disease (135–137). This phenotype is linked to persistent MuNoV infection, an association that was revealed when researchers re-derived mice into a barrier facility free of endemic MuNoV and observed a loss of the intestinal abnormalities, particularly in Paneth cells, previously detected in the *Atg16L1*-deficient mice (136). Aberrant Paneth cell morphology (e.g., in granule packaging) was fully restored upon experimental MNV-CR6, but not MNV-1, infection (137). Moreover, intestinal abnormalities in MNV-CR6-infected *Atg16L1*-deficient mice were dependent on commensal bacteria and the inflammatory cytokines TNF α and IFN γ . A model emerges whereby *Atg16L1* deficiency causes a reduction in the amount of antimicrobial peptides produced in, or packaged into granules of, Paneth cells while increasing inflammatory cytokine production in response to a viral infection (138). This is a perfect example of how endemic infections in colonies of research animals can confound data interpretation and introduce lab-to-lab variability within the same model system. It is also, however, an elegant and powerful example of how animal models can facilitate the dissection of complex multifactorial disease processes that are responsible for many inflammatory and autoimmune disorders common in the human population. Future studies are needed to investigate how the newly recognized interdependence between the microbiota, MuNoV infection and the mucosal immune system (34, 84, 85) contribute to these pathologies.

Other models of IBD showed less dramatic effects, underscoring the conditional nature of MuNoV effects on inflammatory disease models. *Helicobacter* infection induces IBD in certain knockout strains of mice, and MNV-4 infection exacerbated this condition in *Mdr1a*^{-/-} mice (139). However, MNV-4 failed to exacerbate disease in *Helicobacter*-infected *Smad3*^{-/-} or *IL-10*^{-/-} mice (128, 140). *Salmonella typhimurium* infection of wild-type mice treated with streptomycin is used as a model of intestinal fibrosis, a complication in IBD, but acute MNV-1 or persistent MNV-4 infections did not alter major features of this model (141).

Potential for murine norovirus infection to shape the intestinal microbiota

Endemic viral infections may influence the composition of the intestinal microbiota, which strongly shape host mucosal immunity (142–144). Conflicting reports exist whether MuNoV infection alters the composition of the microbiota in immunocompetent mice. No major changes were observed in the intestinal microbiota of inbred C57BL/6 or outbred Swiss Webster mice infected with MNV-1, MNV-4, or CR6 (145). In another study, a significant reduction in the Bacteroidetes/Firmicutes ratio was observed in MNV-1-infected C57BL/6 mice compared to mock-inoculated controls (101). The reason for these discordant results is unclear but institutional variability may play a role. Reduced Bacteroidetes and increased Proteobacteria were observed in a minority of HuNoV-infected subjects (146), although very limited health data and inability for repeated sampling confound the observation. These limited data suggest that NoV infections may alter the microbiota under certain conditions.

Murine noroviruses provide beneficial effects of commensal bacteria in germ-free mice

The beneficial effect of a “healthy” microbiota is widely recognized (e.g. (147, 148)). Remarkably, one study recently demonstrated that a single MuNoV could restore beneficial

functions attributed typically to commensal bacteria in germ-free mice despite reduced viral loads (85). Germ-free mice infected with different strains of MuNoV (CR6, MNV-1, and MNV-SKI) resembled bacterially-colonized mice in terms of villus width, lymphocyte number and function, antibody levels, and basal suppression of innate lymphoid cell expansion. These phenotypes were dependent on type I IFN signaling since MNV-CR6 was not able to restore intestinal health in *Ifnar1*^{-/-} germ-free mice. MuNoV infection was also protective against infectious and chemical forms of intestinal injury in Abx-treated mice. These results suggest that under certain conditions MuNoVs may be considered commensals in nature, shifting the perception that viral infections are strictly pathogenic. However, as this applied only to a bacteria-free environment, the translatability to other NoV infections is unclear.

Conclusions

The discovery of MNV-1 and subsequent development of cell culture and reverse genetics systems have provided important tools to the NoV field to study molecular aspects of NoV biology. Combined with advances in HuNoV research, these studies have yielded important insights into unifying principles underlying NoV replication mechanisms. Further comparative studies between NoVs from different host species will likely reveal additional shared mechanisms that deepen our understanding of these viruses. In addition, the study of MuNoVs as a model for enteric virus infections has yielded exciting new findings regarding the role of commensal bacteria during acute and persistent viral infections, effects on host physiology, and innate immunity in the intestine. Since these studies are in their infancy, much remains to be discovered regarding the host-microbiota-virus interplay.

Complicating but also enriching these studies is the phenotypic diversity of MuNoV strains. Although the strains are genetically highly related and segregate into a single cluster within genogroup V, there are striking differences in the duration of infections, virulence, cytopathogenicity in cultured cells, immune responses, and effects on other inflammatory conditions. Even for the strains preferentially studied in the MuNoV field (i.e. MNV-1 as the prototype acute MuNoV strain, and MNV-3, MNV-4, MNV-CR3, and MNV-CR6 as representative persistent strains), viral determinants regulating their varied outcomes of infection are only beginning to be defined. HuNoVs exhibit much greater genetic diversity between different clusters and genogroups. Thus, one implication for HuNoV research is the existence of a variety of unrecognized differential infection outcomes and immune responses. Dissecting and defining these differences and developing the tools to do so remains a major challenge for the future. Knowledge from studies of MuNoVs may help guide such efforts. It may be in this capacity that a common contaminant to mouse research facilities makes its greatest contribution to improving human health.

Acknowledgments

C.E.W. is a recipient of the Friedrich Wilhelm Bessel award from the Alexander von Humboldt Foundation. Work in her laboratory was supported by NIH grants AI102106, AI080611, AI103961 and DARPA grant HR0011-11-C-0093. S.M.K. is funded by University of Florida (UF) Emerging Pathogens Institute Seed Fund 13-1; UF Opportunity Fund 00115487; and Center for Produce Safety award 2014-355. We thank Dr. T. Smith (University of Texas, Galveston, USA) for Fig. 1C, and Luisa Ströh (University of Tübingen, Germany), for helpful discussions and help with Pymol.

Acronyms and Definitions

Enteric virus

a virus, which infects its host via the gastrointestinal tract

Peyer's patch

lymphoid follicles present in the lamina propria of the small intestine

Histo-blood group antigens (HBGA)

neutral carbohydrates attached to lipids or proteins presented on some cell types, including intestinal epithelial cells, and in body secretions like saliva

Secretor/Non-secretor

an individual that has/does not have a functional copy of α 1,2 fucosyltransferase 2 (FUT2) to synthesize the H antigen and secrete/does not secrete blood group A, B, or AB antigens in saliva and gastric mucus

Genogroup/genotype

related viruses within a genus; can be further subdivided into genotype

Germ-free mice

animals born and raised under sterile conditions

Commensal bacteria

bacteria living on another host without causing harm or benefit

Virome

Collection of genes from viruses or viruses living in a given environment, e.g. gastrointestinal tract.

Microbiome/microbiota

Collection of genes from microorganisms / collection of microorganisms (herein used to describe bacteria) living in a given environment, e.g. gastrointestinal tract.

Literature Cited

- 1Mumphrey SM, Changotra H, Moore TN, Heimann-Nichols ER, Wobus CE, et al. Murine norovirus 1 infection is associated with histopathological changes in immunocompetent hosts, but clinical disease is prevented by STAT1-dependent interferon responses. *J Virol.* 2007; 81:3251–63. [PubMed: 17229692]
- 2Karst SM, Wobus CE, Lay M, Davidson J, Virgin HW IV. STAT1-dependent innate immunity to a Norwalk-like virus. *Science.* 2003; 299:1575–8. [PubMed: 12624267]
- 3Hsu CC, Wobus CE, Steffen EK, Riley LK, Livingston RS. Development of a microsphere-based serologic multiplexed fluorescent immunoassay and a reverse transcriptase PCR assay to detect murine norovirus 1 infection in mice. *Clin Diagn Lab Immunol.* 2005; 12:1145–51. [PubMed: 16210475]
- 4Henderson KS. Murine norovirus, a recently discovered and highly prevalent viral agent of mice. *Lab Anim (NY).* 2008; 37:314–20. [PubMed: 18568010]
- 5Xiang Z, Tian S, Tong W, Chang H, Su J, et al. MNV primarily surveillance by a recombination VP1-derived ELISA in Beijing area in China. *J Immunol Methods.* 2014; 408:70–7. [PubMed: 24857757]

- 6Mahler M, Kohl W. A serological survey to evaluate contemporary prevalence of viral agents and *Mycoplasma pulmonis* in laboratory mice and rats in western Europe. *Lab Anim* (NY). 2009; 38:161–5. [PubMed: 19384313]
- 7Pritchett-Corning KR, Cosentino J, Clifford CB. Contemporary prevalence of infectious agents in laboratory mice and rats. *Lab Anim*. 2009; 43:165–73. [PubMed: 19015179]
- 8Cadwell K. Expanding the Role of the Virome: Commensalism in the Gut. *J Virol*. 2014
- 9Green KY. Caliciviridae: The Noroviruses. In: Knipe DM, Howley PM, Cohen JI, Griffin DI, Lamb RA, Martin MA, Racaniello VR, Roizman B, editors *Fields Virology* Philadelphia: Lippincott Williams & Wilkins, a Wolters Kluwer Business; 2013 582608
- 10Vongpunswad S, Venkataram Prasad BV, Estes MK. Norwalk Virus Minor Capsid Protein VP2 Associates within the VP1 Shell Domain. *J Virol*. 2013; 87:4818–25. [PubMed: 23408637]
- 11Glass PJ, White LJ, Ball JM, Leparc-Goffart I, Hardy ME, Estes MK. Norwalk virus open reading frame 3 encodes a minor structural protein. *J Virol*. 2000; 74:6581–91. [PubMed: 10864672]
- 12McFadden N, Bailey D, Carrara G, Benson A, Chaudhry Y, et al. Norovirus regulation of the innate immune response and apoptosis occurs via the product of the alternative open reading frame 4. *PLoS Pathog*. 2011; 7:e1002413. [PubMed: 22174679]
- 13Thackray LB, Wobus CE, Chachu KA, Liu B, Alegre ER, et al. Murine noroviruses comprising a single genogroup exhibit biological diversity despite limited sequence divergence. *J Virol*. 2007; 81:10460–73. [PubMed: 17652401]
- 14Kroneman A, Vega E, Vennema H, Vinje J, White PA, et al. Proposal for a unified norovirus nomenclature and genotyping. *Arch Virol*. 2013
- 15Vinje J. Advances in Laboratory Methods for Detection and Typing of Norovirus. *J Clin Microbiol*. 2014
- 16Siebenga JJ, Vennema H, Zheng DP, Vinje J, Lee BE, et al. Norovirus illness is a global problem: emergence and spread of norovirus GII.4 variants, 2001–2007. *J Infect Dis*. 2009; 200:802–12. [PubMed: 19627248]
- 17Vega E, Barclay L, Gregoricus N, Shirley SH, Lee D, Vinje J. Genotypic and epidemiologic trends of norovirus outbreaks in the United States, 2009 to 2013. *J Clin Microbiol*. 2014; 52:147–55. [PubMed: 24172151]
- 18Ahmed SM, Hall AJ, Robinson AE, Verhoef L, Premkumar P, et al. Global prevalence of norovirus in cases of gastroenteritis: a systematic review and meta-analysis. *Lancet Infect Dis*. 2014; 14:725–30. [PubMed: 24981041]
- 19Patel MM, Widdowson MA, Glass RI, Akazawa K, Vinje J, Parashar UD. Systematic literature review of role of noroviruses in sporadic gastroenteritis. *Emerg Infect Dis*. 2008; 14:1224–31. [PubMed: 18680645]
- 20Hall AJ, Lopman BA, Payne DC, Patel MM, Gastanaduy PA, et al. Norovirus disease in the United States. *Emerg Infect Dis*. 2013; 19:1198–205. [PubMed: 23876403]
- 21Becker-Dreps S, Bucardo F, Vilchez S, Zambrana LE, Liu L, et al. Etiology of childhood diarrhea after rotavirus vaccine introduction: a prospective, population-based study in Nicaragua. *Pediatr Infect Dis J*. 2014; 33:1156–63. [PubMed: 24879131]
- 22Koo HL, Neill FH, Estes MK, Munoz FM, Cameron A, et al. Noroviruses: The Most Common Pediatric Viral Enteric Pathogen at a Large University Hospital After Introduction of Rotavirus Vaccination. *J Pediatric Infect Dis Soc*. 2013; 2:57–60. [PubMed: 23687584]
- 23Hemming M, Rasanen S, Huhti L, Paloniemi M, Salminen M, Vesikari T. Major reduction of rotavirus, but not norovirus, gastroenteritis in children seen in hospital after the introduction of RotaTeq vaccine into the National Immunization Programme in Finland. *Eur J Pediatr*. 2013; 172:739–46. [PubMed: 23361964]
- 24Scharff RL. Health Related Costs from Foodborne Illness in the United States Washington D.C: Georgetown University; 2010
- 25Hoffmann S, Batz MB, Morris JG Jr. Annual cost of illness and quality-adjusted life year losses in the United States due to 14 foodborne pathogens. *J Food Prot*. 2012; 75:1292–302. [PubMed: 22980013]
- 26Lopman B, Gastanaduy P, Park GW, Hall AJ, Parashar UD, Vinje J. Environmental transmission of norovirus gastroenteritis. *Curr Opin Virol*. 2012; 2:96–102. [PubMed: 22440972]

- 27Cannon JL, Papafragkou E, Park GW, Osborne J, Jaykus LA, Vinje J. Surrogates for the study of norovirus stability and inactivation in the environment: A comparison of murine norovirus and feline calicivirus. *J Food Prot.* 2006; 69:2761–5. [PubMed: 17133824]
- 28Manuel CA, Hsu CC, Riley LK, Livingston RS. Soiled-bedding sentinel detection of murine norovirus 4. *J Am Assoc Lab Anim Sci.* 2008; 47:31–6.
- 29Teunis PF, Moe CL, Liu P, Miller SE, Lindesmith L, et al. Norwalk virus: how infectious is it? *J Med Virol.* 2008; 80:1468–76. [PubMed: 18551613]
- 30Atmar RL, Opekun AR, Gilger MA, Estes MK, Crawford SE, et al. Determination of the 50% human infectious dose for Norwalk virus. *J Infect Dis.* 2014; 209:1016–22. [PubMed: 24253285]
- 31Teunis PF, Sukhrie FH, Vennema H, Bogerman J, Beersma MF, Koopmans MP. Shedding of norovirus in symptomatic and asymptomatic infections. *Epidemiol Infect.* 2014:1–8.
- 32Mathijs E, Stals A, Baert L, Botteldoorn N, Denayer S, et al. A review of known and hypothetical transmission routes for noroviruses. *Food Environ Virol.* 2012; 4:131–52. [PubMed: 23412887]
- 33Bitler EJ, Matthews JE, Dickey BW, Eisenberg JN, Leon JS. Norovirus outbreaks: a systematic review of commonly implicated transmission routes and vehicles. *Epidemiol Infect.* 2013; 141:1563–71. [PubMed: 23433247]
- 34Jones MK, Watanabe M, Zhu S, Graves CL, Keyes LR, et al. Enteric bacteria promote human and mouse norovirus infection of B cells. *Science.* 2014; 346:755–9. [PubMed: 25378626]
- 35Wobus CE, Karst SM, Thackray LB, Chang KO, Sosnovtsev SV, et al. Replication of Norovirus in cell culture reveals a tropism for dendritic cells and macrophages. *PLoS Biol.* 2004; 2:e432. [PubMed: 15562321]
- 36Ward VK, McCormick CJ, Clarke IN, Salim O, Wobus CE, et al. Recovery of infectious murine norovirus using pol II-driven expression of full-length cDNA. *Proc Natl Acad Sci U S A.* 2007; 104:11050–5. [PubMed: 17581883]
- 37Arias A, Urena L, Thorne L, Yunus MA, Goodfellow I. Reverse genetics mediated recovery of infectious murine norovirus. *J Vis Exp.* 2012
- 38Chaudhry Y, Skinner MA, Goodfellow IG. Recovery of genetically defined murine norovirus in tissue culture by using a fowlpox virus expressing T7 RNA polymerase. *J Gen Virol.* 2007; 88:2091–100. [PubMed: 17622609]
- 39Wobus CE, Thackray LB, Virgin HW IV. Murine norovirus: a model system to study norovirus biology and pathogenesis. *J Virol.* 2006; 80:5104–12. [PubMed: 16698991]
- 40Karst SM, Wobus CE, Goodfellow IG, Green KY, Virgin HW. Advances in norovirus biology. *Cell Host Microbe.* 2014; 15:668–80. [PubMed: 24922570]
- 41Thorne LG, Goodfellow IG. Norovirus gene expression and replication. *J Gen Virol.* 2014; 95:278–91. [PubMed: 24243731]
- 42Lay MK, Atmar RL, Guix S, Bharadwaj U, He H, et al. Norwalk virus does not replicate in human macrophages or dendritic cells derived from the peripheral blood of susceptible humans. *Virology.* 2010; 406:1–11. [PubMed: 20667573]
- 43Papafragkou E, Hewitt J, Park GW, Greening G, Vinje J. Challenges of culturing human norovirus in three-dimensional organoid intestinal cell culture models. *PLoS One.* 2013; 8:e63485. [PubMed: 23755105]
- 44Duizer E, Schwab KJ, Neill FH, Atmar RL, Koopmans MP, Estes MK. Laboratory efforts to cultivate noroviruses. *J Gen Virol.* 2004; 85:79–87. [PubMed: 14718622]
- 45Herbst-Kralovetz MM, Radtke AL, Lay MK, Hjelm BE, Bolick AN, et al. Lack of norovirus replication and histo-blood group antigen expression in 3-dimensional intestinal epithelial cells. *Emerg Infect Dis.* 2013; 19:431–8. [PubMed: 23622517]
- 46Takanashi S, Saif LJ, Hughes JH, Meulia T, Jung K, et al. Failure of propagation of human norovirus in intestinal epithelial cells with microvilli grown in three-dimensional cultures. *Arch Virol.* 2014; 159:257–66. [PubMed: 23974469]
- 47Ward JM, Wobus CE, Thackray LB, Erexson CR, Faucette LJ, et al. Pathology of immunodeficient mice with naturally occurring murine norovirus infection. *Toxicol Pathol.* 2006; 34:708–15. [PubMed: 17074739]

- 48Bok K, Prikhodko VG, Green KY, Sosnovtsev SV. Apoptosis in murine norovirus-infected RAW264.7 cells is associated with downregulation of survivin. *J Virol.* 2009; 83:3647–56. [PubMed: 19211757]
- 49Bok K, Parra GI, Mitra T, Abente E, Shaver CK, et al. Chimpanzees as an animal model for human norovirus infection and vaccine development. *Proc Natl Acad Sci U S A.* 2011; 108:325–30. [PubMed: 21173246]
- 50Chan MC, Ho WS, Sung JJ. In vitro whole-virus binding of a norovirus genogroup II genotype 4 strain to cells of the lamina propria and Brunner's glands in the human duodenum. *J Virol.* 2011; 85:8427–30. [PubMed: 21680503]
- 51Taube S, Kolawole AO, Hohne M, Wilkinson JE, Handley SA, et al. A mouse model for human norovirus. *MBio.* 2013:4.
- 52Gonzalez-Hernandez MB, Liu T, Blanco LP, Auble H, Payne HC, Wobus CE. Murine norovirus transcytosis across an in vitro polarized murine intestinal epithelial monolayer is mediated by M-like cells. *J Virol.* 2013; 87:12685–93. [PubMed: 24049163]
- 53Gonzalez-Hernandez MB, Liu T, Payne HC, Stencel-Baerenwald JE, Ikizler M, et al. Efficient Norovirus and Reovirus Replication in the Mouse Intestine Requires Microfold (M) Cells. *J Virol.* 2014; 88:6934–43. [PubMed: 24696493]
- 54Miller H, Zhang J, Kuolee R, Patel GB, Chen W. Intestinal M cells: the fallible sentinels? *World J Gastroenterol.* 2007; 13:1477–86. [PubMed: 17461437]
- 55Neutra MR, Frey A, Kraehenbuhl JP. Epithelial M cells: gateways for mucosal infection and immunization. *Cell.* 1996; 86:345–8. [PubMed: 8756716]
- 56Basic M, Keubler LM, Buettner M, Achard M, Breves G, et al. Norovirus triggered microbiota-driven mucosal inflammation in interleukin 10-deficient mice. *Inflamm Bowel Dis.* 2014; 20:431–43. [PubMed: 24487272]
- 57Makino A, Shimojima M, Miyazawa T, Kato K, Tohya Y, Akashi H. Junctional adhesion molecule 1 is a functional receptor for feline calicivirus. *J Virol.* 2006; 80:4482–90. [PubMed: 16611908]
- 58Taube S, Jiang M, Wobus CE. Glycosphingolipids as receptors for non-enveloped viruses. *Viruses.* 2010; 2:1011–49. [PubMed: 21994669]
- 59Ruvoen-Clouet N, Belliot G, Le Pendu J. Noroviruses and histo-blood groups: the impact of common host genetic polymorphisms on virus transmission and evolution. *Rev Med Virol.* 2013; 23:355–66. [PubMed: 23959967]
- 60Tan M, Jiang X. Norovirus-host interaction: multi-selections by human histo-blood group antigens. *Trends Microbiol.* 2011; 19:382–8. [PubMed: 21705222]
- 61Zakhour M, Ruvoen-Clouet N, Charpilienne A, Langpap B, Poncet D, et al. The alphaGal epitope of the histo-blood group antigen family is a ligand for bovine norovirus Newbury2 expected to prevent cross-species transmission. *PLoS Pathog.* 2009; 5:e1000504. [PubMed: 19578439]
- 62Caddy S, Breiman A, le Pendu J, Goodfellow I. Genogroup IV and VI canine noroviruses interact with histo-blood group antigens. *J Virol.* 2014; 88:10377–91. [PubMed: 25008923]
- 63Singh BK, Glatt S, Ferrer JL, ADK, Leuthold MM, et al. Structural analysis of a feline norovirus protruding domain. *Virology.* 2014; 474C:181–85.
- 64Tamura M, Natori K, Kobayashi M, Miyamura T, Takeda N. Genogroup II noroviruses efficiently bind to heparan sulfate proteoglycan associated with the cellular membrane. *J Virol.* 2004; 78:3817–26. [PubMed: 15047797]
- 65Rydell GE, Nilsson J, Rodriguez-Diaz J, Ruvoen-Clouet N, Svensson L, et al. Human noroviruses recognize sialyl Lewis x neoglycoprotein. *Glycobiology.* 2009; 19:309–20. [PubMed: 19054801]
- 66Bally M, Rydell GE, Zahn R, Nasir W, Eggeling C, et al. Norovirus GII.4 virus-like particles recognize galactosylceramides in domains of planar supported lipid bilayers. *Angew Chem Int Ed Engl.* 2012; 51:12020–4. [PubMed: 23097253]
- 67Taube S, Perry JW, McGreevy E, Yetming K, Perkins C, et al. Murine noroviruses bind glycolipid and glycoprotein attachment receptors in a strain-dependent manner. *J Virol.* 2012; 86:5584–93. [PubMed: 22438544]
- 68Taube S, Perry JW, Yetming K, Patel SP, Auble H, et al. Ganglioside-linked terminal sialic acid moieties on murine macrophages function as attachment receptors for murine noroviruses. *J Virol.* 2009; 83:4092–101. [PubMed: 19244326]

- 69Tan M, Jiang X. Histo-blood group antigens: a common niche for norovirus and rotavirus. *Expert Rev Mol Med*. 2014; 16:e5. [PubMed: 24606759]
- 70Taube S, Rubin JR, Katpally U, Smith TJ, Kendall A, et al. High-resolution x-ray structure and functional analysis of the murine norovirus 1 capsid protein protruding domain. *J Virol*. 2010; 84:5695–705. [PubMed: 20335262]
- 71Kolawole AO, Li M, Xia C, Fischer AE, Giacobbi NS, et al. Flexibility in surface-exposed loops in a virus capsid mediates escape from antibody neutralization. *J Virol*. 2014; 88:4543–57. [PubMed: 24501415]
- 72Chen Z, Sosnovtsev SV, Bok K, Parra GI, Makiya M, et al. Development of Norwalk virus-specific monoclonal antibodies with therapeutic potential for the treatment of Norwalk virus gastroenteritis. *J Virol*. 2013; 87:9547–57. [PubMed: 23785216]
- 73Le Pendu J. Histo-blood group antigen and human milk oligosaccharides: genetic polymorphism and risk of infectious diseases. *Adv Exp Med Biol*. 2004; 554:135–43. [PubMed: 15384573]
- 74Guix S, Asanaka M, Katayama K, Crawford SE, Neill FH, et al. Norwalk virus RNA is infectious in mammalian cells. *J Virol*. 2007; 81:12238–48. [PubMed: 17855551]
- 75Rydell GE, Dahlin AB, Hook F, Larson G. QCM-D studies of human norovirus VLPs binding to glycosphingolipids in supported lipid bilayers reveal strain-specific characteristics. *Glycobiology*. 2009; 19:1176–84. [PubMed: 19625485]
- 76Rydell GE, Svensson L, Larson G, Johannes L, Romer W. Human GII.4 norovirus VLP induces membrane invaginations on giant unilamellar vesicles containing secretor gene dependent alpha1,2-fucosylated glycosphingolipids. *Biochim Biophys Acta*. 2013; 1828:1840–5. [PubMed: 23528203]
- 77Oriol R, Le Pendu J, Sparkes RS, Sparkes MC, Crist M, et al. Insights into the expression of ABH and Lewis antigens through human bone marrow transplantation. *Am J Hum Genet*. 1981; 33:551–60. [PubMed: 6167165]
- 78Wilks J, Golovkina T. Influence of microbiota on viral infections. *PLoS Pathog*. 2012; 8:e1002681. [PubMed: 22615558]
- 79Robinson CM, Pfeiffer JK. Viruses and the Microbiota. *Annu Rev Virol*. 2014; 1:55–69. [PubMed: 25821837]
- 80Kuss SK, Best GT, Etheredge CA, Pruijssers AJ, Frierson JM, et al. Intestinal microbiota promote enteric virus replication and systemic pathogenesis. *Science*. 2011; 334:249–52. [PubMed: 21998395]
- 81Kane M, Case LK, Kopaskie K, Kozlova A, MacDermid C, et al. Successful transmission of a retrovirus depends on the commensal microbiota. *Science*. 2011; 334:245–9. [PubMed: 21998394]
- 82Uchiyama R, Chassaing B, Zhang B, Gewirtz AT. Antibiotic treatment suppresses rotavirus infection and enhances specific humoral immunity. *J Infect Dis*. 2014; 210:171–82. [PubMed: 24436449]
- 83Robinson CM, Jesudhasan PR, Pfeiffer JK. Bacterial lipopolysaccharide binding enhances virion stability and promotes environmental fitness of an enteric virus. *Cell Host Microbe*. 2014; 15:36–46. [PubMed: 24439896]
- 84Baldridge MT, Nice TJ, McCune BT, Yokoyama CC, Kambal A, et al. Commensal microbes and interferon-lambda determine persistence of enteric murine norovirus infection. *Science*. 2014
- 85Kernbauer E, Ding Y, Cadwell K. An enteric virus can replace the beneficial function of commensal bacteria. *Nature*. 2014; 516:94–8. [PubMed: 25409145]
- 86Miura T, Sano D, Suenaga A, Yoshimura T, Fuzawa M, et al. Histo-blood group antigen-like substances of human enteric bacteria as specific adsorbents for human noroviruses. *J Virol*. 2013; 87:9441–51. [PubMed: 23804639]
- 87Karst SM, Wobus CE. A Working Model of how Noroviruses Infect the Intestine. *PLoS Pathog*. 2015 in press.
- 88Atmar RL, Opekun AR, Gilger MA, Estes MK, Crawford SE, et al. Norwalk virus shedding after experimental human infection. *Emerg Infect Dis*. 2008; 14:1553–7. [PubMed: 18826818]
- 89Murata T, Katsushima N, Mizuta K, Muraki Y, Hongo S, Matsuzaki Y. Prolonged norovirus shedding in infants <or=6 months of age with gastroenteritis. *Pediatr Infect Dis J*. 2007; 26:46–9. [PubMed: 17195705]

- 90Bok K, Green KY. Norovirus gastroenteritis in immunocompromised patients. *N Engl J Med*. 2012; 367:2126–32. [PubMed: 23190223]
- 91Siebenga JJ, Beersma MF, Vennema H, van Biezen P, Hartwig NJ, Koopmans M. High Prevalence of Prolonged Norovirus Shedding and Illness among Hospitalized Patients: A Model for In Vivo Molecular Evolution. *J Infect Dis*. 2008; 198:994–1001. [PubMed: 18774885]
- 92Bull RA, Eden JS, Luciani F, McElroy K, Rawlinson WD, White PA. Contribution of intra- and interhost dynamics to norovirus evolution. *J Virol*. 2012; 86:3219–29. [PubMed: 22205753]
- 93Arias A, Bailey D, Chaudhry Y, Goodfellow I. Development of a reverse-genetics system for murine norovirus 3: long-term persistence occurs in the caecum and colon. *J Gen Virol*. 2012; 93:1432–41. [PubMed: 22495235]
- 94Nice TJ, Strong DW, McCune BT, Pohl CS, Virgin HW. A single-aminoacid change in murine norovirus NS1/2 is sufficient for colonic tropism and persistence. *J Virol*. 2013; 87:327–34. [PubMed: 23077309]
- 95Tomov VT, Osborne LC, Dolfi DV, Sonnenberg GF, Monticelli LA, et al. Persistent enteric murine norovirus infection is associated with functionally suboptimal virus-specific CD8 T cell responses. *J Virol*. 2013; 87:7015–31. [PubMed: 23596300]
- 96Nice TJ, Baldrige MT, McCune BT, Norman JM, Lazear HM, et al. Interferon lambda cures persistent murine norovirus infection in the absence of adaptive immunity. *Science*. 2014
- 97Haque R, Snider C, Liu Y, Ma JZ, Liu L, et al. Oral polio vaccine response in breast fed infants with malnutrition and diarrhea. *Vaccine*. 2014; 32:478–82. [PubMed: 24300591]
- 98Qadri F, Bhuiyan TR, Sack DA, Svennerholm AM. Immune responses and protection in children in developing countries induced by oral vaccines. *Vaccine*. 2013; 31:452–60. [PubMed: 23153448]
- 99von Bubnoff A. A gut response to vaccines. *IAVI Rep*. 2011; 15:12–4.
- 100Beck MA, Handy J, Levander OA. Host nutritional status: the neglected virulence factor. *Trends Microbiol*. 2004; 12:417–23. [PubMed: 15337163]
- 101Hickman D, Jones MK, Zhu S, Kirkpatrick E, Ostrov DA, et al. The effect of malnutrition on norovirus infection. *MBio*. 2014; 5:e01032–13. [PubMed: 24595373]
- 102Osborne LC, Monticelli LA, Nice TJ, Sutherland TE, Siracusa MC, et al. Coinfection. Virus-helminth coinfection reveals a microbiota-independent mechanism of immunomodulation. *Science*. 2014; 345:578–82. [PubMed: 25082704]
- 103Chang KO, Sosnovtsev SV, Belliot G, King AD, Green KY. Stable expression of a Norwalk virus RNA replicon in a human hepatoma cell line. *Virology*. 2006; 353:463–73. [PubMed: 16843517]
- 104Arias A, Emmott E, Vashist S, Goodfellow I. Progress towards the prevention and treatment of norovirus infections. *Future Microbiol*. 2013; 8:1475–87. [PubMed: 24199805]
- 105Rocha-Pereira J, Neyts J, Jochmans D. Norovirus: targets and tools in antiviral drug discovery. *Biochem Pharmacol*. 2014; 91:1–11. [PubMed: 24893351]
- 106Zeitler CE, Estes MK, Venkataram Prasad BV. X-ray crystallographic structure of the Norwalk virus protease at 1.5-Å resolution. *J Virol*. 2006; 80:5050–8. [PubMed: 16641296]
- 107Ng KK, Pendas-Franco N, Rojo J, Boga JA, Machin A, et al. Crystal structure of norwalk virus polymerase reveals the carboxyl terminus in the active site cleft. *J Biol Chem*. 2004; 279:16638–45. [PubMed: 14764591]
- 108Rocha-Pereira J, Jochmans D, Dallmeier K, Leyssen P, Cunha R, et al. Inhibition of norovirus replication by the nucleoside analogue 2'-C-methylcytidine. *Biochem Biophys Res Commun*. 2012; 427:796–800. [PubMed: 23063849]
- 109Costantini VP, Whitaker T, Barclay L, Lee D, McBrayer TR, et al. Antiviral activity of nucleoside analogues against norovirus. *Antivir Ther*. 2012; 17:981–91. [PubMed: 22910194]
- 110Rocha-Pereira J, Jochmans D, Debing Y, Verbeken E, Nascimento MS, Neyts J. The Viral Polymerase Inhibitor 2'-C-Methylcytidine Inhibits Norwalk Virus Replication and Protects against Norovirus-Induced Diarrhea and Mortality in a Mouse Model. *J Virol*. 2013; 87:11798–805. [PubMed: 23986582]
- 111Rocha-Pereira J, Jochmans D, Neyts J. Prophylactic treatment with the nucleoside analogue 2'-C-methylcytidine completely prevents transmission of norovirus. *J Antimicrob Chemother*. 2014

- 112 Toniutto P, Fabris C, Bitetto D, Fornasiere E, Rapetti R, Pirisi M. Valopicitabine dihydrochloride: a specific polymerase inhibitor of hepatitis C virus. *Curr Opin Investig Drugs*. 2007; 8:150–8.
- 113 Sidwell RW, Barnard DL, Day CW, Smee DF, Bailey KW, et al. Efficacy of orally administered T-705 on lethal avian influenza A (H5N1) virus infections in mice. *Antimicrob Agents Chemother*. 2007; 51:845–51. [PubMed: 17194832]
- 114 Gowen BB, Wong MH, Jung KH, Sanders AB, Mendenhall M, et al. In vitro and in vivo activities of T-705 against arenavirus and bunyavirus infections. *Antimicrob Agents Chemother*. 2007; 51:3168–76. [PubMed: 17606691]
- 115 Morrey JD, Taro BS, Siddharthan V, Wang H, Smee DF, et al. Efficacy of orally administered T-705 pyrazine analog on lethal West Nile virus infection in rodents. *Antiviral Res*. 2008; 80:377–9. [PubMed: 18762216]
- 116 Smither SJ, Eastaugh LS, Steward JA, Nelson M, Lenk RP, Lever MS. Post-exposure efficacy of oral T-705 (Favipiravir) against inhalational Ebola virus infection in a mouse model. *Antiviral Res*. 2014; 104:153–5. [PubMed: 24462697]
- 117 Rocha-Pereira J, Jochmans D, Dallmeier K, Leyssen P, Nascimento MS, Neyts J. Favipiravir (T-705) inhibits in vitro norovirus replication. *Biochem Biophys Res Commun*. 2012; 424:777–80. [PubMed: 22809499]
- 118 Arias A, Thorne L, Goodfellow I. Favipiravir elicits antiviral mutagenesis during virus replication in vivo. *Elife*. 2014; 3:e03679. [PubMed: 25333492]
- 119 Jin Z, Smith LK, Rajwansi VK, Kim B, Deval J. The ambiguous base-pairing and high substrate efficiency of T-705 (Favipiravir) Ribofuranosyl 5'-triphosphate towards influenza A virus polymerase. *PLoS One*. 2013; 8:e68347. [PubMed: 23874596]
- 120 Furuta Y, Gowen BB, Takahashi K, Shiraki K, Smee DF, Barnard DL. Favipiravir (T-705), a novel viral RNA polymerase inhibitor. *Antiviral Res*. 2013; 100:446–54. [PubMed: 24084488]
- 121 Kapuria V, Peterson LF, Fang D, Bornmann WG, Talpaz M, Donato NJ. Deubiquitinase inhibition by small-molecule WP1130 triggers aggresome formation and tumor cell apoptosis. *Cancer Res*. 2010; 70:9265–76. [PubMed: 21045142]
- 122 Perry JW, Ahmed M, Chang KO, Donato NJ, Showalter HD, Wobus CE. Antiviral activity of a small molecule deubiquitinase inhibitor occurs via induction of the unfolded protein response. *PLoS Pathog*. 2012; 8:e1002783. [PubMed: 22792064]
- 123 Hayashimoto N, Morita H, Ishida T, Uchida R, Tanaka M, et al. Microbiological survey of mice (*Mus musculus*) purchased from commercial pet shops in Kanagawa and Tokyo, Japan. *Exp Anim*. 2014
- 124 Kastenmayer RJ, Perdue KA, Elkins WR. Eradication of murine norovirus from a mouse barrier facility. *J Am Assoc Lab Anim Sci*. 2008; 47:26–30. [PubMed: 18210995]
- 125 Larsson E, Tremaroli V, Lee YS, Koren O, Nookaew I, et al. Analysis of gut microbial regulation of host gene expression along the length of the gut and regulation of gut microbial ecology through MyD88. *Gut*. 2012; 61:1124–31. [PubMed: 22115825]
- 126 Paik J, Fierce Y, Mai PO, Phelps SR, McDonald T, et al. Murine norovirus increases atherosclerotic lesion size and macrophages in *Ldlr*($-/-$) mice. *Comp Med*. 2011; 61:330–8. [PubMed: 22330248]
- 127 Paik J, Fierce Y, Drivdahl R, Treuting PM, Seamons A, et al. Effects of murine norovirus infection on a mouse model of diet-induced obesity and insulin resistance. *Comp Med*. 2010; 60:189–95. [PubMed: 20579433]
- 128 Lencioni KC, Drivdahl R, Seamons A, Treuting PM, Brabb T, Maggio-Price L. Lack of effect of murine norovirus infection on a mouse model of bacteria-induced colon cancer. *Comp Med*. 2011; 61:219–26. [PubMed: 21819691]
- 129 Compton SR, Paturzo FX, Macy JD. Effect of murine norovirus infection on mouse parvovirus infection. *J Am Assoc Lab Anim Sci*. 2010; 49:11–21. [PubMed: 20122310]
- 130 Hensley SE, Pinto AK, Hickman HD, Kastenmayer RJ, Bennink JR, et al. Murine norovirus infection has no significant effect on adaptive immunity to vaccinia virus or influenza A virus. *J Virol*. 2009

- 131Doom CM, Turula HM, Hill AB. Investigation of the impact of the common animal facility contaminant murine norovirus on experimental murine cytomegalovirus infection. *Virology*. 2009; 392:153–61. [PubMed: 19647849]
- 132Ammann CG, Messer RJ, Varvel K, Debuysscher BL, Lacasse RA, et al. Effects of acute and chronic murine norovirus infections on immune responses and recovery from Friend retrovirus infection. *J Virol*. 2009; 83:13037–41. [PubMed: 19812147]
- 133Hampe J, Franke A, Rosenstiel P, Till A, Teuber M, et al. A genome-wide association scan of nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. *Nat Genet*. 2007; 39:207–11. [PubMed: 17200669]
- 134Rioux JD, Xavier RJ, Taylor KD, Silverberg MS, Goyette P, et al. Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nat Genet*. 2007; 39:596–604. [PubMed: 17435756]
- 135Cadwell K, Patel KK, Komatsu M, Virgin HW, Stappenbeck TS. A common role for Atg16L1, Atg5 and Atg7 in small intestinal Paneth cells and Crohn disease. *Autophagy*. 2009; 5:250–2. [PubMed: 19139628]
- 136Cadwell K, Liu JY, Brown SL, Miyoshi H, Loh J, et al. A key role for autophagy and the autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature*. 2008; 456:259–63. [PubMed: 18849966]
- 137Cadwell K. Crohn's disease susceptibility gene interactions, a NOD to the newcomer ATG16L1. *Gastroenterology*. 2010; 139:1448–50. [PubMed: 20875485]
- 138Cadwell K, Stappenbeck TS, Virgin HW. Role of autophagy and autophagy genes in inflammatory bowel disease. *Curr Top Microbiol Immunol*. 2009; 335:141–67. [PubMed: 19802564]
- 139Lencioni KC, Seamons A, Treuting PM, Maggio-Price L, Brabb T. Murine norovirus: an intercurrent variable in a mouse model of bacteria-induced inflammatory bowel disease. *Comp Med*. 2008; 58:522–33. [PubMed: 19149409]
- 140Hsu CC, Paik J, Treuting PM, Seamons A, Meeker SM, et al. Infection with murine norovirus 4 does not alter Helicobacter-induced inflammatory bowel disease in Il10(–/–) mice. *Comp Med*. 2014; 64:256–63. [PubMed: 25296012]
- 141Higgins PD, Johnson LA, Sauder K, Moons D, Blanco L, et al. Transient or persistent norovirus infection does not alter the pathology of Salmonella typhimurium induced intestinal inflammation and fibrosis in mice. *Comp Immunol Microbiol Infect Dis*. 2011; 34:247–57. [PubMed: 21237511]
- 142Abt MC, Artis D. The intestinal microbiota in health and disease: the influence of microbial products on immune cell homeostasis. *Curr Opin Gastroenterol*. 2009; 25:496–502. [PubMed: 19770652]
- 143Brestoff JR, Artis D. Commensal bacteria at the interface of host metabolism and the immune system. *Nat Immunol*. 2013; 14:676–84. [PubMed: 23778795]
- 144Kau AL, Ahern PP, Griffin NW, Goodman AL, Gordon JI. Human nutrition, the gut microbiome and the immune system. *Nature*. 2011; 474:327–36. [PubMed: 21677749]
- 145Nelson AM, Elftman MD, Pinto AK, Baldrige M, Hooper P, et al. Murine norovirus infection does not cause major disruptions in the murine intestinal microbiota. *Microbiome*. 2013; 1:7. [PubMed: 24451302]
- 146Nelson AM, Walk ST, Taube S, Taniuchi M, Houghton ER, et al. Disruption of the Human Gut Microbiota following Norovirus Infection. *PLoS ONE*. 2012; 7:e48224. [PubMed: 23118957]
- 147Power SE, O'Toole PW, Stanton C, Ross RP, Fitzgerald GF. Intestinal microbiota, diet and health. *Br J Nutr*. 2014; 111:387–402. [PubMed: 23931069]
- 148Schippa S, Conte MP. Dysbiotic Events in Gut Microbiota: Impact on Human Health. *Nutrients*. 2014; 6:5786–805. [PubMed: 25514560]
- 149Cao S, Lou Z, Tan M, Chen Y, Liu Y, et al. Structural basis for the recognition of blood group trisaccharides by norovirus. *J Virol*. 2007; 81:5949–57. [PubMed: 17392366]
- 150Hao N, Chen Y, Xia M, Tan M, Liu W, et al. Crystal structures of GI.8 Boxer virus P dimers in complex with HBGAs, a novel evolutionary path selected by the Lewis epitope. *Protein Cell*. 2014

- 151 Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol.* 2011; 7:539. [PubMed: 21988835]
- 152 Shanker S, Czako R, Sankaran B, Atmar RL, Estes MK, Prasad BV. Structural analysis of determinants of histo-blood group antigen binding specificity in genogroup I noroviruses. *J Virol.* 2014; 88:6168–80. [PubMed: 24648450]

Summary Points

1. The MuNoV system has made significant contributions to a greater understanding of NoV biology.
2. MuNoV studies have driven the identification of B cells as NoV targets and commensal bacteria as co-factors for NoV infection.
3. While the specific glycan structures used during NoV infection display a wide degree of virus strain variability, carbohydrate usage by NoVs is highly conserved.
4. NoVs have evolved specific features to successfully infect and spread in the intestinal ecosystem dominated by bacteria.
5. MuNoV infections have a wide range of effects on biomedical research models.

Future Issues

1. MuNoV and HuNoV share many similarities but additional comparative studies are needed to better characterize the extent of similarities and differences and identify the unifying principles in NoV biology.
2. Studies investigating the mechanisms of NoV disease pathogenesis are needed.
3. The field has a need for improved animal models of HuNoV disease that incorporate the microbiota and a full complement of immune functions.
4. Future research should aim to gain a mechanistic understanding regarding the role(s) of carbohydrates and enteric bacteria during NoV infection.
5. Studies are needed to identify HuNoV target cells *in vivo* and to develop new or more robust cell culture models.

Fig. 1B

Figure 1. Norovirus genomic organization and capsid structure. A)

NoV genomic organization. Open reading frame (ORF) 1–3 are found in all NoVs, while ORF4 is only found in MuNoVs. **B)** Cryo-electron microscopy image reconstruction of MNV-1. Reconstructions are colored according to the radial position. View is into an icosahedral five-fold axis. P2 subdomain, P1 subdomain, and shell domain are colored blue, green and yellow/red, respectively.

Fig. 2A-D

Fig. 2E

[illegible]

Fig. 2F

Figure 2. Common features of murine and human NoV glycan binding site. A-D)

The norovirus glycan binding site location is conserved. Surface representations of P dimers from **(A)** MNV-1 (PDB 3LQ6), **(B)** GII.4 VA387 (PDB 2OBS), **(C)** GI.2 FUV258 (PDB 3ASP), and **(D)** GI.7 TCH060 (PDB 4P12) are shown. The monomeric units are colored in shades of grey. The glycan binding site is shown in red. The functional MNV-1 glycan binding site is based on (67), the VA387 glycan binding site is from (149), and the GI binding sites are from Figure 2 of (150). **E–F** Antigenic loops of noroviruses are highly flexible. **E** A multiple sequence alignment of P domain sequences from representative NoVs was generated with ClustalOmega (151). Numbers represent amino acid numbers of the P domain. P domain structures of MNV-1 (PDB 3LQ6), GII.4 VA387 (PDB 2OBS), GI.2 FUV258 (PDB 3ASP), GI.7 TCH060 (PDB 4P12) and Norwalk virus VP1 (PDB 1IHM) were viewed in PyMol (PyMOL Molecular Graphics System, Version 1.3 Schrödinger,

LLC.). Residues forming loops are highlighted in yellow for MNV-1 and green for HuNoVs and corresponding loop names are shown on the top or bottom, respectively. Loop nomenclature is based on (70) and (152) for MNV- 1 and HuNoVs, respectively. For GI HuNoVs, T and U loops form one large unstructured region in Pymol (light and dark green shaded sequences). **F)** Comparison of open versus closed loop conformation. The open conformation is shown on the left with the indicated loops from MNV-1 (PDB 3LQ6, A chain) in turquoise and HuNoV GI.7 TCH 060 (PDB 4P12, C chain) in light green. The closed conformation is shown on the right with MNV-1 (PDB 3LQ6, B chain) in orange, and HuNoVs GII.4 VA387 (PDB 2OBS) in purple and GI.2 FUV258 (PDB 3ASP, A chain) in grey.

Figure 3A

Figure 3B **Figure 3. Commensal bacteria promote norovirus infections. A)**

MuNoV infection in Abx-treated or germ-free mice (right panel) is significantly reduced compared to infection of microbially colonized mice (left panel), demonstrating that commensal bacteria stimulate NoV infections *in vivo* by an unknown mechanism. Because infection of germ-free mice is not completely ablated, bacteria-independent mechanisms exist to breach the intestinal epithelium and access target immune cells. This is represented by free virus interacting with target cells. **B)** A model for bacterial stimulation of HuNoV B cell infection is based on Jones et al. (34). In this model, (1) a HuNoV binds commensal bacteria expressing the appropriate HBGA in the gut lumen; (2) the virus-bacteria complex is co-transcytosed across the polarized epithelium; (3) and the bacterial glycan stimulates viral attachment to permissive B cells. This model is based on data showing filtration-based removal of bacteria from a virus-positive stool inoculum significantly reduced HuNoV infection of B cells in both direct B cell infection and in a co-culture system where the viral inoculum (placed into the apical supernatant) and target B cells (cultured in the basolateral chamber) were separated by a layer of polarized IECs.

Figure 4A **Figure 4B** **Figure 4. Commensal bacterial interactions may drive the immune response during norovirus infections. A)**

Commensal bacteria stimulate a tolerogenic IL-10-dependent microenvironment that suppresses inflammation to MuNoV infection. Basic et al. (56) demonstrated that MuNoV infection is inflammatory in IL-10^{-/-}, but not wild-type mice; and that inflammation in the absence of IL-10 was dependent on commensal bacteria since it did not occur in germ-free IL-10^{-/-} mice. Together with the observations of Jones et al. (34), Baldrige et al. (84), and Kernbauer et al. (85) revealing that commensal bacteria stimulate MuNoV infection, we propose the following model. (i) In wild-type (WT) mice with a normal intestinal microbiota, commensal bacteria stimulate MuNoV infection of immune cells (1);

commensal bacteria-specific regulatory T cells (Tregs) recognize bacterial antigens (2) and secrete IL-10, which prevents virus-induced inflammation (3). There is currently no evidence that Tregs are producers of IL-10 in this model, and future studies will be required to determine the cellular source of this immunosuppressive cytokine. (ii) In microbially colonized mice deficient in IL-10 expression, viral infection proceeds normally since stimulatory commensal bacteria are present (1). However because Tregs cannot secrete IL-10 in response to bacterial antigen recognition (2), immune responses to the virus are not suppressed and instead induce an inflammatory response (3). (iii) In germ-free IL-10^{-/-} mice, viral infection is significantly reduced due to the absence of stimulatory commensal bacteria (1); thus no inflammation is observed. **B)** Based on findings by Nice et al. (96), we propose a model whereby acute NoV strains, such as MNV-1, induce IFN λ (1); IFN λ signals to IECs or other non-immune cell (2); this indirectly reduces viral replication and prevents persistence (3). Persistent MuNoV strains such as MNV-CR6 fail to induce significant IFN λ , likely a key strategy for persistence establishment.