Research Article

Periodic limb movements in tetraplegia

Annemieke Emma Josina Peters^{1,2}, Lysanne van Silfhout^{1,2}, Marnie Graco ¹, Rachel Schembri¹, Dick Thijssen¹, David J. Berlowitz¹

¹Institute for Breathing and Sleep, Austin Health, Melbourne, Australia, ²Department of Physiology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

Objective: To establish the prevalence of Periodic Limb Movements during Sleep (PLMS) in patients with tetraplegia, controlling for obstructive sleep apnea. To explore whether demographic and injury characteristics affect PLMS.

Study Design: Retrospective cohorts.

Setting and Participants: One hundred seventy-three participants with acute (< 12 months) and 92 with chronic (> 12 months) tetraplegia who underwent full overnight diagnostic sleep studies.

Interventions and outcome measures: Two hundred sixty-two sleep study recordings were included. A randomly selected subgroup of 21 studies was assessed for PLM during wakefulness. Data were analysed according to the current American Academy of Sleep Medicine guidelines.

Results: Of the participants, 41.6% (43(15.7) years and 14.9% female) had a motor and sensory complete lesion. Sleep was poor with both OSA (87.8% with apnea hypopnoea index ≥ 5) and PLMS (58.4% with PLMS per hour PLMSI > 15) highly prevalent. There was no difference in the PLMSI between those with OSA (36.3(39.8)) or without (42.2(37.7), $P = 0.42$). PLMS were evident during REM and NREM sleep in all of the 153 patients with PLMSI > 15. All 21 participants in the subgroup of studies analysed for the PLM during quiet wakefulness, exhibited limb movements. None of the modelled variables (injury completeness, gender, OSA severity or time since injury) significantly predicted a PLMSI > 15 ($P = 0.343$).

Conclusion: In conclusion, this study confirms the high prevalence of PLM in tetraplegia and the presence of leg movements in NREM and REM sleep along with wakefulness after controlling for OSA. No associations between the presence of PLMS and patient characteristics or injury specific aspects were found.

Keywords: Sleep, Periodic limb movements of sleep, Cervical spinal cord injury, Obstructive sleep apnea, Tetraplegia

Introduction

Sleep problems are frequently reported in people with a spinal cord injury (SCI).^{1–3} It appears that obstructive sleep apnea (OSA) is the predominant disorder in tetraplegia,² whereas muscle spasms and somatic disturbances such as pain and paraesthesia additionally contribute to sleep disturbance in paraplegia and incomplete lesions.¹

Periodic Limb Movements are a sleep related movement disorder first described by Lugaresi *et al.* in 1972.⁴ The movements are characterized by episodes of repetitive and highly stereotyped periodic limb movements that occur during sleep. These movements may be related to

arousals, obstructive sleep apnea (OSA) and awakenings⁵ and can therefore negatively impact sleep efficiency.

Periodic Limb Movements of Sleep (PLMS) are indexed per hour to provide the periodic leg movement of sleep index (PLMSI). Excluding those with significant OSA, the prevalence of an elevated PLMSI in the general population is 7.6%.⁶ In the able-bodied population, PLMS typically occur during Non Rapid Eye Movement (NREM) sleep and have a clear periodicity.⁷ This contrasts with SCI where PLMS are highly prevalent in those with lesions above T10,^{8–10} not always clearly periodic and observed throughout quiet wakefulness and REM.¹¹ These SCI data are however limited in their sample sizes and inconsistent in their sampling and findings.^{7,10}

Confounding these previous data is the issue of OSA in SCI. Leg movements are frequently associated with apnea termination and arousal from sleep. To address this, the American Association of Sleep Medicine

Correspondence to: David J. Berlowitz, Institute for Breathing and Sleep, Austin Health, Studley Rd, Heidelberg 3084, Victoria, Australia. Email: david.berlowitz@austin.org.au

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/yscm.

(AASM) scoring rules exclude events that occur during a period from 0.5 seconds prior to an apnea or hypopnoea to 0.5 seconds following a scored respiratory event.¹² Obstructive sleep apnea is apparent within weeks of cervical SCI (tetraplegia),^{13,14} is 2–5 times more prevalent than in the general population^{2,15} and is associated with significant symptoms and reduced quality of life.³ Some PLMS in SCI studies have excluded OSA,¹⁰ others have included participants with OSA and used the AASM rules to exclude PLMS adjacent to respiratory events,¹¹ while some authors make no reference to possible co-existent OSA.^{8,9} To address these limitations in the previous literature and to explore factors that are potentially related to PLMS in tetraplegia we analysed a large sample of polysomnographic records collected in previous sleep research in tetraplegia.^{2,16} This analysis aims to: characterise PLMS in a large sample of people with traumatic tetraplegia resulting from a cervical SCI who have undergone polysomnography (PSG), to confirm PLMS across REM, NREM and wake in tetraplegia and to explore whether the presence of OSA, demographic and injury characteristics were related to the prevalence or severity of PLMS.

Subjects and methods

Subjects

Polysomnographic sleep recordings collected from 92 people with chronic (>12 months) tetraplegia who participated in previous, community based research between July 2007 and March 2008 at Austin Health² and 173 people with acute (<12 months) tetraplegic injuries, previously enrolled in the ‘Continuous positive airway pressure for obstructive sleep apnea in quadriplegia’ (COAQP) trial¹⁶ between September 2009 and February 2013. Both studies were approved by the Austin Health Human Research Ethics Committee. The polysomnographic data were collected for acute and chronic participants after an average of 0.28 (SD=0.16) and 12.32 (7.76) years, respectively, from the SCI. The full description of the sampling strategy and inclusion, exclusion criteria can be found in the original references.^{2,16} For this current analysis, participants were excluded if there was technical failure of the limb movement signal on the PSG or if demographic data were unavailable. Demographic data were obtained from clinical trial and hospital medical records. The level and completeness of the lesion were assessed by the International Standards for Neurological Classification of SCI (ISNCSCI).

Polysomnographic sleep recording

Each participant underwent overnight PSG (Compumedics SomtePSG, Abbotsford, Australia)

carried out in patients’ homes (chronic) or on inpatient wards (acute) as per original study protocols. The PSG consisted of two central electroencephalographic (EEG) channels, bilateral electrooculography (EOG), submentalis muscle electromyogram (EMG), electrocardiogram (ECG), nasal cannulae for flow, abdominal and thoracic bands for respiratory effort. Limb movements were detected using a piezoelectric sensor on the left and right dorsum of each foot and recorded via a single, combined channel. The leg sensors detect movement *per se*, not muscle EEG activity, whereas in-laboratory PSG typically employs direct surface EMG measurement of the tibialis anterior. No nasal thermistor was used. Data were analysed using Compumedics ProfusionPSG 3 software (Compumedics Ltd, Abbotsford, Victoria, Australia).

Scoring of the PSG and the detection of the periodic limb movements

The PLMS were scored and summarized per hour of sleep according to the American Academy of Sleep Medicine (AASM) criteria to generate the PLMSI.¹² Specifically, PLMS were scored when at least four consecutive leg movements during a period of sleep were identified with each movement lasting 0.5–10 seconds and an onset to onset interval of 5–90 seconds. Consecutive leg movement recordings within 0.5 seconds were counted as one. A PLMS was not scored if it occurred within 0.5 seconds of the commencement or the termination of an apnea or a hypopnoea. Sleep was staged, respiratory events and arousals scored according to current AASM guidelines.^{12,17} A PLMSI of greater than 15 was considered pathological.⁶ To determine if there was any effect of the progression of sleep on PLMSI, the PLMSI was calculated separately for each quartile of the night in a randomly selected subset of 29 participants.

The leg movement indices were also calculated during wake in a second, randomly selected subgroup of 21 participants who fulfilled the criteria of having more than 10 minutes of quiet resting prior to sleep onset and after awakening. The periodic leg movement index during wakefulness (PLMWI) was obtained from the time-weighted averages of three segments of PSG; 30 minutes before the first sleep onset, 30 minutes after sleep offset (morning awakening) and 30 minutes within the longest period of wake after sleep onset. If the wake after sleep onset segment was less than 30 minutes, the longest available segment was scored provided at least 10 minutes of data were available.

Statistical analyses

Descriptive statistics (mean and standard deviations (SD)) were calculated for all groups. parametric

(Student *t*-test) and non-parametric (χ^2 test) analyses were performed to assess differences between the acute and chronic participants and between those with and without OSA. A conservative apnea hypopnoea index (AHI) of less than five events per hour was used as the initial cut-off for categorization of “no OSA”. A less conservative cut-off of AHI < 15 was also calculated to illustrate possible differences with the prevalence of PLMSI > 15/hour in the general, able-bodied population.⁶ The population data were plotted alongside the mean and 95% CI of the SCI sample and the SCI sample proportions were compared with the chi-squared statistic. No statistical comparisons were made between the SCI and able-bodied data due to differences in study methodologies. Multivariate logistic regression analysis was performed to test for associations between the presence of PLMSI and sex, AHI, age, injury severity (motor and sensory complete versus incomplete) and time since injury. The distribution of PLMSI across the night (quartiles) were compared using a univariate repeated measures analysis of variance and univariate Pearson *r* correlations were performed to examine the relationships between the arousal index (AI), PLMSI and AHI.

Results

All of the chronic participants' studies were able to be fully analysed, but three of the acute studies were excluded due to technical failures of the leg recording leads. The spinal lesion levels of injury ranged from C1 to T1 and injury severity from ISNCSCI (International Standards for Neurological Classification of SCI) A to D.

Those with chronic injuries had their injuries on average over 12 years earlier than the acute cohort, were older, twice as likely to be women and more likely to have complete injuries (Table 1). No significant

Table 1 Demographic and objective data for the SCI groups.

	Acute (n=170)	Chronic (n=92)	P- value
Mean (SD) age	41.6 (13.0)	46.0 (16.8)	0.03
% female	10.0%	24.2%	0.002
% Complete (ISNCSCI A)	37.0%	50.0%	0.04
Mean time since injury (years)	0.28 (0.16)	12.2 (7.74)	<0.000
Mean PLMSI	36.2 (36.8)	38.4 (44.2)	0.67
% PLMSI > 15	58.8%	57.6%	0.85
Mean AHI	29.7 (23.3)	29.5 (24.3)	0.93
% OSA (AHI > 5)	88.2%	87.0%	0.76

ISNCSCI A - international standards for neurological classification of SCI, motor and sensory complete lesion; AHI = apnea hypopnoea index.

differences were observed in PLMSI, proportions with PLMSI > 15 events per hour, OSA severity or prevalence between the acute and chronic samples. Obstructive sleep apnea prevalence (AHI ≥ 5) was extremely high at over 85%. The significant demographic differences were expected as a consequence of the samples pooled for this study. The outcome variables of primary interest (PLMSI and OSA/AHI) were not different across the acute and chronic cohorts and as such, data were pooled for subsequent analyses.

Sleep was poor in the sample overall (Table 2) with both OSA and PLMS highly prevalent. At least mild OSA (AHI ≥ 5) was present in 87.8%, moderate OSA (AHI ≥ 15) in 64.9%, PLMS (PLMSI > 15) in 58.4% and severe PLMS (PLMSI > 29) in 43.9% of the cohort. Overall sleep architecture was poor with a preponderance (almost 50% in both groups) of Stage 2 NREM sleep, and REM sleep was reduced overall. Ninety seven patients (36.7%) spent less than 15% of TST in REM and in 34 patients, (12.9%) the REM sleep was less than 10% of TST. The average arousal index was mildly elevated at 22.4 events per hour.¹⁸

There was no difference in the PLMSI between those with OSA (AHI ≥ 5, *n* = 230, mean PLMSI (SD) = 36.3(39.8)) or without (AHI < 5, *n* = 32, 42.2(37.7), *P* = 0.42). Similarly, OSA had no effect on the proportion of people with a PLMSI > 15 (OSA *n* = 230, 60%; no OSA *n* = 32, 50%, *P* = 0.30). The overall prevalence of PLMSI > 15 in the tetraplegia cohort was 58.4% (Fig. 1).

No variation in PLM severity was observed over the course of the night. (Quartile 1 average=49.6

Table 2 Sleep summary statistics.

	Mean	SD	Min-Max
Total sleep time (min)	358.2	96.5	34.5-583.5
Wake after sleep onset (min)	105.9	74.6	5-434.5
AHI	29.7	23.6	0-122.7
PLMSI <i>n</i> = 262	37.0	39.5	0-234.9
PLMSI NREM <i>n</i> = 262	38.4	42.9	0-253.7
PLMSI REM <i>n</i> = 262	28.4	34.3	0-166.8
PLMSI (AHI ≤ 5) <i>n</i> = 32	42.2	37.7	0-156.2
PLMSI (AHI ≤ 15) <i>n</i> = 92	37.7	35.6	0-156.2
AI	22.4	15.2	2.1-107.8
REM latency (min)	101.5	69.8	1-464.5
N1 (% TST)	11.1	9.5	0.42-60.2
N2 (% TST)	48.3	11.9	18.4-87.4
N3&N4 (% TST)	23.5	12.8	0-56.8
REM (% TST)	17.2	7.2	0-41.2

AHI, apnea hypopnoea index; REM, rapid eye movement sleep; NREM, non-rapid eye movement sleep; N1 = stage one sleep; N2 = stage two sleep; N3&N4 = stage three and four (slow wave) sleep; REM latency = time to first REM after sleep onset, TST, total sleep time; OSA, obstructive sleep apnea; PLMSI, periodic leg movement sleep index (events per hour).

Figure 1 Illustration of periodic leg movements during sleep index (PLMSI) prevalence in the able-bodied and people with tetraplegia. Proportions and 95% confidence intervals for the able bodied (from Scofield 2008, $n = 593$)⁶ and all of the tetraplegia sample ($n = 262$) and those with an AHI < 15 ($n = 170$). Able-bodied participants with sleep apnea as defined by AHI ≥ 15 were excluded from the Scofield publication. Note that differences in PLM measurement methodologies between the current data and the able-bodied prevalence do not allow for direct comparisons.

(SD = 54), Q2 46.4(53), Q3 55.3(57), Q4 46.1(46.3), $F = 0.44$, $P = 0.72$. Of the 153 patients with PLMSI > 15 , PLMS were evident during REM and NREM sleep in all. All 21 participants in the subgroup of studies analysed for the PLM during resting wakefulness, exhibited limb movements that could be classified according to the criteria detailed in the methods. The average PLM index during wake was 92.9 (SD = 52.2) events per hour (Figures 2, 3 & 4).

None of the modelled variables (injury completeness, gender, AHI or time since injury) significantly predicted a PLMSI > 15 (model significance $P = 0.343$). The variable coefficients (and associated P value) were; motor and sensory completeness -0.18 , ($P = 0.5$); AHI 0.002 (0.72); age (years) 0.01 (0.24); sex (male = yes) 0.68 (0.057); time since injury (years) 0.0058 (0.75). There was a significant correlation between the AHI and the frequency of arousals from sleep (AI, $r = 0.61$, $P < 0.001$) but no relationship between PLMSI and the AI ($r = -0.07$, $P = 0.24$) or between the PLMSI and the AHI ($r = -0.04$, $P = 0.50$).

Discussion

The present study aimed to document the prevalence and severity of PLM during sleep in a large cohort of

people with tetraplegia and to examine whether PLM were related to time since injury, obstructive sleep apnea, lesion level or injury severity. This study confirmed that PLMS are highly prevalent in tetraplegia and further, the current study controlled for the highly prevalent confounder of OSA in this same population.¹⁰ The current study found that the PLMSI was not affected by OSA severity and that the arousal index was associated with the AHI but not the PLMSI.

Leg movements were observed in people with tetraplegia during REM and wake. This finding supports prior studies that have documented leg movements during sleep in patients with SCI that continued during wakefulness.^{10,11,19} These findings raise the question of whether the leg movements occurring during wake could be classified as Periodic Limb Movements, given that the AASM defines Periodic Limb Movements as occurring only during sleep.¹⁷ In addition, the studies from which these data were derived did not assess restless leg syndrome specific symptoms and as such we are unable to meaningfully differentiate the observed PLM during wake from the restless leg syndrome.

Although the aetiology of PLMS remains unclear, the findings of PLMS in NREM, REM and wakefulness may suggest that PLMS activity is not exclusively triggered by hypothalamic sleep-wake rhythms in patients with SCI, as is suggested in the able-bodied. While in the able-bodied, PLMS only appear during NREM, the current finding in SCI supports the suggestion that the movement itself may be derived from spinal cord central pattern generation rather than solely a brain derived signal.¹⁹ In SCI the reflex movements are clearly no longer controlled by the descending axonal pathway inhibitory stimuli, which potentially cause a motor neuron overexpression during all stages of sleep and wakefulness. This is consistent with an inhibitory effect controlled by the hypothalamic sleep-wake rhythm, which in able-bodied, with intact descending axonal pathway, suppresses PLMS during wake and REM.¹⁹

Findings in radiology and pharmacology have provided indirect evidence of the involvement of the dopaminergic system in the pathogenesis of PLMS and restless leg syndrome.²⁰ Iron is an essential co-factor for the formation of L-dopa, the precursor of dopamine. Iron deficiency in the multiregional brain, mainly in the substantia nigra, has been shown to correlate with the presence of restless legs,²¹ however differences in dopamine and iron-related markers (specifically, a low ferritin) have also been found in the cerebrospinal fluid of restless leg syndrome patients.^{22,23} Iron deficient mice and knock

out iron-deficient spinal dopamine D(3)-receptor mice support this theory by showing increased locomotor activity resembling restless or PLMS.²⁴ Whether this mechanism may also play a role in the aetiology of PLMS is still unknown. The high prevalence of PLMS in patients with SCI in this study may support this theory, due to a high prevalence of malnutrition, including iron deficiencies, in patients with SCI.²⁵ The current paper was not designed to directly assess this hypothesis however, and as such this remains speculation.

This study has not been able to find a pattern in the appearance of PLMS over the course of the night, supporting previous data that clear periodicity is not always observed in people with SCI and PLM.⁷ Additionally, there were no apparent relationships between demographic factors such as gender, age or injury (AIS, severity and time since injury) and the presence of PLMS. This further suggests that the cervical spinal injury *per se* is the causative factor in the tetraplegia population.

Several studies have reported that people with SCI experience substantially worse subjective sleep than their able-bodied counterparts^{1,3} and in this study the percentage of wakefulness after sleep onset was

approximately 30% of total sleep time compared to 5% in the general population.²⁶ Without a matched control group and identical methods of assessment however, it is not possible to precisely estimate the difference between the SCI sample and the able-bodied population. Further, the majority of able-bodied population PLMS prevalence data were generated using EMG assessment of tibialis anterior activity compared with movement of the foot as in the current dataset.^{6,10} Only one study has directly compared the performance of EMG versus piezo-electric movement sensors and while the PLMS indices were comparable, methodological issues preclude direct comparisons.²⁷ While the confidence intervals on the data in Figure 1 suggests that the PLMS prevalence is higher for tetraplegia than the able-bodied, imprecision in the prevalence estimate comparisons made inferential statistical comparisons inappropriate. The primary potential confounder of the study results remains the up to 78% prevalence of OSA in patients with tetraplegia¹⁶ and associated scoring and attribution of association of the observed PLMS. In reviewing the raw data traces for this project, it was apparent that many of the leg movements

Figure 2 Illustration of PLMS during REM sleep. The top pane of the polysomnography is a 30 second time base and the lower 5 minutes. The fine red vertical line indicates the same point on the recording on the upper and lower panels. Note the regular leg movements seemingly unrelated to the apneic events and the proximity of a number of leg movements to the start or the end of scored respiratory events (obstructive apneas). These PLM events would not be scored or included in the PLMSI.

Figure 3 Illustration of PLMS during stage 2 NREM sleep.

Figure 4 Illustration of PLMS during REM and wakefulness. Note the regular PLM that do not appear modified by sleep state (wake or REM) nor by respiratory events.

were coincident with snoring and respiratory events. As such, these leg movements were unable to be scored as PLMS according to the AASM rules.¹⁷ This would be expected to underestimate the PLMSI, yet the PLMSI was markedly elevated and the PLMSI > 15 category was highly prevalent. The high PLMSI persisted in the subset of participants without OSA; in fact, the PLMSI was higher in those without OSA suggesting that the AASM scoring rules lowered the PLMSI in those with OSA (Figs. 2, 3, and 4).

Participants from this study were initially recruited for studies investigating OSA, and the sleep studies were performed without CPAP or other treatments. As such, the clinical significance and impact on daytime function of PLMS in SCI was not able to be addressed directly. The relationships between daytime function and sleep related movement, breathing and circadian disorders are complex in the able-bodied and are only recently being explored after SCI.^{3,28,29} Given the multi-factorial nature of sleep disturbances in SCI, it is likely that future research on PLMS should consider specific treatment studies as a method for “unpacking” symptomology. For example repeat studies on continuous positive airway pressure treatment for OSA in tetraplegia may reveal the contribution of an elevated PLMSI to sleep-related daytime symptoms.³⁰ Future research could also take the opportunity to examine the effect of electrode type and location on PLMS indices. In the current study, we took the opportunity to add to the literature data from a large cohort of sleep studies, however the original samples were not collected with PLMS prevalence estimation as the primary outcome. People with paraplegia are also excluded from our findings, despite there being evidence that PLMS may also be prevalent in those with lesions from T1 to T10.^{9,10}

Only one channel was used to document the leg events detected in both legs. No distinction has thus been able to be made between which leg moved and therefore no data detailing whether the PLMS were bi- or unilateral. Different ratios of mono- and bilateral movements are reported in various disorders³¹ and potentially hemispheric incomplete SCI could provide additional information about PLMS aetiology.

Conclusion

In conclusion, this study confirms the high prevalence of PLM in tetraplegia and the presence of leg movements in NREM and REM sleep along with wakefulness after controlling for OSA. No associations between the presence of PLMS and patient characteristics or injury specific aspects were found.

Funding

This study was supported by a writing scholarship from the Institute for Breathing and Sleep, a grant from the Radboud University Nijmegen and a grant from the Radboud University Nijmegen Medical Centre.

Disclosure statement

The authors declare they have neither relevant financial disclosures nor conflicts of interest.

ORCID

Marnie Graco <http://orcid.org/0000-0001-6048-0147>

References

- 1 Biering-Sorensen F, Biering-Sorensen M. Sleep disturbances in the spinal cord injured: an epidemiological questionnaire investigation, including a normal population. *Spinal Cord* 2001;39(10):505–13.
- 2 Berlowitz DJ, Spong J, Gordon I, Howard ME, Brown DJ. Relationships between objective sleep indices and symptoms in a community sample of people with tetraplegia. *Arch Phys Med Rehabil* 2012;93(7):1246–52.
- 3 Spong J, Graco M, Brown DJ, Schembri R, Berlowitz DJ. Subjective sleep disturbances and quality of life in chronic tetraplegia. *Spinal Cord* 2015;53(8):636–40.
- 4 Lugaresi E, Coccagna G, Mantovani M, Lebrun R. Some periodic phenomena arising during drowsiness and sleep in man. *Electroencephalogr Clin Neurophysiol* 1972;32(6):701–5.
- 5 Sforza E, Jouny C, Ibanez V. Time course of arousal response during periodic leg movements in patients with periodic leg movements and restless legs syndrome. *Clin Neurophysiol* 2003;114(6):1116–24.
- 6 Scofield H, Roth T, Drake C. Periodic limb movements during sleep: population prevalence, clinical correlates, and racial differences. *Sleep* 2008;31(9):1221–7.
- 7 Dickel MJ, Renfrow SD, Moore PT, Berry RB. Rapid eye movement sleep periodic leg movements in patients with spinal cord injury. *Sleep* 1994;17(8):733–8.
- 8 de Mello MT, Esteves AM, Tufik S. Comparison between dopaminergic agents and physical exercise as treatment for periodic limb movements in patients with spinal cord injury. *Spinal Cord* 2004;42(4):218–21.
- 9 de Mello MT, Lauro FA, Silva AC, Tufik S. Incidence of periodic leg movements and of the restless legs syndrome during sleep following acute physical activity in spinal cord injury subjects. *Spinal Cord* 1996;34(5):294–6.
- 10 Ferri R, Proserpio P, Rundo F, Lanza A, Sambusida K, Redaelli T, *et al.* Neurophysiological correlates of sleep leg movements in acute spinal cord injury. *Clin Neurophysiol* 2015;126(2):333–8.
- 11 Proserpio P, Lanza A, Sambusida K, Fratticci L, Frigerio P, Sommariva M, *et al.* Sleep apnea and periodic leg movements in the first year after spinal cord injury. *Sleep Med* 2015;16(1):59–66.
- 12 The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications. 1st ed. Westchester, IL: American Association of Sleep Medicine; 2007.
- 13 Berlowitz DJ, Brown DJ, Campbell DA, Pierce RJ. A longitudinal evaluation of sleep and breathing in the first year after cervical spinal cord injury. *Arch Phys Med Rehabil* 2005;86(6):1193–9.
- 14 Tran K, Hukins C, Geraghty T, Eckert B, Fraser L. Sleep-disordered breathing in spinal cord-injured patients: A short-term longitudinal study. *Respirology* 2009;15(2).
- 15 Sankari A, Bascom A, Oomman S, Badr MS. Sleep disordered breathing in chronic spinal cord injury. *J Clin Sleep Med* 2014;10(1):65–72.
- 16 Berlowitz DJ, Ayas N, Barnes M, Brown DJ, Cistulli P, Geraghty T, *et al.* Auto-titrating continuous positive airway pressure treatment for obstructive sleep apnoea after acute quadriplegia (COSAQ): study protocol for a randomized controlled trial. *Trials* 2013;14:181.

- 17 Berry RB, Budhiraja R, Gottlieb DJ, Gozal D, Iber C, Kapur VK, *et al.* Rules for scoring respiratory events in sleep: update of the 2007 AASM Manual for the Scoring of Sleep and Associated Events. Deliberations of the Sleep Apnea Definitions Task Force of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2012;8(5):597–619.
- 18 Bonnet MH, Arand DL. EEG Arousal Norms by Age. *J Clin Sleep Med* 2007;3(3):271–4.
- 19 Yokota T, Hirose K, Tanabe H, Tsukagoshi H. Sleep-related periodic leg movements (nocturnal myoclonus) due to spinal cord lesion. *J Neurol Sci* 1991;104(1):13–8.
- 20 Hening W, Allen R, Earley C, Kushida C, Picchietti D, Silber M. The treatment of restless legs syndrome and periodic limb movement disorder. *An American Academy of Sleep Medicine Review. Sleep*. 1999;22(7):970–99.
- 21 Godau J, Klose U, Di Santo A, Schweitzer K, Berg D. Multiregional brain iron deficiency in restless legs syndrome. *Mov Disord*. 2008;23(8):1184–7.
- 22 Allen RP, Connor JR, Hyland K, Earley CJ. Abnormally increased CSF 3-Ortho-methyldopa (3-OMD) in untreated restless legs syndrome (RLS) patients indicates more severe disease and possibly abnormally increased dopamine synthesis. *Sleep Med* 2009;10(1):123–8.
- 23 Earley CJ, Connor JR, Beard JL, Malecki EA, Epstein DK, Allen RP. Abnormalities in CSF concentrations of ferritin and transferrin in restless legs syndrome. *Neurology* 2000;54(8):1698–700.
- 24 Dowling P, Klinker F, Stadelmann C, Hasan K, Paulus W, Liebetanz D. Dopamine D3 receptor specifically modulates motor and sensory symptoms in iron-deficient mice. *J Neurosci* 2011;31(1):70–7.
- 25 Dionysiotis Y. Malnutrition in spinal cord injury: more than nutritional deficiency. *J Clin Med Res* 2012;4(4):227–36.
- 26 Carskadon MA, Dement WC, Mitler MM, Roth T, Westbrook PR, Keenan S. Guidelines for the multiple sleep latency test (MSLT): a standard measure of sleepiness. *Sleep* 1986;9(4):519–24.
- 27 Terrill PI, Leong M, Barton K, Freakley C, Downey C, Vannierkerk M, *et al.*, editors. Measuring leg movements during sleep using accelerometry: Comparison with EMG and piezo-electric scored events. 2013 35th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC); 3–7 July 2013.
- 28 Gainche L, Berlowitz DJ, LeGuen M, Ruehland WR, O'Donoghue FJ, Trinder J, *et al.* Nasal resistance is elevated in people with tetraplegia and is reduced by topical sympathomimetic administration. *J Clin Sleep Med* 2016;12(11):1487–92.
- 29 Schembri R, Spong J, Graco M, Berlowitz DJ. Neuropsychological function in patients with acute tetraplegia and sleep disordered breathing. *Sleep* 2016; DOI 10.1093/sleep/zsw037.
- 30 Hedli LC, Christos P, Krieger AC. Unmasking of Periodic Limb Movements With the Resolution of Obstructive Sleep Apnea During Continuous Positive Airway Pressure Application. *J Clin Neurophysiol* 2012;29(4):339–44.
- 31 Manconi M, Ferri R, Zucconi M, Fantini ML, Plazzi G, Ferini-Strambi L. Time structure analysis of leg movements during sleep in REM sleep behavior disorder. *Sleep* 2007;30(12):1779–85.