CASE REPORT

Trans-scleral explantation of posteriorly dislocated IOL-CTR complex in a case of anterior megalophthalmos with an unusually thin cornea

Anil Babanrao Gangwe, Priyavrat Bhatia, Deepshikha Agrawal, Samrat Chatterjee

Department of Ophthalmology,
MGM Eye Institute, Raipur,
Chhattisgarh, India

Correspondence to
Dr Priyavrat Bhatia,
gorbi.bhatia@gmail.com

Accepted 12 June 2018

SUMMARY

Anterior megalophthalmos and keratoglobus are characterised by corneal thinning and deep anterior chamber. They are clinically distinguished on the basis of normal to slightly decreased corneal thickness with a large corneal diameter (>13 mm) in case of anterior megalophthalmos, and marked limbus-to-limbus corneal thinning with globular protrusion in keratoglobus. To achieve and maintain a centred and stable intraocular lens (IOL) position in the bag is often difficult in cases of anterior megalophthalmos due to a too large diameter of the capsular bag and ciliary ring. We report a case of a 40-year-old man with features of anterior megalophthalmos with extremely thin cornea. He had spontaneous posterior dislocation of IOL and capsular tension ring (CTR) within the bag after initial successful cataract surgery. The dislocated complex could neither have been repositioned with scleral fixation due to large diameter of ciliary ring nor could it have been explanted through a clear corneal incision due to associated very thin cornea. We performed pars plana vitrectomy and separated the IOL-CTR complex inside the eye and explanted them separately through the sclerotomy, as is done for removal of intraocular foreign body. This avoided incision on the thin cornea. We conclude that this method can be very useful in a similar clinical situation but is complex enough to justify its use in routine cases.

BACKGROUND

Posteriorly dislocated intraocular lens (IOL) is removed conventionally through a clear corneal incision, following a pars plana vitrectomy (PPV). In the present case, removal of the IOL-capsular tension ring (CTR) complex through a clear corneal incision was not justified due to the thin and weak cornea. Therefore, a trans-scleral route was opted for the explantation.

CASE PRESENTATION

A 40-year-old man presented with sudden-onset diminution of vision in right eye since the previous day and a history of large eyes since birth. Two years before, he had undergone phacoemulsification through a scleral tunnel, with implantation of single-piece acrylic IOL having a total length of 13 mm and the optic diameter measuring 6 mm along with a 13 mm CTR in the right eye (figure 1A–C). Pseudophacodonesis was documented during

follow-up. The presenting visual acuity in right eye was 20/300 and in left eye was 20/100. Corrected distance visual acuity in the right eye with +4.00 D was 20/40 and in the left eye with –3.00 D was 20/32. Examination of both eyes revealed horizontal corneal diameter of 15 mm, global corneal thinning with mosaic like changes at the deep stromal level, deep anterior chamber and iridodonesis. Right eye was aphakic with vitreous at pupillary margin and left eye had cataract (nuclear sclerosis grade 3) without phacodonesis. Fundus examination of the right eye revealed a posteriorly dislocated IOL-CTR complex within the capsular bag (figure 1D). Fundus of the left eye was within normal limits. Intraocular pressure (not corrected for corneal thickness) was 3 mm Hg in right eye and 4 mm Hg in left eye.

INVESTIGATIONS

The axial length was 27.66 mm in the right eye and 26.18 mm in left eye. Keratometry for right eye was 44.63 D (129°), 43.18 D (39°); (figure 2) and for left eye was 44.59 D (60°), 43.39 D (150°) (figure 3). Pachymetry using anterior segment optical coherence tomography (AS-OCT) was performed which showed corneal thinning in all zones. The corneal thinning was uniform across the cornea (figure 4). The average central corneal thickness measured by AS-OCT pachymetry was 330 µm in right eye and 326 µm in left eye.

DIFFERENTIAL DIAGNOSIS

Keratoglobus is a rare non-inflammatory corneal thinning disorder characterised by generalised thinning and globular protrusion of the cornea.¹ Based on the findings of clinical examination, we initially assumed that the patient had concomitant anterior megalophthalmos and keratoglobus. Factor which favours the diagnosis of keratoglobus is limbus-to-limbus corneal thinning unlike anterior megalophthalmos where thinning is more prominent at the centre than periphery,² but the absence of a steep cornea or rather the presence of normal keratometry values rules out keratoglobus as the diagnosis.³ Final diagnosis of anterior megalophthalmos with very thin cornea was made. Presence of enlarged corneal diameter and relatively normal keratometry values favours a diagnosis anterior megalophthalmos with very thin cornea rather than keratoglobus.³

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Gangwe AB, Bhatia P, Agrawal D, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224691

Figure 1 (A and B) Preoperative slit-lamp photograph of the right and left eye, respectively, showing megalophthalmos, thin cornea, central corneal haze, deep anterior chamber and cataractous lens. (C) Slit-lamp photograph of the right eye postoperatively in retroillumination showing well-centred intraocular lens inside the capsular bag. (D) Intraoperative photograph of the right eye showing intraocular lens—capsular bag—CTR complex lying on the retina. CTR, capsular tension ring.

TREATMENT

The patient underwent PPV and explantation of CTR and IOL separately (video 1). Intraoperatively, few inferior zonular fibres were found to be attached to the capsular bag. Following initial vitrectomy, the dislocated capsular bag-IOL-CTR complex was freed from its residual zonular attachments. After completion of vitrectomy, the complex was floated using perfluorocarbon liquid (PFCL). IOL and CTR were separated bimanually by cutting the fibrotic adhesions between them, using vitrectomy cutter, intravitreal scissor and forceps. Adhesion between ends of ring was also released. CTR was explanted by grasping one of its free ends with an intravitreal forceps and then pulling it through the sclerotomy. After this, the sclerotomy was enlarged to 4 mm and IOL was also explanted through it using intravitreal forceps. Sclerotomy was sutured and PFCL was removed. Peripheral retina was examined with scleral depression and no iatrogenic retinal tear

Figure 2 Placido disc-based videokeratography of the right eye.

Figure 3 Placido disc-based videokeratography of the left eye.

was identified. Patient was left aphakic due to lack for consent for Artisan iris claw lens.

OUTCOME AND FOLLOW-UP

Postoperative course was uneventful and patient gained visual acuity of 20/32, N6 with aphakic correction of +4.00 D and near addition of +2.50 D, 1 month following surgery.

DISCUSSION

Anterior megalophthalmos is clinically distinct from keratoglobus, which shows generalised corneal ectasia in presence of normal corneal diameter. Patient described in this report had corneal diameter of 15 mm and corneal thickness of 330 μ m which was more or less same all throughout, and a negative family history. Enlarged ciliary ring in megalophthalmos can cause zonular dehiscence leading to subluxation of lens.⁴ Implantation of standard in the bag IOL often results in its decentration due to large diameter of the bag.⁴ Various methods of IOL implantation are described to avoid problem of IOL decentration, like use of anterior chamber iris supported lens,⁵ posterior chamber retro fixed iris claw lens,^{6,7} posterior chamber IOL sutured to the posterior surface of the iris and anterior capsule,⁸ standard posterior chamber IOL in the bag,⁴ and customised posterior chamber IOL with larger diameter.⁹ Dislocation of IOL in the bag occurring 3 months or later following cataract surgery has been termed as late dislocation.⁵ It occurs due to progressive zonular weakness and contraction of capsular bag. Pseudoexfoliation syndrome, trauma, uveitis, previous vitrectomy, anterior megalophthalmos, high axial length and retinitis pigmentosa are the predisposing factors for most of these cases.⁶ The dislocated IOL-CTR complex can be repositioned and fixed in to the sclera or explanted and exchanged with either a scleral fixed¹⁰ or anterior chamber IOL. Repositioning of the complex can be effectively achieved with PPV and use of PFCL¹¹ as well as by anterior approach without vitrectomy.¹² This case had a unique situation where repositioning of the IOL-CTR complex by scleral fixation was not possible due to large ciliary ring diameter. A large corneal incision to deliver 13 mm IOL-CTR complex was avoided due to central as well as peripheral corneal thinning. A clear corneal incision made on the thin cornea would have further added on an area of weakness to it. Sutures through such thin and weak cornea could have resulted in cheese wiring and subsequent increased risk of wound dehiscence or leakage. We

OU: Pachymetry Analysis

OD

OS

Figure 4 Corneal pachymetry using anterior segment optical coherence tomography showing global corneal thinning in both eyes.

Video 1 Video showing the key steps involved in removal of IOL and CTR from the vitreous cavity through the enlarged sclerotomy. CTR, capsular tension ring; IOL, intraocular lens.

preferred separating the IOL and CTR component in vitreous cavity followed by their explantation through sclerotomy.

To the best of our knowledge, this is the first published case of anterior megalophthalmos, where the posteriorly dislocated IOL and CTR complex has been explanted through a sclerotomy rather than a clear corneal incision. In the largest series of patients with keratoglobus using corneal topography, the average steep and flat sim k values in paediatric patients were $57.95 \text{ D} \pm 6.85 \text{ D}$ and $52.15 \text{ D} \pm 6.33 \text{ D}$, respectively. In adults the values were $54.7 \text{ D} \pm 8.29 \text{ D}$ and $49.37 \text{ D} \pm 7.68 \text{ D}$.³ On the other hand, the keratometries in the published cases of anterior megalophthalmos are around the mean of the normal population.¹³ In the present case, corneal topography also showed a central curvature of the cornea within normal limits in both eyes (OD: $43.18/44.63 \text{ D}$ and OS: $43.39/44.59 \text{ D}$), therefore ruling out the presence of a keratoglobus. In a review on this topic, Galvis *et al* explained that unlike megalocornea and megalophthalmos in which the corneas have a normal curvature and normal or mildly decreased thickness, the corneas in keratoglobus are remarkably thin and the enlargement of the corneal diameter is small. It is noteworthy that according to their review only two cases of anterior megalophthalmos have

been previously described with such significant corneal thinning one case with 377 μm and the other with 390 μm .¹³ The present case is therefore the one with the thinnest cornea ever reported (328 and 331 μm).

Learning points

Spontaneous dislocation can occur even years after successful cataract surgery in cases of anterior megalophthalmos, and patient has to be counselled beforehand regarding possible outcomes.

- ▶ Though it is not a routine practice to explant intraocular lens (IOL) or capsular tension ring through a sclerotomy, the method described above can be successfully employed whenever confronted with a similar situation where corneal incision has to be avoided.
- ▶ Though achieving in the bag IOL should be the ultimate goal of cataract surgery, in cases associated with progressive zonular weakness, an uncomplicated anterior chamber iris supported,⁵ posterior chamber retro fixed iris claw lens can be useful.^{6,7}

Contributors ABG, PB, DA and SC contributed for concept, design, manuscript review and editing. ABG and PB contributed in literature search, manuscript preparation.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Wallang BS, Das S. Keratoglobus. *Eye (Lond)*. 2013;27:1004–12.
- 2 Nemeth G, Hassan Z, Berta A, *et al*. Scheimpflug imaging in anterior megalophthalmos. *Indian J Ophthalmol* 2013;61:32–5.
- 3 Rath V, Murthy SI, Bagga B, *et al*. Keratoglobus: An experience at a tertiary eye care center in India. *Indian J Ophthalmol* 2015;63:233–8.
- 4 Javadi MA, Jafarinasab MR, Mirdehghan SA. Cataract surgery and intraocular lens implantation in anterior megalophthalmos. *J Cataract Refract Surg* 2000;26:1687–90.
- 5 Neumann AC. Anterior megalophthalmos and intraocular lens implantation. *J Am Intraocul Implant Soc* 1984;10:220–2.
- 6 Lee GA, Hann JV, Braga-Mele R. Phacoemulsification in anterior megalophthalmos. *J Cataract Refract Surg* 2006;32:1081–4.
- 7 Galvis V, Tello A, Miotto G, *et al*. Artisan aphakic lens for cataract surgery in anterior megalophthalmos. *Case Rep Ophthalmol* 2012;3:428–33.
- 8 Dua HS, Azuara-Blanco A, Pillai CT. Cataract extraction and intraocular lens implantation in anterior megalophthalmos. *J Cataract Refract Surg* 1999;25:716–9.
- 9 Vaz FM, Osher RH. Cataract surgery and anterior megalophthalmos: custom intraocular lens and special considerations. *J Cataract Refract Surg* 2007;33:2147–50.
- 10 Gunenc U, Kocak N, Ozturk AT, *et al*. Surgical management of spontaneous in-the-bag intraocular lens and capsular tension ring complex dislocation. *Indian J Ophthalmol* 2014;62:876–8.
- 11 Gimbel HV, Condon GP, Kohnen T, *et al*. Late in-the-bag intraocular lens dislocation: incidence, prevention, and management. *J Cataract Refract Surg* 2005;31:2193–204.
- 12 Krępszt L, Kuzmienė L, Miliauskas A, *et al*. Possible predisposing factors for late intraocular lens dislocation after routine cataract surgery. *Medicina* 2013;49:37–34.
- 13 Galvis V, Tello A, M Rangel C. Cataract surgery in anterior megalophthalmos: a review. *Med Hypothesis Discov Innov Ophthalmol* 2015;4:101–8.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow