Published in final edited form as:

Epidemiology. 2018 September ; 29(5): 697–706. doi:10.1097/EDE.0000000000000872.

Exposure to community violence and self-harm in California: A multi-level, population-based, case–control study

Ellicott C. Matthey^{1,*}, Kriszta Farkas¹, Jennifer Skeem^{2,3}, and Jennifer Ahern¹

¹Division of Epidemiology, University of California-Berkeley School of Public Health, Berkeley, California, USA

²University of California-Berkeley School of Social Welfare, Berkeley, California, USA

³Goldman School of Public Policy, University of California-Berkeley, Berkeley, California, USA

Abstract

Background—Self-harm is a leading cause of morbidity and mortality. Exposure to community violence is an important and potentially modifiable feature of the social environment that may affect self-harm, but studies to date are limited in the samples and outcomes examined.

Methods—We conducted a population-based, nested case–control study. Cases were all deaths and hospital visits due to self-harm in California, 2006–2013. We frequency-matched California resident population-based controls from the American Community Survey to cases on age, gender, race/ethnicity, and year of survey/injury. We assessed past-year community violence using deaths and hospital visits due to interpersonal violence in the community of residence. We estimated risk-difference parameters that were defined to avoid extrapolation and to capture associations between changes in the distribution of community violence and the population-level risk of self-harm.

Results—After adjustment for confounders, setting past-year community violence to the lowest monthly levels observed within each community over the study period was associated with a 30.1 (95% CI: 29.6 to 30.5) per 100,000 persons per year lower risk of nonfatal self-harm, but no difference in the risk of fatal self-harm. Associations for a parameter corresponding to a hypothetical violence prevention intervention targeting high-violence communities indicated a 5% decrease in self-harm at the population level. In sensitivity analyses, results were robust.

Conclusions—This study strengthens evidence on the relationship between community violence and self-harm. Future research should investigate reasons for differential associations by age and gender and whether community violence prevention programs have meaningful impacts on self-harm.

Corresponding Author: Ellicott C. Matthey, Berkeley Way West, 2121 Berkeley Way, Room 5302, University of California, Berkeley, Berkeley, CA 94720-7360, ematthey@berkeley.edu, 510-642-4937.

Conflicts of interest: The authors declare no conflicts of interest.

Disclaimer: The analyses, interpretations, and conclusions of this paper are attributable to the authors, and not to the California Department of Public Health or the National Institutes of Health.

Keywords

Community violence; self-harm; suicide; social environment; social epidemiology; epidemiologic methods; positivity; case-control

Introduction

Self-harm is a leading cause of morbidity and mortality in the United States (US), accounting for more than 44,000 deaths and 505,000 injuries in 2015.¹ Between 2001 and 2015, rates of fatal self-harm (suicide) increased 24% and rates of nonfatal self-harm increased 39%.¹ The reasons for these increases are not well understood. Rates of self-harm also vary substantially by population subgroup. For example, compared to the general population, fatal self-harm rates are nearly three times higher among older men and nonfatal self-harm rates are four times higher among young women.¹ Rising rates have drawn attention to self-harm as an important population health issue.^{2,3} Additional research is needed to understand the drivers of self-harm and to identify effective interventions.⁴

Aspects of the social environment such as social fragmentation and inequality are key risk factors for self-harm.⁵ Community violence is an important and modifiable feature of the social environment⁶ that may contribute to the burden of self-harm, particularly in the US where levels of community violence are high and rising.⁷ Exposure to community violence, meaning witnessing, hearing about, or directly experiencing violence in one's community,⁸ may increase the risk of self-harm in several ways. Increased stress, depressed mood, anxiety, symptoms of post-traumatic stress, and mental disorders can result from exposure to community violence^{9–14} and are strong risk factors for self-harm.^{15–20} Similarly, exposure to community violence can lead to substance use^{21,22} or social isolation (e.g. staying inside),²³ thereby increasing risk for self-harm.^{24–27} Moreover, exposure to community violence can normalize violence and aggression, another important risk factor for self-harm.^{28,29}

Epidemiologic research on the relationship between community violence and self-harm is limited.^{30–36} Although positive associations between community violence and suicidal ideation or nonfatal suicidal behavior have been observed, existing studies are generally limited to small samples of urban adolescents. To our knowledge, no previous studies have quantified the association of community violence with self-harm in a general population. Moreover, no studies have examined both fatal and nonfatal self-harm in the same population, which is critical because these forms of self-harm appear to differ in their distribution and determinants.^{1,37} Finally, no studies have estimated parameters corresponding to the potential impacts of specific reductions in community violence, which are particularly informative for public health decision-making.

Existing studies also suffer methodologic limitations, making it difficult to draw meaningful conclusions. In particular, community violence is strongly associated with other features of communities that are also associated with self-harm (e.g. economic opportunity). This makes it difficult to disentangle the effects of community violence from such factors.³⁸ When these factors are controlled using standard regression methods, the analysis often relies on extrapolation beyond the observed data, which can bias the results.³⁹ Previous

studies have also relied on self-reported measures of community violence exposure and suicide-related outcomes. This approach can introduce same-source bias, where self-report of both the exposure and outcome leads to spurious associations due to correlated measurement error (for example, due to pessimistic outlook).

In this study, we assessed the association of exposure to community violence with fatal and nonfatal self-harm, overall and by age and gender. We applied a population-based, case-control design to a large dataset including all deaths and hospital visits in California, a state with self-harm trends similar to those nationwide. We estimated risk difference parameters that avoid extrapolation and are relevant to potential public health interventions.

Methods

Data and study design

We compiled data on self-harm and community violence for the period 2005 to 2013 from two sources: deaths records from the California Department of Public Health Vital Records and emergency department and inpatient hospitalization discharge records from California's Office of Statewide Health Planning and Development. Records included all deaths and hospital visits statewide, except active duty military hospitals, and captured the external cause of death or injury, demographic characteristics, and residence of the patient or decedent. External cause of injury coding in California's hospital discharge records is mandatory, subject to ongoing quality assurance measures, and considered 100% complete.⁴⁰ Studies also indicate completeness and validity of external cause of mortality codes for homicide and self-harm in mortality data.⁴¹ Emergency department records are not available prior to 2005.

We treated the residents of California as a cohort and conducted a population-based, nested case-control study.^{42,43} Cases were all deaths and hospital visits due to deliberate self-harm in California, 2006–2013 (ICD-9-CM hospital visit code: E95; ICD-10 death codes: X6–X8). Self-harm outcomes were included starting in 2006 so that data on community violence were available for the relevant pre-injury exposure period (see Exposure assessment). We made efficient use of an existing population-representative sampling frame by sampling population-based controls⁴² from California resident participants in the American Community Survey (ACS). The ACS is an ongoing, nationwide survey conducted by the US Census Bureau. It is designed to generate population-representative small-area estimates of demographic, economic, and social indicators over time. ACS interviews were conducted with between 170,000 and 220,000 Californians annually between 2006 and 2013.

We created a state-representative pseudo-population of control units by duplicating each ACS record by the corresponding ACS person weight⁴⁴ and drew controls from this expanded dataset. For statistical efficiency,⁴³ we matched four controls to each case on confounders strongly associated with self-harm: gender, race/ethnicity, 5-year age group, and year of survey/injury. We used population-based controls to avoid the possibility of Berkson's bias that could result from hospital- or death record-based controls.⁴² We assumed that selected controls were not also cases at the time they were selected as controls; this is reasonable because self-harm risk was low (<0.5% in all matching strata). We

restricted to individuals residing in California at the time of survey/injury and to those aged 15 to 84 years due to small numbers outside that age range.

Exposure assessment

Exposure to community violence was defined as the average of the monthly rate of deaths due to homicide (ICD-10 death codes X85-X99, Y00-Y09, Y35, U01, U02, Y871) and injuries due to assault (ICD-9-CM hospital visit codes E960-E969, E970-E977) in the Consistent Public Use Microdata Area (CPUMA) of residence for the 12 months prior to survey/injury. CPUMAs are mutually exclusive and collectively exhaustive geographic units designated by the US Census Bureau. CPUMAs include at least 100,000 residents and are consistently defined over the study period. There are 110 CPUMAs in California. In urban areas (95% of the California population), CPUMAs correspond to known neighborhoods (e.g. Chinatown in San Francisco). In rural areas, CPUMAs are counties or aggregations of small counties.

Decedent addresses from vital records were geocoded to the CPUMA of residence. Patient zip codes from hospital records were assigned to the corresponding CPUMA of residence using a geographic crosswalk.⁴⁵ We selected CPUMAs, instead of census tracts or zip codes, to define neighborhoods because they are locally recognized places of residence but are large enough for stable estimation of monthly community violence rates. CPUMAs are also the smallest geographic identifier available in the ACS. We used objectively measured rates of community violence because they are strongly correlated with frequency of experiences of direct injury and witnessing violence reported by residents⁴⁶ while avoiding same-source bias. We used census-based population estimates equivalent to the ACS pseudo-population as denominators to calculate rates.

We used the average monthly violence rate over the 12-month period immediately prior to occurrence of self-harm for each case and selection of each corresponding control, because we conceptualize community violence as a chronic predisposing factor that theoretically can interact with acute stressors (e.g., psychosocial crisis) to cause self-harm.⁴⁷ The 12-month time frame is a proxy for longer-term exposure, given its strong association with multiyear measures (e.g. $R > 0.95$ with 36-month measure). A 12-month exposure ensured that seasonality did not impact the results, without extending so far back in time that residential mobility introduced excessive measurement error (within a year, 14% of people move and only 5% change counties of residence).⁴⁸ Crime data may also be used to measure community violence, but differences in reporting practices between jurisdictions and over time may introduce bias.^{49,50}

Confounder assessment

Individual- and community-level confounders were identified *a priori* based on the scientific literature and development of a directed acyclic graph (eAppendix 1). We considered established risk factors for self-harm and factors that affect community violence or share common causes with community violence. Variables controlled in the final analysis depended on availability in death, discharge, and ACS records. Individual-level confounders included in analyses of fatal self-harm were marital status, education, foreign birth, history

of military service, and recent immigration to the United States. Analyses of nonfatal self-harm controlled for individual-level primary language spoken. Sensitivity analyses for nonfatal self-harm also controlled for health insurance type, a proxy for socioeconomic status, which was available in the ACS after 2007. Community-level confounders in all analyses were annual or monthly community measures of sociodemographic composition, economic factors, social cohesion, firearm access, population mental health status, primary care provider density, alcohol outlet density, and weather (see eAppendix 2 for details).

Parameters

We estimated three risk difference parameters that capture how the population risk of self-harm is associated with specific changes in the distribution of community violence. Accurate estimation of these parameters relies on positivity, meaning that individuals in all confounder subgroups have to be observed under the different exposure conditions for which estimates are made. Positivity is a particular concern in studies of community violence, because individuals with certain covariate combinations may only be present in either high-violence or low-violence communities.

To ensure that the risk difference parameters did not rely on extrapolation, we identified the highest and lowest monthly violence rates within each community between 2005 and 2013, and, for each individual, we only estimated the risk difference for reductions/increases in community violence to the minimum/maximum observed in their community. By restricting the predictions to violence levels actually observed within communities, we minimized bias from extrapolating predictions beyond what is supported by the data. Specifically, we estimated:

1. RD_{overall} : the overall population risk difference comparing the estimated risk of self-harm if all individuals were exposed to 12-month average violence rates equal to the *highest* versus the *lowest* monthly violence rate observed within their communities⁵¹
2. RD_{PA} : the population attributable risk difference comparing the *observed* risk of self-harm to the risk of self-harm if all individuals were exposed to 12-month average violence rates equal to the *lowest* monthly violence rate observed within their communities
3. RD_{targeted} : the population risk difference comparing the *observed* risk of self-harm to the risk of self-harm if individuals in the *top quartile* of community violence (i.e. individuals living in high-violence communities) were exposed to 12-month average violence rates equal to the *lowest* monthly violence rate observed within their communities and exposure for all other individuals were left unchanged⁵²

The last parameter corresponds to the expected change in the population-level risk of self-harm under a hypothetical violence-prevention intervention that targets the most violent communities and reduces violence substantially but within the range previously experienced.

Statistical analysis

To estimate these marginal parameters, we used g-computation,⁵¹ which allows estimation of additive scale parameters and summarizes the association between community violence and self-harm for the population overall, rather than within covariate sub-groups, as in typical regression. We used generalized additive models with a logit link to model the risk of self-harm as a function of community violence, frequency matching factors (year, 5-year age group, race/ethnicity, and gender), and the confounders.⁵³ We used cubic smoothing splines⁵³ for all continuous independent variables, including community violence, to capture potential non-linear relationships with self-harm risk. We then used the fitted model to predict the risk of self-harm for each individual under the different exposure scenarios and took the difference of the average estimated risks for the relevant contrasts to estimate the three RD parameters. All analyses were weighted to be population-representative by assigning weights equal to the risk of self-harm within each matching strata (q_0) for cases and weights equal to $(1 - q_0)/J$ to controls, where J is the ratio of controls to cases.⁵⁴ We estimated 95% confidence intervals (CIs) using the nonparametric bootstrap.⁵¹

All analyses were stratified by self-harm type (fatal versus nonfatal) because the distribution and relative impacts of different determinants of self-harm vary by type.^{1,37} We report results for overall associations and for analyses stratified by 5-year age group and gender, because age and gender define the groups most commonly described as high risk,^{1,37} and we hypothesized that these groups would respond differently to community violence.

We excluded case records with incomplete covariate data (2.8%) from analyses, resulting in a final sample of 27,027 self-harm fatalities, and 331,203 nonfatal self-harm injuries. Data analysis was conducted using R 3.2.1 (R Foundation for Statistical Computing, Vienna, Austria; see eAppendix 7 for code), and model fitting and prediction were done using the gam package. This study was approved by the State of California and University of California, Berkeley Committees for the Protection of Human Subjects.

Nonfatal cases include only suicide attempts and self-harm injuries that were sufficiently serious to result in an emergency department visit or hospitalization. To assess the sensitivity of results to the inclusion of less severe cases for whom care-seeking may be optional and dependent on factors potentially associated with community violence (e.g. health insurance), we tested analyses restricted only to those nonfatal cases requiring inpatient hospitalization.

To assess the potential role of confounding due to unmeasured factors, we conducted a quantitative bias analysis. Using the bias equations presented by VanderWeele and Arah,⁵⁵ we estimated the characteristics of an unmeasured confounder that would yield the observed association between community violence and nonfatal self-harm, if the true effect were null.

Results

Table 1 presents the risk of fatal and nonfatal self-harm overall and by study characteristics. The risk of self-harm varied substantially by age group, gender, and type of self-harm and was positively correlated with community violence. Observed 12-month average incidence of community violence ranged between 6.9 and 126.6 per 100,000. The lowest within-

community monthly violence rates ranged from 2.4 to 64.7 per 100,000; the highest ranged from 14.5 to 154.6 per 100,000 (see eFigure 2 for geographic distribution). eAppendix 3 presents the number of cases and controls by age and gender.

Table 2 presents the overall associations between community violence and self-harm, adjusted for observed confounders. There were no associations of community violence with fatal self-harm. For nonfatal self-harm, the RD_{overall} was 62.9 per 100,000 (CI: 62.0, 63.9), or approximately a 27% reduction in self-harm relative to the observed risk. The RD_{PA} was 30.1 per 100,000 (CI: 29.6, 30.5), or a 13% reduction. The RD_{targeted} was 10.8 per 100,000 (CI: 10.6, 11.0), or a 5% reduction. The median differences in community violence for affected communities were 21.2 per 100,000 for the RD_{overall} , 9.9 per 100,000 for the RD_{PA} , and 14.8 per 100,000 RD_{targeted} .

Associations for the overall population masked substantial sub-group heterogeneity. Figure 1 presents the RD_{PA} by age and gender and shows that community violence was associated with increased risk of nonfatal self-harm predominantly among the young and middle-aged groups (ages 15–59), with the strongest relationships for women ages 15–24 and men ages 40–49. Community violence was generally not meaningfully associated with fatal self-harm. RD_{overall} (eFigure 3) and RD_{targeted} (Figure 2) estimates showed similar age and gender patterns to the RD_{PA} , but RD_{overall} were larger in magnitude and RD_{targeted} were smaller.

In sensitivity analyses (Table 2), restricting nonfatal self-harm to 2008–2013 to additionally control for health insurance type slightly attenuated the association of community violence with self-harm. Restriction to only inpatient cases reduced the overall risk of self-harm and showed similarly patterned but attenuated risk differences compared to the main analysis. For inpatient self-harm, the RD_{overall} corresponded to 9% less self-harm, the RD_{PA} corresponded to 4% less self-harm, and RD_{targeted} corresponded to 2% less self-harm.

Results of the bias analysis are presented in eAppendix 6. Briefly, for the association of community violence with nonfatal self-harm to be spurious, there would have to be an unmeasured confounder that is at least 50 percentage points more prevalent in high versus low violence communities and that causes a 100 per 100,000 increase in the risk of nonfatal self-harm (a very large association relative to the observed risk of 240 per 100,000).

Discussion

To our knowledge, this is the first study to examine the relationship of community violence with self-harm in a general population. We found that higher past-year community violence was associated with increased risk of nonfatal self-harm but not fatal self-harm, and that a parameter corresponding to setting community violence to lower levels for the highest-violence communities shows associations indicating meaningful reductions in nonfatal self-harm at the population level. Findings suggest that previously reported associations between community violence and nonfatal self-harm among adolescents^{30,31,33,34} extend statewide to the entire California population. Further, we identified important heterogeneity by age and gender, with the strongest associations for women ages 15–24 and men ages 40–49.

As in all observational studies, there may be residual confounding in the observed associations between community violence and self-harm. Confounding control was limited by the covariates available in death, discharge, and ACS records. The quantitative bias analysis indicates that for the observed association to be spurious, there would have to be an unmeasured factor that very strongly affects both community violence and self-harm. Identifying such a factor is possible. For example, mental disorder strongly increases self-harm risk, and also makes one more likely to live in a high-violence community. Confounders of particular concern include the type, extent, and history of mental and substance use disorders, personality traits, early life adversity, and precipitating life circumstances such as the loss of a loved one. However, exposure to community violence may causally precede these (e.g. incite substance use; contribute to the loss of a loved one). If these factors are on the causal pathway, adjusting for them would be inappropriate. We controlled for a large set of confounders including demographic, socioeconomic, contextual, and health indicators. However, additional research using longitudinal designs, more detailed covariate data on participants, and mediation analyses would help to separate these influences.

The community violence–self-harm association may also be driven by the co-occurrence of self-directed and outward-directed violence among the same individuals. Indeed, studies suggest that perpetration of violence against others (i.e., participating in community violence) is linked with psychiatric disorder, aggression, and other traits predisposing to self-harm, and that violence and suicidality mutually affect one another.^{29,32,56} We did not capture whether cases or controls were also direct contributors to community violence and therefore could not assess the co-occurrence of internally- and externally-directed violence. Further investigation is needed to disentangle these factors, particularly for non-adolescents for whom existing research is limited.

Our finding that community violence is associated with nonfatal self-harm but not fatal self-harm may indicate that nonfatal self-harm is more responsive to community violence. Community violence may induce psychologic distress or other psychologic and behavioral correlates sufficient to provoke expressions of self-harm, but insufficient to induce serious intent to kill oneself. Nonfatal self-harm can be a means of coping with distress,⁵⁷ whereas fatal self-harm may be a means of escaping distressing environments.⁵ These are fundamentally different responses, and community violence may be more likely to prompt one than the other. Differences in the covariates controlled in the analyses of fatal versus nonfatal self-harm or differential effects of residual confounding may also explain the different associations observed for nonfatal and fatal self-harm.

Our finding that the strongest associations were for young women and middle-aged men may be due to differences in vulnerability to stressors. Theory and evidence suggest that young women may be particularly vulnerable to life stressors and depression that can lead to suicidal behavior.^{58,59} There is less research on psychologic vulnerability to stressors among middle-aged men, but this group is less likely to seek or receive needed mental health care.^{60,61} Thus, untreated mental or substance use disorders or psychologic distress precipitated by community violence may be more likely to lead to self-harm in this group. Other work has documented recent increases in suicide among non-Hispanic White middle-aged men

and suggested that rising rates of long-term physical disability and mental and substance use disorders in addition to declining job prospects may contribute.² Given rising rates of community violence,⁷ our study suggests that community violence may contribute or exacerbate the risk of self-harm in this group. We also found minimal associations between community violence and self-harm among some high-risk groups (e.g. fatal self-harm for older men). This may indicate that community violence is not a key contributor to risk in these groups, and other social environment and individual factors would be worth examining.

Unlike previous studies that operationalize community violence as a binary “all or nothing” contrast, we used a continuous measure and estimated the impacts of plausible changes in exposure in an effort to more accurately estimate population-level impacts and better inform public health decision-making.⁵² Differences in the magnitudes of the overall risk difference (RD_{overall}), population attributable risk difference (RD_{PA}), and targeted risk difference (RD_{targeted}) reflect differences in the levels of community violence contrasted and the proportion of people affected. The RD_{overall} intervenes on everyone maximally, the RD_{PA} intervenes on individuals exposed to higher-than-minimum violence to varying degrees, and the RD_{targeted} intervenes most restrictively on only individuals in high-violence communities. The RD_{targeted} may be particularly informative because it corresponds to a hypothetical intervention to reduce violence in the highest-risk communities to achievable levels observed within those communities at some point over the study period. Focused deterrence strategies such as the Cure Violence model^{62,63} and mentoring programs for delinquent youth,⁶⁴ are examples of scientifically supported, locally targeted programs to reduce community violence that would fit this hypothetical scenario.

Our data do not include suicide attempts or other self-harm not resulting in hospital visits or deaths. Thus, we may be underestimating the burden of self-harm associated with community violence. In addition, if cases of self-harm of the same severity are more or less likely to receive care in a hospital depending on whether they live in a more or less violent community, selection bias may result. This pattern might result from less health insurance coverage, lower social support for care-seeking, or less access to emergency medical services in high-violence communities. However, results from sensitivity analyses restricted to the most severe cases for whom receipt of hospital services is unlikely to be optional were consistent, albeit attenuated, with those in the main analysis. Our control of proxy measures of healthcare access and other community-level determinants of care-seeking also help to address this concern.

Several other limitations of this study must be noted. First, records on the cause of death and injury classification are imperfect. However, studies suggest the degree of misclassification is not substantial enough to alter major trends and patterns.^{40,41} Second, we lacked long-term exposure data for cases and controls, and exposure misclassification may occur if study participants did not actually reside at the reported location for the 12 months prior to injury/survey. Third, we used distinct data sources to draw cases and controls, which may generate differences in the measurement of covariates. Finally, we used CPUMAs as a proxy for communities. Although these units are locally recognized places of residence, they may not fully capture the social environments of persons in this study.

Overall, this study strengthens the evidence on the relationship between community violence and self-harm. We used complete, population-wide data that included all deaths and hospital visits due to self-harm in California over an 8-year period, which allowed us to compare rare outcomes among important subgroups for whom previous assessments have been limited. We estimated easily interpretable population-level parameters that avoided extrapolation and made novel and efficient use of an existing population-representative survey to draw controls. This approach could serve as a model for future investigations seeking to reconstruct population exposure and outcome experiences to answer important public health questions using existing big data. This study suggests that lower levels of community violence, even when limited to the highest-violence communities, are associated with lower risk of nonfatal self-harm, particularly among young and middle-aged persons. Future research should strive for greater confounding control, investigate reasons for differential associations by age and gender, and assess whether community violence prevention programs have meaningful impacts on self-harm.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Source of Funding: This work was supported by grant DP2HD080350 from NICHD/NIH Office of the Director, and from the University of California-Berkeley Committee on Research, Robert Wood Johnson Health and Society Scholars Program, Harry Frank Guggenheim Foundation, and University of California-Berkeley Mack Center on Mental Health and Social Conflict.

The authors thank the following funding sources: NICHD/NIH Office of the Director; University of California-Berkeley Committee on Research; Robert Wood Johnson Health and Society Scholars Program; Harry Frank Guggenheim Foundation; University of California-Berkeley Mack Center on Mental Health and Social Conflict.

References

1. Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. [Accessed January 22, 2016] Web-based Injury Statistics Query and Reporting System (WISQARS) www.cdc.gov/injury/wisqars. Published 2005
2. Case A, Deaton A. Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. *Proc Natl Acad Sci*. 2015; 112(49):15078–15083. DOI: 10.1073/pnas.1518393112 [PubMed: 26575631]
3. Matthay EC, Galin J, Ahern J. Changing Patterns in Rates and Means of Suicide in California, 2005 to 2013. *Am J Public Health*. 2017; 107(3):427–429. DOI: 10.2105/AJPH.2016.303624 [PubMed: 28177819]
4. Claassen CA, Pearson JL, Khodyakov D, et al. Reducing the burden of suicide in the U.S.: the aspirational research goals of the National Action Alliance for Suicide Prevention Research Prioritization Task Force. *Am J Prev Med*. 2014; 47(3):309–314. DOI: 10.1016/j.amepre.2014.01.004 [PubMed: 24750971]
5. Durkheim E. *Suicide* London: Routledge and Kegan Paul; 1952
6. Krug EG, Mercy JA, Dahlberg LL, Zwi AB. The world report on violence and health. *The Lancet*. 2002; 360(9339):1083–1088. DOI: 10.1016/S0140-6736(02)11133-0
7. Grinshteyn E, Hemenway D. Violent Death Rates: The US Compared with Other High-income OECD Countries, 2010. *Am J Med*. 2016; 129(3):266–273. DOI: 10.1016/j.amjmed.2015.10.025 [PubMed: 26551975]

8. Suglia SF, Ryan L, Wright RJ. Creation of a community violence exposure scale: accounting for what, who, where, and how often. *J Trauma Stress*. 2008; 21(5):479–486. DOI: 10.1002/jts.20362 [PubMed: 18956446]
9. Margolin G, Gordis EB. The Effects of Family and Community Violence on Children. *Annu Rev Psychol*. 2000; 51(1):445–479. DOI: 10.1146/annurev.psych.51.1.445 [PubMed: 10751978]
10. Clark C, Ryan L, Kawachi I, Canner MJ, Berkman L, Wright RJ. Witnessing Community Violence in Residential Neighborhoods: A Mental Health Hazard for Urban Women. *J Urban Health*. 2007; 85(1):22–38. DOI: 10.1007/s11524-007-9229-8 [PubMed: 17965940]
11. Astell-Burt T, Feng X, Kolt GS, Jalaludin B. Does rising crime lead to increasing distress? Longitudinal analysis of a natural experiment with dynamic objective neighbourhood measures. *Soc Sci Med* 1982. 2015; 138:68–73. DOI: 10.1016/j.socscimed.2015.05.014
12. Mair C, Diez Roux AV, Shen M, et al. Cross-sectional and longitudinal associations of neighborhood cohesion and stressors with depressive symptoms in the multiethnic study of atherosclerosis. *Ann Epidemiol*. 2009; 19(1):49–57. DOI: 10.1016/j.annepidem.2008.10.002 [PubMed: 19064189]
13. Fowler PJ, Tompsett CJ, Braciszewski JM, Jacques-Tiura AJ, Baltes BB. Community violence: a meta-analysis on the effect of exposure and mental health outcomes of children and adolescents. *Dev Psychopathol*. 2009; 21(1):227–259. DOI: 10.1017/S0954579409000145 [PubMed: 19144232]
14. Ahern J, Matthay EC, Goin DE, Farkas K, Rudolph KE. Acute changes in community violence and increase in hospital visits and deaths due to stress-responsive disease. *Epidemiology*. 2018 In press.
15. Bell S, Russ TC, Kivimäki M, Stamatakis E, Batty GD. Dose-Response Association Between Psychological Distress and Risk of Completed Suicide in the General Population. *JAMA Psychiatry*. 2015; 72(12):1254–1256. DOI: 10.1001/jamapsychiatry.2015.2107 [PubMed: 26560886]
16. Cavanagh JTO, Carson AJ, Sharpe M, Lawrie SM. Psychological autopsy studies of suicide: a systematic review. *Psychol Med*. 2003; 33(3):395–405. [PubMed: 12701661]
17. Krysinaka K, Lester D. Post-traumatic stress disorder and suicide risk: a systematic review. *Arch Suicide Res Off J Int Acad Suicide Res*. 2010; 14(1):1–23. DOI: 10.1080/13811110903478997
18. Harris EC, Barraclough B. Suicide as an outcome for mental disorders. A meta-analysis. *Br J Psychiatry J Ment Sci*. 1997; 170:205–228.
19. Khan A, Leventhal RM, Khan S, Brown WA. Suicide risk in patients with anxiety disorders: a meta-analysis of the FDA database. *J Affect Disord*. 2002; 68(2–3):183–190. [PubMed: 12063146]
20. Liu RT, Miller I. Life events and suicidal ideation and behavior: A systematic review. *Clin Psychol Rev*. 2014; 34(3):181–192. DOI: 10.1016/j.cpr.2014.01.006 [PubMed: 24534642]
21. Vermeiren R, Schwab-Stone M, Deboutte D, Leckman PE, Ruchkin V. Violence exposure and substance use in adolescents: findings from three countries. *Pediatrics*. 2003; 111(3):535–540. [PubMed: 12612233]
22. Stockdale SE, Wells KB, Tang L, Belin TR, Zhang L, Sherbourne CD. The importance of social context: Neighborhood stressors, stress-buffering mechanisms, and alcohol, drug, and mental health disorders. *Soc Sci Med*. 2007; 65(9):1867–1881. DOI: 10.1016/j.socscimed.2007.05.045 [PubMed: 17614176]
23. Wolfer TA. Coping With Chronic Community Violence: The Variety and Implications of Women's Efforts. *Violence Vict*. 2000; 15(3):283–301. [PubMed: 11200103]
24. Kaplan MS, McFarland BH, Huguet N, et al. Acute alcohol intoxication and suicide: a gender-stratified analysis of the National Violent Death Reporting System. *Inj Prev*. Jan.2012 injuryprev-2012-040317. doi: 10.1136/injuryprev-2012-040317
25. Nock M, Hwang I, Sampson N, Kessler R. Mental Disorders, Comorbidity and Suicidal Behavior: Results from the National Comorbidity Survey Replication. *Mol Psychiatry*. 2010; 15(8):868–876. DOI: 10.1038/mp.2009.29 [PubMed: 19337207]
26. Trout DL. The Role of Social Isolation in Suicide. *Suicide Life Threat Behav*. 1980; 10(1):10–23. DOI: 10.1111/j.1943-278X.1980.tb00693.x [PubMed: 7361340]

27. Hall-Lande JA, Eisenberg ME, Christenson SL, Neumark-Sztainer D. Social Isolation, Psychological Health, and Protective Factors in Adolescence. *Adolescence*. 2007; 42(166):265–286. [PubMed: 17849936]
28. McGirr A, Renaud J, Bureau A, Seguin M, Lesage A, Turecki G. Impulsive-aggressive behaviours and completed suicide across the life cycle: a predisposition for younger age of suicide. *Psychol Med*. 2008; 38(3):407–417. DOI: 10.1017/S0033291707001419 [PubMed: 17803833]
29. Guerra NG, Rowell Huesmann L, Spindler A. Community Violence Exposure, Social Cognition, and Aggression Among Urban Elementary School Children. *Child Dev*. 2003; 74(5):1561–1576. DOI: 10.1111/1467-8624.00623 [PubMed: 14552414]
30. Vermeiren R, Ruchkin V, Leckman PE, Deboutte D, Schwab-Stone M. Exposure to Violence and Suicide Risk in Adolescents: A Community Study. *J Abnorm Child Psychol*. 2002; 30(5):529–537. DOI: 10.1023/A:1019825132432 [PubMed: 12403155]
31. Lambert SF, Copeland-Linder N, Ialongo NS. Longitudinal associations between community violence exposure and suicidality. *J Adolesc Health*. 2008; 43(4):380–386. DOI: 10.1016/j.jadohealth.2008.02.015 [PubMed: 18809136]
32. Van Dulmen M, Mata A, Claxton S, et al. Longitudinal associations between violence and suicidality from adolescence into adulthood. *Suicide Life Threat Behav*. 2013; 43(5):523–531. DOI: 10.1111/sltb.12036 [PubMed: 23725554]
33. Farrell CT, Bolland JM, Cockerham WC. The Role of Social Support and Social Context on the Incidence of Attempted Suicide Among Adolescents Living in Extremely Impoverished Communities. *J Adolesc Health*. 2015; 56(1):59–65. DOI: 10.1016/j.jadohealth.2014.08.015 [PubMed: 25438969]
34. Thompson R, Litrownik AJ, Isbell P, et al. Adverse experiences and suicidal ideation in adolescence: Exploring the link using the LONGSCAN samples. *Psychol Violence*. 2012; 2(2): 211–225.
35. Zimmerman GM. Does Violence toward Others Affect Violence toward Oneself? Examining the Direct and Moderating Effects of Violence on Suicidal Behavior. *Soc Probl*. 2013; 60(3):357–382. DOI: 10.1525/sp.2013.60.3.357
36. Colson KE, Galin J, Ahern J. Spatial Proximity to Incidents of Community Violence Is Associated with Fewer Suicides in Urban California. *J Urban Health Bull N Y Acad Med*. 2016; 93(5):770–796. DOI: 10.1007/s11524-016-0072-7
37. Silverman MM, Berman AL, Sanddal ND, O'Carroll PW, Joiner TE. Rebuilding the Tower of Babel: A Revised Nomenclature for the Study of Suicide and Suicidal Behaviors Part 2: Suicide-Related Ideations, Communications, and Behaviors. *Suicide Life Threat Behav*. 2007; 37(3):264–277. DOI: 10.1521/suli.2007.37.3.264 [PubMed: 17579539]
38. Oakes JM. Commentary: Advancing neighbourhood-effects research—selection, inferential support, and structural confounding. *Int J Epidemiol*. 2006; 35(3):643–647. DOI: 10.1093/ije/dyl054 [PubMed: 16556642]
39. Petersen ML, Porter KE, Gruber S, Wang Y, van der Laan MJ. Diagnosing and responding to violations in the positivity assumption. *Stat Methods Med Res*. 2012; 21(1):31–54. DOI: 10.1177/0962280210386207 [PubMed: 21030422]
40. Centers for Disease Control and Prevention. Strategies to Improve External Cause-of-Injury Coding in State-Based Hospital Discharge and Emergency Department Data Systems: Recommendations of the CDC Workgroup for Improvement of External Cause-of-Injury Coding. *Morb Mortal Wkly Rep*. 2008; 57(RR-1):1–16.
41. Moyer LA, Boyle CA, Pollock DA. Validity of death certificates for injury-related causes of death. *Am J Epidemiol*. 1989; 130(5):1024–1032. [PubMed: 2683747]
42. Wacholder S, Silverman DT, McLaughlin JK, Mandel JS. Selection of Controls in Case-control Studies: II. Types of Controls. *Am J Epidemiol*. 1992; 135(9):1029–1041. [PubMed: 1595689]
43. Wacholder S, Silverman DT, McLaughlin JK, Mandel JS. Selection of Controls in Case-control Studies: III. Design Options. *Am J Epidemiol*. 1992; 135(9):1042–1050. DOI: 10.1093/oxfordjournals.aje.a116398 [PubMed: 1595690]

44. UC Census Bureau. [Accessed August 2, 2017] American Community Survey Design and Methodology 2014 <https://www.census.gov/programs-surveys/acs/methodology/design-and-methodology.html>
45. Missouri Census Data Center. [Accessed December 15, 2017] MABLE/Geocorr14, Version 1.0: Geographic Correspondence Engine <http://mcdc.missouri.edu/websas/geocorr14.html>. Published 2015
46. Hartinger-Saunders RM, Rine CM, Nochajski T, Wieczorek W. Neighborhood crime and perception of safety as predictors of victimization and offending among youth: A call for macro-level prevention and intervention models. *Child Youth Serv Rev*. 2012; 34(9):1966–1973. DOI: 10.1016/j.childyouth.2012.05.020 [PubMed: 23049152]
47. Mann JJ. Neurobiology of suicidal behaviour. *Nat Rev Neurosci*. 2003; 4(10):819–828. DOI: 10.1038/nrn1220 [PubMed: 14523381]
48. Ihrke D, , Faber C, , Koerber W. Geographical Mobility: 2008–2009. Current Population Reports Washington, DC: US Census Bureau; 2011
49. Barnett-Ryan C, , Griffith EH. The Dynamic Nature of Crime Statistics. In: Cuevas CA, , Rennison C, editors *The Wiley Handbook on the Psychology of Violence* John Wiley & Sons, Ltd; 2016:523
50. Maltz MD, Targonski J. A Note on the Use of County-Level UCR Data. *J Quant Criminol*. 2002; 18(3):297–318. DOI: 10.1023/A:1016060020848
51. Ahern J, Hubbard A, Galea S. Estimating the Effects of Potential Public Health Interventions on Population Disease Burden: A Step-by-Step Illustration of Causal Inference Methods. *Am J Epidemiol*. 2009; 169(9):1140–1147. DOI: 10.1093/aje/kwp015 [PubMed: 19270051]
52. Ahern J, Colson KE, Margerson-Zilko C, Hubbard A, Galea S. Predicting the Population Health Impacts of Community Interventions: The Case of Alcohol Outlets and Binge Drinking. *Am J Public Health*. 2016; 106(11):1938–1943. DOI: 10.2105/AJPH.2016.303425 [PubMed: 27631757]
53. Hastie T, Tibshirani RJ. *Generalized Additive Models* New York, NY: Chapman & Hall; 1990
54. van der Laan MJ. Estimation based on case-control designs with known prevalence probability. *Int J Biostat*. 2008; 4(1) Article 17.
55. Vanderweele TJ, Arah OA. Bias formulas for sensitivity analysis of unmeasured confounding for general outcomes, treatments, and confounders. *Epidemiol Camb Mass*. 2011; 22(1):42–52. DOI: 10.1097/EDE.0b013e3181f74493
56. Swanson JW, Holzer CE, Ganju VK, Jono RT. Violence and Psychiatric Disorder in the Community: Evidence From the Epidemiologic Catchment Area Surveys. *Psychiatr Serv*. 1990; 41(7):761–770. DOI: 10.1176/ps.41.7.761
57. Laye-Gindhu A, Schonert-Reichl KA. Nonsuicidal Self-Harm Among Community Adolescents: Understanding the “Whats” and “Whys” of Self-Harm. *J Youth Adolesc*. 2005; 34(5):447–457. DOI: 10.1007/s10964-005-7262-z
58. Rudolph KD, Hammen C. Age and Gender as Determinants of Stress Exposure, Generation, and Reactions in Youngsters: A Transactional Perspective. *Child Dev*. 1999; 70(3):660–677. DOI: 10.1111/1467-8624.00048 [PubMed: 10368914]
59. Hankin B, Abramson L. Development of gender differences in depression: An elaborated cognitive vulnerability-transactional stress theory. *Psychol Bull*. 2001; 127(6):773–796. [PubMed: 11726071]
60. Tran LD, Ponce NA. Who gets needed mental health care? Use of mental health services among adults with mental health need in California. *Californian J Health Promot*. 2016; 15(1):36–45.
61. Galdas PM, Cheater F, Marshall P. Men and health help-seeking behaviour: literature review. *J Adv Nurs*. 2005; 49(6):616–623. DOI: 10.1111/j.1365-2648.2004.03331.x [PubMed: 15737222]
62. Braga AA, Hureau DM, Papachristos AV. Deterring Gang-Involved Gun Violence: Measuring the Impact of Boston’s Operation Ceasefire on Street Gang Behavior. *J Quant Criminol*. 2014; 30(1):113–139. DOI: 10.1007/s10940-013-9198-x
63. Boyle DJ, Lanterman JL, Pascarella JE, Cheng C-C. The Impact of Newark’s Operation Ceasefire on Trauma Center Gunshot Wound Admissions. *Justice Res Policy*. 2010; 12(2):105–123. DOI: 10.3818/JRP.12.2.2010.105

64. TolanP, , HenryD, , SchoenyM. [Accessed August 9, 2017] Mentoring Interventions to Affect Juvenile Delinquency and Associated Problems: A Systematic Review. Campbell Syst Rev2013<https://search.proquest.com/openview/3215b1a8999163f4330c9ca552bcef0a/1?pq-origsite=gscholar&cbl=2040247>

Figure 1. Adjusted population attributable risk difference for fatal and nonfatal self-harm associated with community violence, by age and gender

Risk differences are presented per 100,000 persons per year. (A) Fatal self-harm. (B) Nonfatal self-harm. The population attributable risk difference compares the observed risk of self-harm to the risk of self-harm if all individuals were exposed to 12-month average violence rates equal to the lowest monthly violence rate observed within their communities. Analyses are adjusted for race/ethnicity, year of injury or survey, and community-level confounders (see Covariate assessment). Analyses of nonfatal outcomes are also adjusted for individual-level primary language spoken. Analyses of fatal outcomes are also adjusted for individual-level marital status, education, foreign birth, military service, and recent immigration to the United States. Confidence intervals for fatal self-harm are asymmetric due to the small sample size of these subgroups and should be interpreted with caution. RD: risk difference. Bars indicate 95% confidence intervals.

^aEstimate for women ages 80–84 is unstable due to small sample size and is not presented.

Figure 2. Adjusted targeted risk difference for fatal and nonfatal self-harm associated with community violence, by age and gender

Risk differences are presented per 100,000 persons per year. (A) Fatal self-harm. (B) Nonfatal self-harm. The targeted risk difference compares the observed risk of self-harm to the risk of self-harm if individuals in the top quartile of community violence were exposed to 12-month average violence rates equal to the lowest monthly violence rate observed within their communities and exposure for all other individuals were left unchanged. Analyses are adjusted for race/ethnicity, year of injury or survey, and community-level confounders (see Covariate assessment). Analyses of nonfatal outcomes are also adjusted for individual-level primary language spoken. Analyses of fatal outcomes are also adjusted for individual-level marital status, education, foreign birth, military service, and recent immigration to the United States. Confidence intervals for fatal self-harm are asymmetric due to the small sample size of these subgroups and should be interpreted with caution. RD: risk difference. Bars indicate 95% confidence intervals.

^aEstimate for women ages 80–84 is unstable due to small sample size and is not presented.

Table 1

Risk of fatal and nonfatal self-harm by study characteristics, per 100,000 persons per year.

Characteristic	Fatal		Nonfatal	
	Female	Male	Female	Male
All	8.8	24.8	279.0	173.4
Quartile of past-year violence in community of residence	Lowest	7.4	21.4	255.5
	2	9.0	24.7	263.3
	3	9.2	25.5	282.3
	Highest	9.5	27.5	315.0
Age group (years)	15–19	3.3	8.9	493.6
	20–24	4.6	15.4	304.8
	25–29	4.7	15.7	237.1
	30–34	5.4	16.9	213.5
	35–39	7.3	19.2	216.2
	40–44	9.3	21.8	219.8
	45–49	10.5	27.7	211.3
	50–54	13.0	30.9	172.6
	55–59	11.6	32.3	121.4
	60–64	9.7	27.8	74.2
	65–69	7.8	25.3	53.1
	70–74	6.4	27.7	38.5
	75–79	7.4	39.1	36.1
	80–84	7.2	43.2	37.7

Estimates of risk of self-harm are weighted to be population representative (see Statistical Analysis).

Table 2
Overall adjusted associations between exposure to community violence and risk of fatal and nonfatal self-harm

Self-harm type	Observed risk	Risk of self-harm if all individuals were exposed to high community violence ^a (95% CI)	Risk of self-harm if all individuals were exposed to low community violence ^a (95% CI)	Overall risk difference (95% CI)	Population attributable risk difference (95% CI)	Targeted risk difference (95% CI)
Fatal	21.0	21.0 (20.1, 22.2)	21.1 (20.0, 22.2)	0.0 (-2.0, 2.1)	0.0 (-1.1, 1.1)	0.2 (-0.2, 0.6)
Nonfatal	234.8	267.7 (267.1, 268.3)	204.7 (204.3, 205.2)	62.9 (62.0, 63.9)	30.1 (29.6, 30.5)	10.8 (10.6, 11.0)
Nonfatal, restricted to 2008–2013, controlling for health insurance type	240.0	266.4 (265.5, 267.3)	214.2 (213.4, 214.9)	52.2 (50.6, 53.8)	25.8 (25.1, 26.5)	9.7 (9.4, 10.0)
Nonfatal, inpatient only	76.5	80.2 (80.0, 80.5)	73.6 (73.3, 73.8)	6.7 (6.2, 7.1)	2.9 (2.7, 3.1)	1.4 (1.3, 1.5)

^aHigh and low violence are defined as the highest and lowest levels of monthly violence observed within the study participants' communities over the study period (2005–2013).

Risks and risk differences are presented per 100,000 persons per year. The overall risk difference compares the estimated risk of self-harm if all individuals were exposed to 12-month average violence rates equal to the highest versus the lowest monthly violence rate observed within their communities. The population attributable risk difference compares the observed risk of self-harm if all individuals were exposed to 12-month average violence rates equal to the lowest monthly violence rate observed within their communities. The targeted risk difference compares the observed risk of self-harm to the risk of self-harm if individuals in the top quartile of community violence were exposed to 12-month average violence rates equal to the lowest monthly violence rate observed within their communities and exposure for all other individuals were left unchanged. Analyses are adjusted for 5-year age group, gender, race/ethnicity, year of injury or survey, and community-level confounders (see Covariate assessment). Analyses of nonfatal outcomes are also adjusted for individual-level primary language spoken, or primary language spoken and insurance type, as indicated. Analyses of fatal outcomes are also adjusted for individual-level marital status, education, foreign birth, military service, and recent immigration to the United States. CI: confidence interval.