Published in final edited form as:

Mycopathologia. 2018 August ; 183(4): 645–658. doi:10.1007/s11046-018-0254-9.

RNA-Seq Profile reveals Th-1 and Th-17 Type of Immune Responses in Mice Infected Systemically with *Aspergillus fumigatus*

Jata Shankar^{1,3,4,¶}, Gustavo C. Cerqueira^{2,¶}, Jennifer R. Wortman², Karl V. Clemons^{3,4,*}, and David A. Stevens^{3,4}

¹Jaypee University of Information Technology, Solan, HP, India

²Broad Institute of MIT and Harvard, Cambridge, MA, United States of America

³California Institute for Medical Research, San Jose, CA, United States of America

⁴Division of Infectious Diseases and Geographic Medicine, Stanford University, Stanford, CA, United States of America

Abstract

With the increasing numbers of immunocompromised hosts, *Aspergillus fumigatus* emerges as a lethal opportunistic fungal pathogen. Understanding innate and acquired immunity responses of the host is important for a better therapeutic strategy to deal with aspergillosis patients. To determine the transcriptome in the kidneys in aspergillosis, we employed RNA-Seq to obtain single 76 base reads of whole genome transcripts of murine kidneys on a temporal basis (days 0; uninfected, 1, 2, 3, and 8) during invasive aspergillosis. A total of 6,284 transcripts were downregulated and 5,602 were upregulated compared to baseline expression. Gene Ontology enrichment analysis identified genes involved in innate and adaptive immune response, as well as iron binding and homeostasis, among others. Our results showed activation of pathogen recognition receptors e.g., β -defensins, C-type lectins (e.g., dectin-1), Toll-like receptors (TLR-2,

*Correspondence: Karl V. Clemons, clemons@cimr.org.

¶These authors contributed equally to this work.

Ethics Approval and consent to participate

All animal studies were using a protocol approved by the Institutional Animal Care and Use Committee of the California Institute for Medical Research (CIMR) under protocol numbers 09-01:3 and 12-01:1. Animal Care and Use protocols followed the Animal Welfare Act (7 U.S.C. 2131 *et seq.*), and the Office of Laboratory Animal Welfare - Public Health Service Policy on Humane Care and Use of Laboratory Animals. CIMR protocols were approved and performed under OLAW assurance number, A3652-01. The NRC Guidelines for the Care and Use of Laboratory Animals, 8th edition, were followed for all care and use of mice. Mice were euthanized using CO₂ narcosis per the 2007 version of the American Veterinary Medical Association (AVMA) guidelines, which was current at the time the animal studies were performed.

Consent for publication

Not applicable

Availability of data and material

Sequencing runs were deposited at Short Read Archive (SRA-NCBI) under the following accession identifiers: SRX1201397, SRX1201396, SRX1201395, SRX1201394, SRX1201392.

Conflict of interest

The authors declare that they have no conflict of interest

Authors' Contributions

KVC and DAS conceived and designed the experiments. JS performed the experiments. GCC and JS analyzed the data. DAS, KVC and JRW contributed reagents/materials/analysis tools. GCC, JS, KVC and DAS contributed to the writing of the manuscript.

TLR-3, TLR-8, TLR-9 and TLR-13), as well as Ptx-3, and C-reactive protein among the soluble receptors. Upregulated transcripts encoding various differentiating cytokines and effector pro-inflammatory cytokines, as well as those encoding for chemokines and chemokine receptors revealed Th-1 and Th-17 type immune responses. These studies form a basic dataset for experimental prioritization, including other target organs, to determine the global response of the host against *Aspergillus* infection.

Keywords

Host-Pathogen; Invasive aspergillosis; *Aspergillus fumigatus*; *Mus musculus*; RNA-Seq; Cytokines; differential expression

1. Introduction

Humans inhale conidia from *Aspergillus* species, daily [1]. *Aspergillus fumigatus* is a ubiquitous filamentous fungus, which can cause a variety of problems ranging from allergic to invasive diseases depending on the individual. Invasive aspergillosis is life-threatening in immunocompromised patients, including those with solid organ transplants and hematologic malignancies [2–3].

Host responses against *A. fumigatus* infection are critical for effective control of the disease [4–5]. Studies in mouse models showed that both innate and acquired immunity of the host determine the protection against aspergillosis [5–7]. With the increasing numbers of at risk patients, *A. fumigatus* has emerged as the most lethal mycosis worldwide [2, 8]. Established infection in these patient groups has proven very difficult to eradicate due to the lack of highly effective and curative antifungal drug treatment [2] and gaps in knowledge regarding the host immune response in normal and immunocompromised individuals.

The exploration of the host transcriptome during the course of *A. fumigatus* infection may indicate which genes play an important role in the host response and help the understanding of the development and the progression of the disease [9–10]. Recently, RNA-Seq has been used to perform transcript profiling of *in vitro* cultured *A. fumigatus* [11–13], *A. oryzae* [14], and *A. flavus* [15]. However, RNA-Seq transcriptome data from *in vivo* or *ex vivo* studies are minimal [16]. Oosthuizen, *et al.* used genome-wide microarrays for fungus and human to examine the transcriptome response of bronchoepithelial cells to infection with *A. fumigatus* conidia, noting up-regulation of IL-6 and innate pathways for the human cells, and up-regulation of iron acquisition, and vacuolar acidification pathways, among others for the fungal cells [17]. Similarly, Morton, *et al.* used microarrays to examine expression profiles of dendritic cells from an *in vitro* model of the alveolar surface and their interaction with *A. fumigatus*, noting changes in various chemokines associated with neutrophil chemotaxis [18]. In the current studies we used RNA-Seq technology to determine the host transcriptome in the kidneys of mice infected intravenously with *A. fumigatus* to better understand the events occurring in the most affected organ during intravenous infection and which may also be a key organ during systemic aspergillosis in humans [19]. Our study demonstrated coordinated expression of genes encoding for interleukins and interferon, interleukin receptors and receptor-associated factors, interferon regulators, interferon

receptor and interferon-induced proteins, chemokines, chemokines receptors, iron homeostasis, and iron binding proteins.

2. Material and methods

2.1. Infection model

Six-week-old female CD-1 mice (Charles River Laboratories, Hollister, CA, USA) weighing approximately 25 grams each were arranged in groups of 5 and provided food and water *ad libitum*.

A. fumigatus strain Af293 [20] conidia for infection were prepared, and infected intravenously with 1.1×10^7 viable conidia in 250 μ l of saline/Tween-80 (0.05%) as described previously [21–23]. Five pre-assigned mice were euthanized on days 0, 1, 3, 5 and 8 to collect the samples. One mouse died on day 8, and thus we considered day 8 as a last time point to collect the samples. Uninfected mice were used as normal controls to provide the baseline murine transcriptome. At each time point, kidneys were removed aseptically from each mouse in the group and pooled before being snap-frozen in liquid N₂. Samples were stored at 80°C until used for total RNA extraction.

2.2. RNA extraction

Kidneys from 5 mice at each point were thawed in Trizol (Invitrogen, Carlsbad, CA, USA) and broken with 0.5 mm zirconia/silica beads (Biospec) using a Bead Beater apparatus (Biospec Products, Inc., Bartlesville, OK, USA) at high-speed setting (3 pulses of 12–15 sec). Total RNA was extracted using TRIzol reagent per the manufacturer's instructions. Contaminating gDNA was removed using RNeasy mini spin column (RNeasy mini kit, Qiagen Science, MD, USA) and RNase-free DNase set (50) (Qiagen GmbH, Germany), according to the manufacturer's instructions. The quantity and quality of the extracted RNA were assessed by A260 nm/A280 nm ratio (NanoDrop 1000 spectrophotometer, Thermo Scientific), as well as by electrophoresis through a 1.5% agarose gel stained with ethidium bromide for the presence of intact 18S and 28S ribosomal RNA bands, visualized by UV transillumination at 302 nm. RNA purity was determined using A260:A280 ratios and the extinction coefficient for RNA to quantify the amount of RNA present [24].

2.3. cDNA library preparation and sequencing

Poly (A) containing mRNA was purified from 8 μ g of total RNA of each sample using Sera-mag Magnetic Oligo (dT) Beads (Illumina RS-100-0801); mRNA molecules were fragmented using divalent cations under elevated temperature followed by first and second strand synthesis. End repair was performed and deoxyadenosine triphosphate (dATP) added to the 3' end of the DNA fragments. Next, adapters were ligated to cDNA fragments, cDNA template purified and finally the cDNA libraries were amplified via PCR. All steps were performed according to manufacturer's instructions. Before sequencing, the quantity and quality of each cDNA library was assessed using a 2100 Bioanalyzer (Agilent Inc.) [14]. Libraries were sequenced as single 76 base reads using the sequencing by synthesis method (Solexa sequencing, Illumina Inc.) at the Stanford Functional Genomic Facility, Stanford University.

2.4 Differential expression analysis

Initially, our intention was to obtain sequence reads for *A. fumigatus* and for the mouse transcripts, to generate two profiles to better understand the host-pathogen interactions. Alignment of sequence reads to the annotated genome of *A. fumigatus* (Af293) resulted in few transcripts that could map to the genome with more than a minimum number of reads (i.e., <10). Thus, *in vivo* expressed transcripts of *A. fumigatus* could be not analyzed. Transcript sequences of *Mus musculus* were downloaded from Ensembl (Build 38, GCA_000001635.2). Functional annotation and Gene Ontology (GO) terms were downloaded from EnsemblBiomart (GRCm38.p1). Original coding sequences (CDS) were complemented by 100 nt of flanking genomic sequence on both ends to enable alignment of short sequencing reads with untranslated regions (UTR). Short reads were aligned against the modified CDS using Bowtie [25]; only the best 20 valid alignments ($-k$ 20) were processed. Quantification of transcript abundance and isoform-adjusted expression was done via RSEM [26]. Normalization across samples was performed using the trimmed mean of M-values method (TMM) implemented by Edge-R [27]. Differential expression was calculated between all pairs of time points using the classical method of Edge-R [28]. Only genes with false discovery rate (FDR) < 0.01 and 16-fold change between a pair of time points were reported. All the applications were executed as part of the differential expression workflow implemented by the Trinity suite [29].

2.5 Clusters of genes with similar expression profile across samples

Clusters with similar expression profile across time points were defined based on the mean centered logarithm on base two of the FPKM values (Fragments per Kilobase of transcript per Million of mapped reads). The mean value was based on all genes with significant differential expression in at least one pair of time points. Similarity of expression profile between genes was computed using Euclidean distance and clusters were defined according to K-mean algorithm.

GO enrichment was defined by Fisher-exact correlation test and corrected for multiple comparisons via FDR, both implemented by scripts developed at the Broad Institute. A GO node was considered over-represented in the case of 3-fold difference between test and background sets and a FDR value lower or equal to 0.05. Non-informational GO terms are not shown in the main manuscript, but listed in supplemental material (Supplementary file 1: Tables 7 and 8). GO terms were summarized by Revigo using *simRel* method and 0.7 maximum similarity allowed between remaining GO terms (*small* threshold) [30].

3. Results

In the murine model of systemic aspergillosis, immunocompetent mice are infected intravenously with conidia to establish infection. However, quantification of the fungal burden of *A. fumigatus* is difficult [31], with the temporal burden of *A. fumigatus* in the kidneys determined by culture of homogenates shows the highest burden at day 0–1, at which time the conidia are morphologically still conidia or have germinated with short hyphae and colony counts determined at later time points (i.e., day 5 and 8) lower because the organism has developed into hyphal forms in the tissues. This can be misinterpreted as a

reduced burden, whereas the use of qPCR to quantify fungal burden demonstrates a progressive increase in the number of organisms present [32]. In addition to that, among the few reads derived from *A. fumigatus*, we observed a progressive increase in abundance of rRNA transcripts (not shown). Thus, we are confident that the model is that of progressive infection and that temporal alterations of gene expression in the kidney are a result of the rise in infectious burden.

Approximately 179 million sequencing reads were generated (Supplementary Table S1), of which 136 million were successfully mapped to mouse transcripts. With respect to transcript coverage, of 83,070 mouse transcripts used as reference sequences, 57,007 (69%) had more than 10 reads mapped and approximately 29,000 (35%) had a normalized FPKM of one or greater. When considering each sample/day separately, at least 26% of transcripts reported a FPKM greater than or equal to one (Supplementary Table S2). Transcripts that were significantly differentially expressed in at least one pairwise comparison between time points accounted for 17% of mouse transcripts (14,397) (Supplementary Table S3). When comparing each time point to day 0 (non-infected animal, baseline), 6,284 transcripts (8%) were downregulated and 5,602 (7%) were upregulated. Despite approximately 2,000 transcripts found to be up-regulated on days 1 and 3 in comparison to day 0 (Fig. 1 and Supplementary Table S4 for the complete list of up-regulated genes in comparison to day 0), there was no consistency in terms of the biological process associated with those genes, as evaluated by GO term enrichment. A concerted positive response only appeared on day 5, consisting of genes associated with leukocyte aggregation, acute inflammatory response, positive regulation of chemokines and other GO terms associated with immune response (set A on Fig. 1; Supplementary Table S7 for the list of enriched GO terms). Down-regulation was maximal on day 1 with twice the number of transcripts (4,295) (set D on Fig. 1 and Supplementary Table S5) than those upregulated at the same day. Genes with lower expression when compared to non-infected mice were consistently enriched for processes and functions associated with iron and heme binding, electron carrier activity and aromatase activity (Fig. 1B and Supplementary Table S7). A detailed inspection of the downregulated genes responsible for the enrichment of these processes and functions indicated that the majority were members of the cytochrome P450 family of proteins (Supplementary Table S10).

Thirty-five clusters with similar expression during the course of infection were defined (Supplementary Table S6). A set of three clusters representing positive, negative and mixed response are shown on Fig. 2A, with three of those having enrichment of GO terms. The positive response (cluster 7) was characterized by genes associated with adaptive and innate immune response; and the negative response (clusters 10 and 18) was enriched with transcripts associated with the assembly and binding of lipoproteins, including apolipoproteins H and A2, previously reported as negative regulator of neutrophils and angiogenesis [33]. The acute inflammatory response and negative regulation of cytokine production were also among the biological processes enriched in genes with lower expression after day 1, cluster 10 and 18 (Fig. 2B and Supplementary Table S8).

A comparison focusing on the dynamics of cytokines and iNOS response between our work and a study on mice infected with *Cryptococcus neoformans* [34] indicated a similar pattern

of robust response on day 5 and up-regulation of TNF-alpha on day 1, in spite of the differences in infected tissue (brain), pathogen and technique (RT-PCR) between the two studies (Supplementary Table S11). A lower correlation and an earlier detection was reported by an ELISA based evaluation of cytokines response in kidneys of mice infected with *A. flavus* [35], possibly different due to the nature of those assays.

3.1. Recruitment of immune cells at the site of infection

We found up-regulated transcripts encoding for chemokines and chemokine receptors. Chemokines (*Ccl4*, *Ccl5*, *Ccl7*, and *Ccl8*) and their receptor, *Ccr1*, present on neutrophils, were upregulated (Supplementary Fig. S1). On monocytes, chemokine receptors *Ccr2* and *Ccr5*, and their ligand chemokines (e.g., *Ccl2*, *Ccl7*, *Ccl8*, and *Ccl4*, *Ccl5*, *Ccl8*, respectively) were also upregulated (Supplementary Fig. S1). Receptors present on macrophages, *Ccr1* and *Cxcr3*, and their corresponding chemokines (i.e., *Ccl3*, *Ccl4*, *Ccl5*, *Ccl7*, *Ccl8* and *Cxcl9*, *Cxcl10*, respectively) were also upregulated (Supplementary Fig. S1). Chemokines *Ccl2*, *Ccl3*, *Ccl4*, *Ccl5*, *Ccl7*, *Ccl8* and the receptors *Ccr1*, *Ccr2* and *Ccr5*, which are specific to dendritic cells, were also upregulated in response to Af293 infection in mice (Supplementary Fig. S1).

Proteins associated with the innate response that we searched included β -defensins, C-type lectins (e.g., dectin-1), TLRs, complement, and adherence factors (e.g., VCAM or selectins) (Fig. 3, Supplementary Fig. S2 and S3). In general, increased duration of infection resulted in increasing number of transcripts for VCAM- and L-selectin, which would correlate with increased influx of host immune cells into the kidneys (Fig. 3). Furthermore, we noted increased expression of chemotactic complement factors, as well as some increased expression of mannose binding lectin (MBL), which can activate complement through an alternate pathway in response to binding of mannose in the fungal cell wall. Pentraxin-3 (Ptx3) and C-reactive protein also can activate the complement pathway and their expression was modulated postinfection (Supplementary Fig. S2). As might be expected, we noted increasing expression of C-type lectins (Fig. 3), and TLRs (Supplementary Fig. S2), which were highest at the day 8 after infection, also reflecting the influx of immune cells.

3.2. Activation of Th-1 and Th-17 cells

Transcripts encoding chemokines, *Ccl2*, *Cxcl9*, *Cxcl10*, *Cxcl11*, *Cxcl13*, and *Ccl20* (Supplementary Fig. S1) were up-regulated, as were the trafficking receptors for chemokines *Ccr5* and *Cxcr3*, associated with Th-1 cells and other trafficking receptors for chemokines, *Ccr2* and *Cxcr4*, which are associated with Th-17 cells.

3.3. Th-cell mediated immune response and interplay of cytokines

We found activation of a Th-1 response mediated by pro-inflammatory cytokines. The genes encoding IFN- γ and IL-27 (IL-12 family) were observed to be steadily up-regulated beginning 3 days postinfection (Supplementary Fig. S4). The effector cytokines, IL-18 and IL-24 were found to be up-regulated beginning 5 days postinfection in this model. IL-18 induces Th-1 response in the presence of IFN- γ and IL-12, whereas IL-24, a member of the IL-10 family, induces a Th-2 type response if differentiating cytokines IL-4 or IL-10 are present. On day 8 postinfection, we found upregulated transcripts encoding for IL-16; IL-16

promotes a Th-1 type response. We also observed activation of Th-17 pathway response genes mediated via pro-inflammatory cytokines IL-6 and IL-1 α on day 5 postinfection. We found an increased number of transcripts encoding for interleukin IL-1 β at day 5 postinfection, followed by a significant increase in transcript copies at day 8 postinfection (Supplementary Fig. 4).

Transcripts encoding IL-33, IL-34, IL-1f6 were upregulated in response to infection in kidney cells of mice. IL-34 transcripts had their highest expression at days 3 and 5 postinfection, but this was followed by a decline in transcript copies at day 8. IL-1f6 transcripts were up-regulated at day 5, and were highest at day 8 postinfection (Supplementary Fig. S4).

3.4. Iron regulation

A consistent enrichment of iron and heme binding, electron carrier activity and aromatase activity was found among downregulated genes (Fig. 1B). This was primarily a result of lower expression of members of the cytochrome P450 family of proteins during infection compared to baseline expression (Supplementary Table S10). However, a more detailed analysis revealed that key components of iron homeostasis had an increased numbers of transcripts in infected mice in comparison to uninfected mice, e.g.: *Nos1* (nitric oxide synthase 1), *Nos2* (nitric oxide synthase 2), lipocalin 2 (*Lcn2*), *p4ha2* (procollagen-proline), *Ogfod1* (2-oxoglutarate and iron dependent oxygenase domain containing 1), *Ltf* (lactotransferrin), *Epb4.2* (erythrocyte protein band 4.2), and *Heph* (hephaestin) (Supplementary Fig. 5). Transcripts that showed a decreased expression compared to uninfected mice were *Tdo2* (tryptophan 2,3-dioxygenase), *Lepre1* (leprecan 1), *Hbq 1a* (hemoglobin theta 1A), *Ttc7* (tetratricopeptide repeat domain 7), and *Hamp* (hepcidin, antimicrobial peptide) (Supplementary Fig. 5).

4. Discussion

The model of intravenous infection with *Aspergillus* was chosen to study the normal immune response to this infection in the absence of various immunosuppressive regimens that could be interposed, each of which would produce different perturbations of the immune response, and thus need each to be studied separately. Such studies can now follow, based on our data. To study the immune response in a setting absent the classical immunosuppressive risk factors also takes on topical relevance, owing to the rising incidence of aspergillosis in the non-immunocompromised [36]. The disease produced in the model also mimics disseminated infection with *Aspergillus*, where the kidney is a target organ, which is seen in one third of cases of aspergillosis [19]. Finally, this model has been used extensively in studies of anti-*Aspergillus* therapy [21] and the assets of this model have been discussed elsewhere [7, 37–38]. A limitation of the model used in this work is that the host is not immunocompromised, whereas most aspergillosis occurs in immunocompromised hosts. In the latter, the usual entry route is the lung, though tissue breaches can also lead to dissemination. Our original intention was to evaluate the interacting transcriptome of host and pathogen along the course of infection. Unfortunately, a very small number of sequencing reads was derived from *A. fumigatus* transcriptome, forcing us to focus our

analysis on the host. For those willing to implement similar experimental design, we would recommend the utilization of either mRNA depletion or enrichment technologies, such as hybrid selection [39].

The first line of defense against *Aspergillus* infections is the macrophages, neutrophils and dendritic cells that are recruited to the site of infection. These cells recognize pathogen associated molecular patterns (e.g. β -glucan, mannans, etc) that are exposed on the surface of germinating conidia or hyphae through pathogen recognition receptors (PRRs) including soluble receptors and cell bound receptors [40, 5, 41]. Among cell bound receptors, our data showed increased transcripts for genes encoding for C-type lectins (e.g., dectin-1), Toll-like receptors (TLR-2, TLR-3, TLR-8, TLR-9 and TLR-13); Ptx-3, and C-reactive protein were among the soluble receptors (Fig. 3 and Supplementary Fig. S2). In addition, we noted the increase of transcripts for β -defensins, which are cationic proteins found in the lysosomal contents of PMNs and contribute to the killing of microorganisms (Supplementary Fig. S2).

During kidney infection *A. fumigatus* cells are recognized by antigen presenting cells (APCs) and resident renal macrophages via PRRs. Furthermore, these immune cells induce the production of differentiating cytokines that act on naive CD4⁺Th-cells [42]. For instance, we observed upregulation of the differentiating cytokine IL-27 (IL-12 family) 3 days postinfection, which induces Th-1 cells, and the cytokine IL-6, 5 days postinfection, which induces Th-17 cells (Supplementary Fig. S4). Subsequently, release of various pro-inflammatory effector cytokines, like IL-1 β , IL-18, IFN- γ , IL-1 α , IL-34, and IL-6, suggests the possible role of Th1 and Th17 cell responses (Fig. 4) [43]. Similar to our observation at 24 h post infection, Armstrong-James, et al., did not observe a significant increase in the level of IFN- γ in pulmonary invasive aspergillosis [44]. In a similar study of invasive renal aspergillosis induced by *A. flavus*, Anand *et al.* observed significant induction of IFN- γ in kidney homogenates after 12 h of infection and levels of IFN- γ were higher along with IL-6, after 12, 72, and 96 hours in infected mice compared to uninfected mice [35]. Furthermore, in *A. flavus* mediated cerebral aspergillosis mouse model studies, brain homogenates showed an increased level of IFN- γ , IL-12p40 and IL-6 at 24 hours postinfection, with a decline in IL-4 and IL-23 cytokines [45]. Interestingly, we did not find signature transcripts encoding for the cytokines IL-4 and IL-10 suggesting the lack of an early Th2 type of response. However, we noted the upregulated transcripts encoding for Ccl11 and IL-33 at later time point during infection (Supplementary Fig. S1 and Fig. S4); both contribute to Th2 type response [46–47]. It should be noted that the effect of IL-33 can be either Th1 or Th2 depending on the disease and model [48–49]. We observed later during infection (i.e., day 5 postinfection) more transcripts for genes encoding for IL-1 β and IL-1 α indicated the possible role of Th17 pathway involvement and suppression of Th2 immune responsive cytokines (Supplementary Fig. S4). Similar to our observations, Caffrey, et al., also observed upregulation of IL-1 β and IL-1 α during the initial days of a pulmonary *A. fumigatus* murine model [50]. In addition, Cortez, et al., observed up-regulated transcripts of IL-1 β during the initial 2 h to 6h of interaction of *A. fumigatus* conidia with human monocytes [51]. IL-1 β also helps in activation of immune system and phagocytosis to decrease the load of *Aspergillus* at the infection site [52]. On the other hand, increased numbers of transcripts of the gene encoding for IFN- γ by day 5 postinfection (Supplementary Fig. S4) indicates earlier activation of the Th1 immune response, and suggests a possible co-evolution of Th1

and Th17 type immune responses during kidney infection. Interestingly, expression of IL-34 throughout infection suggests survival of monocytes and their differentiation into immunosuppressive macrophages at the site of infection [53]. We also observed increased transcripts of genes encoding for the chemokine Cxcl10 and its receptor Cxcr3 (Supplementary Fig. S1), which activates immune cells responsible for IFN- γ secretion and inhibition of Th17 regulating cytokines [54]. The current data showed a notable profile of proinflammatory cytokines (e.g., IL-27, IFN- γ , IL-6, IL-34, IL-16, IL-1 α and IL-1 β). Measurement of selected cytokines in the blood samples of aspergillosis patients may be promising as a tool for the monitoring of treatment responses [55–56].

Transcripts encoding numerous chemokines were upregulated. The trafficking receptor for chemokines Cxcl9, Cxcl10, and Cxcl11, is *Cxcr3*, and for chemokine Cxcl13, it is *Cxcr5*; we found transcripts for both these receptors were upregulated. These receptors are reported to be associated with Th-1 cells [57]. Similarly, we observed up-regulation of transcripts encoding for trafficking receptors *Ccr2* and *Cxcr4*. These receptors are present on Th17 cells and respond to different chemokines (i.e., Ccl2, Cxcl10, Cxcl13, Ccl20) [58]. Consistent with our observation, Cortez, et al., also observed up-regulation of transcripts encoding for the chemokine *Cxcl2*, *Ccl3*, *Ccl4* and *Ccl20* in human monocytes in response to *A. fumigatus* [51]. Chemokines are released at the site of infection by epithelial cells, macrophages and neutrophils and attract an influx of more neutrophils, monocytes and macrophages, thus causing the inflammation at the site of infection [59].

Members of cytochrome P450 family were downregulated on most days postinfection (Supplementary Table S10), recapitulating reports of transcription modulation of those enzymes by cytokines and interferons during hepatic and renal infections [60–63]. The biological underpinnings of this phenomenon are unclear. It has been proposed that the downregulation of cytochrome P450 genes during infection may be an attempt in reducing the damaging effects of oxidizing species derived from uncoupled catalytic turnover of those enzymes when in the presence of either iron or *Nos2* (up-regulated, Supplementary Fig. S5) [64]. Changes in the expression of host cytochrome P450 genes are especially relevant in the context of aspergillosis treatment and disease management. Cytochrome P450 enzymes have key roles in the activation and clearance of many drugs, including those used in the treatment of AIDS patients [65], one of the groups vulnerable to invasive aspergillosis [66].

The consistent downregulation of cytochrome P450 genes postinfection and the association of those genes with iron binding and aromatase activity, according to Gene Ontology, caused an enrichment of those pathways in the downregulated set (Fig. 1B). This may lead to the false interpretation that iron sequestration by the host is downregulated during *A. fumigatus* infection. Actually, the key components of iron homeostasis and regulation were up-regulated after infection (e.g. lipocalin 2, lactotransferrin) (Supplementary Fig. S5).

In response to the use of siderophores by pathogens to acquire iron from host iron binding proteins, the host produces proteins that sequester ferric ion-siderophore complexes away from pathogen siderophore receptors [67]. A member of this functional group, Lipocalin 2, is secreted by the host in response to infection and inflammation, and binds not only to the metabolites of the host but also iron-siderophore complexes, thus sequestering iron from the

pathogen [68]. Lactoferrin or lactotransferrin is a glycoprotein that is secreted by exocrine glands and neutrophils. Lactoferrin has various biological functions, including roles in iron metabolism, cell proliferation and differentiation [69]. The iron binding capacity of lactoferrin is twice that of transferrin (downregulated, Supplementary Fig. S5) [70]. Because of its iron binding properties, it reduces iron availability to any pathogens and also influences the immune system and cells involved in the inflammatory response [71].

Our study indicates downregulation of genes associated with aromatase activity (Fig. 1). Aromatase catalyzes estrogen synthesis from androgens (e.g. testosterone) [72]. Estrogens have both proinflammatory and anti-inflammatory properties [73]. The estrogen 17 β -estradiol, as well as IL-6, IL1 β , and TNF- α , stimulates aromatase activity, possibly in vascular endothelial cells or tissue macrophages [74–76]. Thus, it is possible that downregulated genes encoding for aromatase enzyme may be involved in downregulation of proinflammatory [74] or anti-inflammatory [77] cytokine production.

When an invading microbe interacts with host cells, tissue damage resulting from inflammation, thrombosis, and necrosis decreases available oxygen concentrations at the site of infection. The production of the non-ribosomal peptide gliotoxin, and possibly other secondary metabolites, by *A. fumigatus* contributes to the inhibition of angiogenesis in the lung [78–79]. However, during invasive infection, *A. fumigatus* hyphae invade the blood vessels, which leads to hypoxia at the site of infection and activates angiogenesis [78]. Angiogenesis is an important physiological response to tissue hypoxia and proinflammatory cytokines. At the site of infection, endothelial cells produce proinflammatory cytokines and leukocyte adhesion molecule chemokines, which recruit neutrophils and release H₂O₂ [78]. Among the earliest cues for initiation of the wound repair response is the release of “growth factor”-related cytokines [80]. We observed up-regulation of transcripts encoding for *Vegfa* (vascular endothelial growth factor) by day 3 of infection and *bFGF* (a-fibroblast growth factor) by day 1 infection. We also observed up-regulation of transcripts encoding for chemokines *Cxcl1*, *Cxcl2* and *Cxcl3*, all of which have been reported to be stimulatory factors for angiogenesis [57, 81–82].

Overall, our results show that infection in the kidneys with *A. fumigatus* triggers a host-response in the mouse that includes differential expression of 14,000 or less genes in the mouse. Our results show that genes such as C-type lectins, Toll-like receptors, C' proteins, and iron sequestration come under regulation during infection as part of the innate response. Furthermore, adaptive cellular immune response includes cytokine and chemokines and their receptors are regulated during the first week of infection. Surprisingly, we did not find evidence of a clear Th2 cellular response, with the exception of up-regulation of Ccl11 and IL-33 at later time points. We postulate that a robust response of this type may only occur later in infections that are less severe or more chronic; evidence for that would require supplementary work. Our study is an entry to understanding host responses to aspergillosis in tissue. Target organs more frequently involved in immunocompromised patients, such as the lung and brain need to be studied for similarities and differences. These studies form a basic dataset on which to build as more work, including other target organs, is performed in determining the global response of the host to infection with *Aspergillus*.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

These studies were funded in part by National Institute of Allergy and Infectious Diseases, National Institutes of Health (URL: nih.gov) grant 1R21AI85566-01 to DAS.

We thank Marife Martinez, Vicky Chen and Ana Claudia Maretti-Mira for their assistance in this work.

References

1. VandenBergh MF, Verweij PE, Voss A. Epidemiology of nosocomial fungal infections: invasive aspergillosis and the environment. *Diagn Microbiol Infect Dis*. 1999; 34(3):221–7. S0732889399000267 [pii]. [PubMed: 10403102]
2. Walsh TJ, , Stevens DA. Aspergillosis. In: Goldman L, , AS, editors *Cecil Textbook of Medicine* Vol. Chapter 347. Elsevier; 2011
3. Marr KA, Patterson T, Denning D. Aspergillosis. Pathogenesis, clinical manifestations, and therapy. *Infect Dis Clin North Am*. 2002; 16(4):875–94. vi. [PubMed: 12512185]
4. Park SJ, Mehrad B. Innate immunity to *Aspergillus* species. *Clin Microbiol Rev*. 2009; 22(4):535–51. DOI: 10.1128/CMR.00014-09 [PubMed: 19822887]
5. Romani L. Immunity to fungal infections. *Nat Rev Immunol*. 2011; 11(4):275–88. DOI: 10.1038/nri2939 [PubMed: 21394104]
6. Sales-Campos H, Tonani L, Cardoso CR, Kress MR. The immune interplay between the host and the pathogen in *Aspergillus fumigatus* lung infection. *Biomed Res Int*. 2013; 2013:693023.doi: 10.1155/2013/693023 [PubMed: 23984400]
7. Clemons KV, Stevens DA. The contribution of animal models of aspergillosis to understanding pathogenesis, therapy and virulence. *Med Mycol*. 2005; 43(Suppl 1):S101–10. [PubMed: 16110800]
8. Pagano L, Caira M, Nosari A, Van Lint MT, Candoni A, Offidani M, et al. Fungal infections in recipients of hematopoietic stem cell transplants: results of the SEIFEM B-2004 study-- Sorveglianza Epidemiologica Infezioni Fungine Nelle Emopatie Maligne. *Clin Infect Dis*. 2007; 45(9):1161–70. DOI: 10.1086/522189 [PubMed: 17918077]
9. Nagalakshmi U, Wang Z, Waern K, Shou C, Raha D, Gerstein M, et al. The transcriptional landscape of the yeast genome defined by RNA sequencing. *Science*. 2008; 320(5881):1344–9. DOI: 10.1126/science.1158441 [PubMed: 18451266]
10. Zhang W, Morris QD, Chang R, Shai O, Bakowski MA, Mitsakakis N, et al. The functional landscape of mouse gene expression. *J Biol*. 2004; 3(5):21.doi: 10.1186/jbiol16 [PubMed: 15588312]
11. Gibbons JG, Beauvais A, Beau R, McGary KL, Latge JP, Rokas A. Global transcriptome changes underlying colony growth in the opportunistic human pathogen *Aspergillus fumigatus*. *Eukaryot Cell*. 2012; 11(1):68–78. DOI: 10.1128/EC.05102-11 [PubMed: 21724936]
12. Rokas A, Gibbons JG, Zhou X, Beauvais A, Latge JP. The diverse applications of RNA-seq for functional genomic studies in *Aspergillus fumigatus*. *Ann N Y Acad Sci*. 2012; 1273:25–34. DOI: 10.1111/j.1749-6632.2012.06755.x [PubMed: 23230834]
13. Cerqueira GC, Arnaud MB, Inglis DO, Skrzypek MS, Binkley G, Simison M, et al. The *Aspergillus* Genome Database: multispecies curation and incorporation of RNA-Seq data to improve structural gene annotations. *Nucleic Acids Res*. 2014; 42(Database issue):D705–10. DOI: 10.1093/nar/gkt1029 [PubMed: 24194595]
14. Wang B, Guo G, Wang C, Lin Y, Wang X, Zhao M, et al. Survey of the transcriptome of *Aspergillus oryzae* via massively parallel mRNA sequencing. *Nucleic Acids Res*. 2010; 38(15):5075–87. DOI: 10.1093/nar/gkq256 [PubMed: 20392818]

15. Yu J, Fedorova ND, Montalbano BG, Bhatnagar D, Cleveland TE, Bennett JW, et al. Tight control of mycotoxin biosynthesis gene expression in *Aspergillus flavus* by temperature as revealed by RNA-Seq. FEMS Microbiol Lett. 2011; 322(2):145–9. DOI: 10.1111/j.1574-6968.2011.02345.x [PubMed: 21707733]
16. Cairns T, Minuzzi F, Bignell E. The host-infecting fungal transcriptome. FEMS Microbiol Lett. 2010; 307(1):1–11. DOI: 10.1111/j.1574-6968.2010.01961.x [PubMed: 20557573]
17. Oosthuizen JL, Gomez P, Ruan J, Hackett TL, Moore MM, Knight DA, et al. Dual organism transcriptomics of airway epithelial cells interacting with conidia of *Aspergillus fumigatus*. PLoS One. 2011; 6(5):e20527.doi: 10.1371/journal.pone.0020527 [PubMed: 21655222]
18. Morton CO, Fliesser M, Dittrich M, Mueller T, Bauer R, Kneitz S, et al. Gene expression profiles of human dendritic cells interacting with *Aspergillus fumigatus* in a bilayer model of the alveolar epithelium/endothelium interface. PLoS One. 2014; 9(5):e98279.doi: 10.1371/journal.pone.0098279 [PubMed: 24870357]
19. Wise GJ, Silver DA. Fungal infections of the genitourinary system. J Urol. 1993; 149(6):1377–88. [PubMed: 8501772]
20. Gautam P, Shankar J, Madan T, Sirdeshmukh R, Sundaram CS, Gade WN, et al. Proteomic and transcriptomic analysis of *Aspergillus fumigatus* on exposure to amphotericin B. Antimicrob Agents Chemother. 2008; 52(12):4220–7. DOI: 10.1128/AAC.01431-07 [PubMed: 18838595]
21. Clemons KV, Stevens DA. Comparative efficacies of four amphotericin B formulations--Fungizone, amphotec (Amphocil), AmBisome, and Abelcet--against systemic murine aspergillosis. Antimicrobial agents and chemotherapy. 2004; 48(3):1047–50. [PubMed: 14982807]
22. Clemons KV, Martinez M, Tong AJ, Stevens DA. Resistance of MBL gene-knockout mice to experimental systemic aspergillosis. Immunol Lett. 2010; 128(2):105–7. S0165-2478(10)00005-2 [pii]. DOI: 10.1016/j.imlet.2009.12.021 [PubMed: 20064561]
23. Liu M, Machova E, Nescakova Z, Medovarska I, Clemons KV, Martinez M, et al. Vaccination with mannan protects mice against systemic aspergillosis. Med Mycol. 2012; 50(8):818–28. DOI: 10.3109/13693786.2012.683539 [PubMed: 22587733]
24. Shankar J, Wu TD, Clemons KV, Monteiro JP, Mirels LF, Stevens DA. Influence of 17beta-estradiol on gene expression of *Paracoccidioides* during mycelia-to-yeast transition. PLoS One. 2011; 6(12):e28402.doi: 10.1371/journal.pone.0028402 [PubMed: 22194832]
25. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol. 2009; 10(3):R25.doi: 10.1186/gb-2009-10-3-r25 [PubMed: 19261174]
26. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics. 2011; 12:323.doi: 10.1186/1471-2105-12-323 [PubMed: 21816040]
27. Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of RNA-seq data. Genome Biol. 2010; 11(3):R25.doi: 10.1186/gb-2010-11-3-r25 [PubMed: 20196867]
28. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2010; 26(1):139–40. DOI: 10.1093/bioinformatics/btp616 [PubMed: 19910308]
29. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nat Protoc. 2013; 8(8):1494–512. DOI: 10.1038/nprot.2013.084 [PubMed: 23845962]
30. Supek F, Bosnjak M, Skunca N, Smuc T. REVIGO summarizes and visualizes long lists of gene ontology terms. PLoS One. 2011; 6(7):e21800.doi: 10.1371/journal.pone.0021800 [PubMed: 21789182]
31. Clemons KV, Stevens DA. Conventional or molecular measurement of *Aspergillus* load. Medical mycology. 2009; 47(Suppl 1):S132–7. [PubMed: 18654916]
32. Singh G, Imai J, Clemons KV, Stevens DA. Efficacy of caspofungin against central nervous system *Aspergillus fumigatus* infection in mice determined by TaqMan PCR and CFU methods. Antimicrobial agents and chemotherapy. 2005; 49(4):1369–76. 49/4/1369 [pii]. DOI: 10.1128/AAC.49.4.1369-1376.2005 [PubMed: 15793114]

33. Furlaneto CJ, Ribeiro FP, Hatanaka E, Souza GM, Cassatella MA, Campa A. Apolipoproteins A-I and A-II downregulate neutrophil functions. *Lipids*. 2002; 37(9):925–8. [PubMed: 12458630]
34. Maffei CML, Mirels LF, Sobel RA, Clemons KV, Stevens DA. Cytokine and Inducible Nitric Oxide Synthase mRNA Expression during Experimental Murine Cryptococcal Meningoencephalitis. *Infection and immunity*. 2004; 72(4):2338–49. DOI: 10.1128/IAI.72.4.2338-2349.2004 [PubMed: 15039359]
35. Anand R, Shankar J, Singh AP, Tiwary BN. Cytokine milieu in renal cavities of immunocompetent mice in response to intravenous challenge of *Aspergillus flavus* leading to aspergillosis. *Cytokine*. 2013; 61(1):63–70. DOI: 10.1016/j.cyto.2012.08.024 [PubMed: 23063795]
36. Stevens DA, Melikian GL. Aspergillosis in the ‘nonimmunocompromised’ host. *Immunol Invest*. 2011; 40(7–8):751–66. DOI: 10.3109/08820139.2011.614307 [PubMed: 21985304]
37. Clemons KV, Stevens DA. Animal models of *Aspergillus* infection in preclinical trials, diagnostics and pharmacodynamics: What can we learn from them? *Med Mycol*. 2006; 44(Supplement 1):S119–S26. DOI: 10.1080/13693780600871174
38. Mirkov I, Stosic-Grujicic S, Kataranovski M. Host immune defense against *Aspergillus fumigatus*: insight from experimental systemic (disseminated) infection. *Immunol Res*. 2012; 52(1–2):120–6. DOI: 10.1007/s12026-012-8274-x [PubMed: 22388638]
39. Melnikov A, Galinsky K, Rogov P, Fennell T, Van Tyne D, Russ C, et al. Hybrid selection for sequencing pathogen genomes from clinical samples. *Genome biology*. 2011; 12(8):R73. gb-2011-12-8-r73 [pii]. doi: 10.1186/gb-2011-12-8-r73 [PubMed: 21835008]
40. Thakur R, Shankar J. In silico Analysis Revealed High-risk Single Nucleotide Polymorphisms in Human Pentraxin-3 Gene and their Impact on Innate Immune Response against Microbial Pathogens. *Frontiers in microbiology*. 2016; 7:192.doi: 10.3389/fmicb.2016.00192 [PubMed: 26941719]
41. Brown GD. Dectin-1: a signalling non-TLR pattern-recognition receptor. *Nat Rev Immunol*. 2006; 6(1):33–43. DOI: 10.1038/nri1745 [PubMed: 16341139]
42. Thakur R, Anand R, Tiwari S, Singh AP, Tiwary BN, Shankar J. Cytokines induce effector T-helper cells during invasive aspergillosis; what we have learned about T-helper cells? *Front Microbiol*. 2015; 6:429.doi: 10.3389/fmicb.2015.00429 [PubMed: 26029179]
43. Stevens DA. Th1/Th2 in aspergillosis. *Med Mycol*. 2006; 44(s1):S229–S35. DOI: 10.1080/13693780600760773
44. Armstrong-James DP, Turnbull SA, Teo I, Stark J, Rogers NJ, Rogers TR, et al. Impaired interferon-gamma responses, increased interleukin-17 expression, and a tumor necrosis factor-alpha transcriptional program in invasive aspergillosis. *J Infect Dis*. 2009; 200(8):1341–51. DOI: 10.1086/605931 [PubMed: 19754306]
45. Anand R, Shankar J, Tiwary BN, Singh AP. *Aspergillus flavus* induces granulomatous cerebral aspergillosis in mice with display of distinct cytokine profile. *Cytokine*. 2015; 72(2):166–72. DOI: 10.1016/j.cyto.2015.01.006 [PubMed: 25647272]
46. Flaczyk A, Duerr CU, Shourian M, Lafferty EI, Fritz JH, Qureshi ST. IL-33 signaling regulates innate and adaptive immunity to *Cryptococcus neoformans*. *J Immunol*. 2013; 191(5):2503–13. jimmunol.1300426 [pii]. DOI: 10.4049/jimmunol.1300426 [PubMed: 23894196]
47. O’Dea EM, Amarsaikhan N, Li H, Downey J, Steele E, Van Dyken SJ, et al. Eosinophils are recruited in response to chitin exposure and enhance Th2-mediated immune pathology in *Aspergillus fumigatus* infection. *Infect Immun*. 2014; 82(8):3199–205. IAI.01990-14 [pii]. DOI: 10.1128/IAI.01990-14 [PubMed: 24842927]
48. Miller AM. Role of IL-33 in inflammation and disease. *J Inflamm (Lond)*. 2011; 8(1):22.doi: 10.1186/1476-9255-8-22 [PubMed: 21871091]
49. Smithgall MD, Comeau MR, Yoon BR, Kaufman D, Armitage R, Smith DE. IL-33 amplifies both Th1- and Th2-type responses through its activity on human basophils, allergen-reactive Th2 cells, iNKT and NK cells. *Int Immunol*. 2008; 20(8):1019–30. dxn060 [pii]. DOI: 10.1093/intimm/dxn060 [PubMed: 18550585]
50. Caffrey AK, Lehmann MM, Zickovich JM, Espinosa V, Shepardson KM, Watschke CP, et al. IL-1alpha signaling is critical for leukocyte recruitment after pulmonary *Aspergillus fumigatus*

challenge. PLoS Pathog. 2015; 11(1):e1004625.doi: 10.1371/journal.ppat.1004625PPATHOGENS-D-14-01338 [PubMed: 25629406]

51. Cortez KJ, Lyman CA, Kotttilil S, Kim HS, Roilides E, Yang J, et al. Functional genomics of innate host defense molecules in normal human monocytes in response to *Aspergillus fumigatus*. Infect Immun. 2006; 74(4):2353–65. DOI: 10.1128/IAI.74.4.2353-2365.2006 [PubMed: 16552065]
52. Savers A, Rasid OPM, Brock M, Jouvion G, Ryffel B, Cavaillon J-M, et al. Infection-Mediated Priming of Phagocytes Protects against Lethal Secondary *Aspergillus fumigatus* Challenge. PLoS one. 2016; 11(4):e0153829.doi: 10.1371/journal.pone.0153829 [PubMed: 27078879]
53. Foucher ED, Blanchard S, Preisser L, Garo E, Ifrah N, Guardiola P, et al. IL-34 induces the differentiation of human monocytes into immunosuppressive macrophages. antagonistic effects of GM-CSF and IFN γ . PLoS One. 2013; 8(2):e56045.doi: 10.1371/journal.pone.0056045 [PubMed: 23409120]
54. Liu M, Guo S, Hibbert JM, Jain V, Singh N, Wilson NO, et al. CXCL10/IP-10 in infectious diseases pathogenesis and potential therapeutic implications. Cytokine Growth Factor Rev. 2011; 22(3):121–30. DOI: 10.1016/j.cytogfr.2011.06.001 [PubMed: 21802343]
55. Chai L, Netea MG, Teerenstra S, Earnest A, Vonk AG, Schlamm HT, et al. Early proinflammatory cytokines and C-reactive protein trends as predictors of outcome in invasive Aspergillosis. J Infect Dis. 2010; 202(9):1454–62. DOI: 10.1086/656527 [PubMed: 20879853]
56. Rodland EK, Ueland T, Bjornsen S, Sagen EL, Dahl CP, Naalsund A, et al. Systemic biomarkers of inflammation and haemostasis in patients with chronic necrotizing pulmonary aspergillosis. BMC Infect Dis. 2012; 12:144.doi: 10.1186/1471-2334-12-144 [PubMed: 22731696]
57. Bendall L. Chemokines and their receptors in disease. Histol Histopathol. 2005; 20(3):907–26. [PubMed: 15944942]
58. Lim HW, Lee J, Hillsamer P, Kim CH. Human Th17 cells share major trafficking receptors with both polarized effector T cells and FOXP3+ regulatory T cells. J Immunol. 2008; 180(1):122–9. [PubMed: 18097011]
59. Borroni EM, Mantovani A, Locati M, Bonecchi R. Chemokine receptors intracellular trafficking. Pharmacol Ther. 2010; 127(1):1–8. DOI: 10.1016/j.pharmthera.2010.04.006 [PubMed: 20451553]
60. Richardson TA, Sherman M, Antonovic L, Kardar SS, Strobel HW, Kalman D, et al. Hepatic and renal cytochrome p450 gene regulation during citrobacter rodentium infection in wild-type and toll-like receptor 4 mutant mice. Drug Metab Dispos. 2006; 34(3):354–60. DOI: 10.1124/dmd.105.007393 [PubMed: 16339354]
61. Morgan ET. Impact of infectious and inflammatory disease on cytochrome P450-mediated drug metabolism and pharmacokinetics. Clin Pharmacol Ther. 2009; 85(4):434–8. DOI: 10.1038/clpt.2008.302 [PubMed: 19212314]
62. Morgan ET. Regulation of cytochromes P450 during inflammation and infection. Drug Metab Rev. 1997; 29(4):1129–88. DOI: 10.3109/03602539709002246 [PubMed: 9421688]
63. Theken KN, Deng Y, Kannon MA, Miller TM, Poloyac SM, Lee CR. Activation of the acute inflammatory response alters cytochrome P450 expression and eicosanoid metabolism. Drug metabolism and disposition: the biological fate of chemicals. 2011; 39(1):22–9. dmd.110.035287 [pii]. DOI: 10.1124/dmd.110.035287 [PubMed: 20947618]
64. Morgan ET, Ullrich V, Daiber A, Schmidt P, Takaya N, Shoun H, et al. Cytochromes P450 and flavin monooxygenases--targets and sources of nitric oxide. Drug Metab Dispos. 2001; 29(11):1366–76. [PubMed: 11602511]
65. Walubo A. The role of cytochrome P450 in antiretroviral drug interactions. Expert Opin Drug Metab Toxicol. 2007; 3(4):583–98. DOI: 10.1517/17425225.3.4.583 [PubMed: 17696808]
66. Denning DW, Follansbee SE, Scolaro M, Norris S, Edelstein H, Stevens DA. Pulmonary Aspergillosis in the Acquired Immunodeficiency Syndrome. New England Journal of Medicine. 1991; 324(10):654–62. DOI: 10.1056/NEJM199103073241003 [PubMed: 1994248]
67. Leal SM Jr, Roy S, Vareechon C, Carrion S, Clark H, Lopez-Berges MS, et al. Targeting iron acquisition blocks infection with the fungal pathogens *Aspergillus fumigatus* and *Fusarium oxysporum*. PLoS Pathog. 2013; 9(7):e1003436.doi: 10.1371/journal.ppat.1003436 [PubMed: 23853581]

68. Flo TH, Smith KD, Sato S, Rodriguez DJ, Holmes MA, Strong RK, et al. Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature*. 2004; 432(7019):917–21. nature03104 [pii]. DOI: 10.1038/nature03104 [PubMed: 15531878]
69. Adlerova L, Bartoskova A, Faldyna M. Lactoferrin: a review. *Veterinari Medicina*. 2008; 53(9): 457–68.
70. Aisen P, Leibman A. Lactoferrin and transferrin: a comparative study. *Biochim Biophys Acta*. 1972; 257(2):314–23. [PubMed: 4336719]
71. Legrand D, Ellass E, Carpentier M, Mazurier J. Lactoferrin: a modulator of immune and inflammatory responses. *Cell Mol Life Sci*. 2005; 62(22):2549–59. DOI: 10.1007/s00018-005-5370-2 [PubMed: 16261255]
72. McPhaul MJ, Noble JF, Simpson ER, Mendelson CR, Wilson JD. The expression of a functional cDNA encoding the chicken cytochrome P-450arom (aromatase) that catalyzes the formation of estrogen from androgen. *Journal of Biological Chemistry*. 1988; 263(31):16358–63. [PubMed: 3182796]
73. Marriott I, Huet-Hudson YM. Sexual dimorphism in innate immune responses to infectious organisms. *Immunologic Research*. 2006; 34(3):177–92. DOI: 10.1385/ir:34:3:177 [PubMed: 16891670]
74. Villaggio B, Soldano S, Cutolo M. 1,25-dihydroxyvitamin D3 downregulates aromatase expression and inflammatory cytokines in human macrophages. *Clin Exp Rheumatol*. 2012; 30(6):934–8. [PubMed: 23253631]
75. Simpson E, Rubin G, Clyne C, Robertson K, O'Donnell L, Jones M, et al. The role of local estrogen biosynthesis in males and females. *Trends Endocrinol Metab*. 2000; 11(5):184–8. [PubMed: 10856920]
76. Simpson ER, Zhao Y, Agarwal VR, Michael MD, Bulun SE, Hinshelwood MM, et al. Aromatase expression in health and disease. *Recent Prog Horm Res*. 1997; 52:185–213. discussion -4. [PubMed: 9238853]
77. Cuzzocrea S, Santagati S, Sautebin L, Mazzon E, Calabro G, Serraino I, et al. 17beta-estradiol antiinflammatory activity in carrageenan-induced pleurisy. *Endocrinology*. 2000; 141(4):1455–63. [PubMed: 10746650]
78. Ben-Ami R, Lewis RE, Leventakos K, Kontoyiannis DP. *Aspergillus fumigatus* inhibits angiogenesis through the production of gliotoxin and other secondary metabolites. *Blood*. 2009; 114(26):5393–9. DOI: 10.1182/blood-2009-07-231209 [PubMed: 19843884]
79. Hsu JL, Khan MA, Sobel RA, Jiang X, Clemons KV, Nguyen TT, et al. *Aspergillus fumigatus* invasion increases with progressive airway ischemia. *PloS one*. 2013; 8(10):e77136. [pii]. doi: 10.1371/journal.pone.0077136PONE-D-13-14976 [PubMed: 24155924]
80. Frantz S, Vincent KA, Feron O, Kelly RA. Innate immunity and angiogenesis. *Circ Res*. 2005; 96(1):15–26. DOI: 10.1161/01.RES.0000153188.68898.ac [PubMed: 15637304]
81. Romagnani P, Annunziato F, Lasagni L, Lazzeri E, Beltrame C, Francalanci M, et al. Cell cycle-dependent expression of CXC chemokine receptor 3 by endothelial cells mediates angiostatic activity. *The Journal of clinical investigation*. 2001; 107(1):53–63. DOI: 10.1172/JCI9775 [PubMed: 11134180]
82. Miyake M, Goodison S, Urquidi V, Gomes Giacoia E, Rosser CJ. Expression of CXCL1 in human endothelial cells induces angiogenesis through the CXCR2 receptor and the ERK1/2 and EGF pathways. *Lab Invest*. 2013; 93(7):768–78. labinvest201371 [pii]. DOI: 10.1038/labinvest.2013.71 [PubMed: 23732813]

Figure 1. Genes differentially expressed during the course of infection in comparison to non-infected mouse (day 0)

(A) Colored bars indicate the number of transcripts detected on each day (X-axis) and the day they were first detected according to colors in the legend. Only transcripts with significant difference in expression (FDR lower than 0.01 and 16-fold change) when compared to day 0 were accounted. Bars marked with letters represent sets of transcripts with significant enrichment (Bonferroni adjusted p-value < 0.05) of at least one GO term. Gene Ontology enrichment analysis is described in the Material and Methods section. Bars marked with an asterisk indicate sets with no significant enrichment. (B) Summarized list of GO terms enriched in the sets defined on panel A. GO summarization is described in the material and methods section. Gray blocks indicate the set (table header) in which the GO term has a proportionally higher frequency than expected by chance.

Figure 2. Cluster of transcripts presenting a similar profile of expression during the course of the infection

(A) Graphs depicting the normalized FPKM (Y-axis) of members of each cluster during the 8 days of sample collection (X-axis). Dark blue lines describe the approximate general pattern presented by members of each cluster. Graphs with caption in red indicate clusters with enriched GO terms. (B) Summarized list of GO terms enriched on cluster 7, 10 and 18. Gray blocks indicate the set (table header) in which the GO term has a proportionally higher frequency than expected. GO enrichment analysis and GO terms summarization are described in the Material and Methods section.

Figure 3. Expression profile of differentially expressed C-type lectins and VCAM genes
Only genes differentially expressed in at least one pair of time points are shown. Color indicates the expression level of each transcript in terms of Fragments per Kilobase of transcript per Million of mapped reads (FPKM) on each time point: lowest FPKM values are depicted in white and the highest values are shown in darkest blue, according to the legend. The dendrogram indicates the similarity of expression profile between transcripts.

APCs = Antigen Presenting cells. MAPK (Mitogen activated protein kinases), d- day of infection. IFNAR2; Interferon αβ receptor 2.

Figure 4. Interplay of cytokines in mouse model of invasive aspergillosis

The diagram shows known and postulated immune pathways that play a role in the hosts' response to *A. fumigatus* infection. These include up-regulated (shown in red) and down-regulated (shown in green) gene transcripts of mice, after intravenous inoculation of Af293. The interaction of conidia with kidney cells leads to the release of cytokines and chemokines at the site of infection, which recruit and activate inflammatory and immune cells (e.g. phagocytes either circulating or resident macrophages), antigen presenting cells (APCs; e.g. dendritic cells) and T cells. Phagocytic cells engulf the circulating conidia and destroy them via phagolysosome activity and also release of cytokines. APCs migrate and present *Aspergillus* antigens in the lymph nodes, to activate naïve CD4⁺ T cells via MHC class II molecules. Consequently, activated T cells differentiate into Th-1 cells type via IFN-γ and IL-24 or a Th-17 cells via IL-6 and IL-1α/IL-1β.