Published in final edited form as:

J Dev Orig Health Dis. 2018 August ; 9(4): 401–408. doi:10.1017/S2040174418000107.

Adipose Tissue Uncoupling Protein 1 Levels and Function are Increased in a Mouse Model of Developmental Obesity Induced by Maternal Exposure to High Fat Diet

Egle Bytautiene Prewit¹, Craig Porter², Mauricio La Rosa¹, Nisha Bhattarai², Huaizhi Yin¹, Phyllis Gamble¹, Talar Kechichian¹, and Labros S. Sidossis^{2,3}

¹Department of Obstetrics & Gynecology, The University of Texas Medical Branch at Galveston, TX

²Department of Surgery, The University of Texas Medical Branch at Galveston, Shriners Hospitals for Children-Galveston, TX

³Department of Kinesiology and Health Department of Medicine, Rutgers University, New Brunswick, NJ

Abstract

With brown adipose tissue (BAT) becoming a possible therapeutic target to counteract obesity, the prenatal environment could represent a critical window to modify BAT function and browning of white AT. We investigated if levels of Uncoupling Protein 1 (UCP1) and UCP1-mediated thermogenesis are altered in offspring exposed to prenatal obesity. Female CD-1 mice were fed a high-fat (HF) or standard fat (SF) diet for 3 months before breeding. After weaning, all pups were placed on SF. UCP1 mRNA and protein levels were quantified using q RT-PCR and Western blot analysis, respectively, in brown (BAT), subcutaneous (SAT), and visceral (VAT) adipose tissues at 6 months of age. Total and UCP1-dependent mitochondrial respiration were determined by high-resolution respirometry. A Student *t* test and Mann-Whitney test were used (significance: $P < .05$). UCP1 mRNA levels were not different between the HF and SF offspring. UCP1 protein levels, total mitochondrial respiration, and UCP1-dependent respiration were significantly higher in BAT from HF males ($P = .02$, $P = .04$, $P = .005$, respectively) and females ($P = .01$, $P = .04$, $P = .02$, respectively). In SAT, the UCP1 protein were significantly lower in HF females ($P = .03$), and the UCP1-dependent thermogenesis was significantly lower from HF males ($P = .04$). In VAT, UCP1 protein levels and UCP1-dependent respiration were significantly lower only in HF females ($P = .$

Corresponding author: Egle Bytautiene Prewit, MD, PhD, 301 University Blvd, Galveston, TX 77555-1062, Phone: 409-747-5139, Fax: 409-772-2261, egbytaut@utmb.edu.

Author Contributions: EB conceived the idea for this project, obtained animals and samples, analyzed the results, and completed the manuscript. CP interpreted mitochondria respiratory experiments, significantly contributed to results interpretation, and prepared the manuscript. MLR, NB, HY, PG, and TK developed and performed experiments and obtained the data. LS provided critical guidance on the project, helped interpret the findings, and prepared the manuscript.

CONFLICTS OF INTEREST

None

ETHICAL STANDARDS

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national guides on the care and use of laboratory animals (National Institutes of Health) and have been approved by the institutional committee (the Animal Care and Use Committee at The University of Texas Medical Branch at Galveston, Texas).

03, $P = .04$, respectively). There were no differences in total respiration in SAT and VAT. Prenatal exposure to maternal obesity leads to significant increases in UCP1 levels and function in BAT in offspring with little impact on UCP1 levels and function in SAT and VAT.

Keywords

brown adipose tissue; UCP1; developmental programming; obesity

INTRODUCTION

According to the National Health and Nutrition Examination Survey (NHANES, 2003–2006), 34% of adults in the US meet the criteria for metabolic syndrome (MetS), a cluster of major cardiovascular risk factors related to obesity and insulin resistance.¹ Obesity is one of the key risk factors in the development of MetS. Of the 3 most common chronic conditions associated with MetS—namely, hypertension, hypercholesterolemia, and diabetes—45% of adults are reported to have at least 1, 13% to have 2, and 3% to have all 3 conditions. Moreover, an estimated 15% of US adults have 1 or more of these conditions without a diagnosis.²

Traditionally, research into the causes of obesity has focused on lifestyle factors in childhood and adulthood, but recently the role of pregnancy in fueling the obesity epidemic has begun to be recognized. The proportion of women who are obese has doubled since the 1960s.³ In 2003, the prevalence of overweight and obese women reached 61.6%.⁴ In the US, almost one-third of women of reproductive age (20–39 years) are obese.⁵ Furthermore, the prevalence of obesity in pregnancy in the US ranges between 18.5% and 38.3%, and these rates continue to rise.^{6–8}

Epidemiological studies have shown that children born of pregnancies complicated by obesity are at increased risk of obesity and other features of MetS.⁹ After controlling for confounders, the relative risk of childhood obesity associated with maternal obesity in the first trimester of pregnancy has been found to increase 2-fold at 2–3 years of age and 2.3-fold at 4 years.¹⁰ MetS is more common in adolescents who were large-for-gestational-age at birth or who were born to obese mothers.¹¹ Maternal BMI was found to be one of the strongest predictors of excessive weight and obesity in sons at 18 years.¹² Adults born to obese mothers were more insulin resistant than those born to nonobese mothers.¹³

Thus, maternal obesity is a significant factor contributing to increased risk of obesity in offspring. While epidemiological studies provide important insight into the origins of obesity in humans, animal models remain essential for confirming the epidemiological associations, studying the mechanisms involved, and developing novel therapies.

Multiple animal studies have shown that offspring born of pregnancies complicated by obesity are at increased risk of obesity, increased blood pressure, and other features of MetS.^{9–18} The mechanisms by which offspring develop cardiovascular disease have been linked to epigenetic changes, endothelial dysfunction, inflammation, oxidative stress, lipogenesis,

nonalcoholic fatty liver disease, adipocyte hypertrophy, pancreatic abnormalities, and alterations in hypothalamic satiety signaling.^{16,18-21}

Brown adipose tissue (BAT) has recently emerged as a possible therapeutic target to counteract obesity and insulin resistance in humans. BAT is present in small amounts in the fetus. After birth, BAT is thought to be gradually replaced with white adipose tissue (WAT). However, recent data have demonstrated that adult humans have small (~50-100 g) BAT depots. Moreover, in rodents and humans, WAT can undergo a process called “browning,” in which brown or beige adipocytes appear within larger WAT depots.²³ Brown and beige adipocytes are characterized by multilocular lipid droplets in adipocytes and an abundance of mitochondria enriched with uncoupling protein 1 (UCP1). BAT and beige adipocytes in WAT may play an important role in nonshivering thermogenesis in humans.²⁴

Thermogenesis in UCP1-positive adipocytes can be stimulated by exposure to cold, β -adrenergic receptor agonists, and diet. BAT activation can increase energy expenditure and alter fat and glucose metabolism in animals, including humans. Indeed, there is a significant negative relationship between increasing BMI and BAT volume in humans, supporting a putative role for BAT in human obesity. However, whether maternal obesity affects the function of BAT and browning of WAT in offspring has not been addressed. The objectives of this study were to determine the UCP1 expression and to investigate total and UCP1-dependent thermogenesis by using mitochondrial respiration as a proxy in BAT and visceral (VWAT) and subcutaneous (SWAT) white adipose tissue from mice offspring born to obese mothers.

METHOD

This work was performed in line with guidelines set out by the National Institutes of Health for the care and use of laboratory animals. The study protocol and all related procedures were reviewed and approved by the Animal Care and Use Committee at The University of Texas Medical Branch at Galveston, Texas. The mice were maintained in the animal care facility at The University of Texas Medical Branch. They were housed separately in temperature (22°C) and humidity-controlled quarters with constant light:dark cycles of 12 h: 12 h. Animals were provided with food and water ad libitum. Mice were fed either a standard-fat diet (SF group, 5.8 gm%; Teklad 7012: Harlan Teklad LM-485 Mouse/Rat Sterilizable Diet, Harlan Teklad, Madison, WI) or a high-fat diet (HF group, 34.9 gm%; D12492, Research Diets, Inc, New Brunswick, NJ). The source of fat in the HF diet is lard, while the SF chow does not contain animal product; the source of fat in the SF show is soybean oil.

Study design

Female CD-1 mice were obtained from Charles River Laboratories at approximately 4-5 weeks of age. Animals were randomly assigned to either the SF-diet group (control mice, SF group) or to the HF-diet group (diet-induced obesity mice, HF group). After 3 months on the assigned diet, mice were mated with a CD-1 male, maintained on standardized diet. Female mice randomized to the HF diet for 12-14 weeks weighed significantly more than those

randomized to the SF group (HF 44.8± 1.7 gm, n=11 vs SF 34.8± 1.0 gm, n=16, $P<.001$). Therefore, we categorized HF-diet mothers as obese during the gestation period.

All pups were weighed at 1 day of age. During lactation, mothers were fed the diet they were originally assigned (ie, SF or HF). At 21 days post partum, offspring from all mothers were weaned onto a SF diet, containing 5.8 gm% fat. At 6 months of age, both female and male offspring were studied for the manifestations of MetS.

Two additional cohorts of offspring obtained according to the protocol above were used for mitochondrial respirometry experiments and to determine UCP1 mRNA expression. In these cohorts, animals were sacrificed at 6 month of age, and BAT, SWAT, and VWAT (including mesenteric, perirenal, and gonadal fat) were collected. Tissues from one cohort were immediately transferred to an ice-cold preservation buffer and stored on ice as described previously.²⁵ Mitochondrial respirometry experiments were performed on this fresh tissue. Approximately 50-100 mg of collected tissues from the other cohort were immediately frozen in liquid nitrogen and stored in -80°C until further analysis for mRNA expression and protein levels.

In Vivo telemetric blood pressure measurement experiments

To implant the internal blood pressure transducers, the mice were anesthetized with a mixture of ketamine (80-100 mg/kg, Ketalar, Parke-Davis, Morris Plains, NJ) and xylazine (5–10 mg/kg, Gemini, Rugby, Rockville Center, NY) as described previously.²⁶ A vertical midline skin incision along the neck was made, and the submaxillary glands were gently separated. The left common carotid artery located next to the trachea was carefully isolated. The catheter (diameter 0.4 mm) was then introduced into the carotid artery through a small incision in the vessel wall, while the body of the transducer (PA-C10 model, Data Systems International [DSI], Overland Park, KS) was secured in a subcutaneous pouch along the animal's right flank through the same ventral neck incision. The neck incision was closed using 6-0 silk. Mice were kept warm on a heating pad and monitored closely until full recovery from anesthesia.

Blood pressure recordings of male and female offspring began 48-72 hours after surgical implantation of the pressure transducer and were continuously monitored for 5 consecutive days using RLA 1020 telemetry receivers (DSI). The information was entered in the data acquisition and recording system, Dataquest software (A.R.T.3.1; Gartner Dataquest, Stamford, CT). Then, the mice were sacrificed, and blood and tissues were collected for later analysis.

Serological studies

Glucose levels were measured in whole blood using a OneTouch Ultra, a blood glucose monitoring system (LifeScan, CA), after mice fasted overnight (16-18 hours). Commercially available kits were used according to the manufacturer's instructions to determine serum levels of cholesterol (Cayman, Ann Arbor, MC), triglycerides (BioAssay Systems, Hayward, Ca), insulin, leptin (both using magnetic bead MILLIPLEX map kit, cat# MBN2B-41K, minimum detect concentration for insulin 14.4 pg/mL, leptin 4.4 pg/mL, Millipore, Billerica, MA), and free fatty acids (BioVision, Mountain View, CA).

RNA isolation and reverse-transcriptase reaction

BAT, SWAT, and VWAT samples from each mouse were analyzed individually. Samples were homogenized utilizing the Bullet Blender from Next Advance (Averill Park, NY). Total RNA was isolated using Trizol and Zymo RNA isolation kits (Ambion, Austin, TX, and Zymo Research Corporation, Irvine, CA) according to manufacturers' instructions. Quantification of RNA was performed by measuring the absorbance of RNA sample solutions at 260 nm. For the reverse transcription reaction, a High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA) was used according to the manufacturer's instructions (25°C for 10 minutes → 37°C for 120 minutes → 85°C for 5 minutes → 4°C for infinity).

Quantitative real-time PCR

mRNA expression of *Ucp1* was quantified using TaqMan Gene Expression Assays (Mm01244861_m1, Life Technologies, Grand Island, NY) according to the manufacturer's instructions for quantitative real-time PCR performed in Real-Time PCR System 7500 Fast (Applied Biosystems). Gene expression was calculated as the mRNA of the targeted gene relative to β -actin mRNA levels (Mm00607939_s1, Life Technologies) in each specific sample (relative unit). Each reaction was carried out in triplicate. Relative quantification was determined using 7500 software, Version 2.06 (Applied Biosystems).

Western blot analysis

Proteins from BAT, SWAT, and VWAT were extracted after homogenization in a RIPA (Radio-immunoprecipitation assay) buffer with freshly-added protease inhibitor cocktail and PMSF (phenylmethylsulfonyl fluoride) and were quantified using the BCA Protein Assay kit (Pierce, Rockford, IL). The protein extracts were electrophoretically separated by SDS/PAGE on a 4%-15% running gel, transferred to PVDF membranes, blocked, and incubated overnight with primary polyclonal antibodies detecting UCP1 (1:500 dilution, 32 kDa, ab10983, Abcam Inc, Cambridge, MA). The blots were then incubated for 1 hour at room temperature with a rabbit peroxidase-labelled antibody (1:10000 dilution, NA934VS, GE, Pittsburg, PA). The bands were then exposed using the Luminata Crescendo Western HRP Substrate (Millipore, Billerica, MA) and captured on autoradiography film. The obtained bands were scanned and analyzed using ImageJ 1.42q software (NIH). Band signal intensity was measured by densitometry, and intensity values for the UCP1 were normalized against those for β -actin (42 kDa, 1:20,000, Sigma- Aldrich, St. Louis, MO).

Mitochondrial respirometry

Mitochondrial respiratory function was determined in digitonin-permeabilized adipose tissue samples on the same day of sample collection. Tissue respiratory rates were quantified using an Oroboros Oxygraph-O2K™ respirometer as detailed previously with minor modifications.^{25,27} Briefly, respiration in the leak state was achieved after titration of pyruvate (5mM), octanoyl-l-carnitine (0.07mM), malate (2mM), and glutamate (10mM) into the O2K-chamber. Total leak respiration was determined before and after the addition of the UCP1 inhibitor guanosine diphosphate (GDP, 20mM). The reduction in respiration after

GDP titration is equal to UCPI-dependent thermogenesis. This approach represents a direct quantification of UCPI function.²⁸⁻³⁵

Data analysis

Systolic and diastolic blood pressure data for each animal were obtained from the telemetry system as a 24-hour average for the last 4 days of the recording period. Then, the overall average for each mouse was calculated, and data were analyzed using the GraphPad Prism 5 software version 5.04 for Windows (GraphPad Software, La Jolla, CA). All other results were expressed as the mean $\pm$ SEM and compared between the groups. For statistical analyses, a Student *t* test and Mann-Whitney test were applied as appropriate (GraphPad Prism 5). A probability value (*P*value) of .05 was considered statistically significant.

RESULTS

A mouse model of developmental MetS

Maternal characteristics before and during pregnancy—Compared with female SF mice, the HF group gained significantly more weight during the 3-month pre-pregnancy period (HF 44.8 ± 1.7 gm vs SF 34.8 ± 1.0 gm, $P < .001$) and had significantly higher cholesterol levels (HF -172.6 ± 15.4 mg/dL, $n=6$, vs SF 98.0 ± 18.0 mg/dL, $n=7$, $P = .01$).

Offspring characteristics at 6 months of age (adulthood)—The developmental programming of MetS was characterized in 1-2 offspring randomly selected from each litter. The results are shown in Table 1. Males and females born to obese mothers were significantly heavier ($P = .03$ and $P = .04$, respectively) and had a significantly higher percentage of VWAT ($P = .03$ and $P = .02$, respectively) than their counterparts born to SF mothers.

Systolic and diastolic blood pressure in males ($P = .04$ and $P = .03$, respectively) and females ($P = .0001$ and $P = .02$, respectively) born to obese mothers were significantly higher than in offspring of control dams.

Exposure to maternal pre-pregnancy obesity led to significantly higher levels of fasting glucose ($P = .007$), total cholesterol ($P = .01$), triglycerides ($P = .02$), free fatty acids ($P = .03$), and leptin ($P = .0002$) in males, compared with those born to SF mothers. In HF female offspring, significantly higher levels of cholesterol ($P = .04$), leptin ($P = .02$), and insulin ($P = .04$) were determined.

Collectively, in this model of maternal obesity, offspring from obese mothers exhibited indicators of MetS—obesity, dyslipidemia, and elevated blood pressure—at 6 months of age. The development of some of the metabolic parameters was also sex specific.

UCP1 mRNA expression, protein levels, thermogenesis—In BAT, UCP1 mRNA levels were not different between the groups (Fig 1a). UCP1 protein levels were significantly higher in BAT from HF males ($P = .04$) and HF females ($P = .04$) when compared to SF (Figs 1b, c). Total leak respiration and UCPI-dependent respiration were significantly higher

in BAT from HF males ($P = .005$, $P = .02$, respectively) and HF females ($P = .02$, $P = .01$, respectively) when compared to SF (Figs 2*a*, *b*).

In SAT, UCP1 mRNA expression was not different between the offspring born to HF and SF mothers (Fig 3*a*). UCP1 protein levels were significantly lower in HF females ($P = .03$, Figs 3*b*, *c*) when compared to SF females. The total leak respiration was not different between the groups (Fig 4*a*). UCP1-dependent thermogenesis in SAT was significantly lower only in HF males ($P = .04$, Fig 4*b*) when compared to SF group.

In VWAT, UCP1 mRNA was not different between the groups (Fig 5*a*). And UCP1 protein levels were significantly lower in HF females ($P = .03$, Fig 5*b*, *c*). Total leak respiration did not differ between the offspring groups (Fig 6*a*). UCP1-dependent thermogenesis was significantly lower in HF females ($P = .04$, Fig 6*b*), with no differences between male groups.

DISCUSSION

Our study confirmed the development of MetS in 6-month-old offspring born to obese mothers, despite being fed a regular diet after weaning. We determined that in adulthood, BAT thermogenic capacity is significantly modified by maternal obesity. UCP1 protein levels in BAT correspond with the direction of UCP1-dependent thermogenesis. Interestingly, the mRNA signal for UCP1 was not different between groups, suggesting that mRNA is a poor proxy of UCP1 protein and function in long-term studies. Prenatal obesity had a much less pronounced effect of UCP1 levels and function in WAT. In SWAT and VAT, alterations in thermogenesis and UCP1 protein levels were sex-dependent. Overall, our data indicate that the thermogenic capacity of BAT is altered in the offspring of obese mothers.

Multiple animal studies have shown that offspring born of pregnancies complicated by obesity are at increased risk of obesity and other features of MetS.^{9-18,36} Similarly, in our model, offspring born to obese mothers were significantly heavier and had a significantly higher percentage of adipose tissue compared to mice born to nonobese mothers, despite being fed a regular-fat diet after weaning. In line with previous reports, we demonstrated sex-related cardiovascular dysfunction in this study. Systolic and diastolic BPs in females and systolic BP in males were significantly higher in mice born to obese mothers. Exposure to maternal obesity led to significantly higher levels of fasting glucose, triglycerides, free fatty acids, and leptin in males exposed to maternal obesity, compared with those born to normal weight mothers. In female offspring born to obese mothers, significantly higher levels of cholesterol, leptin, and insulin were found. Thus, by 6 months of age, offspring exhibited indicators of MetS—obesity, dyslipidemia, and elevated blood pressure—if their mothers were obese before pregnancy, indicating the suitability of the model to study the developmental programming of MetS. Moreover, in line with other models of developmental MetS, our mouse model exhibited gender differences.

Several animal models have been used to investigate BAT function in offspring. Maternal caloric-, protein-, or food-restricted diets during pregnancy have led to a reduction in the expression of UCP1 and depression of the thermogenic capacity in BAT and negatively

affected the browning of WAT in offspring.^{37–40} Others have reported increased UCP1 expression in late-gestation ovine fetuses after prenatal exposure to hypoxia.⁴¹ Overfeeding rodents in the first few weeks of life have resulted in lower levels of UCP1 and changes in BAT structure in adulthood.^{42,43} Supplementation of obese ewes with L-arginine have led to a significant increase in fetal BAT, among other metabolic improvements, while maternal diet supplementation with olive oil, protein, or fiber has increased UCP1 expression in the BAT of rat pups.^{40,44,45} Only one of these studies reported gender differences in UCP1 expression.³⁷ Ours is the first study to demonstrate that UCP1-dependent mitochondrial thermogenesis and UCP1 levels are altered in offspring born to obese mothers, regardless of being fed a regular diet postnatally. Thus, in the current study, we fill an important knowledge gap regarding offspring's BAT function after exposure to maternal obesity. All the aforementioned studies, including our own, provide evidence that the in utero environment plays an important role in the development of BAT and its function in offspring.

An intriguing result in our current study is the higher levels of UCP1-dependent mitochondrial thermogenesis and UCP1 protein expression in the BAT of offspring born to obese mothers when compared to controls. We hypothesize that this increased nonshivering thermogenesis in BAT is a counteractive process against obesity in offspring. However, what is most interesting is that while both male and female offspring exposed to prenatal obesity developed BAT with greater UCP1 levels and thermogenic function, these animals were still bigger, had a greater fat mass percentage, were hypertensive, and had altered systemic lipid metabolism when compared to control animals. Thus, while increased BAT function in animals exposed to prenatal obesity may be a countermeasure to prevent adiposity, it is a response that falls short of being fully successful. That said, it does not mean that the increase in BAT thermogenic capacity seen here is not physiologically significant. However, these data do suggest a disassociation between BAT function and metabolic health in this model.

While exposure to prenatal obesity had a large impact on BAT UCP1 function in both male and female offspring, we observed a far less pronounced effect in white adipose depots. There was a sexual dimorphism in the response of SAT and VAT to prenatal obesity in terms of UCP1 protein levels, where maternal obesity reduced UCP1 protein in female but not male offspring. However, this effect only translated to lower UCP1 function in VAT in female offspring. Moreover, in SWAT, we found that UCP1-dependent thermogenesis, but not UCP1 protein levels, were modified in males exposed to prenatal obesity. Importantly, it should be noted that UCP-dependent thermogenesis was ~50-100-fold greater in interscapular BAT vs SWAT or VWAT. Indeed, a clear concordance between UCP protein levels and UCP-dependent thermogenesis was seen in BAT only. It may be that the vastly lower levels of UCP1 and capacity for UCP1-dependent respiration in WAT depots compared to BAT may limit our ability to demonstrate clear associations between UCP1 abundance and function in WAT depots. We found no difference in UCP1 gene expression in any of white fat depots investigated, while UCP1 protein levels were differentially regulated. These findings raise the possibility that UCP1 translation is epigenetically regulated and that exposure to maternal obesity induces negative translational regulation, mediated by miRNAs in the offspring's BAT and WAT. Micro-RNAs are short (~20-23 nucleotides in length)

endogenous noncoding RNA molecules that inhibit translation. It had been established that multiple miRNAs are involved in the development and differentiation of adipose tissue, including BAT and BWAT.^{46,47} Another explanation could be as follows—increased thermogenic capacity in BAT brought about by chronic cold exposure first causes a big increase in UCP1 mRNA, which is eventually caught up by higher UCP1 protein. So, perhaps we are looking at the wrong time point to see differences in UCP1 mRNA. That said, UCP1 mRNA does not produce heat, only the protein does, and we observed significant differences in BAT UCP1 protein and function in the current study, despite an absence in mRNA levels.

Several limitations of our study should be acknowledged. We investigated only 1 age group—the adult mice. We did not evaluate homeostasis model assessment of insulin resistance (HOMA-IR) since fasting glucose and insulin were not measured in the same animals. We did not expose our mice to a cold temperature or to β -adrenergic receptor agonists, which are known to induce nonshivering thermogenesis, though it has been shown that an animal house temperature of 22°C is significantly below the mouse thermoneutral temperature (28°C–30°C).^{29,48} Also, the question of whether altered thermogenesis in BAT is a consequence of exposure to maternal obesity or a consequence of the offspring's own obesity needs to be answered in the future. Moreover, teasing out the effects of obesity vs high fat diet exposure during pregnancy, during gestation and lactation needs to be performed.

Our study offers proof of concept that UCP1 protein levels and function are altered in BAT (and to a lesser degree WAT) of offspring exposed to prenatal obesity. With increasing interest in BAT as a possible therapeutic target to counteract obesity and MetS, we have demonstrated that an in utero environment could represent a critical window to modify BAT function.

Acknowledgments

Language editing and figure layout assistance were provided by the Publication, Grant, and Media Support Office funded by the Department of Obstetrics & Gynecology at The University of Texas Medical Branch at Galveston.

FINANCIAL SUPPORT

Funding for this study was provided by the Shriner's Hospital for Children grants 84090, 85310, and 85410; NIH/NICHD grant K12HD052023; and the Department of Surgery, UTMB.

References

- Writing G, Roger VrL, Go AS, et al. Executive Summary: Heart Disease and Stroke Statistics—2011 Update. *Circ*. 2011; 123(4):459–463.
- Fryar CD, Hirsch R, Eberhardt MS, Yoon SS, Wright JD. Hypertension, high serum total cholesterol, and diabetes: racial and ethnic prevalence differences in U.S. adults, 1999–2006. *NCHS Data Brief*. 2010; 36:1–8.
- American Heart Association. Statistical fact Sheet. Women and Cardiovascular Diseases: Statistics2006<http://www.americanheartorg/downloadable/heart/1136818052118Females06pdf>
- Thom T, Haase N, Rosamond W, et al. Heart Disease and Stroke Statistics—2006 Update: A Report From the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circ*. 2006; 113(6):e85–151.

5. Huda SS, Brodie LE, Sattar N. Obesity in pregnancy: prevalence and metabolic consequences. *Semin Fetal Neonatal Med.* 2010; 15(2):70–76. [PubMed: 19896913]
6. Galtier-Dereure F, Boegner C, Bringer J. Obesity and pregnancy: complications and cost. *American J Clin Nutr.* 2000; 71(5 Suppl):1242S–1248S.
7. Kim SY, Dietz PM, England L, Morrow B, Callaghan WM. Trends in Pre-pregnancy Obesity in Nine States, 1993–2003. *Obesity.* 2007; 15(4):986–993. [PubMed: 17426334]
8. Lu GC, Rouse DJ, DuBard M, Cliver S, Kimberlin D, Hauth JC. The effect of the increasing prevalence of maternal obesity on perinatal morbidity. *Am J Obstet Gynecol.* 2001; 185(4):845–849. [PubMed: 11641663]
9. Catalano PM, Ehrenberg HM. The short- and long-term implications of maternal obesity on the mother and her offspring. *BJOG.* 2006; 113(10):1126–1133. [PubMed: 16827826]
10. Whitaker RC. Predicting preschooler obesity at birth: the role of maternal obesity in early pregnancy. *Pediatrics.* 2004; 114(1):e29–e36. [PubMed: 15231970]
11. Boney CM, Verma A, Tucker R, Vohr BR. Metabolic syndrome in childhood: association with birth weight, maternal obesity, and gestational diabetes mellitus. *Pediatrics.* 2005; 115(3):e290–e296. [PubMed: 15741354]
12. Koupil I, Toivanen P. Social and early-life determinants of overweight and obesity in 18-year-old Swedish men. *Int J Obes.* 2008; 32(1):73–81.
13. Mingrone G, Manco M, Valera Mora ME, et al. Influence of Maternal Obesity on Insulin Sensitivity and Secretion in Offspring. *Diabetes Care.* 2008; 31(9):1872–1876. [PubMed: 18535193]
14. Desai M, Jellyman JK, Han G, Beall M, Lane RH, Ross MG. Maternal obesity and high-fat diet program offspring metabolic syndrome. *Am J Obstet Gynecol.* 2014; 211(3):237.e231–237.e237. [PubMed: 24631702]
15. Forsen T, Eriksson JG, Tuomilehto J, Teramo K, Osmond C, Barker DJ. Mother's weight in pregnancy and coronary heart disease in a cohort of finnish men: follow up study. *BMJ.* 1997; 315(7112):837–840. [PubMed: 9353502]
16. Khan IY, Dekou V, Douglas G, et al. A high-fat diet during rat pregnancy or suckling induces cardiovascular dysfunction in adult offspring. *Am J Physiol Regul Integr Comp Physiol.* 2005; 288(1):R127–R133. [PubMed: 15308487]
17. Khan IY, Taylor PD, Dekou V, et al. Gender-Linked Hypertension in Offspring of Lard-Fed Pregnant Rats. *Hypertension.* 2003; 41(1):168–175. [PubMed: 12511548]
18. Samuelsson AM, Matthews PA, Argenton M, et al. Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance: a novel murine model of developmental programming. *Hypertension.* 2008; 51(2):383–392. [PubMed: 18086952]
19. Oben JA, Mouralidarane A, Samuelsson AM, et al. Maternal obesity during pregnancy and lactation programs the development of offspring non-alcoholic fatty liver disease in mice. *J Hepatol.* 52(6):913–920.
20. Oben JA, Patel T, Mouralidarane A, et al. Maternal obesity programmes offspring development of non-alcoholic fatty pancreas disease. *Biochem Biophys Res Commun.* 2010; 394(1):24–28. [PubMed: 20170634]
21. Suter MA, Ma J, Vuguin PM, et al. In utero exposure to a maternal high-fat diet alters the epigenetic histone code in a murine model. *Am J Obstet Gynecol.* 2014; 210(5):463–463. [PubMed: 24793723]
22. Persichetti A, Sciuto R, Rea S, et al. Prevalence, Mass, and Glucose-Uptake Activity of 18F-FDG-Detected Brown Adipose Tissue in Humans Living in a Temperate Zone of Italy. *PLoS One.* 2013; 8(5):e63391. [PubMed: 23667608]
23. Lo Kinyui A, Sun L. Turning WAT into BAT: a review on regulators controlling the browning of white adipocytes. *Biosci Rep.* 2013; 33(5):e00065. [PubMed: 23895241]
24. Fenzl A, Kiefer FW. Brown adipose tissue and thermogenesis. *Horm Mol Biol Clin Investig.* 2014; 19(1):25–37.
25. Porter C, Herndon DN, Bhattarai N, et al. Severe Burn Injury Induces Thermogenically Functional Mitochondria in Murine White Adipose Tissue. *Shock.* 2015; 44(3):258–264. [PubMed: 26009824]

26. Bytautienė E, Tamayo E, Kechichian T, et al. Prepregnancy obesity and sFlt1-induced preeclampsia in mice: developmental programming model of metabolic syndrome. *Am J Obstet Gynecol.* 2011; 204(5):398.e391–398.e398. [PubMed: 21444063]
27. Porter C, Hurren NM, Cotter MV, et al. Mitochondrial respiratory capacity and coupling control decline with age in human skeletal muscle. *Am J Physiol Endocrinol Metab.* 2015; 309(3):E224–E232. [PubMed: 26037248]
28. Cannon B, Nedergaard J. Respiratory and thermogenic capacities of cells and mitochondria from brown and white adipose tissue. *Methods Mol Biol.* 2001; 155:295–303. [PubMed: 11293080]
29. Cannon B, Nedergaard J. Nonshivering thermogenesis and its adequate measurement in metabolic studies. *J Exp Biol.* 2010; 214(2):242–253.
30. Cannon B, Nedergaard J. Studies of thermogenesis and mitochondrial function in adipose tissues. *Methods Mol Biol.* 2008; 456:109–121. [PubMed: 18516556]
31. Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev.* 2004; 84(1):277–359. [PubMed: 14715917]
32. Matthias A, Ohlson KB, Fredriksson JM, Jacobsson A, Nedergaard J, Cannon B. Thermogenic responses in brown fat cells are fully UCP1-dependent. UCP2 or UCP3 do not substitute for UCP1 in adrenergically or fatty acid-induced thermogenesis. *J Biol Chem.* 2000; 275(33):25073–25081. [PubMed: 10825155]
33. Nedergaard J, Cannon B. UCP1 mRNA does not produce heat. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids.* 2013; 1831(5):943–949. [PubMed: 23353596]
34. Shabalina IG, Ost M, Petrovic N, Vrbicky M, Nedergaard J, Cannon B. Uncoupling protein-1 is not leaky. *Biochim Biophys Acta.* 2010; 1797(6–7):773–784. [PubMed: 20399195]
35. Shabalina IG, Petrovic N, de Jong JM, Kalinovich AV, Cannon B, Nedergaard J. UCP1 in Brite/Beige Adipose Tissue Mitochondria Is Functionally Thermogenic. *Cell Reports.* 2013; 5(5):1196–1203. [PubMed: 24290753]
36. Elahi MM, Cagampang FR, Mukhtar D, Anthony FW, Ohri SK, Hanson MA. Long-term maternal high-fat feeding from weaning through pregnancy and lactation predisposes offspring to hypertension, raised plasma lipids and fatty liver in mice. *Br J Nutr.* 2009; 102(04):514–519. [PubMed: 19203419]
37. Watkins AJ, Lucas ES, Wilkins A, Cagampang FRA, Fleming TP. Maternal Periconceptional and Gestational Low Protein Diet Affects Mouse Offspring Growth, Cardiovascular and Adipose Phenotype at 1 Year of Age. *PLoS One.* 2011; 6(12):e28745. [PubMed: 22194901]
38. Ojha S, Robinson L, Yazdani M, Symonds ME, Budge H. Brown adipose tissue genes in pericardial adipose tissue of newborn sheep are downregulated by maternal nutrient restriction in late gestation. *Pediatr Res.* 2013; 74(3):246–251. [PubMed: 23788058]
39. Felipe A, Villarroya F, Mampel T. Effects of maternal hypocaloric diet feeding on neonatal rat brown adipose tissue. *Biol Neonate.* 1988; 53(2):105–112. [PubMed: 2833318]
40. Priego T, Sanchez J, Garcia A, Palou A, Pico C. Maternal Dietary Fat Affects Milk Fatty Acid Profile and Impacts on Weight Gain and Thermogenic Capacity of Suckling Rats. *Lipids.* 2013; 48(5):481–495. [PubMed: 23417844]
41. Myers DA, Hanson K, Mlynarczyk M, Kaushal KM, Ducas CA. Long-term hypoxia modulates expression of key genes regulating adipose function in the late-gestation ovine fetus. *Am J Physiol Regul Integr Comp Physiol.* 2008; 294(4):R1312–R1318.
42. de Almeida DL, Fabricio GS, Trombini AB, et al. Early Overfeed-Induced Obesity Leads to Brown Adipose Tissue Hypoactivity in Rats. *Cell Physiol Biochem.* 2013; 32(6):1621–1630. [PubMed: 24335411]
43. Xiao XQ, Williams SM, Grayson BE, et al. Excess Weight Gain during the Early Postnatal Period Is Associated with Permanent Reprogramming of Brown Adipose Tissue Adaptive Thermogenesis. *Endocrinology.* 2007; 148(9):4150–4159. [PubMed: 17525123]
44. Carey Satterfield M, Dunlap K, Keisler D, Bazer F, Wu G. Arginine nutrition and fetal brown adipose tissue development in diet-induced obese sheep. *Amino Acids.* 2012; 43(4):1593–1603. [PubMed: 22327565]
45. Maurer AD, Reimer RA. Maternal consumption of high-prebiotic fibre or -protein diets during pregnancy and lactation differentially influences satiety hormones and expression of genes

involved in glucose and lipid metabolism in offspring in rats. *Br J Nutr.* 2011; 105(03):329–338. [PubMed: 21129233]

46. Arias N, Aguirre L, Fernandez-Quintela A, et al. MicroRNAs involved in the browning process of adipocytes. *J Physiol Biochem.* 2015:1–13.
47. Guller I, McNaughton S, Crowley T, et al. Comparative analysis of microRNA expression in mouse and human brown adipose tissue. *BMC Genomics.* 2015; 16(1):1–11. [PubMed: 25553907]
48. Quevedo S, Roca P, Pico C, Palou A. Sex-associated differences in cold-induced UCP1 synthesis in rodent brown adipose tissue. *Pflugers Arch.* 436(5):689–695. [PubMed: 9716701]

Fig 1.

(a) UCP1 mRNA (for both sexes SF=6, HF=5), (b) protein levels (SF males n=3, HF males n=5, SF females n=5, HF females n=3), and (c) Western (for both sexes SF n=5, HF n=6) blots in BAT from offspring at 6 months of age born to control (SF) and obese (HF) mice. Values are presented as mean $\pm$ SD.

Fig 2.

(a) Total leak respiration (n=6 in each group) and (b) UCPI-dependent respiration (SF males n=6, HF males n=7, SF females n=7, HF females n=7) in BAT from offspring at 6 months of age born to control (SF) and obese (HF) mice. Values are presented as mean $\pm$ SD.

Fig 3. (a) UCP1 mRNA (SF males n=4, HF males n=3, SF females n=5, HF females n=3), (b) protein levels (SF males n=4, HF males n=3, SF females n=4, HF females n=5), and (c) Western blots (for both sexes SF n=5, HF n=6) in SAT from offspring at 6 months of age born to control (SF) and obese (HF) mice. Values are presented as mean $\pm$ SD.

Fig 4.

(a) Total leak respiration (SF males $n=5$, HF males $n=8$, SF females $n=6$, HF females $n=8$) and (b) UCPI-dependent respiration (SF males $n=4$, HF males $n=6$, SF females $n=6$, HF females $n=8$) in SAT from offspring at 6 months of age born to control (SF) and obese (HF) mice. Values are presented as mean $\pm$ SD.

Fig 5. (a) UCP1 mRNA (SF males n=5, HF males n=5, SF females n=5, HF females n=4), (b) protein levels (SF males n=5, HF males n=5, SF females n=4 HF females n=4), and (c) Western blots (for both sexes SF n=5, HF n=6) in VBAT from offspring at 6 months of age born to control (SF) and obese (HF) mice. Values are presented as mean ± SD.

Fig 6.

(a) Total leak respiration (SF males n=7, HF males n=9, SF females n=5, HF females n=7) and (b) UCPI-dependent respiration (SF males n=7, HF males n=8, SF females n=5, HF males n=5) in VWAT from offspring at 6 months of age born to control (SF) and obese (HF) mice. Values are presented as mean $\pm$ SD.

Table 1

Offspring characteristics at 6 months of age.

Parameters	MALES			FEMALES		
	SF group (n)	HF group (n)	P value	SF group (n)	HF group (n)	P value
Body weight (gm)	46.4± 1.0 (15)	50.4± 1.4 [*] (12)	.03	41.8 ±2.1 (11)	48.4 ±1.2 [*] (6)	.04
Adiposity (%)	4.1 ±0.5 (10)	5.7 ± 0.5 [*] (11)	.03	6.9 ±1.1 (11)	10.7 ±0.5 [*] (6)	.02
Systolic BP (mmHg)	128.8 ±3.3 (12)	140.4 ±4.1 [*] (12)	.04	116.3 ±2.3 (14)	129.2 ± 1.5 [*] (12)	.0001
Diastolic BP (mmHg)	104.2± 3.5 (12)	118.7 ±5.1 (11)	.03	95.3 ±2.1 (14)	102.4 ± 2.5 [*] (13)	.02
Fasting glucose (mg/dL)	120.7 ± 6.6 (15)	193.1 ±31.9 [*] (8)	.007	116.0 ± 6.0 (6)	120.6 ±2.4 (5)	.52
Cholesterol (mg/dL)	116.2 ± 11.9 (10)	180.2 ± 20.3 [*] (10)	.01	94.93 ±12.61 (6)	141.8 ± 15.10 [*] (6)	.04
Triglycerides (mg/mL)	126.6± 13.9 (9)	195.7 ± 26.0 [*] (5)	.02	146.1 ±18.7 (4)	186.9 ± 73.7 (4)	.49
Free fatty acids	29.1 ±4.3 (10)	42.3 ±3.7 [*] (10)	.03	35.6 ± 5.7 (12)	39.6 ± 3.5 (12)	.55
Leptin (ng/mL)	5.0 ±0.3 (5)	14.8 ± 1.4 [*] (6)	.0002	9.8 ±3.0 (5)	19.2 ± 1.9 [*] (6)	.02
Insulin (ng/mL)	3.7 ±1.1 (4)	7.0 ±1.0 (5)	.06	3.6 ±0.6 (6)	6.5 ±1.2 [*] (4)	.04

* indicates statistically significant differences; SF—standard fat; HF—high fat; BP—blood pressure

Mean ± SEM (number of mice).