CORRESPONDENCE

The Risks Associated with Obesity in Pregnancy

by PD Dr. med. Johannes Stubert, Dr. med. Steffi Hartmann, PD Dr. med. Frank Reister, and Prof. Dr. med. Wolfgang Janni in issue 16/2018

Benefits of Breastfeeding

The prevalence of obesity has increased in Germany and is continuing to rise. For this reason, the topic is of great relevance in the setting of pregnancy care. The authors write that interventions have not been found to be successful and that therefore preventive measures are gaining in importance. In this context, it is important not only to consider pregnancy, but also the actual birth and breastfeeding.

The long-term effects for mothers are positive: on the one hand, if they breastfeed for more than three months, this leads to long-term changes in their metabolism and helps prevent type 2 diabetes. On the other hand, breastfeeding also protects mothers from cardiovascular disorders. Both these facts are important, especially in obese women, because they often develop gestational diabetes during their pregnancies (1–3).

The authors also described (4) effects on the newborns of overweight mothers. Postnatally, the risk for hypoglycemia is raised. Early colostrum feeding, 30 minutes postpartum in the delivery suite, is the best prevention (AWMF [Association of the Scientific Medical Societies in Germany] guideline on caring for healthy and mature neonates born to diabetic mothers).

There is hardly any primary prevention method that yields a comparable success to breastfeeding for children regarding overweight when they are older.

As obesity impairs lactogenesis phase II, these women breastfeed less and their breastfeeding period is shortened, the topic of breastfeeding should be discussed with those women during their pregnancies. Gynecologists have a crucial role in this. It may be valuable in terms of supporting health promotion in women and their babies to recommend that they book an antenatal appointment with an International Board Certified Lactation Consultant (IBCLC) in the Professional Association of Lactation Consultants in Germany.

DOI: 10.3238/arztebl.2018.0453a

References

1. Much D, Beyerlein A, Kindt A, et al.: Lactation is associated with altered metabolomic signatures in women with gestational diabetes. *Diabetologia* 2016; 59: 2193–202.
2. WHO (eds.): Impact of breastfeeding on maternal and child health. *Acta paediatrica* 2015; 104 (Issue Supplement S46): 1–134. <https://onlinelibrary.wiley.com/toc/16512227/104/S467> (last accessed on 14 June 2018).
3. Hansen K: Breastfeeding: a smart investment in people and in economies. *The Lancet* 2016; 387: 10017.
4. Stubert J, Reister F, Hartmann S, Janni W: The risks associated with obesity in pregnancy. *Dtsch Arztebl Int* 2018; 115: 276–83.

Gudrun von der Ohe

Physician and International Board Certified Lactation Consultant (IBCLC)
Hamburg
postfach@stillberatung.info

Conflict of interest statement

The author is on the staff of the European Institute for Breastfeeding and Lactation (www.stillen-institut.com) and is an expert adviser with the WHO/ UNICEF initiative "Verein Babyfreundlich" (www.babyfreundlich.org).

Migration Background Should Be Considered

A substantial proportion of pregnant women in the maternity facilities in large German cities and conurbations have a migration background. In order to ensure that high quality healthcare can be delivered, it is important to consider this fact when discussing overweight and pregnancy risks.

In a study funded by the German Research Foundation (DFG), we investigated peripartum health outcomes in 7100 pregnant women at three large maternity hospitals in Berlin, whose parturition course we followed prospectively (1, 2). Induced births were more common in overweight and obese women, but no differences were seen between women with and without a migration background. Similarly, in the context of secondary cesarean sections, we observed an association only with overweight, not with a migration background. This indicates that peripartum care and support are of a high quality (1, 2).

We used Cox regression models to compare rates of large-for-gestational-age (LGA) neonates, small-for-gestational-age (SGA) neonates, and premature babies. A total of 2586 women with a migration background were included, the largest subgroups were from Turkey and Lebanon. The comparison group consisted of 2676 women without a migration background. Of the women of Turkish origin, 53.6% were overweight or obese; compared to 33.4% of the women without a migration background. The greatest risks were seen for LGA babies, both in women with and without a migration background. SGA babies were less frequent in the group of overweight women. The attributable risks (proportion of preventable cases in the population under study) for LGA neonates, preterm births, and extreme preterm births were 16.3%, 3.6%, and 16.5%, respectively (3). In the group of women with a migration background, the attributable risks owing to overweight/obesity were thus higher than those due to smoking (3). Women with a migration background therefore require particular attention with regard to the risk factor overweight.

DOI: 10.3238/arztebl.2018.0453b

References

1. Reiss K, Breckenkamp J, Borde T, et al.: The association of pre-pregnancy overweight and obesity with delivery outcomes: a comparison of immigrant and non-immigrant women in Berlin, Germany. *Int J Public Health* 2016; 61: 455–63.
2. David M, Borde T, Brenne S, Henrich W, Breckenkamp J, Razum O: Caesarean section frequency among immigrants, second- and third-generation women, and non-immigrants: prospective study in Berlin/Germany. *PLoS ONE* 2015; 10: e0127489.
3. Reiss K, Breckenkamp J, Borde T, Brenne S, David M, Razum O: Contribution of overweight and obesity to adverse pregnancy outcomes among immigrant and non-immigrant women in Berlin, Germany. *Eur J Public Health* 2015; 25: 239–44.
4. Stubert J, Reister F, Hartmann S, Janni W: The risks associated with obesity in pregnancy. *Dtsch Arztebl Int* 2018; 115: 276–83.

Prof. Dr. med. Oliver Razum

Abteilung Epidemiologie und International Public Health
Fakultät für Gesundheitswissenschaften
Universität Bielefeld, oliver.razum@uni-bielefeld.de

Prof. Dr. med. Matthias David

Charité Berlin, Campus Virchow-Klinikum
Klinik für Gynäkologie

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We wholeheartedly agree with Gudrun von der Ohe's response to our article. In the context of breastfeeding advice in general, the benefits for the neonate should be

discussed, but so should the benefits for the mothers themselves. Emphasizing the positive effects of breastfeeding on normalizing weight and metabolism is also a way to improve women's willingness to breastfeed.

Professor Razum and Professor David underline the associations between obesity and pregnancy that we identified in our article for a population subgroup. The prevalence of obesity in women with a migration background reflects—among other things—the situation in their countries of origin (1). The particular attention our colleagues demand is to be supported, but in all experience, any pregnancy advice or instruction is particularly challenging in these women, owing to language and cultural barriers.

DOI: 10.3238/arztebl.2018.0454a

References

1. Global Health Observatory (GHO) data: Prevalence of obesity among adults. www.who.int/gho/ncd/risk_factors/overweight_obesity/obesity_adults/en/ (last accessed on 29 May 2018).
2. Stubert J, Reister F, Hartmann S, Janni W: The risks associated with obesity in pregnancy. *Dtsch Arztebl Int* 2018; 115: 276–83.

On behalf of the authors

PD Dr. med. habil. Johannes Stubert

Universitätsfrauenklinik und Poliklinik
am Klinikum Südstadt Rostock
johannes.stubert@uni-rostock.de

Conflict of interest statement

The author declares that no conflict of interest exists.

CLINICAL SNAPSHOT

Arthralgia and an Unusual Radiological Finding

An 18-year-old man complained of pain in an ankle of three days' duration, without any antecedent trauma, and presented for radiological examination, accompanied by his mother. The image revealed multiple, irregular, rounded, oval foci of diminished transparency. The patient was of Turkish ancestry; the bone changes were typical of osteopoikilosis, a very rare, benign bone malformation first described in 1915, with familial clustering in eastern Anatolia and along the shores of the Black Sea. This is a hereditary condition with an autosomal dominant inheritance pattern. The skeletal changes are usually discovered as an incidental finding in childhood or adolescence; as the individual grows, they may increase or decrease in size or disappear entirely. No treatment is needed. This patient's symptoms resolved after treatment with anti-inflammatory drugs and were presumably due to a nonspecific arthropathy or tendinopathy.

Dr. med. Dietmar Häussler, chirurgische Praxis Laupheim, Praxis für Chirurgie und Unfallchirurgie, Illerkirchberg, dietmarhaeusler@web.de

Conflict of interest statement: The authors state that they have no conflict of interest.

Translated from the original German by Ethan Taub, M.D.

Cite this as: Häussler D: Arthralgia and an unusual radiological finding. *Dtsch Arztebl Int* 2018; 115: 454.
DOI: 10.3238/arztebl.2018.0454b