RESEARCH ARTICLE

[View Article Online](#)
[View Journal](#) | [View Issue](#)

Cite this: *Med. Chem. Commun.*,
2017, 8, 1946

An agent for optical imaging of TrkC-expressing, breast cancer†

Anyanee Kamkaew,^{ab} Feng Li,^c Zheng Li^{*c} and Kevin Burgess^{id} ^{*a}

Tropomyosin receptor kinases receptor C is expressed at high levels on the surface of tumors from metastatic breast cancer, metastatic melanoma, glioblastoma, and neuroblastoma. Previous studies have shown synthetic TrkC ligands bearing agents for photodynamic therapy could be used to completely ablate 4T1 metastatic breast tumors and suppress metastatic spread *in vivo*. Modification of these probes (**A** in the text) to make them suitable for near infrared optical imaging *in vivo* would require a substantial increase in molecular mass (and hence increased vulnerability to undesirable absorption, metabolism and immunogenicity effects), or significant changes to the probe design which might compromise binding to TrkC in histochemical studies and on live cells. The research featured here was undertaken to investigate if the second strategy could be achieved without compromising binding to TrkC-expressing tissues. Specifically, an “aza-BODIPY” probe was synthesized to replace a spacer fragment in the original probe **A**. In the event, this new probe design (**1a** in the text) binds TrkC⁺ breast cancer in live cell cultures, in histochemical studies and in an *in vivo* murine model. Probe **1a** binds TrkC⁺ tissues with good contrast with respect to healthy tissues, and much more strongly than an isomeric, non-TrkC binding, probe (**1b**) prepared as a negative control.

Received 28th June 2017,
Accepted 17th August 2017

DOI: 10.1039/c7md00328e

rsc.li/medchemcomm

Introduction

Tropomyosin receptor kinases (Trk's) are cell-surface receptors that bind the neurotrophin growth hormones.¹ Docking neurotrophin-3 (NT3) with TrkC, for instance, triggers intracellular phosphorylation then a network of cell signaling processes that lead to cell growth and differentiation.^{2,3} Overexpression of Trk receptors is associated with several forms of cancer, including neuroblastoma,^{4–16} glioblastoma,^{14,17–21} metastatic breast cancer^{22–28} and metastatic melanoma.^{29–37} Consequently, functionalized Trk ligands may be prepared as possible imaging agents for cancer.^{28,38–40} Synthetic TrkB⁴¹ and pan-Trk-binding agents¹⁶ functionalized with radionuclei have been explored for imaging with positron emission tomography (PET), but, to the best of our knowledge, no synthetic imaging agents have been reported for selective optical-imaging of TrkC *in vivo*.

Prior research from these laboratories featured the TrkC-homing ligand in structure **A**. Probe **A** was shown to be internalized by TrkC-expressing cells and it could also be used as a staining agent for metastatic breast cancer and melanoma tissue, both of which are associated with TrkC overexpression. Moreover, one 10 mg kg^{−1} dose of **A** coupled to a BODIPY-based PDT agent was shown to almost completely ablate a 4T1 primary tumor, and suppress metastatic spread.⁴²

Probe **A** is not ideal for optical imaging because it absorbs light optimally at a wavelength (around 500 nm) that is too short for efficient penetration of more than a few mm in tissue.^{43,44} To circumvent this, the most obvious solution would be to prepare similar molecules with a near-IR-absorbing probe directly replacing the BODIPY. However, modifications to fluors to make them near-IR absorbing invariably increase their molecular size; this is undesirable because larger structures tend to be more vulnerable to undesirable absorption, metabolism, and excretion effects. We reasoned that one way to reconcile these opposing design criteria is to replace the black part of structure **A** with a near-IR fluor giving the generic structures **B** (Fig. 1). Designs **B** eliminate the triazine core which serves no purpose other than as a scaffold-spacer. One possibility would be to replace it with an aza-BODIPY dye; these dyes tend to have significantly longer absorption and emission wavelength maxima than their BODIPY analogs.^{45–47}

^a Department of Chemistry, Texas A & M University, Box 30012, College Station, TX 77842, USA. E-mail: burgess@tamu.edu

^b School of Chemistry, Institute of Science, Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand

^c Center for Bioenergetics, Houston Methodist Research Institute, Houston, TX 77030, USA. E-mail: zli@houstonmethodist.org

† Electronic supplementary information (ESI) available: Scheme and full details of compounds syntheses and characterizations, cell toxicity of **1a** and TrkC antibody stained on breast tissues. See DOI: 10.1039/c7md00328e

Fig. 1 First generation TrkC targeting ligand-conjugate **A** and the design concept in this paper **B**.

Based on the considerations outlined above, we designed aza-BODIPY dyes **1a** and **b** (Fig. 2). Probe **1a** was appealing because the sulfonic acid groups would enhance the hydrophilicity of the molecule and their negative charge might reduce non-specific binding to cells. Conversely, **1b** has the advantages of smaller size, and more facile synthetic access. We anticipated the best of these two probe candidates, based on solubility and spectroscopic experiments, would be tested for: (i) binding and internalization by TrkC-expressing cells; (ii) staining TrkC-expressing fixed tissue (histochemistry); and, (iii) comparative distribution in healthy mice and ones bearing TrkC-expressing breast cancer.

Results and discussion

Probe syntheses

Coupling of the azido acid from L-tyrosine⁴⁸ with the disulfonated aza-BODIPY **2**, prepared as described,⁴⁹ gave the diazide **3** (Scheme 1). A copper-mediated cycloaddition^{50,51} of **3** with the alkyne derived from L-isoleucine⁴⁸ gave the desired product **1a**. Probe **1b** was prepared *via* an analogous route, but begin-

Fig. 2 Second generation TrkC targeting ligands **1a** and **1b** featured in this work.

ning with a diamine aza-BODIPY precursor to **2**; a scheme and full details of that synthesis are given in the ESI.[†]

Distribution of probes **1a** and **b** in TrkC⁺ murine breast cancer cells

TrkC-expressing, 4T1 murine breast cancer cells were treated with probes **1a** and **b**. Confocal microscopy (Fig. 3a) showed the sulfonated system **1a** was efficiently internalized whereas the more lipophilic system **1b** had poor solubility and appeared to give significantly more cell-surface binding (Fig. 3b). For both probes, the material that was internalized overlaid with a lysosome tracker, indicating it accumulated in that organelle. These data support the assertion that the sulfonic acid groups of **1a** reduce non-specific binding to TrkC⁺ cells relative to the neutral probe **1b**, as expected because of repulsion between the negatively charged sulfonic acids and polar head groups at the cell surface.⁵² Probe **1a** showed no sign of toxicity to 4T1 cells at concentrations up to and including 25 μ M (Fig. S1[†]). All the subsequent work in this research was focused on **1a**, and studies of **1b** were discontinued, because these data indicate **1a** may have better characteristics for non-specific binding.

Histochemistry

Commercial samples of fixed human tissue from patients with stage 1, 2, and 3 malignant breast cancer were

Scheme 1 Preparation of TrkC-targeted near IR probe **1a**.

purchased (US Biomax). Probe **1a** was used to treat these samples and the staining was compared to that observed for normal breast tissue proximal to the tumor, obtained from the same patient in each case; illustrative data are shown in Fig. 4. In all cases the tumor tissue stained strongly whereas the healthy samples showed much less fluorescence. We were encouraged by the selectivity, but slightly surprised that the healthy tissue showed faint staining. It appears that this is genuine binding to the TrkC receptor in the healthy tissue (rather than non-specific binding or fluorescence leakage into the red channel) because fluorescently labeled TrkC mAb also stained weakly the normal tissue (Fig. S2†). This may mean the tumor environment is causing the surrounding normal tissue to upregulate TrkC expression.

Optical imaging with **1a** *in vivo*

A 4T1 breast cancer xenograft model was used (female BALB/c nu/nu mice, 8–10 mm diameter 4T1 tumors, *iv* administration) to evaluate **1a** as an *in vivo* optical imaging probe. Fig. 5 showed the NIR fluorescence images of nude mice bearing orthotopic 4T1 tumors after intravenous injection of 10 nmol of **1a**. The 4T1 tumors were clearly visualized with good tumor to background contrast at all time points (6, 24 and 48 h). As reported, Trk receptors are a family of tyrosine kinases that regulates synaptic strength and plasticity in the mammalian nervous system. TrkC is ordinarily activated by binding

with NT-3 and is expressed by proprioceptive sensory.⁵³ We observed high uptake of **1a** in central nervous system (CNS) which correlated with the expression of TrkC receptor.

To validate the TrkC targeting specificity of **1a**, the scrambled negative control, **1a-control**, was prepared (structure in Fig. 6 and Scheme S2†) and tested for *in vivo* imaging and *ex vivo* studies (Fig. 5 and 6). 4T1 tumor uptake of the negative control was much lower at all time points comparing to **1a**, which confirmed the specific TrkC targeting. As shown in Fig. 5b, the tumor/normal tissue fluorescent signal ratios in the imaging group were 3.06 ± 0.88 at 6 h, 3.01 ± 0.85 at 24 h, and 2.54 ± 0.55 at 48 h p.i., respectively, which is 2–3 times higher than that of the control group (1.44 ± 0.09 at 6 h, 1.26 ± 0.08 at 24 h, and 1.21 ± 0.19 at 48 h p.i.).

Ex vivo imaging was performed to acquire the fluorescent images of internal organs for both imaging and control groups as shown in Fig. 6. The tumor fluorescent intensities in the imaging group (7.29 ± 0.67) are significantly higher than in the control group (1.52 ± 0.29), also confirming the specificity of the probe **1a** for TrkC. The average radiant efficiency of other organs in control group was similar to that of the imaging group.

Conclusion

Compounds **1a** and **1b** are sufficiently different to the known probes **A** that it was not certain that these would also bind

Fig. 3 Imaging with 4T1 cells. a Probe **1a** was internalized; whereas, b much more of **1b** accumulated in the cell walls (overlay images of fluorescent probes and bright field imaging). c LysoTracker and probe **1a** (d) colocalized as shown in e.

TrkC. In the event, **1a** was selectively internalized into TrkC expressing cells, and stained human metastatic breast tumor tissues from cancer patients. *In vivo*, **1a** accumulated in a TrkC-expressing (4T1) breast cancer tumor model to a much higher degree than the isomeric control that does not bind TrkC (**1a-control**). Accumulation of both **1a** and **1a-control** in the liver was also observed, but this is to be expected for such a lipophilic compound (aza-BODIPY cores are very hydrophobic). The key probe **1a** did not accumulate in other organs to any extent that would preclude optical imaging of a primary breast cancer tumor.

Materials and methods

Cell culture and imaging

Murine breast cancer 4T1 cells were obtained from American Type Culture Collection and cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS) at 37 °C in humidified atmosphere with 5% CO₂. Sub-cellular localization was measured on living 4T1 cells using

Fig. 4 Histochemistry on human breast tissue array. Probe **1a** stained on breast cancer tissue (b, d, f) much brighter than matched adjacent breast tissue control (a, c, e), (example from 3 cases).

an Olympus FV1000 Confocal Microscope. Throughout, digital images were captured with a 100×/1.4 oil objective with the following filter sets: for LysoTracker Green: excitation 488 nm; for probes **1a** and **1b**: excitation 633 nm. Sequential optical sections (Z-stacks) from the basal-to-apical surfaces of the cell were also acquired.

Intracellular localization

4T1 cells were incubated with **1a**, 1 μM, for 12 h at 37 °C. After the cells were washed with PBS, LysoTracker Green (Life Technology, 500 nM) was added and the cells were incubated for 30 min at 37 °C. The cells were washed again with PBS before imaging.

Histochemistry

Two slides of human breast cancer tissue microarrays (BRC962) containing 36 cases of breast cancers and 12 cases

Fig. 5 a *In vivo* NIR fluorescence images (at 6 h, 24 h, and 48 h p.i.) of 4T1 tumor-bearing mice injected (iv) with **1a**. Arrows indicate the tumors. b Representative NIR fluorescent images of mice injected with **1a** (left) or **1a-control** compound at 48 h p.i. (right). The arrows indicate the tumors (top); an ROI study of tumor/normal tissue (muscle) ratio of **1a** and **1a-control** in the imaging groups ($n = 3$).

Fig. 6 a Structure of **1a-control**. b Representative *ex vivo* fluorescent images of organs (1 heart, 2 liver, 3 lungs, 4 spleen, 5 tumor, 6 kidneys, 7 muscle) collected from the mice injected with **1a** and **1a-control** at 48 h p.i.; and, quantification of fluorescence in those organs ($n = 3$).

of normal, reactive and benign tumour tissues of the breast in duplicates were purchased from US Biomax, Inc. **1a** and anti-TrkC antibody – N-terminal (ab188592, Abcam U.S.A.) were used as staining materials and the histology protocol was performed according to a published procedure.⁴² All tissues were visualized using an Olympus FV1000 confocal microscope.

Animal model

Murine breast cancer 4T1 cells were obtained from American Type Culture Collection and cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS) at 37 °C in humidified atmosphere with 5% CO₂. Animal procedures were performed according to a protocol approved by Houston Methodist Research Institute Animal Care and Use Committee. Murine 4T1 cells (5×10^6) were suspended in 100 μ L PBS and injected orthotopically into the lower left mammary fat pad of 6 weeks female BALB/c nu/nu mice (Charles River). Mice were subjected to the *in vivo* imaging studies when tumors reached 8–10 mm in diameter.

In vivo fluorescence imaging and *ex vivo* biodistribution study

In vivo and *ex vivo* fluorescence imaging was performed with IVIS200 imaging system, and quantified by Living Imaging software (Xenogen, CA). Excitation and emission filters were set at 640 nm and 680 nm respectively as suggested by the system for image acquisition. For *in vivo* studies ($n = 3$), stock solutions (~ 1 mM) of **1a** and **1a-control** conjugate in DMSO was diluted with PBS for injection. Each tumor-bearing mouse was injected intravenously with 10 nmol of **1a**. Images were acquired at 6 h, 24 h and 48 h post-injection (p.i.). For control studies ($n = 3$), mice were injected with 10 nmol of **1a-control** following the same procedures described above. Regions of interest (ROI) were drawn over the tumor and adjacent normal tissue to calculate the *in vivo* tumor/normal tissue ratio. In the ROI analysis, mean fluorescence intensities of the tumor area (tumor) defined as fluorescent radiant efficiency and of the area at the adjacent normal tissue (muscle) were calculated. After the 48 h imaging, mice were sacrificed, and tumors, muscles and major organs (*i.e.*, heart, lung, liver, spleen and kidney) were excised for *ex vivo* imaging acquisition. ROI analysis and quantification of the fluorescence signals ($n = 3$) were performed with Living Imaging (Xenogen, CA).

Notes

The authors declare no competing financial interests.

Conflicts of interest

The authors declare no competing interests.

Acknowledgements

We thank DoD BCRP Breakthrough Award (BC141561), CPRIT (RP150559 and RP170144), The Robert A. Welch Foundation (A-1121), and High Impact Research (HIR (UM.C/625/1/HIR/MOHE/MED/17 & UM.C/625/1/HIR/MOHE/MED/33) from the Ministry of Higher Education, Malaysia, for financial support. The NMR instrumentation at Texas A&M University was supported by a grant from the National Science Foundation (DBI-9970232) and the Texas A&M University System. The Olympus FV1000 confocal microscope acquisition was supported by the Office of the Vice President for Research at Texas A&M University. We thank the preclinical imaging core facility of Houston Methodist Research Institute for imaging support. Procedures for animal housing, maintenance, and euthanization were performed according to the American Veterinary Medical Association guidelines, and the IACUC approval from Methodist Hospital.

References

- 1 A. Patapoutian and L. F. Reichardt, *Curr. Opin. Neurobiol.*, 2001, **11**, 272–280.
- 2 E. J. Huang and L. F. Reichardt, *Annu. Rev. Biochem.*, 2003, **72**, 609–642.
- 3 M. V. Chao, *Nat. Rev. Neurosci.*, 2003, **4**, 299–309.
- 4 L. Tessarollo, P. Tsoulfas, D. Martin-Zanca, D. J. Gilbert, N. A. Jenkins, N. G. Copeland and L. F. Parada, *Development*, 1993, **118**, 463–475.
- 5 M. Ryden, R. Sehgal, C. Dominici, F. H. Schilling, C. F. Ibanez and P. Kogner, *Br. J. Cancer*, 1996, **74**, 773–779.
- 6 D. J. Yamashiro, A. Nakagawara, N. Ikegaki, X. G. Liu and G. M. Brodeur, *Oncogene*, 1996, **12**, 37–41.
- 7 G. M. Brodeur, A. Nakagawara, D. J. Yamashiro, N. Ikegaki, X. G. Liu, C. G. Azar, C. P. Lee and A. E. Evans, *J. Neuro-Oncology*, 1997, **31**, 49–55.
- 8 D. J. Yamashiro, X. G. Liu, C. P. Lee, A. Nakagawara, N. Ikegaki, L. M. McGregor, S. B. Baylin and G. M. Brodeur, *Eur. J. Cancer*, 1997, **33**, 2054–2057.
- 9 M. Encinas, M. Iglesias, N. Llecha and J. X. Comella, *J. Neurochem.*, 1999, **73**, 1409–1421.
- 10 M. A. Grotzer, A. J. Janss, K. M. Fung, J. A. Biegel, L. N. Sutton, L. B. Rorke, H. Zhao, A. Cnaan, P. C. Phillips, V. M. Y. Lee and J. Q. Trojanowski, *J. Clin. Oncol.*, 2000, **18**, 1027–1035.
- 11 M. E. Evangelopoulos, J. Weis and A. Kruttgen, *J. Neuro-Oncology*, 2004, **66**, 101–110.
- 12 P. Laneve, L. Di Macrotullio, U. Gioia, M. E. Fiori, E. Ferretti, A. Gulino, I. Bozzoni and E. Caffarelli, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 7957–7962.
- 13 G. M. Brodeur, J. E. Minturn, R. Ho, A. M. Simpson, R. Iyer, C. R. Varela, J. E. Light, V. Kolla and A. E. Evans, *Clin. Cancer Res.*, 2009, **15**, 3244–3250.
- 14 M. Bassili, E. Birman, N. F. Schor and H. U. Saragovi, *Cancer Chemother. Pharmacol.*, 2010, **65**, 1047–1056.
- 15 L. Harel, B. Costa and M. Fainzilber, *Dev. Neurobiol.*, 2010, **70**, 298–303.

- 16 V. Bernard-Gauthier, A. Aliaga, A. Aliaga, M. Boudjemeline, R. Hopewell, A. Kostikov, P. Rosa-Neto, A. Thiel and R. Schirmmacher, *ACS Chem. Neurosci.*, 2015, **6**, 260–276.
- 17 S. Kumar and J. de Vellis, *J. Neurosci. Res.*, 1996, **44**, 490–498.
- 18 Y. Wang, C. Hagel, W. Hamel, S. Muller, L. Kluwe and M. Westphal, *Acta Neuropathol.*, 1998, **96**, 357–364.
- 19 C. Calatuzzolo, A. Salmaggi, B. Pollo, F. L. Sciacca, M. Lorenzetti, A. Franzini, A. Boiardi, G. Broggi and C. Marras, *Neurol. Sci.*, 2007, **28**, 304–310.
- 20 M. Assimakopoulou, M. Kondyli, G. Gatzounis, T. Maraziotis and J. Varakis, *BMC Cancer*, 2007, **7**, 202–210.
- 21 S. Lawn, A. Chan Jennifer, N. Krishna, A. Pisklakova, X. Qu, A. Fenstermacher David, M. Fournier, D. Vrionis Frank, N. Tran, S. Kenchappa Rajappa and A. Forsyth Peter, *J. Biol. Chem.*, 2014, **290**, 3814–3824.
- 22 P. Stephens, S. Edkins, H. Davies, C. Greenman, C. Cox, C. Hunter, G. Bignell, J. Teague, R. Smith, C. Stevens, S. O'Meara, A. Parker, P. Tarpey, T. Avis, A. Barthorpe, L. Brackenbury, G. Buck, A. Butler, J. Clements, J. Cole, E. Dicks, K. Edwards, S. Forbes, M. Gorton, K. Gray, K. Halliday, R. Harrison, K. Hills, J. Hinton, D. Jones, V. Kosmidou, R. Laman, R. Lugg, A. Menzies, J. Perry, R. Petty, K. Raine, R. Shepherd, A. Small, H. Solomon, Y. Stephens, C. Tofts, J. Varian, A. Webb, S. West, S. Widaa, A. Yates, F. Brasseur, C. S. Cooper, A. M. Flanagan, A. Green, M. Knowles, S. Y. Leung, L. H. J. Looijenga, B. Malkowicz, M. A. Pierotti, B. Teh, S. T. Yuen, A. G. Nicholson, S. Lakhani, D. F. Easton, B. L. Weber, M. R. Stratton, P. A. Futreal and R. Wooster, *Nat. Genet.*, 2005, **37**, 590–592.
- 23 S. V. Ivanov, A. Panaccione, B. Brown, Y. Guo, C. A. Moskaluk, M. J. Wick, J. L. Brown, A. V. Ivanova, N. Issaeva, A. K. El-Naggar and W. G. Yarbrough, *Oncogene*, 2013, **32**, 3698–3710.
- 24 W. Jin, G.-M. Kim, M.-S. Kim, M.-H. Lim, C.-H. Yun, J. Jeong, J.-S. Nam and S.-J. Kim, *Carcinogenesis*, 2010, **31**, 1939–1947.
- 25 W. Jin, C. Yun, M. K. Kwak, T. A. Kim and S. J. Kim, *Oncogene*, 2007, **26**, 7684–7691.
- 26 W. Jin, C. Yun, J. Jeong, Y. Park, H.-D. Lee and S.-J. Kim, *J. Biol. Chem.*, 2008, **283**, 1391–1400.
- 27 M. J. Blasco-Gutierrez, I. J. San Jose-Crespo, E. Zozaya-Alvarez, R. Ramos-Sanchez and N. Garcia-Atares, *Cancer Invest.*, 2007, **25**, 405–410.
- 28 H. Hondermarck, *Cytokine Growth Factor Rev.*, 2012, **23**, 357–365.
- 29 X. Xu, S. R. Tahan, T. L. Pasha and P. J. Zhang, *J. Cutaneous Pathol.*, 2003, **30**, 318–322.
- 30 A. Marconi, M. Terracina, C. Fila, J. Franchi, F. Bonte, G. Romagnoli, R. Aurelli, C. M. Failla, M. Dumas and C. Pincelli, *J. Invest. Dermatol.*, 2003, **121**, 1515–1521.
- 31 D. Marchetti, B. Murry, J. Galjour and A. Wilke-Greiter, *J. Cell. Biochem.*, 2003, **88**, 865–872.
- 32 M. J. C. Hendrix, E. A. Seftor, D. A. Kirschmann, V. Quaranta and R. E. B. Seftor, *Ann. N. Y. Acad. Sci.*, 2003, **995**, 151–161.
- 33 A. Marconi, M. C. Panza, M. Bonnet-Duquennoy, K. Lazou, R. Kurfurst, F. Truzzi, R. Lotti, G. De Santis, M. Dumas, F. Bonte and C. Pincelli, *Int. J. Cosmet. Sci.*, 2006, **28**, 255–261.
- 34 F. Truzzi, A. Marconi, R. Lotti, K. Dallaglio, L. E. French, B. L. Hempstead and C. Pincelli, *J. Invest. Dermatol.*, 2008, **128**, 2031–2040.
- 35 V. A. Botchkarev, M. Yaar, E. M. J. Peters, S. P. Raychaudhuri, N. V. Botchkareva, A. Marconi, S. K. Raychaudhuri, R. Paus and C. Pincelli, *J. Invest. Dermatol.*, 2006, **126**, 1719–1727.
- 36 D. Marchetti, Y. Denkins, J. Reiland, A. Greiter-Wilke, J. Galjour, B. Murry, J. Blust and M. Roy, *Pathol. Oncol. Res.*, 2003, **9**, 147–158.
- 37 Y. Denkins, J. Reiland, M. Roy, N. D. Sinnappah-Kang, J. Galjour, B. P. Murry, J. Blust, R. Aucoin and D. Marchetti, *Neuro-Oncology*, 2004, **6**, 154–165.
- 38 A. Vaishnavi, T. Le Anh and C. Doebele Robert, *Cancer Discovery*, 2015, **5**, 25–34.
- 39 F. Truzzi, A. Marconi and C. Pincelli, *Derm.-Endocrinol.*, 2011, **3**, 32–36.
- 40 L. Dolle, E. Adriaenssens, I. E. Yazidi-Belkoura, X. L. Bourhis, V. Nurcombe and H. Hondermarck, *Curr. Cancer Drug Targets*, 2004, **4**, 463–470.
- 41 V. Bernard-Gauthier, M. Boudjemeline, P. Rosa-Neto, A. Thiel and R. Schirmmacher, *Bioorg. Med. Chem.*, 2013, **21**, 7816–7829.
- 42 S. C. Kue, A. Kamkaew, H. B. Lee, L. L. Chung, L. V. Kiew and K. Burgess, *Mol. Pharmaceutics*, 2015, **12**, 212–222.
- 43 A. Kamkaew, H. Lim Siang, B. Lee Hong, V. Kiew Lik, Y. Chung Lip and K. Burgess, *Chem. Soc. Rev.*, 2012, **42**, 77–88.
- 44 S. G. Awuah and Y. You, *RSC Adv.*, 2012, **2**, 11169–11183.
- 45 R. Ziesel, C. Goze and G. Ulrich, *Synthesis*, 2007, **6**, 936–949.
- 46 A. Loudet and K. Burgess, BODIPY Dyes and Their Derivatives: Syntheses and Spectroscopic Properties, *Handbook of Porphyrin Science: With Applications to Chemistry, Physics, Materials Science, Engineering, Biology and Medicine*, ed. K. Kadish, K. Smith and R. Guilard, World Scientific, 2010, p. 203.
- 47 H. Lu, J. Mack, Y. Yang and Z. Shen, *Chem. Soc. Rev.*, 2014, **43**, 4778–4823.
- 48 Y. Angell, D. Chen, F. Brahimi, H. U. Saragovi and K. Burgess, *J. Am. Chem. Soc.*, 2008, **130**, 556–565.
- 49 A. Kamkaew and K. Burgess, *Chem. Commun.*, 2015, **51**, 10664–10667.
- 50 M. Meldal and C. W. Tornøe, *Chem. Rev.*, 2008, **108**, 2952–3015.
- 51 P. Wu and V. V. Fokin, *Aldrichimica Acta*, 2007, **40**, 7–17.
- 52 G. A. Johnson, N. Muthukrishnan and J.-P. Pellois, *Bioconjugate Chem.*, 2013, **24**, 114–123.
- 53 R. A. Segal, *Annu. Rev. Neurosci.*, 2003, **26**, 299–330.