RESEARCH ARTICLE

[View Article Online](#)
[View Journal](#) | [View Issue](#) Cite this: *Med. Chem. Commun.*,
2017, **8**, 2258Synthesis and antifungal activity of novel
oxazolidin-2-one-linked 1,2,3-triazole derivatives†Alejandra Ramírez-Villalva,^a Davir González-Calderón, ^a Roxana I. Rojas-García,^a
Carlos González-Romero, ^a Joaquín Tamariz-Mascarúa,^c
Macario Morales-Rodríguez,^b Nieves Zavala-Segovia^d and Aydeé Fuentes-Benites ^{*a}

Novel oxazolidin-2-one-linked 1,2,3-triazole derivatives (**4a–k**) were synthesized by straightforward and versatile azide–enolate (3 + 2) cycloaddition. The series of compounds was screened for antifungal activity against four filamentous fungi as well as six yeast species of *Candida* spp. According to their efficiency and breadth of scope, they can be ordered as **4k** > **4d** > **4h** > **4a**, especially in relation to the activity displayed against *Candida glabrata* ATCC-34138, *Trichosporon cutaneum* ATCC-28592 and *Mucor hiemalis* ATCC-8690, i.e. compounds **4d**, **4h** and **4k** showed excellent activity against *C. glabrata* (MIC 0.12, 0.25 and 0.12 $\mu\text{g mL}^{-1}$, respectively), better than that of itraconazole (MIC 1 $\mu\text{g mL}^{-1}$). The activity of compound **4d** (MIC = 2 $\mu\text{g mL}^{-1}$) was higher than that observed for the standard antifungal drug (MIC = 8 $\mu\text{g mL}^{-1}$) against *Trichosporon cutaneum*, while compound **4k** displayed an excellent antimycotic activity against *Mucor hiemalis* (MIC = 2 $\mu\text{g mL}^{-1}$ vs. 4 $\mu\text{g mL}^{-1}$ for itraconazole). In addition, we describe herein a novel mild and eco-friendly synthetic protocol for obtaining β -ketosulfones (adducts to afford compounds **4a–k**) from α -brominated carbonyls in an aqueous nanomicellar medium at room temperature.

Received 28th August 2017,
Accepted 14th October 2017

DOI: 10.1039/c7md00442g

rsc.li/medchemcomm

1. Introduction

Both oxazolidin-2-one¹ and 1,2,3-triazole² cores are very well-recognized pharmacophores in the literature. They have become extremely versatile in medicinal chemistry, featuring a number of clinically used drugs (e.g. Linezolid **1** or Tazobactam **2**, Fig. 1). Consequently, several antimicrobials based on one or the other scaffold have been reported in the literature.³ On the other hand, the study of *both cores in the same molecule* (**3**) as an antibacterial agent has been reported⁴ and patented (e.g. **3a** (ref. 5) and **3b** (ref. 6)). This year – 2017 – Melinta Therapeutics Inc. announced the acceptance of FDA of the Investigational New Drug (IND) application for topical

Radezolid **3c**, a second-generation oxazolidinone/triazole compound discovered by Melinta scientists as a novel antibiotic to treat serious bacterial infections.⁷ Curiously, antifungal activity has not been demonstrated for these kinds of compounds to the best of our knowledge.

Even though the mechanism of action of oxazolidinone⁸ and triazole⁹ cores has been described, the inhibition of monoamine oxidase A (MAO-A)^{4a} as well as the RNA-binding process^{4g} has been proposed as the mode of action for oxazolidin-2-one-linked 1,2,3-triazole.

Due to the strong resistance of fungi to current drugs, there is a continuous search for antifungal agents. As part of our ongoing research, we herein describe the synthesis and biological evaluation of novel oxazolidin-2-one-linked 1,2,3-triazole derivatives (**4**).

2. Chemistry

Our initial study began by obtaining the starting materials, azide/oxazolidin-2-one **8** and ketones **11** (Scheme 1). Firstly, chlorine derivative **7** was synthesized (61%) by coupling between 2-chloroethyl isocyanate **6** and benzoin **5**.¹⁰ Subsequently, the nucleophilic substitution of **7** by the azide ion furnished compound **8** (58%).

Ketones **11a–11g** were synthesized since, unlike **11i–11j**, they are not commercially available (ketone **11h** was kindly donated by Syntex-La Roche). For these purposes, we have developed a novel organic solvent-free synthesis of

^a Departamento de Química Orgánica, Facultad de Química, Universidad Autónoma del Estado de México, Paseo Colón/Paseo Tollocan s/n, Toluca, Estado de México, 50120, Mexico. E-mail: mpagfuentesb@uaemex.mx; Fax: +52 722 217 3890; Tel: +52 722 217 5109x113

^b Departamento de Microbiología, Facultad de Química, Universidad Autónoma del Estado de México, Paseo Colón/Paseo Tollocan s/n, Toluca, Estado de México, 50120, Mexico

^c Departamento de Química Orgánica, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Prol. Carpio y Plan de Ayala, Ciudad de México, 11340, Mexico

^d Centro Conjunto de Investigación en Química Sustentable UAEM-UNAM, Carretera Toluca-Aticomulco Km 14.5, Toluca, 52000, Mexico

† Electronic supplementary information (ESI) available: Supplementary data (experimental procedures, characterization data of all compounds and copies of ¹³C, ¹H, and spectra, as well as the study on the synthesis of β -ketosulfones). See DOI: 10.1039/c7md00442g

U.S. FDA approved antimicrobial pharmaceuticals:

Oxazolidin-2-ones linked-1,2,3-triazoles:

Fig. 1 The novel series of compounds **4** involves the extremely important oxazolidin-2-one and 1,2,3-triazole pharmacophoric cores. A similar feature is found in antimicrobial oxazolidin-2-one-linked 1,2,3-triazole **3**, including lead compounds **1** and **2**.

β -ketosulfones in an aqueous nanomicellar medium by the concomitant use of the surfactant reagent SPGS-550-M or 'Nok' (now available at Sigma-Aldrich¹¹) and sodium *p*-toluenesulfonate from α -bromoketones. The surfactant nature of NOK (a third generation surfactant) allows it to act as an efficient nanoreactor in catalytic amounts. The study and scope of this novel methodology are described in the ESI† in greater detail. Therefore, we decided to apply such a methodology in the synthesis of compound **11** from **10**, which in turn was accomplished by reported protocols¹² with *N*-bromosuccinimide (NBS) as the bromine source.

Although Cu-catalyzed azide-alkyne cycloaddition (CuAAC) is the conventional method for obtaining 1,2,3-triazole moieties,¹³ other strategies have emerged as alternatives for such a purpose. We previously reported¹⁴ a novel synthetic protocol to achieve the efficient assembly of 1,4,5-trisubstituted 1,2,3-triazole cores through azide-enolate cycloaddition. In consequence, oxazolidin-2-one-linked 1,2,3-triazole derivatives (**4**) were synthesized by the efficient 1,3-dipolar cycloaddition of azide/oxazolidin-2-one in the presence of enolates prepared *in situ* from ketones **11** activated by DBU as the base. Table 1 summarizes these outcomes.

3. Microbiology

Compounds **4a–k** were tested for their *in vitro* activity against four filamentous fungi: *Aspergillus fumigatus* ATCC-16907, *Trichosporon cutaneum* ATCC-28592, *Rhizopus oryzae* ATCC-10329 and *Mucor hiemalis* ATCC-8690. These compounds were also evaluated, employing standardized microbiological methods developed by the CLSI, against six yeast specimens: *Candida albicans* ATCC-10231, *Candida utilis* ATCC-9226, *Candida tropicalis* ATCC-13803, *Candida parapsilopsis* ATCC-22019,

Scheme 1 Reagents and conditions: (i) N₂, 180 °C, 24 h. (ii) NaN₃ (1.1 eq.), DMF anhyd., 60 °C, 12 h, N₂. (iii) NBS (1.1 eq.), TsOH·H₂O (1.0 eq.), MeCN, 60 °C, 4 h. (iv) SPGS-550-M aq. (2% w/w), *p*-Tol-SO₂Na (1.5 eq.), r.t., 12 h.

Table 1 Synthesis of oxazolidin-2-one-linked 1,2,3-triazoles **4a–k** from azide **8** by coupling with active ketones **11**

Entry ^a	Ketone	Triazole ^b (yield%) ^c
1	11a : R ² = Ph, R ¹ = SO ₂ - <i>p</i> -Tol	4a (68)
2	11b : R ² = <i>p</i> -CH ₃ Ph, R ¹ = SO ₂ - <i>p</i> -Tol	4b (61)
3	11c : R ² = <i>p</i> -NO ₂ -Ph, R ¹ = SO ₂ - <i>p</i> -Tol	4c (74)
4	11d : R ² = <i>m</i> -NO ₂ Ph, R ¹ = SO ₂ - <i>p</i> -Tol	4d (70)
5	11e : R ² = <i>p</i> -Cl-Ph, R ¹ = SO ₂ - <i>p</i> -Tol	4e (72)
6	11f : R ² = thiophen-2-yl, R ¹ = SO ₂ - <i>p</i> -Tol	4f (68)
7	11g : R ² = 5-chlorothiophen-2-yl, R ¹ = SO ₂ - <i>p</i> -Tol	4g (64)
8	11h : R ² = pentyl, R ¹ = SO ₂ Ph	4h (71)
9	11i : R ² = CH ₃ , R ¹ = COCH ₃	4i (73)
10	11j : R ² = Ph, R ¹ = COPh	4j (71)
11	11k : R ² = Ph, R ¹ = CN	4k (67)

^a Reaction conditions: To a solution of compound **8** (1.0 eq.) and **11** (1.0 eq.) in DMF anhyd., DBU (2.0 eq.) was added. The reaction mixture was stirred at 50–60 °C for 12–24 h. ^b Confirmed by ¹H-NMR, ¹³C-NMR and MS. ^c Yields refer to chromatographically pure isolated compounds.

Candida glabrata ATCC-34138, and *Candida krusei* ATCC-14243. Then, the sensitivity of the filamentous microorganisms was determined by the microdilution M38-A method,¹⁵ and that of the yeast fungi with the M27-A3 method.¹⁶

Such antifungal activity was compared to that of itraconazole as the standard antifungal drug. The minimum inhibitory concentration (MIC) values of the standard and compounds **4a–k**, expressed in micrograms per milliliter, were determined in 96-well plates using MOPS (3-[*N*-morpholino]propanesulfonic acid buffered RPMI-1640 medium, Sigma-Aldrich).

4. Results and discussion

The antifungal activity of the test compounds is summarized in Table 2. Compounds **4d**, **4h** and **4k** showed excellent activity against *C. glabrata* (MIC 0.12, 0.25 and 0.12 μg mL⁻¹, respectively), better than that of

itraconazole (MIC 1 μg mL⁻¹). The activity of compound **4d** (MIC = 2 μg mL⁻¹) was higher than that observed for the standard antifungal drug (MIC = 8 μg mL⁻¹) against *Trichosporon cutaneum*, while compound **4k** displayed an excellent antimycotic activity against *Mucor hiemalis* (MIC = 2 μg mL⁻¹ vs. 4 μg mL⁻¹ for itraconazole). Compounds **4a**, **4c** and **4e** proved to be moderate antifungal agents against *Aspergillus fumigatus* strains. The current results suggest that the presence of either the *a*-CN or *p*-NO₂Ph group at position 5 of the triazole cores increases the biological activity of these compounds in both yeast and filamentous fungi.

These outcomes can also be described by the ‘sensitivity’ parameters of yeasts, according to the breakpoints described in the M27-A3 document (Table 3). In general, *C. glabrata*, *C. krusei* and *C. parapsilosis* showed some susceptibility to the test compounds, whereas *C. albicans*, *C. tropicalis* and *C. utilis* were resistant to all of them.

Table 2 *In vitro* antifungal activities of the synthesized compounds (MIC, μg mL⁻¹)

Compound	<i>C. alb</i>	<i>C. trop</i>	<i>C. uti</i>	<i>C. kru</i>	<i>C. gla</i>	<i>C. par</i>	<i>M. hie</i>	<i>A. fum</i>	<i>T. cut</i>	<i>R. ory</i>
4a	8	8	8	0.5	1	8	4	2	8	16
4b	1	8	4	8	2	0.25	16	8	8	16
4c	4	8	4	0.25	4	8	16	2	8	16
4d	8	8	8	8	0.12	0.5	16	16	2	16
4e	8	8	8	8	4	8	16	4	8	16
4f	8	8	8	8	1	8	16	8	8	16
4g	8	8	8	2	2	4	16	16	8	16
4h	8	8	8	8	0.25	8	16	16	8	16
4i	8	8	8	8	2	4	16	16	8	8
4j	8	8	8	8	2	8	16	16	8	16
4k	8	8	8	0.5	0.12	8	2	16	8	16
Standard ^a	0.03	0.06	0.25	0.25	1	0.06	4	1	8	1

Abbreviations: *C. alb.*, *Candida albicans*; *C. trop.*, *Candida tropicalis*; *C. uti.*, *Candida utilis*; *C. kru.*, *Candida krusei*; *C. gla.*, *Candida glabrata*, *C. par.*, *Candida parapsilosis*; *M. hie.*, *Mucor hiemalis*; *A. fum.*, *Aspergillus fumigatus*; *T. cut.*, *Trichosporon cutaneum*; *R. ory.*, *Rhizopus oryzae*.^a Itraconazole.

Table 3 Determination of the sensitivity of yeast (according to document M27-A3): Susceptible (S), dose-dependent sensitive (SDD) and resistant (R)

Compound	<i>C. alb</i>	<i>C. trop</i>	<i>C. uti</i>	<i>C. kru</i>	<i>C. gla</i>	<i>C. par</i>
4a	R	R	R	SDD	R	R
4b	R	R	R	R	R	SDD
4c	R	R	R	SDD	R	R
4d	R	R	R	R	S	SDD
4e	R	R	R	R	R	R
4f	R	R	R	R	R	R
4g	R	R	R	R	R	R
4h	R	R	R	R	SDD	R
4i	R	R	R	R	R	R
4j	R	R	R	R	R	R
4k	R	R	R	SDD	S	R
Standard ^a	S	S	SDD	SDD	R	S

Abbreviations: *C. alb.*, *Candida albicans*; *C. trop.*, *Candida tropicalis*; *C. uti.*, *Candida utilis*; *C. kru.*, *Candida krusei*; *C. gla.*, *Candida glabrata*; *C. par.*, *Candida parapsilosis*. ^a Itraconazole. Interpretive criteria: breakpoints (MIC, $\mu\text{g mL}^{-1}$) = 0.12 [S], 0.25–0.5 [SDD], 1 [R].

5. Conclusion

In summary, eleven oxazolidin-2-one-linked 1,2,3-triazole derivatives (4a–k) were synthesized in good yields based on azide–enolate 1,3-dipolar cycloaddition. *In vitro* assays demonstrated that compound 4k is the most efficient antimicrobial agent, since it was either better than or comparable to itraconazole against three species (*C. glabrata*, *M. hiemalis* and *T. cutaneum*). The second best antimicrobial activity was exhibited by compound 4d, which was much better than the reference drug against two species (*C. glabrata* and *T. cutaneum*). In consequence, these compounds can be considered as drug candidates for future complementary biological studies. In addition, we have developed a novel organic solvent-free synthesis of β -ketosulfones in an aqueous nanomicellar medium. The surfactant nature of SPGS-550-M or 'NOK' (a third generation surfactant) allows it to act as an efficient nanoreactor in catalytic amounts. The notable advantages of this methodology over those previously reported include its simplicity of handling, mild conditions, high yields, cheap reagents and great tolerance of functional groups.

Conflicts of interest

The authors declare no competing interest.

Acknowledgements

We gratefully acknowledge financial support from the Secretaría de Investigación y Estudios Avanzados/UAEMéx (project no. 3804/2014/CID) and CONACYT-Mexico (postgraduate scholarship no. 227581 and no. 273644). The authors would also like to thank the referees for their valuable comments and suggestions, Signa S.A. de C.V. for kindly donating some solvents and reagents, and L. Triana-Cruz (CCIQS UAEMéx-UNAM) for technical support.

References

- (a) N. Pandit, R. K. Singla and B. Shrivastava, *Int. J. Med. Chem.*, 2012, **2012**, 24; (b) K. J. Shaw and M. R. Barbachyn, *Ann. N. Y. Acad. Sci.*, 2011, **1241**, 48–70.
- (a) A. Massarotti, S. Aprile, V. Mercalli, E. Del Grosso, G. Grosa, G. Sorba and G. C. Tron, *ChemMedChem*, 2014, **9**, 2497–2508; (b) S. G. Agalave, S. R. Maujan and V. S. Pore, *Chem. – Asian J.*, 2011, **6**, 2696–2718; (c) R. Kharb, P. C. Sharma and M. S. Yar, *J. Enzyme Inhib. Med. Chem.*, 2011, **26**, 1–21.
- Oxazolidinone scaffolds: (a) O. A. Phillips and L. H. Sharaf, *Expert Opin. Ther. Pat.*, 2016, **26**, 591–605; (b) K. Michalska, I. Karpiuk, M. Król and S. Tyski, *Bioorg. Med. Chem.*, 2013, **21**, 577–591; (c) K. Michalska, I. Karpiuk, M. Król and S. Tyski, *Bioorg. Med. Chem.*, 2013, **21**, 577–591; (d) 1,2,3-Triazole scaffolds: F. de Carvalho da Silva, M. F. do Carmo Cardoso, P. Garcia-Ferreira and V. F. Ferreira, *Biological properties of 1H-1,2,3- and 2H-1,2,3-triazoles*, in *Chemistry of 1,2,3-triazoles*, ed. W. Dehaen and V. A. Bakulev, Springer International Publishing, Switzerland, 2015, pp. 117–165, DOI: 10.1007/7081_2014_124.
- (a) M. V. Nora de Souza, *Expert Opin. Ther. Pat.*, 2008, **18**, 1101–1105; (b) F. Reck, F. Zhou, M. Girardot, G. Kern, C. J. Eyermann, N. J. Hales, R. R. Ramsay and M. B. Gravestock, *J. Med. Chem.*, 2005, **48**, 499–506; (c) J. A. Demaray, J. E. Thuener, M. N. Dawson and S. J. Sucheck, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 4868–4871; (d) O. A. Phillips, E. E. Udo and S. M. Samuel, *Eur. J. Med. Chem.*, 2008, **43**, 1095–1104; (e) O. A. Phillips, E. E. Udo, M. E. Abdel-Hamid and R. Varghese, *Eur. J. Med. Chem.*, 2009, **44**, 3217–3227; (f) H. Fan, Y. Chen, Z. Jiang, S. Zhang, D. Zhong, R. Ji and Y. Yang, *Eur. J. Med. Chem.*, 2008, **43**, 1706–1714; (g) G. Acquah-Harrison, S. Zhou, J. V. Hines and S. C. Bergmeier, *J. Comb. Chem.*, 2010, **12**, 491–496.
- M. B. Gravestock, M. J. Betts, D. A. Griffin and I. R. Matthews, Oxazolidinone derivatives with antibacterial activity, PCT Int. Appl. WO0181350, Astrazeneca U.K. Limited, Astrazeneca AB, Sweden, 2001.
- A. L. Choy and V. P. V. N. Josyula, Benzisoxazole oxazolidinones as antibacterial agents, PCT Int. Appl. WO2007088438A2, Pfizer Products Inc., US, 2007.
- <http://melinta.com/>.
- (a) B. Bozdogan and P. C. Appelbaum, *Int. J. Antimicrob. Agents*, 2004, **23**, 113–119; (b) A. Chan, J. Cross, Y. He, B. Lippa and D. Ryan, *Antibacterial Drugs*, in *Drug Discovery: Practices, Processes, and Perspectives*, ed. J. J. Li and E. J. Corey, John Wiley & Sons, Inc., Hoboken, New Jersey, 2013, ch. 10, pp. 412–413, DOI: 10.1002/9781118354483.ch10.
- (a) F. Schiaffella, A. Macchiarulo, L. Milanese, A. Vecchiarelli, G. Costantino and D. Pietrella, *J. Med. Chem.*, 2005, **48**, 7658–7666; (b) J. Y. Choi, L. M. Podust and W. R. Roush, *Chem. Rev.*, 2014, **114**, 11242–11271.
- B. M. Santoyo, C. González-Romero, O. Merino, R. Martínez-Palou, A. Fuentes-Benites, H. A. Jiménez-Vázquez, F. Delgado and J. Tamariz, *Eur. J. Org. Chem.*, 2009, **2009**, 2505–2518.

- 11 Sigma-Aldrich catalog number: 776033.
- 12 J. C. Lee, Y. H. Bae and S. K. Chang, *Bull. Korean Chem. Soc.*, 2003, **24**, 407–408.
- 13 (a) N. Jung and S. Bräse, Click Reactions: Azide-Alkyne Cycloaddition, *Kirk-Othmer Encycl. Chem. Technol.*, 2013, 1–43, DOI: 10.1002/0471238961.clicjung.a01; (b) *Click Reactions in Organic Synthesis*, ed. S. Chandrasekaran, Wiley-VCH, Weinheim, Germany, 2016, DOI: 10.1002/9783527694174.
- 14 D. González-Calderón, I. Santillán-Iniesta, C. A. González-González, A. Fuentes-Benites and C. González-Romero, *Tetrahedron Lett.*, 2015, **56**, 514–516.
- 15 (a) Clinical and Laboratory Standards Institute (CLSI), *Document M38-A2: Reference Method for Broth Dilution Antifungal Susceptibility Testing of Filamentous Fungi*, Approved Standard, Clinical and Laboratory Standards Institute, Wayne, PA, 2nd edn, 2002; (b) A. L. Barry, An overview of the Clinical and Laboratory Standards Institute (CLSI) and its impact on antimicrobial susceptibility tests, in *Antimicrobial Susceptibility Testing Protocols*, ed. R. Schwalbe, L. Steele-Moore and A. C. Goodwin, CRC Press Taylor & Francis Group, Florida, 2007, pp. 1–6, ISBN: 9780824741006; (c) A. Espinel-Ingroff and E. Canton, Antifungal susceptibility testing of filamentous fungi, in *Antimicrobial Susceptibility Testing Protocols*, ed. R. Schwalbe, L. Steele-Moore and A. C. Goodwin, CRC Press Taylor & Francis Group, Florida, 2007, pp. 209–241, ISBN: 9780824741006.
- 16 (a) Clinical and Laboratory Standards Institute (CLSI), *M27-A3: Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts*, Approved Standard, Clinical and Laboratory Standards Institute, Wayne, PA, 3rd edn, 2008, ISBN: 1-56238-666-2; (b) A. Espinel-Ingroff and E. Canton, Antifungal susceptibility testing of yeasts, in *Antimicrobial Susceptibility Testing Protocols*, ed. R. Schwalbe, L. Steele-Moore and A. C. Goodwin, CRC Press Taylor & Francis Group, Florida, 2007, pp. 173–208, ISBN: 9780824741006; (c) A. W. Fothergill, *Antifungal Susceptibility Testing: Clinical Laboratory and Standards Institute (CLSI) methods*, in *Interactions of Yeasts, Moulds, and Antifungal Agents, How to Detect Resistance*, ed. G. S. Hall, Springer Science-Business Media, 2012, pp. 65–74, DOI: 10.1007/978-1-59745-134-5_2; (d) M. A. Pfaller and D. J. Diekema, *J. Clin. Microbiol.*, 2012, **50**, 2846–2856; (e) M. C. Arendrup, G. Garcia-Effron, C. Lass-Flörl, A. G. López, J. L. Rodríguez-Tudela, M. Cuenca-Estrella and D. S. Perlin, *Antimicrob. Agents Chemother.*, 2010, **54**, 426–439.