what's your diagnosis?

Diagnosis: Benign bowel distribution of lanthanum carbonate

Hsin-Chi Wu, Shih-Hua Lin

From the Division of Nephrology, Department of Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan, ROC

Correspondence and reprints: Shih-Hua Lin, MD · Division of Nephrology, Department of Medicine, Tri-Service General Hospital · Number 325, Section 2, Cheng-Kung Road, Neihu 114, Taipei, Taiwan ROC · T: +886-2-87927213 F: +886-2-87927134 · shihhualin@yahoo.com · Accepted for publication August 2008

Ann Saudi Med 2008; 28(6): __

Hyperphosphatemia in uremic patients is associated with metastatic calcification such as tumoral calcinosis and vascular calcification, and an increased cardiovascular mortality and morbidity.¹ To avoid hyperphosphatemia-related complications, the use of oral phosphate-binding agents is universal to reduce phosphate absorption in the gastrointestinal tract. "Old generation" phosphate binders with aluminum or calcium-based agents tend to cause

neurological toxicity and exacerbation of calcium overloading and soft tissue or cardiovascular calcification, respectively.² The "new generation" of aluminum or calcium-free phosphate binders has been shown to effectively control serum phosphate concentration with the attenuation of cardiovascular and soft tissue calcification.

Lanthanum carbonate ($\text{La}_2(\text{CO}_3)_3$), is a novel calcium- and aluminum-free phosphate binder for the

Figure 1. Radiopaque lanthanum carbonate tablets milled in the gastrointestinal distribution. Starry opacities also seen.

treatment of hyperphosphatemia in ESRD patients. The element, lanthanum (La, atomic number 57, atomic weight 138.9) is a rare-earth natural metal belonging to group 3B of the periodic table and causes radiopacities in the roentgenogram, as does barium (atomic number 56) (Figure 1). The half-life of lanthanum elimination from plasma is 53 hours though the plasma level is extremely low and elimination half-life from bone is 2.0 to 3.6 years.³ The affinity of lanthanum compound for phosphate is >97% and is more potent than calcium or magnesium, and similar to aluminum.⁴ Lanthanum binds phosphate optimally at pH 3~5 and retains binding activity at pH 1~7.⁵ It is virtually non-absorbed (0.00005%) and chelates dietary phosphate

into insoluble lanthanum phosphate by ionized form. Lanthanum phosphate and carbonate in the gut may form numerous radiopaque spots on the abdominal x-ray, which is not reported with other phosphate binders.^{6,7}

These unexpected starry opacities disappear completely after the withdrawal of lanthanum carbonate. The patient continued lanthanum carbonate treatment because of no abdominal symptoms and improved pain. Due to this radiopaque feature, one can evaluate the compliance to this drug. Given the increasing usage of lanthanum carbonate in patients with ESRD, every physician must understand this typical benign radiopaque finding to avoid unnecessary examination and inappropriate management.

REFERENCES

1. Coladonato JA. Control of hyperphosphatemia among patients with ESRD. *J Am Soc Nephrol* 2005;16 Suppl 2:S107-114.
2. Amann K, Gross ML, London GM, Ritz E. Hyperphosphataemia--a silent killer of patients with renal failure? *Nephrol Dial Transplant* 1999;14:2085-2087.
3. Behets GJ, Verberckmoes SC, D'Haese PC, De Broe ME. Lanthanum carbonate: a new phosphate binder. *Curr Opin Nephrol Hypertens* 2004;13:403-409.
4. Fricker SP. The therapeutic application of lanthanides. *Chem Soc Rev* 2006;35:524-33.
5. Albaaj F, Hutchison AJ. Lanthanum carbonate (Fosrenol): a novel agent for the treatment of hyperphosphataemia in renal failure and dialysis patients. *Int J Clin Pract* 2005;59:1091-1096.
6. Savica V, Calò LA, Monardo P, Santoro D, Bellinghieri G. Phosphate binders and management of hyperphosphataemia in end-stage renal disease. *Nephrol Dial Transplant* 2006;21:1065-1068.
7. de Freitas D, Donne RL, Hutchison AJ. Lanthanum carbonate--a first line phosphate binder? *Semin Dial* 2007;20:325-328.