ERRATA

The November 1, 2008, commentary by Kvols, entitled, “Revisiting C.G. Moertel’s Land of Small Tumors” (J Clin Oncol 26:5005-5007, 2008) contained an error.

In the Interferon section, the second sentence of the first paragraph should have indicated that more than 500 patient cases have been reported worldwide, as follows:

“Since then, more than 500 patient cases have been reported worldwide and have been published in the literature.”

The online version has been corrected in departure from the print.

DOI: 10.1200/JCO.2009.22.0533

The January 1, 2009, article by Slotman et al entitled, “Prophylactic Cranial Irradiation in Extensive Disease Small-Cell Lung Cancer: Short-Term Health-Related Quality of Life and Patient Reported Symptoms—Results of an International Phase III Randomized Controlled Trial by the EORTC Radiation Oncology and Lung Cancer Groups” (J Clin Oncol 27:78-84, 2009) contained errors.

In the Patients and Methods section, under Study Design, a reference to Figure 1 should have been included in the first sentence, as follows:

“This was an international multicenter phase III randomized controlled trial (RCT; **Fig 1**).”

In the Results section, the third-to-last and second-to-last sentences of the second-to-last paragraph referred to Table 5 and Figure 1, respectively, whereas it should have been Table 4 and Figure 2, respectively, as follows:

“The impact of PCI on functioning scales was similar to the one on global health status—more limited (**Table 4**). For hair loss, the difference between the two arms was larger, as there was an improvement in the control arm (**Fig 2**).”

Also in the Results section, the first sentence of the last paragraph referred to Figure 2, whereas it should have been Figure 3, as follows:

“Our exploratory analysis of the remaining HRQOL and symptom scales (ie, not preselected for the analysis) showed statistically significant ($P < .01$) and clinically significant (or near to) mean differences between the two arms for appetite loss, constipation, nausea/vomiting, social functioning, future uncertainty, headaches, motor dysfunction, and weakness of the legs at 6 weeks and/or at 3 months (**Fig 3**).”

The online version has been corrected in departure from the print.

DOI: 10.1200/JCO.2009.22.0566