Published in final edited form as:

Curr Opin Chem Biol. 2018 August ; 45: 148–156. doi:10.1016/j.cbpa.2018.05.009.

Advances in bioluminescence imaging: New probes from old recipes

Zi Yao^{1, #}, Brendan S. Zhang^{1, #}, and Jennifer A. Prescher^{1, 2, 3}

¹Departments of Chemistry, University of California, Irvine, Irvine, California 92697 USA

²Molecular Biology & Biochemistry, University of California, Irvine, Irvine, California 92697 USA

³Pharmaceutical Sciences, University of California, Irvine, Irvine, California 92697 USA

Abstract

Bioluminescent probes are powerful tools for visualizing biology in live tissues and whole animals. Recent years have seen a surge in the number of new luciferases, luciferins, and related tools available for bioluminescence imaging. Many were crafted using classic methods of optical probe design and engineering. Here we highlight recent advances in bioluminescent tool discovery and development, along with applications of the probes in cells, tissues, and organisms. Collectively, these tools are improving *in vivo* imaging capabilities and bolstering new research directions.

Introduction

Bioluminescent enzymes (luciferases) are among the most sensitive probes for imaging in thick tissues and whole organisms.[1] Luciferases catalyze light emission via the oxidation of small molecule substrates (luciferins). Since no external light is required, the background emission is virtually zero, enabling sensitive imaging *in vivo*. Bioluminescence has long been used to track cells, gene expression, and other biological features in tissues and whole organisms.[2] The emitted light is inherently weak, though, compared to conventional fluorescent tools. For this reason, luciferases are typically used in conjunction with fluorescent reporters. The bioluminescent enzymes survey processes on the macro scale and in heterogeneous environments. The fluorescent probes capture events at the micro scale or *ex vivo* – environments where excitation light is more efficiently delivered.

Historically, the most popular bioluminescent reporter for imaging *in vivo* has been firefly luciferase (Fluc). This enzyme emits the largest percentage of tissue-penetrant light with its cognate luciferin (D-luciferin, Figure 1A).[3] Other luciferases, including *Renilla* luciferase (Rluc) and *Gaussia* luciferase (Gluc) have also found broad utility in biological research.[4] These enzymes oxidize coelenterazine and emit blue light in the process. Rluc and Gluc

[#]These authors contributed equally

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

require no additional cofactors (other than oxygen), making them well suited for extracellular work. Compared to their fluorescent protein counterparts, though, luciferases have been less frequently employed in bioimaging studies. Fewer bioluminescent probes have been developed and even fewer have been optimized for application *in vivo*. There is a constant demand for more bioluminescent colors, improved enzymes, and more biocompatible substrates.

Advances in protein engineering and chemical syntheses are addressing voids in the bioluminescent toolbox. The past few years, in particular, have seen an uptick in the number of sensitive and substrate-selective luciferases available for use. Much of the progress mirrors trends in fluorescent protein development, including identifying mechanistically distinct probes in nature and subsequently evolving for new function.[5] Systematic efforts to engineer fluorescent probes for altered colors of emission, photo-switching capabilities, and other features ultimately enabled new studies in biology. This iterative cycle of tool development and biological discovery is similarly driving the field of bioluminescence. Below we highlight recent efforts to discover and evolve new bioluminescent tools, and showcase their application to biological sensing.

Discovering new luciferases and luciferins

Thousands of luminescent species exist in the natural world, but only a fraction of the associated luciferases and luciferins have been characterized in detail.[4,6] Even fewer have been coopted for use in heterologous systems.[1] Continued efforts to mine new luciferase and luciferin architectures from natural sources are expanding the number of available tools. For example, the luciferase gene from *Photinus scintillans* was recently cloned.[7] *P. scintillans* emits predominantly orange light, in contrast to the well-known North American firefly (which emits predominantly yellow-green light). The unique spectrum was traced to a single amino acid change (Y255F) in the luciferase structure. In 2016, Sharpe and colleagues reported the isolation of bioluminescent, crystalline protein assemblies in the Japanese firefly squid.[8] The crystals comprise three different—but homologous—proteins that catalyze light emission with coelenterazine-disulfate and ATP (Figure 1B).

New luciferin scaffolds and light-emitting mechanisms have also been elucidated in recent years (Figure 1B). One example includes the peptide-like luciferin from *Fridericia heliota*. Yampolsky and colleagues speculated that this molecule undergoes oxidative decarboxylation in the light-emitting reaction (similar to D-luciferin), despite its highly divergent structure.[9] This same group also discovered a new bioluminescent mechanism operative within glowing fungi. Some species convert 3-hydroxyhispidin to a putative endoperoxide en route to light emission[10*]. This scaffold is distinct from dioxetanones and other intermediates observed in classic bioluminescent reactions. Such unique luciferins and light-emitting mechanisms are potentially useful for multi-component imaging.[11]

The majority of luciferin biosyntheses remain unknown, but advances in genome sequencing are beginning to shed light on these historical mysteries. Unraveling the biosynthetic pathways would be a huge boon to *in vivo* imaging efforts: both enzyme production and substrate generation could potentially be genetically encoded in mammalian cells. Cells

would thus be autoluminescent without the need for exogenous substrate (similar to the *lux* operon for bacterial imaging).[12] Weng and coworkers have taken key steps toward this goal by harvesting the light-emitting organs from fireflies and other insects. Transcriptome profiling revealed multiple conserved genes that are likely involved in the de novo synthesis of D-luciferin.[13] Related efforts to elucidate coelenterazine biosynthesis have also been reported.[14]

Generating a palette of bioluminescent probes

The discovery and characterization of native bioluminescent systems, while important, have often not kept pace with the demand for user-friendly imaging tools. Thus, efforts to engineer bioluminescent probes with desirable properties have been critical to fill voids in the imaging toolbox. Many of the approaches have mirrored those in fluorescent protein development: mutagenesis and screening for desired properties such as thermostability, turnover, and color. Some of the most impactful luciferase engineering work in recent years has centered on NanoLuciferase (Nluc).[15] Nluc is a small (16 kDa) luciferase derived from the luminous sea shrimp *Oplophorus gracilirostris*. Nluc was evolved to process a stabilized coelenterazine analog (furimazine) in the light-emitting reaction. The Nluc-furimazine pair has been widely adopted for imaging studies in diverse fields, due to its brightness and stability. Split versions of Nluc have also been reported.[16] Like other split reporters, these tools have been useful for analyzing protein-protein interactions in cells[16] and screening inhibitors.[17]

Nluc has also proven to be a versatile platform for broadening the palette of bioluminescent probes. Much like fluorescent proteins, distinct bioluminescent reporters are desirable for applications in multicellular imaging. An enhanced set of colors can be achieved via bioluminescence resonance energy transfer (BRET, Figure 2). BRET involves luminescent reactions that excite acceptor fluorophores, resulting in altered emission spectra. The process is analogous to Förster resonance energy transfer (FRET), where energy transfer processes between two fluorophores can tune emission spectra. Nagai and colleagues recently generated a set of Nluc-fluorescent protein conjugates for BRET imaging. These chimeras were inspired by earlier Rluc-fluorescent protein conjugates (i.e., the “Nano-lanterns”). [18,19] The cyan- and green-emitting Nluc lanterns exhibited quantum yields on par with (or exceeding) Nluc itself and enabled real-time colorimetric imaging.[20*] Fluorescent dyes are also suitable BRET acceptors. Johnsson and coworkers pioneered a strategy to append different fluorophores to Nluc using SNAPtag and HaloTag technologies (Figure 2B). The suite of resulting probes provided a bioluminescent portrait reminiscent of the famous fluorescent protein collection.[21] The Nluc chimeras were also shown to be well suited for multi-component imaging in cells.[22**]

Pushing the frontiers in noninvasive imaging, the Lin lab reported an Nluc BRET construct (Antares) suitable for *in vivo* work.[23**] Antares comprises Nluc flanked by 2 copies of an orange fluorescent protein (CYOFP1). This construct produces ~20-fold more tissue-penetrant photons (>600 nm) compared to Nluc, enabling sensitive imaging in rodents. Further engineering and analog optimization yielded a second-generation reporter: Nluc (teLuc) that uses a modified furimazine analog (DTZ, Figure 2C). teLuc and DTZ provide

improved spectral overlap with CYOFP1. The optimized BRET construct (termed Antares2) exhibited enhanced red-shifted light emission and more robust bioluminescence in deep tissues.[24] Related BRET probes with Rluc have similarly provided bioluminescent probes that emit in the near-infrared regime.[25]

Efforts to produce multi-spectral tools have historically focused on the luciferase enzyme, although modifying the luciferin architecture offers another viable route. Changes to the luciferin chromophore can directly impact the color of light released. For example, extending the conjugation of the luciferin pi system or altering heteroatom substituents can alter emission wavelengths.[26–28] Both blue- and red-shifted analogs have been synthesized in recent years, although most remain weak emitters with native luciferases.[26,27,29] Engineering enzymes to better process the modified analogs—and thus recover light intensity—has been successful in some cases.[27,29,30]

Engineering orthogonal luciferase-luciferin pairs

Discriminating among wavelengths *in vivo* is challenging, as the perceived color changes with depth. Multi-component bioluminescence imaging has thus been most often achieved using *substrate*-resolved luciferases versus spectrally resolved pairs. For example, Fluc and Rluc oxidize completely different luciferins and can therefore be readily distinguished in two-component assays.[31] The Fluc/Rluc combination has further inspired the expansion of orthogonal bioluminescent tools. Unique patterns of substrate use, rather than color, can serve as diagnostic fingerprints for collections of cells or other features (Figure 3A). In our own lab, we synthesized dozens of chemically distinct luciferins and screened them against a panel of Fluc mutants.[32,33] A computer algorithm was used to identify orthogonal enzyme-substrate pairs. Substrate selectivity was maintained in both mammalian cells and in mouse models, enabling multi-cellular imaging *in vivo* (Figure 3B). Additional screening analysis further revealed triplet sets and higher-order orthogonal combinations.[33**] Simultaneous engineering of enzymes and substrates has also been applied to luciferases that use coelenterazine.[34]

Luciferase-like enzymes are further expanding the number of orthogonal probes. Luciferases belong to the ANL (Acyl-CoA synthetases, NRPS adenylation domains, and Luciferase enzymes) superfamily; these enzymes use a common mechanism to activate carboxylates as adenylates. These intermediates can be displaced with biological thiols (e.g. CoA) or, in the case of firefly luciferase, react with molecular oxygen.[35] Most ANL enzymes do not catalyze light-emitting reactions with their cognate substrates. Excitingly, though, Miller showed that some fatty acyl-CoA synthetases exhibit “latent” luciferase activity when supplied with a luminogenic substrate.[36] For example, AbLL (a synthetase from the non-luminous beetle *Agrypnus binodulus*) catalyzed light emission with a panel of synthetic luciferin analogs (Figure 3C).[36**] Many of the latent luciferases exhibit unique patterns of substrate use, expanding the number of new and orthogonal tools.

The functional and sequence similarity between luciferase and other ANL enzymes is further enabling luciferase-engineering efforts. ANL enzymes are known to be promiscuous,[35] and can potentially serve as starting points to identify new luciferase-luciferin pairs.

The Leconte lab used the homologous enzymes in combination with a previously developed bioinformatic method, statistical coupling analysis (SCA)[37,38] to guide the design of mutant luciferase libraries. SCA was used to analyze amino acid positions that were mutable and functionally important, along with networks of potentially synergistic interactions.[39*] In a single round of selection, mutants with desirable emission spectra and improved thermostability were identified. Mutants with >50-fold changes in specificity for modified luciferins were also found.

Identifying new enzyme-substrate pairs requires rapid access to diverse collections of luciferins. Such molecules have historically been difficult to synthesize from common routes. Recent advances in luciferin chemistry, though, are beginning to address this issue. Modular coupling reactions to outfit D-luciferin with diverse steric modifications have been reported.[32,40*] Ring-closing metathesis and carbene insertions have also been used to produce a series of conformationally restricted and pi-extended coelenterazines.[41–43] Many these probes exhibited red-shifted emission or other desirable photophysical properties.

Monitoring new facets of biology

Advances in luciferase engineering have ushered in a flurry of new sensors for metabolites and enzyme activities.[44–47] Many of the probes have parallels to classic fluorescent sensors, but are more tailored for *in vivo* work. A notable example is CalfluxVTN, a BRET-based calcium sensor comprising Nluc and Venus fluorescent protein (Figure 4A). In the absence of Ca^{2+} , Nluc emission is observed. Upon Ca^{2+} binding, the sensor undergoes a conformational change and BRET is observed. CalfluxVTN enabled sensitive imaging of calcium flux in response to stimulation of a rhodopsin photoreceptor.[48] Such measurements were refractory to FRET, as external light interfered with receptor activation. Johnsson and coworkers further developed a universal BRET sensor platform for analyte detection. They fused various antibody fragments to an Nluc-fluorophore pair; upon binding of a complementary analyte, a conformational change was induced, accompanied by a change in emission color.[49] The modularity of this system could enable point-of-care diagnosis for a variety of antigens.

Advances in luciferin synthesis have also enabled access to new probes of cellular function, including “caged” luciferins. These molecules typically contain a bulky group (i.e. “cage”) that renders the molecule non-emissive with luciferase. Upon removal of the cage (typically from an enzymatic reaction), an active luciferin is revealed and available for light emission. “Caged” luciferins have recently been used to detect biologically relevant metal ions and other species.[50–53**] (Figure 4B). Some have also been used to profile cell-cell interactions[54] and improve delivery.[55] The caged luminophore concept has recently been expanded to craft novel chemiluminescent sensors.[56] Some of these probes comprise embedded dioxetanes that are cleavable—and thus emit light—in response to a variety of triggers.[57*,58] Unlike canonical caged luciferins, these reporters do not require a luciferase to produce light. Such probes further diversify the portfolio of tools for biological imaging.

Conclusions and Future Directions

Many advances in bioluminescent probe technology have mirrored trends in fluorescent probe development. Dozens of luciferases have been evolved for new functions via iterative mutagenesis and screening. Collections of robust and structurally distinct luciferins have also synthesized. A variety of unique bioluminescent mechanisms have further been uncovered in the natural world, providing platforms from which to craft new tools. The continued discovery and development of bioluminescence probes, like other optical imaging agents, promises to expand what researches can “see” in cells and tissues.

Acknowledgments

This work was supported by a grant from the National Institutes of Health (R01-GM107630 to J.A.P.). B.S.Z. was supported by a GAANN Fellowship and Z.Y. was supported by the BEST IGERT program (National Science Foundation DGE-1144901). We thank members of the Prescher lab for helpful discussions.

References

Papers of particular interest, published within the period of review, have been highlighted as:

* of special interest

** of outstanding interest

1. Rathbun CM, Prescher JA. Bioluminescent probes for imaging biology beyond the culture dish. *Biochemistry*. 2017; 56:5178–5184. [PubMed: 28745860]
2. Paley MA, Prescher JA. Bioluminescence: A versatile technique for imaging cellular and molecular features. *MedChemComm*. 2014; 5:255–267. [PubMed: 27594981]
3. Zhao H, Doyle TC, Coquoz O, Kalish F, Rice BW, Contag CH. Emission spectra of bioluminescent reporters and interaction with mammalian tissue determine the sensitivity of detection *in vivo*. *J Biomed Opt*. 2005; 10:41210.doi: 10.1117/1.2032388 [PubMed: 16178634]
4. Kaskova ZM, Tsarkova AS, Yampolsky IV. 1001 lights: Luciferins, luciferases, their mechanisms of action and applications in chemical analysis, biology and medicine. *Chem Soc Rev*. 2016; 45:6048–6077. [PubMed: 27711774]
5. Rodriguez EA, Campbell RE, Lin JY, Lin MZ, Miyawaki A, Palmer AE, Shu XK, Zhang J, Tsien RY. The growing and glowing toolbox of fluorescent and photoactive proteins. *Trends Biochem Sci*. 2017; 42:111–129. [PubMed: 27814948]
6. Martini S, Haddock SH. Quantification of bioluminescence from the surface to the deep sea demonstrates its predominance as an ecological trait. *Sci Rep*. 2017; 7:45750.doi: 10.1038/srep45750 [PubMed: 28374789]
7. Branchini BR, Southworth TL, Fontaine DM, Murtiashaw MH, McGurk A, Talukder MH, Qureshi R, Yetil D, Sundlov JA, Gulick AM. Cloning of the orange light-producing luciferase from *Photinus scintillans*-a new proposal on how bioluminescence color is determined. *Photochem Photobiol*. 2017; 93:479–485. [PubMed: 27861940]
8. Gimenez G, Metcalf P, Paterson NG, Sharpe ML. Mass spectrometry analysis and transcriptome sequencing reveal glowing squid crystal proteins are in the same superfamily as firefly luciferase. *Sci Rep*. 2016; 6:27638.doi: 10.1038/srep27638 [PubMed: 27279452]
9. Dubinnyi MA, Kaskova ZM, Rodionova NS, Baranov MS, Gorokhovatsky AY, Kotlobay A, Solntsev KM, Tsarkova AS, Petushkov VN, Yampolsky IV. Novel mechanism of bioluminescence: Oxidative decarboxylation of a moiety adjacent to the light emitter of *Fridericia* luciferin. *Angew Chem Int Ed Engl*. 2015; 54:7065–7067. [PubMed: 25913753]
- 10*. Kaskova ZM, Dorr FA, Petushkov VN, Purtov KV, Tsarkova AS, Rodionova NS, Mineev KS, Guglya EB, Kotlobay A, Baleeva NS, Baranov MS, Arseniev AS, Gitelson JI, Lukyanov S, Suzuki Y, Kanie S, Pinto E, Di Mascio P, Waldenmaier HE, Pereira TA, Carvalho RP, Oliveira

AG, Oba Y, Bastos EL, Stevani CV, Yampolsky IV. Mechanism and color modulation of fungal bioluminescence. *Sci Adv.* 2017; 3:e1602847. The oxidized product of fungal bioluminescence was detected by mass spectrometry. Based on $^{18}\text{O}_2$ labeling studies, the luminescent reaction was hypothesized to involve decomposition of an endoperoxide intermediate. doi: 10.1126/sciadv.1602847 [PubMed: 28508049]

11. Maguire CA, Bovenberg MS, Crommentuijn MH, Niers JM, Kerami M, Teng J, Sena-Esteves M, Badr CE, Tannous BA. Triple bioluminescence imaging for *in vivo* monitoring of cellular processes. *Mol Ther Nucleic Acids.* 2013; 2:e99. doi: 10.1038/mtna.2013.25 [PubMed: 23778500]
12. Gregor C, Gwosch KC, Sahl SJ, Hell SW. Strongly enhanced bacterial bioluminescence with the *ilux* operon for single-cell imaging. *Proc Natl Acad Sci U S A.* 2018; 115:962–967. [PubMed: 29339494]
13. Fallon TR, Lower Sarah E, Chang Ching-Ho, Bessho-Uehara Manabu, Martin Gavin J, Bewick Adam J, Behringer Megan, Debat Humberto J, Wong Isaac, Day John C, Suvorov Anton, Silva Christian J, Hall David W, Schmitz Robert J, Nelson David R, Lewis Sara, Shigenobu Shuji, Bybee Seth M, Larracuente Amanda M, Oba Yuichi, Weng Jing-Ke. Firefly genomes illuminate the origin and evolution of bioluminescence. *bioRxiv.* 2017.
14. Francis WR, Shaner NC, Christianson LM, Powers ML, Haddock SH. Occurrence of isopenicillin-N-synthase homologs in bioluminescent ctenophores and implications for coelenterazine biosynthesis. *PLoS One.* 2015; 10:e0128742. doi: 10.1371/journal.pone.0128742 [PubMed: 26125183]
15. Hall MP, Unch J, Binkowski BF, Valley MP, Butler BL, Wood MG, Otto P, Zimmerman K, Vidugiris G, Machleidt T, Robers MB, Benink HA, Eggers CT, Slater MR, Meisenheimer PL, Klaubert DH, Fan F, Encell LP, Wood KV. Engineered luciferase reporter from a deep sea shrimp utilizing a novel imidazopyrazinone substrate. *ACS Chem Biol.* 2012; 7:1848–1857. [PubMed: 22894855]
16. Dixon AS, Schwinn MK, Hall MP, Zimmerman K, Otto P, Lubben TH, Butler BL, Binkowski BF, Machleidt T, Kirkland TA, Wood MG, Eggers CT, Encell LP, Wood KV. NanoLuc complementation reporter optimized for accurate measurement of protein interactions in cells. *ACS Chem Biol.* 2016; 11:400–408. [PubMed: 26569370]
17. Hayes MP, Soto-Velasquez M, Fowler CA, Watts VJ, Roman DL. Identification of FDA-approved small molecules capable of disrupting the calmodulin-adenylyl cyclase 8 interaction through direct binding to calmodulin. *ACS Chem Neurosci.* 2018; 9:346–357. [PubMed: 28968502]
18. Saito K, Chang YF, Horikawa K, Hatsugai N, Higuchi Y, Hashida M, Yoshida Y, Matsuda T, Arai Y, Nagai T. Luminescent proteins for high-speed single-cell and whole-body imaging. *Nat Commun.* 2012; 3:1262. doi: 10.1038/ncomms2248 [PubMed: 23232392]
19. Takai A, Nakano M, Saito K, Haruno R, Watanabe TM, Ohyanagi T, Jin T, Okada Y, Nagai T. Expanded palette of Nano-lanterns for real-time multicolor luminescence imaging. *Proc Natl Acad Sci U S A.* 2015; 112:4352–4356. [PubMed: 25831507]
- 20*. Suzuki K, Kimura T, Shinoda H, Bai G, Daniels MJ, Arai Y, Nakano M, Nagai T. Five colour variants of bright luminescent protein for real-time multicolour bioimaging. *Nat Commun.* 2016; 7:13718. New Nano-lantern conjugates were produced. The probes were used for multicolor imaging and to create a new Ca^{2+} sensor. doi: 10.1038/ncomms13718 [PubMed: 27966527]
21. Tsien lab website. <http://www.tsienlab.ucsd.edu/>
- 22**. Hiblot J, Yu Q, Sabbadini MDB, Reymond L, Xue L, Schena A, Sallin O, Hill N, Griss R, Johnsson K. Luciferases with tunable emission wavelengths. *Angew Chem Int Ed Engl.* 2017; 56:14556–14560. A palette of Nluc-fluorophore chimeras was prepared using self-labeling proteins (SNAP-tag or Halo-tag). These reporters were used for multicomponent imaging and to develop luciferase-based indicators of drugs (LUCIDs). [PubMed: 28941028]
- 23**. Chu J, Oh Y, Sens A, Ataie N, Dana H, Macklin JJ, Laviv T, Welf ES, Dean KM, Zhang F, Kim BB, Tang CT, Hu M, Baird MA, Davidson MW, Kay MA, Fiolka R, Yasuda R, Kim DS, Ng HL, Lin MZ. A bright cyan-excitable orange fluorescent protein facilitates dual-emission microscopy and enhances bioluminescence imaging *in vivo*. *Nat Biotechnol.* 2016; 34:760–767. Nluc was fused to two copies of CYOFPI, an engineered orange-emitting fluorescent protein. The resulting BRET reporter enabled improved imaging in mouse models. [PubMed: 27240196]

24. Yeh HW, Karmach O, Ji A, Carter D, Martins-Green MM, Ai HW. Red-shifted luciferase-luciferin pairs for enhanced bioluminescence imaging. *Nat Methods*. 2017; 14:971–974. [PubMed: 28869756]
25. Rumyantsev KA, Turoverov KK, Verkhusha VV. Near-infrared bioluminescent proteins for two-color multimodal imaging. *Sci Rep*. 2016; 6:36588.doi: 10.1038/srep36588 [PubMed: 27833162]
26. Anderson JC, Grounds H, Jathoul AP, Murray JAH, Pacman SJ, Tisi L. Convergent synthesis and optical properties of near-infrared emitting bioluminescent infra-luciferins. *RSC Adv*. 2017; 7:3975–3982. [PubMed: 28496975]
27. Hall MP, Woodroffe CC, Wood MG, Que I, Van't Root M, Ridwan Y, Shi C, Kirkland TA, Encell LP, Wood KV, Lowik C, Mezzanotte L. Click beetle luciferase mutant and near infrared naphthyl-luciferins for improved bioluminescence imaging. *Nat Commun*. 2018; 9:132.doi: 10.1038/s41467-017-02542-9 [PubMed: 29317625]
28. Kuchimaru T, Iwano S, Kiyama M, Mitsumata S, Kadonosono T, Niwa H, Maki S, Kizaka-Kondoh S. A luciferin analogue generating near-infrared bioluminescence achieves highly sensitive deep-tissue imaging. *Nat Commun*. 2016; 7:11856.doi: 10.1038/ncomms11856 [PubMed: 27297211]
29. Zhang BS, Jones KA, McCutcheon DC, Prescher JA. Pyridone luciferins and mutant luciferases for bioluminescence imaging. *ChemBioChem*. 2018; 19:470–477. [PubMed: 29384255]
30. Mofford DM, Reddy GR, Miller SC. Aminoluciferins extend firefly luciferase bioluminescence into the near-infrared and can be preferred substrates over D-luciferin. *J Am Chem Soc*. 2014; 136:13277–13282. [PubMed: 25208457]
31. Chan CT, Reeves RE, Geller R, Yaghoubi SS, Hoehne A, Solow-Cordero DE, Chiosis G, Massoud TF, Paulmurugan R, Gambhir SS. Discovery and validation of small-molecule heat-shock protein 90 inhibitors through multimodality molecular imaging in living subjects. *Proc Natl Acad Sci U S A*. 2012; 109:E2476–E2485. [PubMed: 22895790]
32. Jones KA, Porterfield WB, Rathbun CM, McCutcheon DC, Paley MA, Prescher JA. Orthogonal luciferase-luciferin pairs for bioluminescence imaging. *J Am Chem Soc*. 2017; 139:2351–2358. [PubMed: 28106389]
- 33**. Rathbun CM, Porterfield WB, Jones KA, Sagoe MJ, Reyes MR, Hua CT, Prescher JA. Parallel screening for rapid identification of orthogonal bioluminescent tools. *ACS Cent Sci*. 2017; 3:1254–1261. Orthogonal luciferase-luciferin pairs were identified via parallel screening of mutant enzymes and substrate analogs. These tools were applied in mouse models. [PubMed: 29296665]
34. Nishihara R, Abe M, Nishiyama S, Citterio D, Suzuki K, Kim SB. Luciferase-specific coelenterazine analogues for optical contamination-free bioassays. *Sci Rep*. 2017; 7:908.doi: 10.1038/s41598-017-00955-6 [PubMed: 28424463]
35. Gulick AM. Conformational dynamics in the acyl-CoA synthetases, adenylation domains of non-ribosomal peptide synthetases, and firefly luciferase. *ACS Chem Biol*. 2009; 4:811–827. [PubMed: 19610673]
- 36**. Mofford DM, Liebmann KL, Sankaran GS, Reddy G, Reddy GR, Miller SC. Luciferase activity of insect fatty acyl-CoA synthetases with synthetic luciferins. *ACS Chem Biol*. 2017; 12:2946–2951. A fatty-acyl CoA synthase originating from the non-luminous beetle *Agrypnus binodulus* was shown to have latent luciferase activity. The enzyme catalyzed light emission with various designer luciferin analogs. [PubMed: 29073357]
37. Socolich M, Lockless SW, Russ WP, Lee H, Gardner KH, Ranganathan R. Evolutionary information for specifying a protein fold. *Nature*. 2005; 437:512–518. [PubMed: 16177782]
38. Smock RG, Rivoire O, Russ WP, Swain JF, Leibler S, Ranganathan R, Gierasch LM. An interdomain sector mediating allostery in Hsp70 molecular chaperones. *Mol Syst Biol*. 2010; 6:414.doi: 10.1038/msb.2010.65 [PubMed: 20865007]
- 39*. Liu MD, Warner EA, Morrissey CE, Fick CW, Wu TS, Ornelas MY, Ochoa GV, Zhang BS, Rathbun CM, Porterfield WB, Prescher JA, Leconte AM. Statistical coupling analysis-guided library design for the discovery of mutant luciferases. *Biochemistry*. 2018; 57:663–671. Statistical coupling analysis (SCA) was used to construct mutant luciferase libraries. Upon screening, enzymes with red-shifted emission and improved stability were identified. Mutants with enhanced specificity for modified luciferins were also identified. [PubMed: 29224332]

- 40*. Sharma DK, Adams ST Jr, Liebmann KL, Miller SC. Rapid access to a broad range of 6'-substituted firefly luciferin analogues reveals surprising emitters and inhibitors. *Org Lett*. 2017; 19:5836–5839. Functionalization of D-luciferin was achieved via Buchwald-Hartwig amination. This chemistry enabled access to a collection of structurally diverse analogs. [PubMed: 29039673]
41. Nishihara R, Suzuki H, Hoshino E, Suganuma S, Sato M, Saitoh T, Nishiyama S, Iwasawa N, Citterio D, Suzuki K. Bioluminescent coelenterazine derivatives with imidazopyrazinone C-6 extended substitution. *Chem Commun*. 2015; 51:391–394.
- 42*. Shakhmin A, Hall MP, Machleidt T, Walker JR, Wood KV, Kirkland TA. Coelenterazine analogues emit red-shifted bioluminescence with NanoLuc. *Org Biomol Chem*. 2017; 15:8559–8567. Several red-shifted luciferin analogs were accessed by functionalization of the furimazine core. Red-shifted BRET constructs utilizing the luciferin analogs were developed. [PubMed: 28972606]
43. Hosoya T, Iimori R, Yoshida S, Sumida Y, Sahara-Miura Y, Sato J, Inouye S. Concise synthesis of ν -coelenterazines. *Org Lett*. 2015; 17:3888–3891. [PubMed: 26194904]
44. Shigeto H, Ikeda T, Kuroda A, Funabashi H. A BRET-based homogeneous insulin assay using interacting domains in the primary binding site of the insulin receptor. *Anal Chem*. 2015; 87:2764–2770. [PubMed: 25655236]
45. den Hamer A, Dierickx P, Arts R, de Vries J, Brunsveld L, Merckx M. Bright bioluminescent BRET sensor proteins for measuring intracellular caspase activity. *ACS Sens*. 2017; 2:729–734. [PubMed: 28670623]
46. Takenouchi O, Kanno A, Takakura H, Hattori M, Ozawa T. Bioluminescent indicator for highly sensitive analysis of estrogenic activity in a cell-based format. *Bioconjug Chem*. 2016; 27:2689–2694. [PubMed: 27690388]
47. Aper SJ, Dierickx P, Merckx M. Dual readout BRET/FRET sensors for measuring intracellular zinc. *ACS Chem Biol*. 2016; 11:2854–2864. [PubMed: 27547982]
48. Yang J, Cumberbatch D, Centanni S, Shi SQ, Winder D, Webb D, Johnson CH. Coupling optogenetic stimulation with NanoLuc-based luminescence (BRET) Ca^{++} sensing. *Nat Commun*. 2016; 7:13268.doi: 10.1038/ncomms13268 [PubMed: 27786307]
49. Xue L, Yu Q, Griss R, Schena A, Johnsson K. Bioluminescent antibodies for point-of-care diagnostics. *Angew Chem Int Ed Engl*. 2017; 56:7112–7116. [PubMed: 28510347]
50. Ke B, Wu W, Liu W, Liang H, Gong D, Hu X, Li M. Bioluminescence probe for detecting hydrogen sulfide *in vivo*. *Anal Chem*. 2016; 88:592–595. [PubMed: 26634959]
51. Kojima R, Takakura H, Kamiya M, Kobayashi E, Komatsu T, Ueno T, Terai T, Hanaoka K, Nagano T, Urano Y. Development of a sensitive bioluminescent probe for imaging highly reactive oxygen species in living rats. *Angew Chem Int Ed Engl*. 2015; 54:14768–14771. [PubMed: 26474404]
52. Heffern MC, Park HM, Au-Yeung HY, Van de Bittner GC, Ackerman CM, Stahl A, Chang CJ. *In vivo* bioluminescence imaging reveals copper deficiency in a murine model of nonalcoholic fatty liver disease. *Proc Natl Acad Sci U S A*. 2016; 113:14219–14224. [PubMed: 27911810]
- 53*. Aron AT, Heffern MC, Lonergan ZR, Vander Wal MN, Blank BR, Spangler B, Zhang Y, Park HM, Stahl A, Renslo AR, Skaar EP, Chang CJ. *In vivo* bioluminescence imaging of labile iron accumulation in a murine model of *Acinetobacter baumannii* infection. *Proc Natl Acad Sci U S A*. 2017; 114:12669–12674. A caged luciferin reporter for Fe^{2+} was developed. This probe was used to monitor iron homeostasis in response to infection. [PubMed: 29138321]
54. Porterfield WB, Prescher JA. Tools for visualizing cell-cell 'interactomes'. *Curr Opin Chem Biol*. 2015; 24:121–130. [PubMed: 25461730]
55. Mofford DM, Adams ST Jr, Reddy GS, Reddy GR, Miller SC. Luciferin amides enable *in vivo* bioluminescence detection of endogenous fatty acid amide hydrolase activity. *J Am Chem Soc*. 2015; 137:8684–8687. [PubMed: 26120870]
56. Hananya N, Shabat D. A glowing trajectory between bio- and chemiluminescence: From luciferin-based probes to triggerable dioxetanes. *Angew Chem Int Ed Engl*. 2017; 56:16454–16463. [PubMed: 28967167]
- 57*. Green O, Gnaïm S, Blau R, Eldar-Boock A, Satchi-Fainaro R, Shabat D. Near-infrared dioxetane luminophores with direct chemiluminescence emission mode. *J Am Chem Soc*. 2017;

139:13243–13248. Red-shifted chemiluminescent probes were developed for *in vivo* imaging. These scaffolds were used as sensors for peroxide and β -galactosidase activity. [PubMed: 28853880]

58. A chemiluminescent probe for cellular peroxynitrite using a self-immolative oxidative decarbonylation reaction. *Chem Sci*. 2018; 9:2552–2558. [PubMed: 29732134]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- Advances in bioluminescent tool production that mirror trends in fluorescent probe development.
- New bioluminescent platforms based on naturally occurring luciferases and luciferins.
- Engineered probes that enable deep tissue and multi-component bioluminescence imaging.
- Applications of bioluminescent tools to cellular biosensing

Figure 1. Luciferase-luciferin pairs in nature

(A) Beetle luciferases oxidize D-luciferin using ATP and O₂, generating primarily yellow-green light. Marine luciferases release blue photons via the oxidation of imidazopyrazinone analogs. (B) Recently characterized luciferases exploit unique molecules and mechanisms to produce light. The relevant luciferins and oxyluciferin products are shown. Orange dots mark the sites of oxidation.

Figure 2. Expanded palette of bioluminescent probes

(A) Resonance energy transfer can tune optical emission spectra. In FRET, donor fluorophores (e.g., CFP) are excited with external light. Emission from the acceptor fluorophore (e.g., YFP) is observed. In BRET, luminescent reactions can excite acceptor fluorophores, resulting in altered emission spectra. A sample BRET construct (ReNL, pairing Nluc with tdTomato) is shown. (B) Nluc-fluorophore chimeras expand the palette of bioluminescent probes. Fluorescent molecules were appended to Nluc via SNAPtag ligation (left), generating a colorful array of S-Luc tags (right). S-Luc images were reproduced with permission from ref. 22. (C) Far red-emitting BRET constructs enable sensitive imaging *in vivo*. Antares comprises Nluc and two copies of a fluorescent protein (CyOPF1). Red-shifted light is produced upon luciferin administration. Antares and a related construct (Antares2) were expressed in mice following hydrodynamic transfection. Light emission was observed upon furimazine or DTZ administration ($3.3 \mu\text{mol i.p.}$). Mouse images were reproduced with permission from ref 24.

Figure 3. Substrate-selective luciferases for multi-component imaging

(A) Orthogonal luciferases were identified via parallel screening of luciferase mutants and luciferin analogs. (B) Dual imaging with engineered luciferase-luciferin pairs. DB7 cells expressing orthogonal mutants (37 and 81) were inoculated in opposing flanks. The populations were readily distinguished upon administration of the complementary luciferins. Bioluminescence images were reproduced with permission from ref. 33. (C) Fatty acyl-CoA synthetases from non-luminous organisms (e.g., CG6178 from *D. melanogaster* and AbLL from *A. binodulus*) exhibit luciferase-like behavior with synthetic luciferin analogs. Bar graph was reproduced with permission from ref. 36.

Figure 4. Visualizing cellular species with bioluminescent sensors

(A) CalfluxVTN comprises a calcium-binding protein (TroponinC) flanked by Nluc and YFP (left). Ca^{2+} binding induces a conformational change in the sensor, resulting in BRET. ClafluxVTN was expressed in neurons and used to monitor Ca^{2+} flux following photoreceptor firing (right, scale bar = 20 μm). Cellular images were reproduced with permission from ref. 48. **(B)** Caged luciferins and dioxetanes can report on cellular activities. Selective removal of the caging groups provides an active luminophore. Light emission via luciferase oxidation (top) or direct chemiluminescence (bottom) thus provides a readout on the uncaging enzyme or analyte of interest.