Published in final edited form as:

J Fish Dis. 2018 May ; 41(5): 805–816. doi:10.1111/jfd.12784.

Construction and evaluation of type III secretion system mutants of the catfish pathogen *Edwardsiella piscicida*

Asmaa Edrees¹, Hossam Abdelhamed², Seong-Won Nho², Seong bin Park³, Attila Karsi², Frank W. Austin², Manal Essa¹, Tibor Pechan⁴, and Mark L. Lawrence²

¹Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef, Egypt Mississippi State University, Mississippi State, MS, USA

²College of Veterinary Medicine, Mississippi State University, Mississippi State, MS, USA

³Department of Animal and Dairy Science, Mississippi State University, Mississippi State, MS, USA

⁴Institute for Genomics, Biocomputing and Biotechnology, Mississippi State University, Mississippi State, MS, USA

Abstract

Catfish is the largest aquaculture industry in the United States. Edwardsiellosis is considered one of the most significant problems affecting this industry. *Edwardsiella piscicida* is a newly described species within the genus *Edwardsiella*, and it was previously classified as *Edwardsiella tarda*. It causes gastrointestinal septicemia, primarily in summer months, in farmed channel catfish in the southeastern U.S. In the current study, we adapted gene deletion methods used for *Edwardsiella* to *E. piscicida* strain C07–087, which was isolated from a disease outbreak in a catfish production pond. Four genes encoding structural proteins in the type III secretion system (T3SS) apparatus of *E. piscicida* were deleted by homologous recombination and allelic exchange to produce in-frame deletion mutants (*Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT*). The mutants were phenotypically characterized, and virulence and vaccine efficacy were evaluated. Three of the mutants, *Ep ssaV*, *Ep yscR*, and *Ep esaM*, were significantly attenuated compared to the parent strain ($P < 0.05$), but *Ep escT* strain was not. Vaccination of catfish with the four mutant strains (*Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT*) provided significant protection when subsequently challenged with wild type strain. In conclusion, we report methods for gene deletion in *E. piscicida* and development of vaccine candidates derived from a virulent catfish isolate.

Keywords

Edwardsiella piscicida; type III secretion system; virulence; catfish; aquaculture

---

Correspondence to: Mark L. Lawrence.

Correspondence M L Lawrence, College of Veterinary Medicine, Mississippi State University, Mississippi State, MS 39762, USA, (lawrence@cvm.msstate.edu).

Introduction

Edwardsiella piscicida is one of five species in the genus *Edwardsiella*. Prior to 2012, the genus consisted of 3 species: *E. ictaluri*, *E. hoshinae*, and *E. tarda*. However, based on collective evidence from genome sequencing and phenotypic comparisons, isolates from fish previously identified as *E. tarda* were reclassified into three different genetic groups: *E. tarda*, *E. piscicida*, and *E. anguillarum*. All three species can grow in a temperature range from 25–37°C; however, the optimum temperature for growth of *E. piscicida* is 28–30°C (Abayneh et al., 2013). Analysis of “*E. tarda*” isolates from channel catfish (*Ictalurus punctatus*) aquaculture in the southeast U.S. revealed that strains associated with gastrointestinal septicemia, especially in summer months, are correctly classified as *E. piscicida* (Griffin et al., 2014). A quantitative PCR method specific for *E. piscicida* was developed and used to quantify the pathogen in various organs in both clinically and subclinically infected fish (Reichley et al., 2015).

E. piscicida was also isolated from farmed whitefish, *Coregonus lavaretus* (L.), in Finland during disease outbreaks in 2000, 2002, and 2013. The species was identified by biochemical tests, a species-specific PCR, enterobacterial repetitive intergenic consensus PCR (ERIC-PCR), and multilocus sequence analysis (MLSA) (Shafiei et al., 2016). *E. piscicida* was reported as the causative agent of Edwardsiellosis in cage-cultured sharpsnout sea bream (*Diplodus puntazzo*), indicating that *E. piscicida* is a possible serious threat for this Mediterranean aquaculture industry (Katharios et al., 2015).

Like many Gram-negative bacteria, *E. piscicida* utilizes a type III secretion system (T3SS) to modulate host cell functions (Blocker et al., 2008). Type III secretion system (T3SS) is a needle-like structure composed of several structural proteins encoded by genes typically located in a chromosome or plasmid locus. It is an important component in the pathogenesis of many Gram-negative bacteria, including several fish pathogens, and its function is to inject virulence proteins directly into host cells to disrupt normal functions (Tan et al., 2005; Xie et al., 2014).

In aquaculture, bacterial infections are often treated with antimicrobial therapy. However, this strategy can result in the development of antimicrobial resistant *Edwardsiella* strains and subsequent treatment failures (Heuer et al., 2009; Wang et al., 2009; Lo et al., 2014). Furthermore, losses have already occurred when medicated feeds are applied; therefore, prevention is a more desirable strategy for farmers. Thus, there is increased interest in vaccination as a method to prevent fish diseases (Marsden et al., 1996; Temprano et al., 2005). The objective of the current study was to construct T3SS *E. piscicida* mutant strains by deleting genes that encode four structural proteins (AGH72974, AGH72975, AGH72976, and AGH72978) to determine the effects on phenotype, and to evaluate their applicability as an attenuated vaccine against *E. piscicida* infection in catfish.

Materials and methods

Bacterial strains, plasmids, and growth conditions.

Bacterial strains and the plasmids used in this work are listed in Table 1. *E. piscicida* C07–087 and mutant strains were cultured in brain heart infusion (BHI) agar and broth (Becton Dickinson, Sparks, MD, USA) and incubated at 30 °C. *Escherichia coli* strains were cultured in Luria–Bertani (LB) agar and broth (Becton Dickinson) and incubated at 37 °C. *E. coli* CC118 λ pir strain was used for cloning and maintenance of pMEG-375 plasmids, and *E. coli* BW19851 and SM10 λ pir strains were used for transferring plasmids into *E. piscicida* by conjugation. Ampicillin was used at 100 µg/ml to select for integration of pMEG-375 derivatives in the *E. piscicida* chromosome or for maintenance of pAKgfplux1. Colistin was used at 12.5 µg/ml for counterselection against *E. coli* following conjugation.

Construction of in-frame deletion mutant strains

Four in-frame deletion mutant strains (*Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT*) were constructed through allelic exchange as described for *E. ictaluri* (Abdelhamed et al., 2013; Dahal et al., 2013). All primers used in mutant construction are given in Table 2. Upstream and downstream fragments for each gene deletion were amplified using primer pair A and B and primer pair C and D, respectively, from each target gene. Fusion of the two fragments was done by overlap extension PCR using the mixed upstream and downstream products as a template DNA with primer pair A and D. The resulting in-frame deleted fragments were purified using Gel/PCR DNA Fragment Extraction Kit (IBI Scientific, Peosta, IA, USA). Plasmid pMEG-375 was purified from *E. coli* CC118 using Hi-Speed Mini Plasmid Kit (IBI Scientific), and the plasmid and overlap extension PCR amplicons were digested with *SacI* and *XbaI* (New England Biolabs, Ipswich, MA, USA) at 37 °C for 8 hours. Amplicons were ligated into pMEG-375 by T4 ligase (New England Biolabs) at 16°C, generating *pEp ssaV*, *pEp esaM*, *pEp yscR*, and *pEp escT* (Table 1).

The resulting plasmids were transferred into *E. coli* BW19851 by electroporation and subsequently mobilized to *E. piscicida* C07–087 by filter conjugation (Karsi and Lawrence, 2007). Selection was on BHI agar plates with colistin and ampicillin. Colonies with ampicillin resistance were streaked on BHI plates with 5% sucrose and 0.35% mannitol (Sigma-Aldrich, St. Louis, MO, USA) and grown for 2–3 days to allow curing pMEG-375 vector from the chromosome using *sacB* counterselection (Santander et al., 2011). After this, resulting colonies were inoculated in BHI broth and tested for ampicillin sensitivity to ensure loss of plasmid. Gene deletions were confirmed by PCR and sequencing using primers listed in Table 1. After confirmation, *Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT* mutants were labeled with bioluminescence using pAKgfplux1 plasmid (Karsi and Lawrence, 2007).

Determination of growth kinetics

The growth kinetics of *E. piscicida* C07–087 and four mutant strains were determined in BHI broth (Liu et al., 2012). Briefly, overnight cultures in BHI were normalized based on optical density at 600 nm (OD600), and pellets were washed twice in phosphate buffered saline (PBS) before being subcultured at 1:1000 dilution into 20 ml BHI. OD600 was

measured at 30 min intervals until the onset of stationary phase. All growth kinetics experiments were performed four times from two independent starter cultures.

Biochemical characterization

E. piscicida C07–087 and mutant strains were suspended in PBS for Gram stain, catalase, oxidase, and motility tests. Other biochemical tests were performed using the API 20E kit (BioMerieux Inc., Durham, NC, USA) according to the manufacturer's instructions. Pure cultures were transferred aseptically to Lysine Iron Agar slant (Remel, Lenexa, KS, USA) and incubated for 24 hours at 35°C to determine H₂S production.

Biofilm assay

Biofilm formation was determined as described (Wakimoto et al., 2004) with minor modifications. Briefly, overnight cultures of *E. piscicida* C07–087 and mutant strains were diluted 1:100 in BHI supplemented with 1% glucose (Sigma-Aldrich), and 100 µl of the bacterial suspension was transferred into the wells of a 96-well microtiter plate (Coster, USA). Plates were incubated at 30°C for 72 hours, wells were gently washed two times with PBS, and they were stained with 0.1% crystal violet (Sigma-Aldrich) for 10 min at room temperature. Plates were rinsed with 100 µl PBS and dried at room temperature, and 100 µl of 70% ethanol was used for destaining. Finally, biofilm formation was quantified by measuring absorbance at 538 nm with a SpectraMax M5 ELISA reader (Molecular Devices, Sunnyvale, CA, USA). Biofilm formation was determined five times for each strain.

Hemolysin activity

Hemolysin activity was determined in triplicate as described (Xu et al., 2010) with minor modifications. Sheep erythrocytes (Innovative Research, Novi, MI, USA) were washed three times with hemolysis buffer [0.85% physiological saline, 0.01 M CaCl₂ (Sigma-Aldrich)] and resuspended in the same buffer to make a final concentration of 3% sRBCs (Watson and White, 1979). *E. piscicida* C07–087 and mutant strains were cultured in BHI for 24 hours at 30°C with shaking, diluted 1:100 in BHI, and incubated at 30°C for 2, 4, 6, 8, 10 and 12 hours. Bacteria were harvested by centrifugation and washed two times in hemolysis buffer. Assay tubes containing 0.2 ml sRBCs, 250 µl bacteria, and 1.55 ml hemolysis buffer were incubated at 30°C for one hour and then pelleted by centrifugation. Absorbance of the supernatant at 560 nm was measured to determine hemolytic activity. Assay tubes containing 1.8 ml hemolysis buffer and 0.2 ml sRBCs was included as negative control, and positive control consisted of 1.8 ml sterile water and 0.2 ml sRBCs. Percent hemolysis was calculated as follows:

$$\frac{[(\text{Absorbance of sample} - \text{Absorbance of negative control}) / \text{Absorbance of positive control}]}{X} \times 100.$$

Extracellular proteins

Extracellular proteins (ECPs) from *E. piscicida* C07–087 and mutant strains were extracted as described (Wang et al., 2010b) with some modifications. One hundred microliters of cultured bacteria were spread on BHI agar medium and incubated at 30 °C for 48 hours. Bacteria were collected, suspended in PBS, and pelleted by centrifugation. Supernatant was

filtered through a 0.22 µm syringe filter (VWR, Radnor, PA, USA). ECPs were precipitated with trichloroacetic acid (TCA, Sigma-Aldrich) and dissolved in PBS. After centrifugation, supernatants were kept, and protein concentrations were determined using Bradford assay (Bio-Rad). One half volume of tricine sample buffer (Bio-Rad, Hercules, CA, USA) was added, and mixture was sonicated by Fisher Scientific Sonic Dismembrator (Waltham, MA, USA). A total of 100 µg of ECPs was loaded on pre-cast gels (Bio-Rad) for SDS-PAGE, and gels were stained with Coomassie Brilliant Blue (Bio Rad).

In-gel digestion was performed as described (Nho et al., 2011) with minor modifications. Briefly, sliced protein bands were completely destained in a mixture of equal volumes of 75 mM ammonium bicarbonate (ABC) and 40% ethanol with vortexing. Gel strips were reduced with 10 mM dithiothreitol (DTT) in 25 mM ammonium bicarbonate (ABC) for 1 h and alkylated by incubating with 55 mM iodoacetamide (IAA) in 25 mM ABC for 45 min in the dark. Supernatants were discarded, and 100 mM ABC was added to the gel bands, followed by dehydration with 100% acetonitrile (ACN). This process was repeated once, and then gels were dried using a vacuum dryer. Trypsin solution (0.02 µg/ml; Promega, Madison, WI, USA) was added to the dried gel particles. After incubating on ice for 40 min, 25 mM ABC was added, and samples were incubated at 37°C for 14 h. Fresh 25 mM ABC was added, samples were vortexed for 30 min, and supernatants were transferred to fresh tubes. Gels were re-extracted by adding extraction solution (0.1% [v/v] formic acid in 25% [v/v] ACN) and vortexing for 30 min. Supernatants were transferred to the tubes containing tryptic peptide mixture from the first extraction, and samples were stored at –20°C until LC-MS/MS analysis.

Liquid Chromatography-Tandem Mass Spectrometry

Mass spectra were collected and analyzed (Takac et al., 2016) using an UltiMate 3000 nano-HPLC system linked to a LTQ Orbitrap Velos mass spectrometer (ThermoFisher Scientific, Waltham, MA, USA). Protein tryptic digest was separated on an Acclaim PepMap C18 column (75µm x 150mm, ThermoFisher Scientific) in 60 min linear gradient of acetonitrile (2%–55% for 35 min, 95% for 10 min, 2% for 15 min), with a flow rate of 300 nl/min. Intact and fragmented peptide ions were measured utilizing the linear trap mass detector in data dependent acquisition mode with dynamic exclusion being applied.

The raw mass spectral data were matched to *Edwardsiella piscicida* protein database using the SEQUEST algorithm of Proteome Discoverer 1.1 (Thermo Fisher Scientific). The target database (7,667 referenced sequences as of March 2016) was obtained from National Center for Biotechnology Information (www.ncbi.nlm.nih.gov). The reversed copy (created automatically by Proteome Discoverer) served as a decoy database to calculate false discovery rate (FDR). To obtain high confidence protein identification, results were filtered with FDR value of less than 1.0%.

Bioluminescence imaging

Bioluminescence imaging (BLI) was conducted using an IVIS Imaging System (IVIS Lumina XRMS In Vivo Imaging System, PerkinElmer, Waltham, MA, USA) (Karsi et al., 2006). Five fish per bacterial strain were anesthetized and injected IP with 3.7×10^6 CFU/

fish. For imaging, fish were anesthetized and transferred to the imaging chamber; total photon emissions from the whole fish body were collected using an exposure time of one minute. Following BLI, fish were returned to well-aerated water for recovery. BLI was conducted at 0, 2.5, 6, 12, 24, and 36 h post-infection.

Virulence/vaccine efficacy of *E. piscicida* T3SS mutants

All fish experiments were approved by the Institutional Animal Care and Use Committee (IACUC) at Mississippi State University. Specific-pathogen-free (SPF) channel catfish fingerlings (19 ± 2.04 g, 11 ± 0.67 cm) were obtained from the SPF Channel Catfish Laboratory at Mississippi State University's College of Veterinary Medicine. Fish were maintained in aerated tanks supplied with a continuous flow of recirculating water (flow rate: 1-L/min) at water temperature $28 \pm 1^\circ\text{C}$, and they were fed twice daily with commercial channel catfish feed. Anesthesia was provided using tricaine methane sulfonate (MS-222) when needed.

We first determined the 50 percent lethal dose (LD50) of *E. piscicida* C07–087. Eight treatment groups were included, each with 3 tanks with 10 fish in each. Overnight cultures of *E. piscicida* C07–087 strain in BHI broth were harvested by centrifugation, washed twice with sterile PBS, and then serially diluted. Fish were injected intraperitoneally (IP) with 0.1 ml of 1.9×10^3 , 9.3×10^3 , 4.7×10^4 , 2.3×10^5 , 1.2×10^6 , 5.8×10^6 , 2.9×10^7 , and 1.5×10^8 CFU/fish of *E. piscicida* C07–087 strain. One more group was used as a negative control group that was handled similarly but injected IP with sterile PBS. Fish were observed for 7 days, and any fish that died were removed for routine bacteriological examination. The LD50 was calculated (Reed and Muench, 1938).

Based on the LD50 determination, a virulence/vaccine trial was conducted. For the virulence determination, about 180 six-month-old SPF channel catfish fingerlings were stocked into 18 tanks at a rate of 10 fish/tank. Each treatment had three replicate tanks. Treatments consisted of *E. piscicida* C07–087, *Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT*, and negative control (sterile PBS). Catfish were experimentally infected by IP injection (0.1 ml per fish). Injection doses were approximately 9×10^5 , 7.5×10^5 , 1.14×10^6 , 9×10^5 , and 1.6×10^6 , and CFU/fish for *E. piscicida* C07–087, *Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT*, respectively. Mortalities were recorded for 21 days. Following the virulence trial (21 days post-exposure from the first challenge), all animals from the mutant-exposed and negative control treatment groups were experimentally infected by IP injection with *E. piscicida* C07–087 (1.2×10^6 CFU/fish). Mortalities were recorded for 7 days, and any fish that died were removed for routine bacteriological examination.

Statistical analysis

Linear regression analysis in PROC MIXED of SAS for Windows 9.4 (SAS Institute, Inc., Cary, NC, USA) was used for growth kinetics, biofilm, hemolysis, and bioluminescence assays where the mutant strains were compared to *E. piscicida* strain C07–087. Separate models were developed for each analytical test using manual forward selection, and each included mutant type and hour as fixed effects. Rep was included as a random effect.

Adjustment for multiple comparisons were made using the SIMULATE option. Results are reported as least square means $\pm$ standard error.

Exact logistic regression was used in PROC LOGISTIC of SAS for Windows 9.4 (SAS Institute, Inc., Cary, NC, USA) to evaluate percent mortalities during the pathogenicity and vaccine trials. The outcome was the number of deaths over the total number of trials. For the challenge analysis, comparisons were made to the negative control, and separate models were developed for low and high doses. Odds ratios and 95% Confidence Limits are reported. For the pathogenicity analysis, comparisons were made to strain C07–087.

Results

Construction of *E. piscicida* mutant strains

Genes *ssaV*, *esaM*, *yscR*, and *escT* encode structural proteins that contribute to formation of the T3SS apparatus of *E. piscicida* C07–087 (Table 3). These four genes were deleted to obtain four mutant strains: *Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT* (Table 1). All mutant strains were ampicillin sensitive, and each was confirmed by PCR (Figure 1) and sequencing.

Growth kinetics

Growth kinetics of *E. piscicida* C07–087 and mutant strains were determined in BHI broth in quadruplicate. No significant difference in growth kinetics was detected between the parent wild type strain and any of the mutants (data not shown).

Biochemical characterization

E. piscicida C07–087 and mutant strains were all cytochrome oxidase negative and catalase positive, and all had the same pattern in API20E (positive reaction on ortho nitrophenyl- β -D-galactopyranosidase, ornithine decarboxylase, indole production, citrate utilization, and glucose utilization; negative reaction on gelatin liquefaction, mannose, sucrose, arabinose, sorbitol, inositol, rhamnose, and melibiose utilization). For H₂S production, all strains showed positive reaction except *Ep esaM* strain, which had a weak positive reaction. This finding was confirmed by streaking an isolated colony from each mutant and the parent strain C07–087 on Lysine Iron Agar slant with incubation at 35 °C for 24 hours, where *Ep esaM* showed the lowest H₂S production.

Biofilm formation

None of the wild type or mutant strains had significant biofilm formation at 24 or 48 hours. All four mutants and parent strain C07–087 had significant biofilm formation at 72 hours ($P < .05$), but there were no significant differences in biofilm formation between strains (Figure 2).

Hemolysin activity

Hemolysin activity was low for all strains at 2 hours incubation and was near 100% for all strains at 6 hours incubation. No significant differences between strains were detected at these time points. However, at 4 hours incubation, mutant strains *Ep esaM* and *Ep escT*

had significantly higher level of hemolytic activity than wild type strain C07–087 ($P=0.0002$) (Figure 3).

Extracellular proteins

SDS-PAGE of ECPs from *E. piscicida* C07–087 and mutant strains indicated a band with molecular weight of approximately 150 kDa was present in *E. piscicida* C07–087, *Ep ssaV*, and *Ep yscR*, but it was missing in *Ep escT* and *Ep esaM* (Figure 4). For comparison, the band from C07–087 and the same gel region from strain *Ep escT* were excised and analyzed by the same method. Mass spectrometry results are shown in Table 4. Parent strain C07–087 exhibited a dominant presence of flagellin protein as indicated by a high number of spectra (>200) matched to peptides belonging to four individual protein entries in the database. On the contrary, only two flagellin peptides were detected in mutant strain *Ep escT*. RNA polymerase subunit was also detected in both strains.

Bioluminescence imaging

Bioluminescence (quantified as photon counts) from the catfish infected with each of the T3SS mutants (*Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT*) was significantly lower than that of strain C07–087 for at least one time point (Figure 5). Strain *Ep ssaV* was the most attenuated and had significantly lower bioluminescence at every time point after 2.5 h post-infection. Strain *Ep esaM* had significantly lower bioluminescence at early time points (6 and 12 h post-infection), and strain *Ep escT* was similar (significantly lower at 12 h post-infection). Strain *Ep yscR* had a different pattern with significantly lower bioluminescence at later time points (24 and 36 h post-infection).

Virulence/vaccine efficacy of *E. piscicida* T3SS mutants

The LD50 of *E. piscicida* strain C07–087 was 3.3×10^5 CFU/fish, and infected fish died within 6 days with typical signs of hemorrhagic septicemia. In the virulence comparison, significantly ($P < 0.05$) higher mortalities occurred in fish infected with parent strain C07–087 (26%) compared with mortalities in fish infected with strains *Ep ssaV* (0%), *Ep yscR* (0%), and *Ep esaM* (6%), (Figure 6A). By contrast, there was no significant difference ($P > 0.05$) in mortalities between fish injected with *Ep escT* (16%) and parent strain C07–087. To determine potential vaccine efficacy of *E. piscicida* T3SS mutants, fish were challenged with *E. piscicida* C07–087 by IP injection three weeks post-immunization. Fish were protected significantly ($P < 0.001$) after vaccination with all four mutants. After the second challenge, no mortalities occurred in fish vaccinated with *Ep esaM*, *Ep yscR*, and *Ep escT*, whereas *Ep ssaV*-vaccinated group had some mortalities (6%) compared with 100% mortalities in sham-vaccinated group (Figure 6B).

Discussion

The newly described *Edwardsiella* species *E. piscicida* has been isolated from diseased fish of multiple species (Griffin et al., 2014; Katharios et al., 2015; Shafiei et al., 2016). T3SS is a nanosyringe that forms a channel across the bacterial membrane and the host cell membrane to transport proteins into the host cell. There are four types of T3SS proteins: apparatus, effectors, chaperons, and regulators. The apparatus is complex and typically

comprised of almost 25 proteins (Hueck, 1998). SsaV, YscR, EsaM, and EscT are four T3SS apparatus proteins in the inner membrane that are collectively necessary for organization of the inner membrane rings. The individual functions of these proteins are still unclear (Galán et al., 2014; Notti and Stebbins, 2016).

In *E. piscicida* C07–087, the *esaM* gene was annotated as encoding a hypothetical protein, but it showed 99.9% identity with a *E. tarda* EIB202 T3SS apparatus gene. The four genes encoding these proteins are located in adjacent divergently transcribed operons with *ssaV* and *esaM* located in one operon and *yscR* and *escT* located in another. These genes were selected for deletion to determine the effects of inner membrane rings on T3SS function and virulence. To our knowledge, these proteins have not been characterized in any *Edwardsiella* species, and in general they are not well studied. The current study also demonstrates that gene deletion methods developed for *E. ictaluri* are effective in *E. piscicida*.

Similar to previous studies on the *E. tarda* T3SS, disruption of the *E. piscicida* T3SS did not interfere with growth kinetics in BHI at 30 °C (Lan et al., 2007; Wang et al., 2010a). Disruption of genes encoding T3SS structural proteins also does not affect growth of plant pathogens on plant cells (Sun et al., 2011). Using API20E strips, the four T3SS mutants were also biochemically homogenous and similar to parent strain C07–087.

The one exception to the biochemical homogeneity is that strain *Ep esaM* showed lower production of H₂S, which was confirmed in Lysine Iron Agar. The H₂S production test measures the ability of bacteria to reduce sulfur-containing compounds to sulfides during metabolism. For example, certain genera of Enterobacteriaceae are able to reduce tetrathionate during respiration as a terminal electron acceptor (as an alternative to oxygen). This is controlled by *ttr* genes, which in *Salmonella* are located within pathogenicity island 2 (SPI-2). SPI-2 also encodes a T3SS that facilitates pathogenesis of *Salmonella* (Figueira and Holden, 2012). Transcription of the *ttr* operon is controlled by a TtrS sensor and TtrR response regulator (Hensel et al., 1999). It is possible that the difference in H₂S production in *Ep esaM* is not due to a difference in effects on T3SS inner membrane ring function; it may be due to a secondary function of EsaM. Interestingly, although EsaM has no similarity with proteins encoded by any known *ttrRSBCA* genes, it has 33.7% identity with putative transcriptional regulator MerR in *Mycobacterium* sp. JLS. If EsaM has a secondary function as a transcriptional regulator, it may have a regulatory effect on the *ttr* operon.

Biofilm formation is important for bacterial survival in adverse environmental conditions, including escape from the host immune system (Wakimoto et al., 2004). Therefore, biofilm formation can contribute to disease pathogenesis and persistence in the host (Sundell and Wiklund, 2011). Biofilm formation requires multiple steps (Watnick and Kolter, 2000). In *P. aeruginosa*, T3SS is required for cell-associated aggregation on the surface of polarized epithelial cells, but it is not required for aggregation on abiotic surfaces (Tran et al., 2014). Therefore, it is possible that we did not detect significant differences in biofilm formation in our T3SS mutants because the assays were performed on plates.

Several virulence factors contribute to *E. tarda* pathogenicity including secretion systems, flagella, fimbriae, hemagglutinins, chondroitinases, hemolysins, and iron scavenging system (Wang et al., 2009; Wang et al., 2010b; Park et al., 2012). In the current study, we investigated effects of the selected T3SS apparatus genes on hemolytic activity of *E. piscicida*. Our results revealed that hemolysis requires 6 hours of incubation at 30 °C in *E. piscicida*, which is similar to previously reported results from *E. tarda* (Watson and White, 1979). Interestingly, *Ep esaM* and *Ep escT* had increased hemolytic activity compared to parent strain C07–087. In *E. tarda* EIB202, a *esrB* mutant strain had higher hemolytic activity than its parent strain (Wang et al., 2010b) due to upregulation of EthA, a hemolysis associated protein. Although *E. tarda* EsrB does not have any similarity to *E. piscicida* EsaM or EscT, it is possible that increased hemolysin activity in *Ep esaM* and *Ep escT* is due to a similar effect in *E. piscicida*.

Virulent and avirulent strains of *E. tarda* have differences in extracellular protein expression (Tan et al., 2002). We expected that mutation of T3SS genes in *E. piscicida* could affect ECP expression. On SDS-PAGE of ECPs, *Ep escT* and *Ep esaM* mutants showed a missing protein band with a molecular mass of approximately 150 kDa. Analysis of the 150 kDa region by band excision, in-del digestion, and LC-MS/MS showed high abundance of flagellin proteins in *E. piscicida* C07–087, while only traces of flagellin were detected in *Ep escT*. Theoretical Mr of the flagellins is given as 43.5–43.7 kDa, which is in discrepancy with experimentally observed Mr (~150 kDa). However, unless steps are taken in the protein isolation to disaggregate flagellins into monomers, it is common that flagellin proteins can exist in multimeric form (Simon et al., 2014), so it is plausible that flagellin trimer(s) would be detectable in the 150 kDa range. In *Yersinia*, T3SS apparatus protein YscT is linked with flagella formation (Ghosh, 2004). EscT is homologous to YscT, so it is not surprising that EscT would participate in *E. piscicida* flagellar formation.

Although *E. piscicida* C07–087 and T3SS mutants display close phenotypic characteristics, virulence comparison in channel catfish showed significant attenuation of *Ep ssaV* and *Ep yscR* compared to the parent strain. Mortalities for the fish challenged with strain C07–087 were lower than expected (26%) despite using a dose higher than our calculated LD₅₀, but both *Ep ssaV* and *Ep yscR* caused no mortalities. Moreover, bioluminescence imaging illustrated that *Ep ssaV* and *Ep yscR* were able to invade, establish infection, and then be cleared without causing mortalities. Low mortalities were observed in catfish injected with strains *Ep esaM* (6% mortality) and *Ep escT* (16% mortality) compared to parent strain C07–087 (26% mortality). However, bioluminescence imaging showed that these two mutations have significant impact on *E. piscicida* during earlier stages of infection, but by 24 and 36 h post-infection, these mutants had similar tissue quantities as the parent strain. Finally, our second challenge with parent strain C07–087 indicated that the four mutant strains provided significant protection against wild-type infection. Altogether, *Ep ssaV*, *Ep esaM*, and *Ep yscR* are safe and provided protection, suggesting they may be viable live attenuated vaccine candidates.

In conclusion, four T3SS mutants (*Ep ssaV*, *Ep esaM*, *Ep yscR*, and *Ep escT*) were successfully constructed by homologous recombination and allelic exchange, illustrating that the methods used for deletion mutation of *E. ictaluri* are effective in *E. piscicida*.

Phenotypic comparisons showed some minor differences in the T3SS mutants in H₂S production, hemolysin activity, and flagellin expression. Importantly, all four mutants provided significant protection against infection with the virulent strain C07–087. Strains *Ep ssaV* and *Ep yscR* were the safest and caused no vaccination mortalities, suggesting they may have potential as live attenuated vaccines.

Acknowledgements

This work was supported by the Egyptian Government, by the Mississippi State University College of Veterinary Medicine, and by INBRE-NIH award 4P20GM103476–15. The proteomics-mass spectrometry analysis was performed at the Institute for Genomics, Biocomputing and Biotechnology, Mississippi State University, with partial support by the Mississippi Agricultural and Forestry Experiment Station. Bioluminescence imaging work was partially supported by USDA-ARS Biophotonics Initiative #58–6402-3–018. We also thank Liesel Schneider and Min Wang for providing statistical analysis.

REFERENCES

- Abayneh T , Colquhoun DJ , Sorum H , 2013 *Edwardsiella piscicida* sp. nov., a novel species pathogenic to fish. *Journal of applied microbiology* 114, 644–654.23167785
- Abdelhamed H , Lu J , Shaheen A , Abbass A , Lawrence ML , Karsi A , 2013 Construction and evaluation of an *Edwardsiella ictaluri* fhuC mutant. *Vet Microbiol* 162, 858–865.23201245
- Blocker AJ , Deane JE , Veenendaal AK , Roversi P , Hodgkinson JL , Johnson S , Lea SM , 2008 What's the point of the type III secretion system needle? *Proceedings of the National Academy of Sciences of the United States of America* 105, 6507–6513.18458349
- Dahal N , Abdelhamed H , Lu J , Karsi A , Lawrence ML , 2013 Tricarboxylic acid cycle and one-carbon metabolism pathways are important in *Edwardsiella ictaluri* virulence. *PloS one* 8, e65973.23762452
- Dozois CM , Daigle F , Curtiss R , 2003 Identification of pathogen-specific and conserved genes expressed in vivo by an avian pathogenic *Escherichia coli* strain. *Proceedings of the National Academy of Sciences of the United States of America* 100, 247–252.12506201
- Figueira R , Holden DW , 2012 Functions of the *Salmonella* pathogenicity island 2 (SPI-2) type III secretion system effectors. *Microbiology* 158, 1147–1161.22422755
- Galán JE , Lara-Tejero M , Marlovits TC , Wagner S , 2014 Bacterial type III secretion systems: specialized nanomachines for protein delivery into target cells. *Annual review of microbiology* 68, 415.
- Ghosh P , 2004 *Process of Protein Transport by the Type III Secretion System*, Vol 68 American Society for Microbiology.
- Griffin MJ , Ware C , Quiniou SM , Steadman JM , Gaunt PS , Khoo LH , Soto E , 2014 *Edwardsiella piscicida* identified in the Southeastern USA by *gyrB* sequence, species-specific and repetitive sequence-mediated PCR. *Diseases of aquatic organisms* 108, 23–35.24492051
- Hensel M , Hinsley AP , Nikolaus T , Sawers G , Berks BC , 1999 The genetic basis of tetrathionate respiration in *Salmonella typhimurium*. *Mol Microbiol* 32, 275–287.10231485
- Herrero M , de Lorenzo V , Timmis KN , 1990 Transposon vectors containing non-antibiotic resistance selection markers for cloning and stable chromosomal insertion of foreign genes in gram-negative bacteria. *Journal of bacteriology* 172, 6557–6567.2172216
- Heuer OE , Kruse H , Grave K , Collignon P , Karunasagar I , Angulo FJ , 2009 Human Health Consequences of Use of Antimicrobial Agents in Aquaculture. *Clinical Infectious Diseases* 49, 1248–1253.19772389
- Hueck CJ , 1998 Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiology and molecular biology reviews* : MMBR 62, 379–433.9618447
- Karsi A , Lawrence ML , 2007 Broad host range fluorescence and bioluminescence expression vectors for Gram-negative bacteria. *Plasmid* 57, 286–295.17207855

- Karsi A , Menanteau-Ledouble S , Lawrence ML , 2006 Development of bioluminescent *Edwardsiella ictaluri* for noninvasive disease monitoring. *FEMS microbiology letters* 260, 216–223.16842347
- Katharios P , Kokkari C , Dourala N , Smyrli M , 2015 First report of *Edwardsiellosis* in cage-cultured sharpnose sea bream, *Diplodus puntazzo* from the Mediterranean. *BMC veterinary research* 11, 155.26193880
- Lan MZ , Peng X , Xiang MY , Xia ZY , Bo W , Jie L , Li XY , Jun ZP , 2007 Construction and characterization of a live, attenuated *esrB* mutant of *Edwardsiella tarda* and its potential as a vaccine against the haemorrhagic septicaemia in turbot, *Scophthalmus maximus* (L.). *Fish & shellfish immunology* 23, 521–530.17478097
- Liu W , Dong N , Zhang XH , 2012 Overexpression of *mltA* in *Edwardsiella tarda* reduces resistance to antibiotics and enhances lethality in zebra fish. *Journal of applied microbiology* 112, 1075–1085.22443589
- Lo DY , Lee YJ , Wang JH , Kuo HC , 2014 Antimicrobial susceptibility and genetic characterisation of oxytetracycline-resistant *Edwardsiella tarda* isolated from diseased eels. *The Veterinary record* 175, 203.24958553
- Marsden MJ , Vaughan LM , Foster TJ , Secombes CJ , 1996 A live (*delta aroA*) *Aeromonas salmonicida* vaccine for furunculosis preferentially stimulates T-cell responses relative to B-cell responses in rainbow trout (*Oncorhynchus mykiss*). *Infection and immunity* 64, 3863–3869.8751940
- Metcalf WW , Jiang W , Wanner BL , 1994 Use of the rep technique for allele replacement to construct new *Escherichia coli* hosts for maintenance of R6K gamma origin plasmids at different copy numbers. *Gene* 138, 1–7.8125283
- Miller VL , Mekalanos JJ , 1988 A novel suicide vector and its use in construction of insertion mutations: osmoregulation of outer membrane proteins and virulence determinants in *Vibrio cholerae* requires *toxR*. *Journal of bacteriology* 170, 2575–2583.2836362
- Nho SW , Hikima J , Cha IS , Park SB , Jang HB , del Castillo CS , Kondo H , Hirono I , Aoki T , Jung TS , 2011 Complete genome sequence and immunoproteomic analyses of the bacterial fish pathogen *Streptococcus parauberis*. *Journal of bacteriology* 193, 3356–3366.21531805
- Notti RQ , Stebbins CE , 2016 The Structure and Function of Type III Secretion Systems. *Microbiology spectrum* 4, 10.1128/microbiolspec.VMBF-0004–2015.
- Park SB , Aoki T , Jung TS , 2012 Pathogenesis of and strategies for preventing *Edwardsiella tarda* infection in fish. *Veterinary research* 43, 67.23035843
- Reed LJ , Muench H , 1938 A simple method of estimating fifty per cent endpoints. *American journal of epidemiology* 27, 493–497.
- Reichley SR , Ware C , Greenway TE , Wise DJ , Griffin MJ , 2015 Real-time polymerase chain reaction assays for the detection and quantification of *Edwardsiella tarda*, *Edwardsiella piscicida*, and *Edwardsiella piscicida*-like species in catfish tissues and pond water. *Journal of veterinary diagnostic investigation : official publication of the American Association of Veterinary Laboratory Diagnosticians, Inc* 27, 130–139.
- Santander J , Mitra A , Curtiss R , 2011 Phenotype, virulence and immunogenicity of *Edwardsiella ictaluri* cyclic adenosine 3',5'-monophosphate receptor protein (Crp) mutants in catfish host. *Fish & shellfish immunology* 31, 1142–1153.22015784
- Shafiei S , Viljamaa-Dirks S , Sundell K , Heinikainen S , Abayneh T , Wiklund T , 2016 Recovery of *Edwardsiella piscicida* from farmed whitefish, *Coregonus lavaretus* (L.), in Finland. *Aquaculture* 454, 19–26.
- Simon R , Curtis B , Deumic V , Nicki J , Tennant SM , Pasetti MF , Lees A , Wills PW , Chacon M , Levine MM , 2014 A scalable method for biochemical purification of *Salmonella* flagellin. *Protein expression and purification* 102, 1–7.25050462
- Sun W , Liu L , Bent AF , 2011 Type III secretion-dependent host defence elicitation and type III secretion-independent growth within leaves by *Xanthomonas campestris* pv. *campestris*. *Molecular Plant Pathology* 12, 731–745.21726374
- Sundell K , Wiklund T , 2011 Effect of biofilm formation on antimicrobial tolerance of *Flavobacterium psychrophilum*. *J Fish Dis* 34, 373–383.21488905

- Takac T , Vadovic P , Pechan T , Luptovciak I , Samajova O , Samaj J , 2016 Comparative proteomic study of *Arabidopsis* mutants mpk4 and mpk6. Scientific reports 6, 28306.27324189
- Tan YP , Lin Q , Wang XH , Joshi S , Hew CL , Leung KY , 2002 Comparative proteomic analysis of extracellular proteins of *Edwardsiella tarda*. Infection and immunity 70, 6475–6480.12379732
- Tan YP , Zheng J , Tung SL , Rosenshine I , Leung KY , 2005 Role of type III secretion in *Edwardsiella tarda* virulence. Microbiology 151, 2301–2313.16000720
- Tekedar HC , Karsi A , Williams ML , Vamenta S , Banes MM , Duke M , Scheffler BE , Lawrence ML , 2013 Complete Genome Sequence of Channel Catfish Gastrointestinal Septicemia Isolate *Edwardsiella tarda* C07–087. Genome announcements 1.
- Temprano A , Riaño J , Yugueros J , González P , de Castro L , Villena A , Luengo JM , Naharro G , 2005 Potential use of a *Yersinia ruckeri* O1 auxotrophic *aroA* mutant as a live attenuated vaccine. Journal of Fish Diseases 28, 419–427.16083447
- Tran CS , Rangel SM , Almblad H , Kierbel A , Givskov M , Tolker-Nielsen T , Hauser AR , Engel JN , 2014 The *Pseudomonas aeruginosa* Type III Translocon Is Required for Biofilm Formation at the Epithelial Barrier. PLoS Pathog 10, e1004479.25375398
- Wakimoto N , Nishi J , Sheikh J , Nataro JP , Sarantuya J , Iwashita M , Manago K , Tokuda K , Yoshinaga M , Kawano Y , 2004 Quantitative biofilm assay using a microtiter plate to screen for enteroaggregative *Escherichia coli*. The American journal of tropical medicine and hygiene 71, 687–690.15569806
- Wang B , Mo ZL , Xiao P , Li J , Zou YX , Hao B , Li GY , 2010a. EseD, a putative T3SS translocon component of *Edwardsiella tarda*, contributes to virulence in fish and is a candidate for vaccine development. Marine biotechnology 12, 678–685.20072793
- Wang Q , Yang M , Xiao J , Wu H , Wang X , Lv Y , Xu L , Zheng H , Wang S , Zhao G , Liu Q , Zhang Y , 2009 Genome sequence of the versatile fish pathogen *Edwardsiella tarda* provides insights into its adaptation to broad host ranges and intracellular niches. PloS one 4, e7646.19865481
- Wang X , Wang Q , Xiao J , Liu Q , Wu H , Zhang Y , 2010b. Hemolysin EthA in *Edwardsiella tarda* is essential for fish invasion in vivo and in vitro and regulated by two-component system EsrA-EsrB and nucleoid protein HhaEt. Fish & shellfish immunology 29, 1082–1091.20832475
- Watnick P , Kolter R , 2000 Biofilm, city of microbes. Journal of bacteriology 182, 2675–2679.10781532
- Watson JJ , White FH , 1979 Hemolysins of *Edwardsiella tarda*. Canadian journal of comparative medicine : Revue canadienne de medecine comparee 43, 78–83.34473
- Xie HX , Lu JF , Rolhion N , Holden DW , Nie P , Zhou Y , Yu XJ , 2014 *Edwardsiella tarda*-Induced cytotoxicity depends on its type III secretion system and flagellin. Infection and immunity 82, 3436–3445.24891103
- Xu L , Wang Q , Xiao J , Liu Q , Wang X , Chen T , Zhang Y , 2010 Characterization of *Edwardsiella tarda* waaL: roles in lipopolysaccharide biosynthesis, stress adaptation, and virulence toward fish. Archives of microbiology 192, 1039–1047.20938646

Figure 1.

Genotypic confirmation of *E. piscicida* mutant strains by PCR using A and D primer pair.

Lane 1: 1KB plus DNA ladder; Lane 2: 4081 bp *E. piscicida* wild type strain *ssaV* gene;

Lane 3: 1922 bp *EP ssaV* mutant; Lane 4: 2198 bp *E. piscicida* wild type strain *esaM* gene;

Lane 5: 1865 bp *Ep esaM* mutant; Lane 6: 2426 bp *E. piscicida* wild type strain *yscR* gene;

Lane 7: 1748 bp *Ep yscR* mutant; Lane 8: 2685 bp *E. piscicida* wild type strain *escT* gene;

Lane 9: 1872 bp *Ep escT* mutant

Figure 2.

Mean biofilm formation of *E. piscicida* C07-087 and mutant strains as measured by absorbance at 538 nm to detect crystal violet staining of biofilm. The data represent means of five replicates plus or minus standard error. *E. piscicida* C07-087 = parent strain, and C-ve = negative control (wells with no bacteria). All strains had significant biofilm formation at 72 h compared to the negative control.

Figure 3. Percent hemolysis of *E. piscicida* C07-087 and mutant strains. Error bars indicate standard deviation for three triplicate samples. * indicates statistically significant difference compared to parent strain C07-087 ($P < 0.05$).

Figure 4. SDS PAGE analysis of ECP of *E. piscicida* strain C07-087 and T3SS mutants. Lane 1: Molecular weight marker; Lane 2: *E. piscicida* C07-087; Lane 3: *Ep escT*; Lane 4: *Ep esaM*; Lane 5: *Ep yscR*; Lane 6: *Ep ssaV*. The arrow refers to the approximately 150 kDa band missing in *Ep escT* and *Ep esaM*.

(A)

(B)

Figure 5.

Bioluminescent imaging of live catfish after IP injection. A, BLI of catfish. B, mean total photon emissions at each time point (S = second; Sr = steradian). *indicates significant difference between *E. piscicida* C07-087 and mutant strains.

Figure 6.

(A) Percent mortalities of catfish fingerlings challenged with *E. piscicida* C07-087 and T3SS mutant strains. (B) Percent mortalities of channel catfish fingerlings vaccinated with the T3SS mutants and challenged with wild-type strain C07-087 at 21 d post-vaccination. Data are presented as means $\pm$ SE. *indicates significant difference compared to parent strain C07-087.

Table 1.**Bacterial strains and plasmids**

Strain or plasmid	Description	Reference
<i>E. piscicida</i>		
C07-087	Isolate from diseased catfish	(Tekedar et al., 2013)
<i>Ep ssaV</i>	<i>E. piscicida</i> ; <i>ssaV</i>	This study
<i>Ep esaM</i>	<i>E. piscicida</i> ; <i>esaM</i>	This study
<i>Ep yscR</i>	<i>E. piscicida</i> ; <i>yscR</i>	This study
<i>Ep escT</i>	<i>E. piscicida</i> ; <i>escT</i>	This study
<i>Escherichia coli</i>		
CC118 λ pir	(<i>ara-leu</i>); <i>araD</i> ; <i>lacX74</i> ; <i>galE</i> ; <i>galK</i> ; <i>phoA20</i> ; <i>thi-I</i> ; <i>rpsE</i> ; <i>rpoB</i> ; <i>argB</i> (Am); <i>recA</i> ; λ pirR6K	(Herrero et al., 1990)
SM10 λ pir	<i>thi</i> ; <i>thr</i> ; <i>leu</i> ; <i>tonA</i> ; <i>lacY</i> ; <i>supE</i> ; <i>recA</i> ; <i>::RP4-2-Tc::Mu</i> ; <i>Kmr</i> ; <i>lpirR6K</i>	(Miller and Mekalanos, 1988)
BW19851	<i>RP4-2 (Km::Tn7, Tc::Mu-1)</i> , <i>uidA3::pir⁺</i> , <i>recA1</i> , <i>endA1</i> , <i>thi-1</i> , <i>hsdR17</i> , <i>creC510</i>	(Metcalf et al., 1994)
Plasmids		
pMEG-375	8142 bp, <i>sacRB</i> <i>mobRP4</i> or <i>R6K</i> . Cm ^R , Ap ^R	(Dozois et al., 2003)
pEp <i>ssaV</i>	<i>ssaV</i> , pMEG-375	This study
pEp <i>esaM</i>	<i>esaM</i> , pMEG-375	This study
pEp <i>yscR</i>	<i>yscR</i> , pMEG-375	This study
pEp <i>escT</i>	<i>escT</i> , pMEG-375	This study
pAK <i>gfpluxI</i>	pBBR1MCS4 with <i>gfpmut3</i> and <i>luxCDABE</i>	(Karsi and Lawrence, 2007)

Table 2.

Primers used to generate and verify in-frame deletion of the *E. piscicida* T3SS apparatus genes.

Primer ID		Sequence (5' - 3') ^b	RE ^a
EpssaVF01	A	AAAGAGCTC ACGGTATGGGTCGAGCGTAT	<i>SacI</i>
EpssaVR87	B	CATGAAAACGGCCAGCAAC	
EpssaVF225	C	<u>GTTGCTGGCCGTTTT</u> CATGAGGCGTTGCTCTCACCTAC	
EpssaVR01	D	AAATCTAGA ATCCTGTCCCGCTCGGTAT	<i>XbaI</i>
EpssaVF01S		AGCAGCCTGACCCTGGAC	
EpesaMF01	A	AAAGAGCTC ACGAGAGGATCAGGACCAG	<i>SacI</i>
EpesaMR	B	GCCCATCAGTCGTAGCCAAAG	
EpesaMF	C	<u>CTTTGGCTACGACTGATGGGCTAGCTGGCTACACAA</u> ACTCACC	
EpesaMR01	D	AAATCTAG ACCGATGAAGATCAGGGTAGG	<i>XbaI</i>
EpesaMF01S		CGGTAACGAGAGGATCAGGA	
EpyscRF01	A	AAAGAGCTCCAGAGCCTTTGGTAATCACG	<i>SacI</i>
EpyscRR48	B	AGTATCTGCCGACAGTGTTC	
EpyscRF48	C	<u>GGAACACTGTGCGCAGATACTGCTATGGGAAA</u> GCACAGCTC	
EpyscRR01	D	AAATCTAG ATACAGCAGGTGCCATAGGTG	<i>XbaI</i>
EpyscRF01S		CTGAAATCCACAGCGCATC	
EpecTF01	A	AAAGAGCTCGTGGAAGTGGGGAACACTG	<i>SacI</i>
EpecTR	B	CTTCCGGCATAGCTAAACAGC	
EpecTF33	C	<u>GCTGTTTAGCTATGCCGGAAGTTGAGCGAGAAA</u> ACGGAACAG	
EpecTR01	D	AAATCTAG ACTGCAGCGATGACTTGGTCT	<i>XbaI</i>
EpecTF01S		GCGATGACTCGCTGCTATTAC	

^aRE: restriction endonuclease site added to the 5' end of the primer sequence.

^bBold letters at the 5' end of the primer sequence represent RE site. AA nucleotides were added to the end of each primer containing a RE site to increase the efficiency of enzyme digestion. Underlined bases indicate reverse complemented primer sequence for overlap extension PCR.

Table 3. Properties of the selected *E. piscicida* T3SS apparatus proteins.

Protein	Locus tag	GI	ORF (bp/aa)
Secretion system apparatus protein SsaV	ETAC_04215	469762556	2058/685
EsaM	ETAC_04220	469762557	336/111
Type III secretion system protein YscR	ETAC_04225	469762558	648/215
Type III secretion apparatus protein EscT	ETAC_04235	469762560	783/260

Table 4.

Extracellular proteins of *Edwardsiella piscicida* identified in 150 kDa region of SDS-PAGE. Two strains (C07–087 and *Ep escT*) are compared. (–) indicated absences of protein in *Ep escT* mutant strain. (aa) is Total amino acids in protein. (pI) is the isoelectric point of the protein.

Protein Characterization					C07–087		<i>Ep escT</i>	
Accession number (gi)	Proteins name	MW [kDa]	pI	aa	Peptides	Coverage	Peptides	Coverage
505274692	Flagellar biosynthesis protein	43.5	5.21	417	55	56.59	2	8.39
505274691	Flagellar biosynthesis protein	43.6	5.30	419	55	56.32	2	2.24
779983474	Flagellin	43.5	5.22	416	48	43.03	-	-
502611981	MULTISPECIES: flagellin	43.7	5.22	416	43	43.51	-	-
779991656	DNA-directed RNA polymerase subunit beta	150.6	5.26	1342	2	1.86	-	-
Total peptides					203		4	