Oligodendrogliomas: a short history of clinical developments

“This special focus issue will give a detailed overview of the current understanding, diagnosis and treatment of oligodendroglial tumors.”

Martin J van den Bent*

Oligodendroglioma: the early days

Oligodendrogliomas have in particular attracted attention once it became clear that these are chemotherapy sensitive tumors. Although since the first description in 1926 and 1929 of oligodendroglial tumors it has been clear that these tumors are a special brand with a relatively good prognosis, they were taken for just another glioma subtype for many years thereafter [1]. Indeed, apart from a relatively more favorable prognosis of oligodendroglioma in comparison to their astrocytic counterparts and their often calcified lesions allowing a first diagnosis even at the time CT scans were not yet available, little pointed toward a specific significance for these tumors. This histological identification was somewhat complicated by the existence of another group with histological characteristics of both astrocytoma and oligodendroglioma, labeled mixed oligoastrocytoma. All these tumors were considered among the grade II and III glioma, and trials on grade II and high-grade glioma usually considered both astrocytoma, oligodendroglioma and mixed oligoastrocytoma. Furthermore, trials on high-grade glioma usually lumped both grade III and grade IV (glioblastoma) under the heading ‘anaplastic glioma’. As a consequence, grade III tumors were in general treated as grade IV tumors, albeit with the clear notion their outcome was better.

Oligodendroglioma: the chemotherapy sensitive glioma

This clinical significance of the diagnosis ‘oligodendroglioma’ radically changed when Cairncross *et al.* in 1988 and again in 1992 reported remarkably high response rates to procarbazine, CCNU (lomustine) and vincristine (PCV) chemotherapy in relapsing anaplastic oligodendroglioma [2]. This led to confirmatory Phase II studies, and subsequently to studies in newly diagnosed anaplastic oligodendroglial tumors [3,4]. At that time, still no distinction was made between pure oligodendroglial tumors and mixed oligodendroglioma. At least in part driven by the wish not to miss oligodendroglial tumors the percentage of patients with oligodendroglioma showed a steep incline, from 5% of all glioma in historical series to no less than 20% of all glioma cases [5]. A clear change in diagnostic criteria was part of that increase, already pointing to the lack of objective criteria for the histological diagnosis of glioma in general but more in particular of oligodendroglial tumors.

Oligodendroglioma: the molecular tumor

From the early nineties on, the advance in molecular techniques began to make contribution to the understanding of oligodendroglioma. In 1994, Reifenberger *et al.* described combined loss of the entire short arm of chromosome 1 and the long arm

KEYWORDS

- anaplastic • IDH
- oligoastrocytoma
- oligodendroglioma
- 1P/19q codeletion
- PCV • temozolomide

“From the early nineties on, the advance in molecular techniques began to make contribution to the understanding of oligodendroglioma.”

*Brain Tumor Center at Erasmus MC Cancer Center, Erasmus MC, Rotterdam, The Netherlands; m.vandenbent@erasmusmc.nl

of chromosome 19 (combined 1p/19q loss) as the most characteristic genetic lesion in oligodendroglioma [6]. In an important next paper, Mainz *et al.* showed that low-grade mixed oligoastrocytoma show either characteristic astrocytic lesions (TP53 mutations) or combined loss of 1p/19q, pointing out that at the molecular level mixed tumors do not exist [7]. Subsequent studies showed that anaplastic oligodendrogliomas showed in addition to 1p/19q loss other lesions (e.g., loss of 9p), but that typical genetic lesions found in glioblastoma were mutually exclusive with combined 1p/19q loss [8,9]. This pointed toward a glioblastoma-like genetic background of many anaplastic mixed oligoastrocytomas, further emphasizing that true mixed oligoastrocytic tumors do not exist.

Oligodendroglioma: 1p/19q loss & chemosensitivity

A second major breakthrough in this field was the observation by Louis and Cairncross that loss of 1p/19q was correlated to a 100% response rate to PCV chemotherapy, as opposed to 20–25% response rate in the absence of 1p/19q loss [10]. Further studies noted that typical oligodendroglial histological features were associated with a greater likelihood of 1p/19q loss, despite the occurrence of 1p/19q loss in tumors with atypical oligodendroglial or even pure astrocytic features [11]. Several series however showed that 1p/19q status was more predictive of outcome to chemotherapy than morphology. This resulted in tendency toward a more restrictive diagnosis of oligodendroglial features, and in many studies into the nature of the sensitivity of chemotherapy of 1p/19q co-deleted oligodendrogliomas [12]. These studies explored in particular the presence of other lesions on 1p and 19q, including epigenetic alterations on these chromosomes. A new findings coming out of these studies was the simultaneous discovery by Burger and Jenkins that the combined loss of 1p/19q in fact represents a balanced translocation, after which a copy of 1p and 19q is lost [13,14]. This translocation has so far not been observed in any other tumor type, and as opposed to other translocations the centromeric fusion of 1q and 19p does not result in an activated oncogene.

Oligodendrogliomas: initial results of the large prospective studies

These genetic studies allowed molecular analysis of the large prospective Phase III trials conducted

by RTOG and EORTC that explored the role of (neo)adjuvant PCV chemotherapy in anaplastic oligodendroglial tumors. These trials, initiated in the mid-nineties were reported for the first time in 2004 and 2005, and although initially failed to show survival benefit from the addition of PCV to radiotherapy they already showed the major prognostic significance of combined 1p/19q loss [15,16]. Side studies showed that as anticipated ~30% of included tumors did not have an oligodendroglial genetic signature, but evidence of alterations commonly observed in glioblastoma [17]. Moreover, a clear interobserver variation with regard to the histopathological diagnosis was present on both studies, even with respect for what is to be called a typical oligodendroglioma [18,19].

Oligodendroglioma: temozolomide

A major question that rose after the pivotal EORTC trial on temozolomide chemoradiotherapy in glioblastoma and the trials on PCV in newly anaplastic oligodendroglial tumors, was why temozolomide improved outcome in glioblastoma, a chemoresistant disease but PCV failed to improve outcome in a chemotherapy sensitive tumor [20]. One possibility was the different administration of chemotherapy in concurrent chemoradiotherapy with temozolomide that had been used in the glioblastoma trial. Here, chemotherapy was combined on a daily basis during the entire radiotherapy phase, which was not the case in the classical (neo)adjuvant PCV studies.

Oligodendroglioma: second-generation trials

In the aftermath of the initial results of EORTC 26951 and RTOG 9402, follow-up studies were designed that no longer embarked on histology *per se*, but on the presence or absence of combined 1p/19q loss in grade III tumors and mark the transition of trials based on histology to trials based on molecular signature. These trials, CATNON and CODEL aimed to define the role of temozolomide in grade III tumors, a question that rose from the outcome of the EORTC trial on chemoradiotherapy with temozolomide in glioblastoma. Similar trials were initiated on low-grade tumors trying to define the role of chemotherapy in low-grade tumors.

Oligodendroglioma: the role of MGMT, IDH & methylation

The observations of the role of MGMT in sensitivity of glioblastoma to temozolomide resulted

in studies of the significance of *MGMT* promoter methylation in the European study on anaplastic oligodendroglioma, but also in the German study ('NOA4') that compared radiotherapy to chemotherapy in anaplastic glioma [21,22]. Surprisingly, both studies found that *MGMT* promoter methylation was correlated with a improved outcome after radiotherapy alone, for which no clear biological explanation was present [22,23]. These and other studies also reported high *MGMT* promoter methylation rates (80% range) in grade II and grade III glioma [24]. This unexpected prognostic finding after radiotherapy alone was sort of resolved with the discovery of *IDH* mutations, which were found in 60–80% of all grade II and III glioma, and in up to 100% in 1p/19q co-deleted tumors in some studies. Not only were *IDH* mutations found to be prognostic, they also strongly correlated to the presence of *MGMT* promoter methylation. First of all, this revealed that the significance of *MGMT* promoter methylation in glioblastoma differed from that in grade III glioma. Genomic wide methylation studies subsequently showed that a CpG Island Methylated Phenotype also exists in glioma, similar to as observed in colon cancer. *In vivo* models have suggested that the metabolic alterations present in *IDH*-mutated cells may induce a hypermethylated state, thus leading to an explanation of some of the findings in the clinical studies. At present, this CpG Island Methylated Phenotype phenotype appears to be a very strong prognostic factor for outcome in grade III tumors, and perhaps predictive for benefit to adjuvant chemotherapy.

Oligodendroglioma: long-term follow-up of the clinical trials

The understanding of the impact of 1p/19q co-deletion was impacted again by the reports in 2012 of the long-term follow-up of the EORTC and RTOG studies on adjuvant PCV chemotherapy [25,26]. Surprisingly, after first negative overall survival results now both studies showed a clear survival benefit of the addition of PCV chemotherapy to radiotherapy in 1p/19q co-deleted tumors, with a separation of the survival curves 5–6 years after randomization. The similar observation in both studies give credit to this finding, even though it was made in subgroups both original studies. On top of this finding, the median survival of more than 12–14 years in 1p/19q co-deleted tumors following RT/PCV exceeds that of their anaplastic astrocytic

counterparts significantly and points to a very significant long-term survivorship leading to new clinical questions. Moreover, an EORTC trial on low-grade glioma now gives evidence that in 1p deleted low-grade glioma, progression-free survival after temozolomide is equal to that after radiotherapy, whereas in 1p intact tumors radiotherapy is superior (Baumert *et al.*, ASCO 2013, oral communication). This adds further to the notion molecular diagnostic are now part of clinical decision making in neuro-oncology.

Oligodendroglioma: next-generation sequencing

Very recently next-generation sequencing has identified novel mutations in 1p/19q co-deleted tumors: in the *CIC* and *FUBP* gene [27]. The clinical significance of these lesions is still unclear, and their functional impact remains speculative. Interest in telomerase activity has also been renewed in neuro-oncology, studies have shown that diffuse glioma are characterized by mutually exclusive *TERT* and *ATRX* mutations [28,29]. Surprisingly, 1p/19q co-deleted tumors have the same abnormality as glioblastoma (*TERT* mutations) whereas diffuse astrocytoma regardless of grade are characterized by *ATRX* mutations. Based on these findings a new classification of diffuse gliomas based on molecular alterations is being considered.

Oligodendroglioma: unanswered questions

Many questions though remain unanswered. It is clear PCV chemotherapy added to radiotherapy improves outcome of 1p/19q co-deleted patients. It is unclear whether that also holds also true for temozolomide. In the past decade temozolomide has replaced PCV because of its better tolerability, but trials on the combination of RT and temozolomide in oligodendroglial tumors are simply unavailable. Several groups have reported on upfront treatment with chemotherapy alone (temozolomide, or combinational regimens with autologous bone marrow transplant) [30,31]. Lack of long-term follow-up and possible bias by patient selection make the interpretation of these series difficult. Although the ongoing CODEL study will give an answer to this question, this answer will not come in the near future. An important question is whether radiotherapy can be safely postponed, whether chemotherapy alone and radiotherapy at progression will be equally effective. If combinational

treatment is indeed making a survival difference this may adversely affect outcome, but delaying radiotherapy in favorable prognosis patients and thus avoiding radiotherapy induced sequelae on cognition may improve quality of survival. Again, no answer from currently available data exists.

On a more basic level, it is still unclear what induces oligodendroglial tumors, and what drives 1p/19q loss. We now know that *IDH* mutations are pivotal in diffuse glioma, but why some develop *TP53* mutations and others 1p/19q co-deletion remains a mystery. The answer to these key questions is hampered by the paucity of 1p/19q loss *in vitro* and *in vivo* models. Although these models do exist, in general these tumors are almost impossible to culture (which also holds for *IDH* mutated tumors). Absence of good models hinder the development of more effective treatments and a better understanding of oncogenesis of 1p/19q co-deleted tumors. We still do not know what other factors (inhibited tumor suppressor genes, oncogenes) drive these tumors. And, perhaps most important, we lack innovative agents. It is good we have a better understanding of the responsiveness to classical radiotherapy and chemotherapy of oligodendroglial tumors. But,

all these treatments were available in the mid-nineties, and since then no other treatment has increased our therapeutic arsenal.

This special focus issue will give a detailed overview of the current understanding, diagnosis and treatment of oligodendroglial tumors. The ins and outs of classical histology are not reviewed here, for that topic we refer our readers to the 2007 edition of the WHO text book on neuro-oncology [32]. Each article in this special issue is written by a specialist in the field, with a track record in that area. We are very happy that these colleagues have been willing to contribute to this special issue, which gives an overview of all aspects of oligodendroglioma and will present the standard of both understanding and clinical care.

Financial & competing interests disclosure

The author has no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

No writing assistance was utilized in the production of this manuscript.

References

- Bailey P, Bucy PC. Oligodendrogliomas of the brain. *J. Pathol.* 32, 735–751 (1929).
- Cairncross JG, Macdonald DR. Successful chemotherapy for recurrent oligodendroglioma. *Ann. Neurol.* 23, 360–364 (1988).
- Cairncross G, Macdonald D, Ludwin S *et al.* Chemotherapy for anaplastic oligodendroglioma. *J. Clin. Oncol.* 12, 2013–2021 (1994).
- van den Bent MJ, Kros JM, Heimans JJ *et al.* Response rate and prognostic factors of recurrent oligodendroglioma treated with procarbazine, CCNU and vincristine chemotherapy. *Neurology* 51, 1140–1145 (1998).
- Coons SW, Johnson PC, Scheithauer BW, Yates AJ, Pearl DK. Improving diagnostic accuracy and interobserver concordance in the classification and grading of primary gliomas. *Cancer* 79, 1381–1391 (1997).
- Reifenberger J, Reifenberger G, Liu L, James CD, Wechsler W, Collins VP. Molecular genetic analysis of oligodendroglial tumors shows preferential allelic deletions on 19q and 1p. *Am. J. Pathol.* 145, 1175–1190 (1994).
- Maintz D, Fiedler K, Koopmann J *et al.* Molecular genetic evidence for subtypes of oligoastrocytomas. *J. Neuropathol. Exp. Neurol.* 56, 1098–1104 (1997).
- Bigner SH, Matthews MR, Rasheed BKA *et al.* Molecular genetic aspects of oligodendrogliomas including analysis by comparative genomic hybridization. *Am. J. Pathol.* 155, 375–386 (1999).
- He J, Mokhtari K, Sanson M *et al.* Glioblastomas with an oligodendroglial component: a pathological and molecular study. *J. Neuropathol. Exp. Neurol.* 60, 863–871 (2001).
- Cairncross JG, Ueki K, Zlatescu MC *et al.* Specific genetic predictors of chemotherapeutic response and survival in patients with anaplastic oligodendrogliomas. *J. Natl. Cancer Inst.* 90, 1473–1479 (1998).
- Smith JS, Perry A, Borell TJ *et al.* Alterations of chromosome arms 1p and 19q as predictors of survival in oligodendrogliomas, astrocytoma, and mixed oligoastrocytomas. *J. Clin. Oncol.* 18, 636–645 (2000).
- Burger PC. What is an oligodendroglioma? *Brain Pathol.* 12, 257–259 (2002).
- Griffin CA, Burger P, Morsberger L *et al.* Identification of der(1;19)(q10;p10) in five oligodendrogliomas suggests mechanism of concurrent 1p and 19q loss. *J. Neuropathol. Exp. Neurol.* 65, 988–994 (2006).
- Jenkins RB, Blair H, Ballman KV *et al.* A t(1;19)(q10;p10) mediates the combined deletions of 1p and 19q and predicts a better prognosis of patients with oligodendroglioma. *Cancer Res.* 66, 9852–9861 (2006).
- Cairncross JG, Berkey B, Shaw E *et al.* Phase III trial of chemotherapy plus radiotherapy (RT) versus RT alone for pure and mixed anaplastic oligodendroglioma (RTOG 9402): an intergroup trial by the RTOG, NCCCTG, SWOG, NCI CTG and ECOG. *J. Clin. Oncol.* 24, 2707–2714 (2006).
- van den Bent MJ, Carpentier AF, Brandes AA *et al.* Adjuvant PCV improves progression free survival but not overall survival in newly diagnosed anaplastic oligodendrogliomas and oligoastrocytomas:

- a randomized EORTC Phase III trial. *J. Clin. Oncol.* 24, 2715–2722 (2006).
- 17 Kouwenhoven MC, Gorlia T, Kros JM *et al.* Molecular analysis of anaplastic oligodendroglial tumors in a prospective randomized study: a report from EORTC study 26951. *Neuro Oncol.* 11, 737–746 (2009).
- 18 Kros JM, Gorlia T, Kouwenhoven MC *et al.* Panel review of anaplastic oligodendroglioma from EORTC trial 26951: assessment of consensus in diagnosis, influence of 1p/19q loss and correlations with outcome. *J. Neuropathol. Exp. Neurol.* 66, 545–551 (2007).
- 19 Giannini C, Burger PC, Berkey BA *et al.* Anaplastic oligodendroglial tumors: refining the correlation among histopathology, 1p 19q deletion and clinical outcome in Intergroup Radiation Therapy Oncology Group Trial 9402. *Brain Pathol.* 18, 360–369 (2008).
- 20 Stupp R, Mason WP, van den Bent MJ *et al.* Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N. Engl. J. Med.* 352, 987–996 (2005).
- 21 Hegi ME, Diserens AC, Gorlia T *et al.* MGMT gene silencing and benefit from temozolomide in glioblastoma. *N. Engl. J. Med.* 352, 997–1003 (2005).
- 22 Wick W, Hartmann C, Engel C *et al.* NOA-04 randomized Phase III trial of sequential radiochemotherapy of anaplastic glioma with procarbazine, lomustine, and vincristine or temozolomide. *J. Clin. Oncol.* 27, 5874–5880 (2009).
- 23 van den Bent MJ, Dubbink HJ, Sanson M *et al.* MGMT promoter methylation is prognostic but not predictive for outcome to adjuvant PCV chemotherapy in anaplastic oligodendroglial tumors: a report from EORTC Brain Tumor Group Study 26951. *J. Clin. Oncol.* 27, 5881–5886 (2009).
- 24 Everhard S, Kaloshi G, Criniere E *et al.* MGMT methylation: a marker of response to temozolomide in low-grade gliomas. *Ann. Neurol.* 60, 740–743 (2006).
- 25 Cairncross G, Wang M, Shaw E *et al.* Phase III trial of chemoradiotherapy for anaplastic oligodendroglioma: long-term results of RTOG 9402. *J. Clin. Oncol.* 31(3), 337–343 (2013).
- 26 van den Bent MJ, Brandes AA, Taphoorn MJ *et al.* Adjuvant procarbazine, lomustine, and vincristine chemotherapy in newly diagnosed anaplastic oligodendroglioma: long-term follow-up of EORTC brain tumor group study 26951. *J. Clin. Oncol.* 31, 344–350 (2013).
- 27 Bettegowda C, Agrawal N, Jiao Y *et al.* Mutations in CIC and FUBP1 contribute to human oligodendroglioma. *Science* 333, 1453–1455 (2011).
- 28 Killela PJ, Reitman ZJ, Jiao Y *et al.* TERT promoter mutations occur frequently in gliomas and a subset of tumors derived from cells with low rates of self-renewal. *Proc. Natl Acad. Sci. USA* 110, 6021–6026 (2013).
- 29 Jiao Y, Killela PJ, Reitman ZJ *et al.* Frequent ATRX, CIC, and FUBP1 mutations refine the classification of malignant gliomas. *Oncotarget* 3, 709–722 (2012).
- 30 Abrey LE, Childs BH, Paleologos N *et al.* High-dose chemotherapy with stem cell rescue as initial therapy for anaplastic oligodendroglioma: long-term follow-up. *Neuro Oncol.* 8, 183–188 (2006).
- 31 Levin N, Lavon I, Zelikovitch B *et al.* Progressive low-grade oligodendrogliomas: response to temozolomide and correlation between genetic profile and O6-methylguanine DNA methyltransferase protein expression. *Cancer* 106, 1759–1765 (2006).
- 32 Louis DN, Ohgaki H, Wiestler OD, Cavenee WK. *WHO Classification of Tumours of the Central Nervous System*. WHO, Geneva, Switzerland (2007).