Published in final edited form as:

J Proteome Res. 2015 June 05; 14(6): 2569–2574. doi:10.1021/acs.jproteome.5b00125.

Metabonomics Reveals Metabolite Changes in Biliary Atresia Infants

Kejun Zhou^{1,2}, Guoxiang Xie^{3,4}, Jun Wang¹, Aihua Zhao³, Jiajian Liu³, Mingming Su⁴, Yan Ni⁴, Ying Zhou¹, Weihua Pan¹, Yanran Che⁵, Ting Zhang⁵, Yongtao Xiao², Yang Wang², Jie Wen², Wei Jia^{3,4,*}, and Wei Cai^{1,2,*}

¹Department of Pediatric Surgery, Xin Hua Hospital, School of Medicine, Shanghai Jiao Tong University (SJTU), Shanghai 200092, China

²Shanghai Key Laboratory of Pediatric Gastroenterology and Nutrition, Shanghai Institute for Pediatric Research, Shanghai 200092, China

³Center for Translational Medicine, Six People's Hospital, SJTU School of Medicine, Shanghai 200230, China

⁴University of Hawaii Cancer Center, Honolulu, HI 96813, United States of America

⁵Department of Infection and Gastroenterology, Shanghai Children's Hospital, SJTU School of Medicine, Shanghai 200040, China

Abstract

Biliary atresia (BA) is a rare neonatal cholestatic disorder caused by obstruction of extra and intra hepatic bile ducts. If untreated, progressive liver cirrhosis will lead to death within two years. Early diagnosis and operation improve the outcome significantly. Infants with neonatal hepatitis syndrome (NHS) present similar symptoms, confounding the early diagnosis of BA. The lack of non-invasive diagnostic methods to differentiate BA from NHS greatly delays the surgery of BA infants, thus deteriorating the outcome. Here, we performed a metabolomics study in plasma of BA, NHS, and healthy infants using gas chromatography-time-of-flight mass spectrometry. Scores plots of orthogonal partial least square discriminant analysis clearly separated BA from NHS and healthy infants. Eighteen metabolites were found to be differentially expressed between BA and NHS, among which seven (L-glutamic acid, L-ornithine, L-isoleucine, L-lysine, L-valine, L-tryptophan, and L-serine) were amino acids. The altered amino acids were quantitatively verified using ultra-performance liquid chromatography-tandem mass spectrometry. Ingenuity pathway analysis revealed the network of "Cellular Function and Maintenance, Hepatic System Development and Function, Neurological Disease" was altered most significantly. This study suggests that plasma metabolic profiling has great potential in differentiating BA from NHS, and amino acid metabolism is significantly different between the two diseases.

Table of Contents Synopsis

*Corresponding author: Wei Cai. 1665 Kong Jiang Road, Shanghai, China. Tel & Fax: 86-21-65791316. caiw1978@163.com, Wei Jia. 701 Ilalo Street, Honolulu, Hawaii, USA. Tel & Fax: 01-808-5645823. wjia@cc.hawaii.edu.

biliary atresia; neonatal hepatitis syndrome; metabonomics; amino acid

Biliary atresia (BA) is the most common cause of chronic cholestasis in neonates which derived from a fibro-inflammatory destruction of extra or intra hepatic bile ducts ^{1, 2}. The incidence ranges from 1 in 5,000 to 1 in 20,000 living births in different areas of the world. It constitutes nearly one-third of all neonatal cholestasis or more than 90% obstructive cholestatic cases ³. If untreated, children will die by age 2 years because of progressive liver cirrhosis.

Neonatal hepatitis syndrome (NHS) is another leading cause of neonatal cholestasis and there is great overlap in the clinical, biochemical, imaging, and histopathological characteristics between NHS and BA ³⁻⁵. BA should be differentiated from NHS as soon as possible, since early Kasai operations greatly improve the outcome of BA infants, while NHS infants don't need surgery and most of which will usually return to normal by one year of age. However, discrimination of BA from NHS is challenging due to the lack of non-invasive diagnostic methods. Currently, clinical determination of BA in an infant with cholestasis usually requires liver biopsy or operative cholangiography. Patients may undergo a wide range of tests before BA or an alternative diagnosis is reached, which delays the time of Kasai operation in BA patients.

Recently, great effort has been made to search for new BA biomarkers. Some serum metabolites or proteins, including hyaluronic acid ⁶, Apolipoprotein C3 (APOC3) ⁷, and interleukin 6 (IL-6) and IL-8 ⁸, are reported altered in BA. Circulating levels of the miR-200b/429 cluster were found elevated in BA infants compared to cholestatic controls ⁹. However, no non-invasive biomarker for BA has been discovered so far. Thus, there is an urgent need for blood biomarkers for BA.

Metabonomics studies the comprehensive small-molecular-weight metabolites and their dynamic changes in biological systems. Mass spectrometry-based metabolic profiling is increasingly used to uncover new biomarkers for diagnosis^{10, 11}, prognosis^{12, 13}, pathogenesis clarification¹⁴, and as potential therapeutic targets for clinical treatment¹⁵. However, no comprehensive metabolic profiling of BA was reported yet. In this study, we

conducted a metabolomics study of 45 BA, 15 NHS, and 6 healthy infants using gas chromatography-time-of-flight-mass spectrometry (GC-TOFMS) to discover the different plasma metabolic compositions. Then, differentially expressed metabolites were verified in plasma of 53 BA and 28 NHS infants using a targeted metabolomics method with ultra-performance liquid chromatography-tandem mass spectrometry (UPLC-MS/MS).

MATERIALS AND METHODS

Chemicals and Reagents

HPLC grade acetonitrile, methanol, and formic acid were purchased from Merck Chemicals (Darmstadt, Germany). Chloroform, pyridine, anhydrous sodium sulfate, BSTFA (1% TMCS), heptadecanoic acid, methoxyamine, and L-2-chlorophenylalanine were purchased from Sigma-Aldrich (St. Louis, MO). Absolute IDQ P180 kit was purchased from Biocrates Life Sciences (Innsbruck, AT) for targeted quantification of amino acids and biogenic amines.

Clinical Samples

Plasma samples were obtained from 45 BA (before operative cholangiography), 15 NHS, and 6 healthy infants for GC-TOFMS analysis at Xin Hua Hospital, Shanghai, China, from Jul 2011 to Jul 2013. Additional 8 BA (before operative cholangiography) and 13 NHS plasma samples were collected from Aug 2013 to May 2014 in the same hospital. Combined with former samples, a total of 53 BA and 28 NHS was used for validation of amino acids expression. All infants with BA were correctly identified based on operative cholangiography and liver pathology. Cholestasis induced by citrin deficiency, α 1-antitrypsin deficiency, Alagille's syndrome, progressive familial intrahepatic cholestasis, parenteral nutrition, and Choledochal cysts were excluded in this study. Plasma samples were kept at -80°C until analysis. Parental consent was provided for all patients to participate in the study. Human sample collection and study protocols were performed in accordance with the guidelines of Xin Hua hospital. All the experimental protocols were approved by the ethics committee of Xin Hua hospital.

Plasma Sample Preparation and Analysis by GC-TOFMS

Plasma metabolites were firstly analyzed by GC-TOFMS following our previously published procedure with minor modifications^{10, 16}. A 50 μL aliquot of plasma sample was spiked with two internal standard solutions (10 μL of L-2-chlorophenylalanine in water, 0.3 mg/mL; 10 μL of heptadecanoic acid in methanol, 1 mg/mL) and vortexed for 10 s. The mixed solution was extracted with 150 μL of methanol/chloroform (3:1) and vortexed for 30 s. After storing for 10 min at -20°C , the samples were centrifuged at 12,000 g for 10 min. An aliquot of the 150 μL supernatant was transferred to a glass sampling vial to vacuum dry at room temperature. Then, a two-step procedure was used for residue derivatization. First, 50 μL of methoxyamine (15 mg/mL in pyridine) was added to the vial and kept at 30°C for 90 min, followed by 50 μL of BSTFA (1% TMCS) at 70°C for 60 min.

Each 1 μL aliquot of the derivatized solution was injected into an Agilent 6890N gas chromatography coupled with a Pegasus HT time-of-flight mass spectrometer (Leco

Corporation, St Joseph, MI) in splitless mode. Metabolites separation was achieved on a DB-5MS capillary column (30 m × 250 µm I.D., 0.25-µm film thickness; (5%-phenyl)-methylpolysiloxane bonded and crosslinked; Agilent J&W Scientific, Folsom, CA). Helium carrier gas was used at a constant flow rate of 1.0 ml/min. The temperature of injection, transfer interface, and ion source was set to 270 °C, 260 °C, and 200 °C, respectively. The GC temperature programming was set to 2 min isothermal heating at 80 °C, followed by 10 °C/min oven temperature ramps to 180 °C, 5 °C/min to 240 °C, and 25 °C/min to 290 °C, and a final 9 min maintenance at 290 °C. MS measurements were implemented with electron impact ionization (70 eV) in the full scan mode (m/z 30_600) and the acquisition rate was 20 spectrum/second in the TOFMS setting.

GC-TOFMS Data Analysis

The acquired MS files from GC-TOFMS analysis were processed by ChromaTOF software (v4.50.8.0, Leco Co., CA). After the pretreatment of baseline correction, de-noising, smoothing, alignment, and de-convolution, raw data containing retention time, intensity, and the mass-to-charge ratio of each peak were obtained. Compounds were identified by comparison with the internal library built with the standard reference compounds and NIST 05 Standard mass spectral databases software in NIST MS search 2.0 (NIST, Gaithersburg, MD). Briefly, the metabolites were annotated initially with NIST 05 (a similarity score over 700) and further validated with the internal metabolite database (over 1000 metabolites) that was previously established using the exactly same analytical protocol. Both mass-spectrum and retention times were used to achieve precise compound annotations. Internal standards and any known artificial peaks, such as peaks caused by noise, column bleed, and BSTFA derivatization procedure, were removed from the data set. Multivariate statistical model of orthogonal partial least square discriminate analysis (OPLS-DA) was constructed with the software SIMCA-P+ (version 13.0, Umetrics, Umea, Sweden). On the basis of a variable importance in the projection (VIP) threshold of 1, a number of metabolites responsible for the difference in the metabolic profiles of different groups could be obtained. In parallel, the metabolites identified by the OPLS-DA model were validated at a univariate level using the non-parametric Mann-Whitney U test (SPSS 21) with the critical p-value set to 0.05.

Molecular Pathway and Network Analysis in Ingenuity Pathway Analysis (IPA)

To systematically understand the metabolism difference between BA and NHS, we uploaded the metabolite lists (with HMDB IDs) and the change folds of the differentially expressed metabolites onto an IPA server¹⁷. Canonical pathways and Chemical-Protein interaction networks were generated based on the knowledge sorted in the Ingenuity Pathway Knowledge Base. A ratio of the number of metabolites that map to the canonical pathway divided by the total number of molecules that map to the pathway was displayed. Fisher's exact test was used to calculate a p-value determining the probability that the association between the metabolites and the canonical pathway was explained by chance alone. The network score was based on the hypergeometric distribution and was calculated with the right-tailed Fisher's Exact Test. The higher a score was, the more relevant the eligible submitted molecules were to the network.

Validation of Deferential Amino Acid Metabolism between BA and NHS

A P180 absolute IDQ kit was used to validate the amino acid expressions in plasma of 53 BA and 28 NHS infants. Forty amino acids and biogenic amines were quantitatively measured according to manufacturer's instruction with minor modifications using an ACQUITY ultra performance liquid chromatography (BEH C18 1.7 μ m 2.1 $\times$ 100 mm column) coupled with Waters Xevo TQ-S triple quadrupole mass spectrometry. In brief, 10 μ L of the internal standard solutions was added to all wells of the plate. Then, 10 μ L plasma or calibration standards was pipetted onto the center of each spot and samples were dried at room temperature under a stream of nitrogen. 50 μ L freshly prepared 5% v/v phenylisothiocyanate solution was immediately added to each dried well. After 20 minutes for derivatization, the plate was dried in the nitrogen evaporator for 60 minutes. 300 μ L extraction solvent (5 mM ammonium acetate in methanol) was added to each well and incubated for 30 minutes. An aliquot of 5 μ L of the derivatized amino acids and biogenic amines was injected for analysis.

RESULTS

Demographic information and clinical manifestations

Demographic and clinical parameters including age, gender, total bilirubin, direct bilirubin, total bile acids, alanine aminotransferase, aspartate aminotransferase, and γ -glutamyltransferase (GGT) were listed in Table 1. Age and gender showed no significant difference among all the groups. Both BA and NHS infants have significantly elevated levels of total bilirubin, direct bilirubin, total bile acids, alanine aminotransferase, and aspartate aminotransferase compared to healthy infants, because of the severe liver damage of BA and NHS infants. However, these liver function parameters showed no significant difference between BA and NHS, except for GGT ($p=0.001$).

Plasma Metabonomic Profiling of BA, NHS and Healthy Infants (HC)

Plasma samples from 45 BA, 15 NHS, and 6 HC infants were conducted to GC-TOFMS metabonomic profiling. After data processing, OPLS-DA was performed on the dataset including 78 identified compounds (the compounds list was shown in supplementary data). Distinct metabolic profiles could be observed from OPLS-DA scores plots between BA and HC (Figure 1A, $R^2X=0.336$, $R^2Y=0.803$, $Q^2Y=0.405$), between NHS and HC (Figure 1B, $R^2X=0.346$, $R^2Y=0.868$, $Q^2Y=0.542$), and between BA and NHS (Figure 1C, $R^2X=0.316$, $R^2Y=0.758$, $Q^2Y=0.389$). On the basis of $VIP>1$ of the OPLS-DA model and $P<0.05$ of nonparametric Mann-Whitney U test, a total of 18 metabolites (L-glutamic acid, L-ornithine, L-isoleucine, 2-hydroxy-3-methylbutyric acid, L-lysine, L-valine, D-mannose, L-tryptophan, gluconolactone, L-serine, phosphate, succinic acid, glycerol 3-phosphate, D-glucose, citric acid, oxalic acid, ribitol, and pentanoic acid) were differentially expressed between BA and NHS (Table 2). There were 7 amino acids (L-Glutamic acid, L-Ornithine, L-Isoleucine, L-Lysine, L-Valine, L-Tryptophan, and L-Serine) among these metabolites.

IPA analysis of the differentially-expressed metabolites between BA and NHS

The total 18 differentially-expressed metabolites were uploaded to IPA website for pathway and network analysis. The most significantly altered network was “Cellular Function and Maintenance, Hepatic System Development and Function, Neurological Disease”, with a score of 28 (Figure 3). The five most significantly different pathways were tRNA Charging, Serine Biosynthesis, Glucose and Glucose-1-phosphate Degradation, Proline Biosynthesis II (from Arginine), TCA cycle II (Table 3). A total of 11 metabolite (L-glutamic acid, L-ornithine, L-lysine, D-mannose, L-tryptophan, L-serine, phosphate, succinic acid, glycerol 3-phosphate, D-glucose, and citric acid) were involved in the network. The amino acids, L-glutamic acid, L-isoleucine, L-lysine, L-serine, L-tryptophan, and L-valine involved in the tRNA Charging were altered most significantly ($p=2.3E-07$).

Validation of Amino Acid metabolism in BA and NHS infants

To further validate the amino acid expressions in BA and NHS, a total of 40 amino acids and biogenic amines were quantitatively measured using a P180 Absolute IDQ kit in 53 BA and 28 NHS infants. Among these 40 compounds, a total of 34 compounds were quantitatively detected. All the 7 differentially expressed amino acids (L-Glutamic acid, L-Ornithine, L-Isoleucine, L-Lysine, L-Valine, L-Tryptophan, and L-Serine) found by GC-TOFMS were significantly different between BA and NHS in the larger sample set (Figure 2). We also found taurine was increased in BA compared to NHS infants (Figure 2).

DISCUSSION

In this study, we systematically profiled total plasma metabolites in BA, NHS, and healthy infants using GC-TOFMS. The OPLS-DA models derived from the metabonomic analysis showed clear separation among BA, NHS, and healthy infants. After the IPA analysis of the differentially-expressed metabolites, UPLC-MS/MS was performed to validate the differentially-expressed amino acids between BA and NHS.

Here, untargeted GC-TOFMS and targeted UPLC-MS/MS were combined to discover and validate the metabonomic differences between BA and NHS. Untargeted GC-TOFMS was used to measure as many features as possible with no bias, and targeted UPLC-MS/MS was used to verify the potential biomarkers in a more reliable and robust manner¹⁸. The comparison showed that the two platforms used in this study were both of great reliability. All of the differentially-expressed amino acids discovered in GC-TOFMS were also found to be significantly altered using UPLC-MS/MS.

IPA analysis revealed that “Cellular Function and Maintenance, Hepatic System Development and Function, Neurological Disease” was the most significantly altered network between BA and NHS (Figure 3). This is reasonable as both BA and NHS have severe liver dysregulation. There were 5 amino acids (L-glutamic acid, L-ornithine, L-lysine, L-tryptophan, and L-serine) involved in this network. The perturbations in amino acid metabolism are very well established in liver diseases although the particular amino acids and directions of change are very complex.^{19, 20} The increased level of seven amino acids in the BA group as compared with the NHS group obtained in our study were in agreement

with the reports that the serum levels of all amino acids were elevated under circumstances of fulminant hepatic necrosis.¹⁹ It was reported that aromatic amino acids (AAA, tyrosine and phenylalanine), taurine, aspartate, threonine, serine, asparagine, methionine, ornithine and histidine were significantly increased in partially hepatectomized rats (65% of liver was removed) while in carbon tetrachloride induced liver cirrhosis, increased levels of AAA, aspartate, asparagine, methionine, ornithine and histidine and decreased levels of branched-chain amino acids (BCAA; i.e. valine, leucine and isoleucine) were observed.¹⁹ L-glutamic acid is the core metabolite in the network and the most significant differentially-expressed amino acid between BA and NHS. Zhao et al also found the higher expression of L-glutamic acid compared to NHS, while both were still lower than healthy controls⁵. It has been found that development of acute liver failure induced a significant reduction in L-glutamic acid levels in pig models²¹. L-glutamic acid was also reported as a protective regulator for deoxynivalenol-induced damage manifested as oxidative stress, intestinal injury, and signaling inhibition²². We found L-tryptophan was significantly elevated in BA compared to NHS while no significant differences of L-tyrosine and L-phenylalanine were observed between the two groups. L-glutamine was found marginally lower in BA compared to NHS ($p=0.075$, data not shown) in our study, which is in agreement with the findings by Zhao et al that L-glutamine was reduced in BA compared to NHS, while both were higher than normal control⁵.

It is worth noting that two of the BCAAs, L-valine and L-isoleucine, were higher expressed in BA than NHS. Their downstream metabolite, 2-hydroxy-3-methylbutyric acid, was significantly decreased. BCAAs are receiving great attention nowadays as their predictive role in the occurrence of type 2 diabetes²³. BCAAs were significantly decreased in carbon tetrachloride induced liver cirrhosis, while significantly elevated in acute carbon tetrachloride induced injury and ischemic damage.¹⁹ A recent article by Holecek et al., on amino acids alterations in liver cirrhosis were reviewed in detail²⁰ and the best-known alteration is the decrease of BACC²⁴. Furthermore, BCAA supplementation reduces oxidative stress and prolongs survival in rats with advanced liver cirrhosis²⁵, and restores the innate immune responses²⁶. Although etiology of BA is unclear, immune dysregulation plays important roles in pathogenesis of the disease²⁷. BA will eventually progress to liver cirrhosis^{1, 28} and the significantly higher expression of γ -glutamyltransferase levels possibly indicates worse hepatic damage in the BA group. The difference in amino acid metabolism may reflect the different causes of liver damages between BA and NHS^{29, 30}.

CONCLUSIONS

This study suggests that plasma metabolic profiling has great potential in differentiating BA from NHS, and amino acid metabolism is significantly different between the two diseases.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

This study is supported by Shanghai Key Laboratory of Pediatric Gastroenterology and Nutrition (14DZ2272400), National Natural Science Youth Foundation of China (81300516), Natural Science Foundation of Shanghai (13ZR1434200), Doctoral Fund of Ministry of Education of China (20130073120013), Medicine-Engineering Cross Research Funding of Shanghai Jiao Tong University (YG2012MS03).

REFERENCES

1. Hartley JL ; Davenport M ; Kelly DA (2009). Biliary atresia. *Lancet*, 374 (9702), 1704–13.19914515
2. Mieli-Vergani G ; Vergani D (2009). Biliary atresia. *Semin Immunopathol*, 31 (3), 371–81.19533128
3. Sira MM ; Taha M ; Sira AM (2014). Common misdiagnoses of biliary atresia. *European journal of gastroenterology & hepatology*, 26 (11), 1300–5.25171020
4. Peng SS ; Hsu WM ; Chen HL ; Mo YH (2014). Using volume index and lateral hepatic angle to differentiate biliary atresia from TPN-associated cholestasis. *Journal of pediatric gastroenterology and nutrition*, 59 (3), 403–8.24821537
5. Zhao D ; Han L ; He Z ; Zhang J ; Zhang Y (2014). Identification of the plasma metabolomics as early diagnostic markers between biliary atresia and neonatal hepatitis syndrome. *PloS one*, 9 (1), e85694 3885726.24416443
6. Ukarapol N ; Wongsawasdi L ; Ong-Chai S ; Riddhiputra P ; Kongtawelert P (2007). Hyaluronic acid: additional biochemical marker in the diagnosis of biliary atresia. *Pediatr Int*, 49 (5), 608–11.17875085
7. Song Z ; Dong R ; Fan Y ; Zheng S (2012). Identification of serum protein biomarkers in biliary atresia by mass spectrometry and enzyme-linked immunosorbent assay. *J Pediatr Gastroenterol Nutr*, 55 (4), 370–5.22569524
8. El-Faramawy AA ; El-Shazly LB ; Abbass AA ; Ismail HA (2011). Serum IL-6 and IL-8 in infants with biliary atresia in comparison to intrahepatic cholestasis. *Trop Gastroenterol*, 32 (1), 50–5.21922857
9. Zahm AM ; Hand NJ ; Boateng LA ; Friedman JR (2012). Circulating microRNA is a biomarker of biliary atresia. *J Pediatr Gastroenterol Nutr*, 55 (4), 366–9 3459263.22732895
10. Chen T ; Xie G ; Wang X ; Fan J ; Qiu Y ; Zheng X ; Qi X ; Cao Y ; Su M ; Wang X ; Xu LX ; Yen Y ; Liu P ; Jia W (2011). Serum and urine metabolite profiling reveals potential biomarkers of human hepatocellular carcinoma. *Molecular & cellular proteomics : MCP*, 10 (7), M110 004945 3134066.
11. Xuan J ; Pan G ; Qiu Y ; Yang L ; Su M ; Liu Y ; Chen J ; Feng G ; Fang Y ; Jia W ; Xing Q ; He L (2011). Metabolomic profiling to identify potential serum biomarkers for schizophrenia and risperidone action. *Journal of proteome research*, 10 (12), 5433–43.22007635
12. Chen WL ; Wang JH ; Zhao AH ; Xu X ; Wang YH ; Chen TL ; Li JM ; Mi JQ ; Zhu YM ; Liu YF ; Wang YY ; Jin J ; Huang H ; Wu DP ; Li Y ; Yan XJ ; Yan JS ; Li JY ; Wang S ; Huang XJ ; Wang BS ; Chen Z ; Chen SJ ; Jia W (2014). A distinct glucose metabolism signature of acute myeloid leukemia with prognostic value. *Blood*, 124 (10), 1645–54.25006128
13. Mathe EA ; Patterson AD ; Haznadar M ; Manna SK ; Krausz KW ; Bowman ED ; Shields PG ; Idle JR ; Smith PB ; Anami K ; Kazandjian DG ; Hatzakis E ; Gonzalez FJ ; Harris CC (2014). Noninvasive urinary metabolomic profiling identifies diagnostic and prognostic markers in lung cancer. *Cancer research*, 74 (12), 3259–70 4100625.24736543
14. Chi Y ; Pei L ; Chen G ; Song X ; Zhao A ; Chen T ; Su M ; Zhang Y ; Liu J ; Ren A ; Zheng X ; Xie G ; Jia W (2014). Metabonomic profiling of human placentas reveals different metabolic patterns among subtypes of neural tube defects. *Journal of proteome research*, 13 (2), 934–45.24397701
15. Spratlin JL ; Serkova NJ ; Eckhardt SG (2009). Clinical applications of metabolomics in oncology: a review. *Clinical cancer research : an official journal of the American Association for Cancer Research*, 15 (2), 431–40 2676437.19147747

16. Qiu Y ; Cai G ; Su M ; Chen T ; Zheng X ; Xu Y ; Ni Y ; Zhao A ; Xu LX ; Cai S ; Jia W (2009). Serum metabolite profiling of human colorectal cancer using GC-TOFMS and UPLC-QTOFMS. *Journal of proteome research*, 8 (10), 4844–50.19678709
17. Sun L ; Li J ; Zhou K ; Zhang M ; Yang J ; Li Y ; Ji B ; Zhang Z ; Zhu H ; Yang L ; He G ; Gao L ; Wei Z ; Wang K ; Han X ; Liu W ; Tan L ; Yu Y ; He L ; Wan C (2013). Metabolomic analysis reveals metabolic disturbance in the cortex and hippocampus of subchronic MK-801 treated rats. *PloS one*, 8 (4), e60598 3618452.23577129
18. Zhu J ; Djukovic D ; Deng L ; Gu H ; Himmati F ; Chiorean EG ; Raftery D (2014). Colorectal cancer detection using targeted serum metabolic profiling. *Journal of proteome research*, 13 (9), 4120–30.25126899
19. Holecsek M ; Mraz J ; Tilser I (1996). Plasma amino acids in four models of experimental liver injury in rats. *Amino Acids*, 10 (3), 229–41.24178537
20. Holecsek M (2015). Ammonia and amino acid profiles in liver cirrhosis: effects of variables leading to hepatic encephalopathy. *Nutrition*, 31 (1), 14–20.25220875
21. Ytrebo LM ; Sen S ; Rose C ; Ten Have GA ; Davies NA ; Hodges S ; Nedredal GI ; Romero-Gomez M ; Williams R ; Revhaug A ; Jalan R ; Deutz NE (2006). Interorgan ammonia, glutamate, and glutamine trafficking in pigs with acute liver failure. *American journal of physiology. Gastrointestinal and liver physiology*, 291 (3), G373–81.16782695
22. Wu M ; Xiao H ; Ren W ; Yin J ; Tan B ; Liu G ; Li L ; Nyachoti CM ; Xiong X ; Wu G (2014). Therapeutic effects of glutamic acid in piglets challenged with deoxynivalenol. *PloS one*, 9 (7), e100591 4077692.24984001
23. Wang TJ ; Larson MG ; Vasan RS ; Cheng S ; Rhee EP ; McCabe E ; Lewis GD ; Fox CS ; Jacques PF ; Fernandez C ; O'Donnell CJ ; Carr SA ; Mootha VK ; Florez JC ; Souza A ; Melander O ; Clish CB ; Gerszten RE (2011). Metabolite profiles and the risk of developing diabetes. *Nature medicine*, 17 (4), 448–53 3126616.
24. Morgan MY ; Marshall AW ; Milsom JP ; Sherlock S (1982). Plasma amino-acid patterns in liver disease. *Gut*, 23 (5), 362–70 1419690.7076013
25. Iwasa M ; Kobayashi Y ; Mifuji-Moroka R ; Hara N ; Miyachi H ; Sugimoto R ; Tanaka H ; Fujita N ; Gabazza EC ; Takei Y (2013). Branched-chain amino acid supplementation reduces oxidative stress and prolongs survival in rats with advanced liver cirrhosis. *PloS one*, 8 (7), e70309 3723692.23936183
26. Nakamura I ; Ochiai K ; Imai Y ; Moriyasu F ; Imawari M (2007). Restoration of innate host defense responses by oral supplementation of branched-chain amino acids in decompensated cirrhotic patients. *Hepatology research : the official journal of the Japan Society of Hepatology*, 37 (12), 1062–7.17608669
27. Feldman AG ; Mack CL (2012). Biliary atresia: cellular dynamics and immune dysregulation. *Seminars in pediatric surgery*, 21 (3), 192–200 3399127.22800972
28. Diniz G ; Tosun Yildirim H ; Calkavur S ; Ecevit C ; Olukman O ; Bekem Soylu O ; Aktas S (2015). Can Neonatal Hepatitis Be More Fatal than Biliary Atresia? *Fetal Pediatr Pathol*.
29. Rastogi A ; Krishnani N ; Yachha SK ; Khanna V ; Poddar U ; Lal R (2009). Histopathological features and accuracy for diagnosing biliary atresia by prelaparotomy liver biopsy in developing countries. *J Gastroenterol Hepatol*, 24 (1), 97–102.19196397
30. El-Guindi MA ; Sira MM ; Sira AM ; Salem TA ; El-Abd OL ; Konsowa HA ; El-Azab DS ; Allam AA (2014). Design and validation of a diagnostic score for biliary atresia. *J Hepatol*, 61 (1), 116–23.24657403

Figure 1.

OPLS-DA score plots of metabolic profiles derived from the data of GC-TOFMS among BA, NHS, and HC groups. A, OPLS-DA score plots between BA and HC. $R^2X=0.336$, $R^2Y=0.803$, $Q^2Y=0.405$. B, OPLS-DA score plots between NHS and HC. $R^2X=0.346$, $R^2Y=0.868$, $Q^2Y=0.542$. C, OPLS-DA score plots between BA and NHS. $R^2X=0.316$, $R^2Y=0.758$, $Q^2Y=0.389$.

Figure 2.

The most significantly differential network between BA and NHS. The network of “Cellular Function and Maintenance, Hepatic System Development and Function, Neurological Disease” was the most significantly altered network between BA and NHS, with a score of 28 and 11 altered metabolites involved. Red means the elevated metabolites in BA compared to NHS.

Figure 3.
The differentially expressed amino acids and biogenic amines validated by UPLC-MS/MS between BA and NHS.

Table 1.

Demographic and clinical investigations of the sample set.

	BA (n=53)	NHS (n=28)	HC (n=6)	P value		
				BA vs HC	NHS vs HC	BA vs NHS
Age (days)	67 (32–145)	63 (33–194)	46.5 (29–113)	0.135	0.392	0.111
Gender (F/M)	16/37	9/19	2/4	0.874	0.890	0.856
Total Bilirubin (μmol/L)	156(80–271)	147 (43–419)	4.7 (4–14)	0.000	0.000	0.727
Direct Bilirubin (μmol/L)	113 (51–181)	99 (32–283)	1.6 (1–6)	0.000	0.000	0.109
Total Bile Acids (μmol/L)	117(34–282)	107 (24–320)	5.6 (3–11)	0.000	0.000	0.798
Alanine Aminotransferase (U/L)	126 (58–742)	111 (31–427)	25 (7–53)	0.000	0.000	0.153
Aspartate Aminotransferase (U/L)	231 (81–1440)	176 (61–651)	28 (9–59)	0.000	0.000	0.096
γ- Glutamyl Transferase (U/L)	670 (123–2074)	186 (41–815)	NA	NA	NA	0.001

BA, biliary atresia; NHS, neonatal hepatitis syndrome; HC, healthy control.

Table 2.

The differentially expressed plasma metabolites between BA and NHS.

Metabolites	VIP	P	Ratio (BA/NHS)
L-glutamic acid	2.24	0.000	2.15
L-ornithine	1.26	0.002	1.51
L-isoleucine	1.85	0.005	1.33
2-hydroxy-3-methylbutyric acid	2.01	0.006	0.57
L-lysine	1.74	0.006	1.17
L-valine	1.72	0.007	1.29
D-mannose	1.55	0.008	1.19
L-tryptophan	1.53	0.008	1.44
Gluconolactone	1.28	0.023	1.42
L-serine	1.52	0.024	1.24
Phosphate	1.26	0.024	1.12
Succinic acid	1.17	0.024	1.14
Glycerol 3-phosphate	1.11	0.024	1.47
D-glucose	1.33	0.032	1.42
Citric acid	1.24	0.034	1.45
Oxalic acid	1.88	0.035	0.82
Ribitol	1.39	0.035	0.75
Pentanoic acid	1.33	0.042	1.42

BA, biliary atresia; NHS, neonatal hepatitis syndrome.

Table 3.

The 5 most significantly altered canonical pathways between BA and NHS revealed by IPA analysis.

Canonical Pathways	P	Differentially expressed metabolites
tRNA Charging	2.3E-07	L-glutamic acid, L-isoleucine, L-lysine, L-serine, L-tryptophan, L-valine
Serine Biosynthesis	1.2E-04	L-glutamic acid, L-serine, phosphate
Glucose and Glucose-1-phosphate Degradation	3.1E-04	D-glucose, gluconolactone, phosphate
Proline Biosynthesis II (from Arginine)	4.0E-04	L-glutamic acid, L-ornithine, phosphate
TCA cycle II	1.1E-03	citric acid, phosphate, succinic acid