Published in final edited form as:

Nano Lett. 2017 December 13; 17(12): 7940–7944. doi:10.1021/acs.nanolett.7b04336.

Single-Molecule Clocks Controlled by Serial Chemical Reactions

Alexander Johnson-Buck^{1,2,3,4} and William M. Shih^{1,2,3,*}

¹Department of Cancer Biology, Dana Farber Cancer Institute, Boston, Massachusetts 02215, United States

²Wyss Institute for Biologically Inspired Engineering, Harvard Medical School, Boston, Massachusetts 02115, United States

³Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts 02115, United States

⁴Present Address: Department of Internal Medicine, The University of Michigan, Ann Arbor, Michigan, 48109, United States

Abstract

Chemical clocks usually achieve well-defined temporal delays through concentration thresholding coupled to the production, degradation, activation, or inhibition of downstream effectors. In this way, the stochastic dynamics of many individual molecules yield essentially deterministic bulk behavior through ensemble averaging. As a result, their temporal evolution is governed by ensemble dynamics rather than by the behavior of an individual molecule or complex. Here, we present a general approach for the design of single-molecule clocks that permits quasi-deterministic control over the lifetime of single molecular interactions without any external synchronization. By coupling the dissociation of a bimolecular complex to a series of irreversible chemical steps, we interpose a well-defined time delay between binding and dissociation. The number and speed of irreversible steps can be varied to systematically tune both the lifetimes of complexes and the precision of the time delay, raising the prospect of localized timekeeping in nanoscale systems and devices.

For Table of Contents Only:

* William_Shih@dfci.harvard.edu; Phone: 617-632-5143; Fax: 617-394-2897.

Supporting Information

Materials and methods including protocols for TIRF microscopy, denaturing polyacrylamide gel electrophoresis assays, kinetic analysis of single-molecule data, and Monte Carlo simulations, as well as supporting references, a list of oligonucleotide sequences used for single-molecule clocks (Table S1), and Figures S1–S7.

Keywords

DNA nanotechnology; dynamic DNA nanotechnology; single-molecule; chemical clock; clock reaction; DNA polymerase

Precise control of the temporal evolution of chemical systems plays important roles in naturally occurring biological systems such as circadian clocks¹, cardiac pacemaker cells², and embryonic development³. Rational engineering of chemical and biochemical kinetics has given rise to systems that mimic some of the features of these natural systems; in addition to classic examples such as the non-oscillatory iodine clock reaction^{4,5} and its oscillatory derivatives⁶, synthetic analogs of biological clocks have been constructed in the form of *in vitro* transcriptional oscillators^{7,8} and synthetic oscillatory networks in bacteria⁹. Atomic clocks¹⁰ permit extremely precise analytic timekeeping by spectroscopic determination of resonant frequencies in atoms or ions in the vapor phase. However, it is challenging to achieve precise local control of the timescale of individual molecular events owing to their stochastic dynamics and the typical lack of internal memory. Such control would be valuable, for instance, in permitting the local timing of proposed autonomous molecular devices such as nanoscale assembly lines^{11–13}. It may also provide new strategies for multiplexing in fluorescence microscopy, where transiently binding probes have been used for super-resolution imaging^{14,15}, high-confidence identification of endogenous nucleic acids¹⁶, and precise determination of stoichiometry within biomolecular complexes and aggregates¹⁷. Here, we outline and experimentally demonstrate a general strategy for precisely controlling the interaction time of single bimolecular complexes. By guiding the complex through a series of several irreversible reactions that terminate in dissociation, it is possible to specify the lifetime of the interaction much more precisely than is achievable for single-step binding or dissociation reactions. We dub such quasi-deterministic devices single-molecule clocks.

The binding equilibrium of a bimolecular complex can usually be approximated as a two-state system whose kinetics are characterized by a bimolecular association rate constant k_0 and a unimolecular dissociation rate constant k_1 (Fig. 1a). Thus, when the reversible binding of a fluorescent probe to an immobilized target¹⁴ is monitored at the single-molecule level, the dwell times in the high-fluorescence (bound) state are exponentially distributed (Fig. 1a). The complex has no memory of how long it has existed, and its probability of dissociating within an interval of time t is independent of its history, resulting in a broad range of complex lifetimes (coefficient of variation equal to 1 in the ideal case). In contrast, single-molecule clocks pass through a series of intermediates *via* irreversible reactions characterized by rate constants $k_1, k_2, \dots, k_N$ before dissociation can occur (Fig. 1b). While the dwell time τ_i of each intermediate is a random value drawn from an exponential distribution, the overall wait time

$$\Delta t = \sum_{i=1}^N \tau_i \quad (1)$$

follows a gamma distribution, with coefficient of variation proportional to $1/\sqrt{N} < 1$ (Fig. 1b)¹⁸. That is, as the number of irreversible steps increases, the lifetime of the complex becomes more narrowly distributed relative to its expectation value, or more deterministic.

Although the single-molecule clock principle might be implemented in various ways, one of the most convenient and compact approaches is to use a processive enzyme such as a DNA polymerase (DNAP). We therefore constructed a system in which a DNAP with strand displacement activity controls the delay between the binding and dissociation of a fluorescent probe (Fig. 1c). An initiator strand *I* binds to a hairpin template *T* (Figure S1), opening the hairpin and exposing a primer binding site on *T*. Subsequently, a fluorophore-labeled primer *P* binds to the complex, resulting in an increase in localized fluorescence. Almost immediately (since [DNAP] > 1 μ M), the DNAP binds and begins elongating the primer. Upon reaching the end of the template, the strand displacement activity of the DNAP causes the fluorescent waste complex *W* to dissociate from *I*, resulting in a loss of fluorescence from the binding site. The dwell time in the high-fluorescence state is thus controlled by the rate of nucleotide addition, and is expected to exhibit a gamma distribution of dwell times whose shape and mean value is dependent on the length of the template and the concentration of dNTPs. Since clock behavior requires that polymerase binding be fast relative to polymerization, we performed a kinetic assay by polyacrylamide gel electrophoresis to confirm this fact (Figure S2).

To characterize clock behavior at the single-molecule level, we used a total internal reflection fluorescence (TIRF) assay in which biotinylated initiator strand *I* was immobilized on a passivated coverslip (Fig. 2b, see Methods for details). In this scheme, repeated cycles of increased and decreased fluorescence intensity are expected to occur at the position of each surface-immobilized copy of *I*, with the duration of each burst controlled by polymerase activity. Such repeated fluorescence bursts are indeed observed (Fig. 2c), and as expected, exhibit a narrow, non-exponential distribution of dwell times (τ) in the bound state (Fig. 2d). Fitting a gamma probability distribution to the τ histogram yields a mean dwell time $\overline{\Delta\tau}$ of 12.7 ± 0.2 s for a 41-nucleotide template (T_{41}). Importantly, the photobleaching lifetime (determined in absence of dNTPs or polymerase) is ~ 83.8 s (Figure S3); thus, only $\sim 15\%$ of primers are expected to photobleach before dissociating under these conditions. Consistent with the hypothesis that dwell time is predominantly controlled by the rate of nucleotide addition, $\overline{\Delta\tau}$ is strongly dependent on the concentration of dNTPs (Fig. 2e), exhibiting saturation behavior with an apparent K_M of 12.6 ± 2.4 μ M with respect to dNTPs, consistent with a previous estimate¹⁹. Notably, a minority of events exhibit much longer dwell times, which may be due to nonspecific binding of probes to the imaging surface or occasional polymerase stalling (Figure S4).

As the number of irreversible steps between binding and dissociation increases, two trends are expected: (i) $\overline{\Delta\tau}$ should increase in proportion to template length, and (ii) dwell time distributions should have a smaller coefficient of variation. To test these predictions, we constructed a set of templates with identical toehold and stem domains but with varying length of the terminal loop, with total lengths ranging from 41 to 153 nucleotides (Fig. 3a–d). For longer loops, short hairpins were intentionally introduced to prevent unwanted base-pairing between the stem and terminal loop, and to reduce the entropic penalty for loop

closure. As expected, in the presence of a saturating concentration (100 μM) of dNTPs, all templates exhibited repeated bursts of fluorescence with a precisely determined τ that increased with template size (Fig. 3a–d). Notably, several of the τ distributions are well separated from one another (Fig. 3e), making it possible in principle to distinguish between templates from the duration of only one binding event. Fitting a gamma distribution to the main peak of each dwell time histogram yields estimates of $\overline{\Delta t}$ that depend linearly on the length of the template (Fig. 3f), providing further evidence that polymerization is the primary rate-limiting process prior to complex dissociation. The slope of a linear fit yields an estimated elongation rate of $\sim 8.3 \text{ nt s}^{-1}$, considerably slower than the maximum elongation rate of 191.2 nt s^{-1} reported for this polymerase at 65°C ²⁰ but much faster than the maximum rate of $0.047\text{--}0.24 \text{ nt s}^{-1}$ reported at room temperature using coumarin-modified dNTPs¹⁹ (all experiments in the present study were performed at room temperature using unmodified dNTPs). In addition to $\overline{\Delta t}$, fitting of a gamma distribution to the experimental τ histograms yields estimates of N_{app} , the apparent number of irreversible steps between binding and dissociation, and $\overline{\Delta t_{step}}$, the apparent average dwell time per step. As expected, N_{app} increases linearly with the length of the template (Figure S5), indicating more deterministic behavior as the number of steps increases. However, contrary to naïve predictions, N_{app} is not equal to the number of nucleotides added, but is related by a proportionality constant of ~ 0.43 , suggesting that not all nucleotide addition steps are equally rate-limiting. Perhaps the simplest explanation for this higher-than-predicted N_{app} is that different dNTPs are incorporated with substantially different kinetics, which has been observed experimentally for KlenTaq polymerase²¹. Using Monte Carlo simulations, we find that a value of $N_{app}/N = 0.43$ is obtained if one of the nucleotides is incorporated at a rate 9-fold lower than the other three nucleotides, for instance (Figure S6). The apparent number of steps could also be diminished by dissociation of the waste complex before the polymerase reaches the end of the template, or by partially rate-limiting polymerase binding/initiation kinetics. On the other hand, $\overline{\Delta t_{step}}$ is essentially constant across all studied template lengths (Figure S5), consistent with expectations that the rate of polymerization is independent of the length of the template. As expected, the dwell times in the unbound state are not clocked, and follow a simple exponential distribution (Figure S7).

We have demonstrated one way of implementing single-molecule clocks, namely, using DNA polymerases with strand displacement activity. We have further shown that the quasi-deterministic temporal behavior of this class of single-molecule clocks can be systematically tuned by varying the length of the template and the rate of polymerization. Historically, the term “chemical clock” has been applied to both non-oscillatory^{4,5,22} systems, in which an initial induction period is followed by a sudden change in reagent concentration²³, and oscillatory⁶ systems, in which the concentration of a reagent undergoes repeated changes in concentration with a well-defined period. While we only demonstrate implementation of a non-oscillatory single-molecule clock here, the principle of generating time delays through serial reactions may be used to design periodic single-molecule clocks in the future, provided that sufficiently well-behaved systems (*i.e.*, devoid of significant kinetic traps) are discovered or designed.

While we have exploited DNA polymerization as the source of serial reactions, in principle any series of irreversible reactions with similar kinetics could be substituted for the polymerization, so long as they control the time interval between two reporting events. For example, DNA-only catalytic cascades²⁴, as well as RNA polymerases, exonucleases, ATP-dependent helicases, and other processive enzymes that act upon templates of well-defined size, may be expected to yield single-molecule clock behavior. In some cases such systems may even function as clocks in their native biological contexts as, for example, has been hypothesized for translational pausing in protein folding²⁵.

In the future, a number of improvements and modifications on this principle may be possible. In our system, a background of exponentially distributed binding events is evident in the dwell time histograms, and becomes more dominant for templates of increasing length (Fig. 3e and Figure S4), possibly due to polymerase stalling, which is expected to occur in a larger fraction of templates as the number of base pairs per template increases. Improved template sequence design and/or alternate polymerases may reduce the influence of polymerase stalling on dwell time distributions, resulting in higher fidelity clocks. While the clocks presented in this manuscript exhibit bound-state lifetimes in the range of ~1–20 s, these may be lengthened (e.g., by lowering dNTP concentration or increasing template length) or shortened (e.g., by lowering monovalent ion concentration to increase the elongation rate of the polymerase) to suit the needs of a given application. In addition, the further development of polymerases that act upon bio-orthogonal nucleotide analogs may ensure greater compatibility with a variety of applications in biological imaging and biomolecular nanotechnology. We anticipate a variety of compelling uses for the deterministic control of kinetics at the single-molecule level.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

The Harvard University Office of Technology Development has filed a provisional patent application on technologies described herein. This work was funded by support from NSF Expeditions Award CCF-1317291, NSF DMREF Award 1435964, ARO MURI Award W911NF-12-1-0420, ONR Award N000141510073, Dana Farber Cancer Institute Claudia Adams Barr Award, Harvard Catalyst Pilot Grant Award, and the Wyss Institute at Harvard Faculty Award to W.M.S.

References

- (1). Harmer SL ; Hogenesch JB ; Straume M ; Chang H-S ; Han B ; Zhu T ; Wang X ; Kreps JA ; Kay SA Science 2000, 290, 2110–2113.11118138
- (2). Glass L Nature 2001, 410, 277–284.11258383
- (3). Pourquié O Science 2003, 301, 328–330.12869750
- (4). Landolt H Berichte Dtsch. Chem. Ges 1886, 19, 1317–1365.
- (5). Oliveira AP ; Faria RB J. Am. Chem. Soc 2005, 127, 18022–18023.16366551
- (6). Briggs TS ; Rauscher WC J. Chem. Educ 1973, 50, 496.
- (7). Kim J ; Winfree E Mol. Syst. Biol 2011, 7.
- (8). Baccouche A ; Montagne K ; Padirac A ; Fujii T ; Rondelez Y Methods 2014, 67, 234–249.24495737

- (9). Elowitz MB ; Leibler S Nature 2000, 403, 335–338.10659856
- (10). Essen L ; Parry JV L. Nature 1955, 176, 280–282.
- (11). He Y ; Liu DR Nat. Nanotechnol 2010, 5, 778–782.20935654
- (12). Gu H ; Chao J ; Xiao S-J ; Seeman NC Nature 2010, 465, 202–205.20463734
- (13). Kassem S ; Lee ATL ; Leigh DA ; Markevicius A ; Solà J Nat. Chem 2015, 8, 138–143.26791896
- (14). Jungmann R ; Steinhauer C ; Scheible M ; Kuzyk A ; Tinnefeld P ; Simmel FC Nano Lett 2010, 10, 4756–4761.20957983
- (15). Jungmann R ; Avendaño MS ; Woehrstein JB ; Dai M ; Shih WM ; Yin P Nat. Methods 2014, 11, 313–318.24487583
- (16). Johnson-Buck A ; Su X ; Giraldez MD ; Zhao M ; Tewari M ; Walter NG Nat. Biotechnol 2015, 33, 730–732.26098451
- (17). Jungmann R ; Avendaño MS ; Dai M ; Woehrstein JB ; Agasti SS ; Feiger Z ; Rodal A ; Yin P Nat. Methods 2016, 13, 439–442.27018580
- (18). Floyd DL ; Harrison SC ; van Oijen AM Biophys. J 2010, 99, 360–366.20643053
- (19). Walsh MT ; Roller EE ; Ko K-S ; Huang X Biochemistry (Mosc.) 2015, 54, 4019–4021.
- (20). Kiefer JR ; Mao C ; Hansen CJ ; Basehore SL ; Hogrefe HH ; Braman JC ; Beese LS Structure 1997, 5, 95–108.9016716
- (21). Montgomery JL ; Rejali N ; Wittwer CT J. Mol. Diagn 2014, 16, 305–313.24607271
- (22). Forbes GS ; Estill HW ; Walker OJ J. Am. Chem. Soc 1922, 44, 97–102.
- (23). Preece SJ ; Billingham J ; King AC J. Math. Chem 1999, 26, 47.
- (24). Yin P ; Choi HMT ; Calvert CR ; Pierce NA Nature 2008, 451, 318–322.18202654
- (25). Marin M Biotechnol. J 2008, 3, 1047–1057.18624343

Figure 1.

Principles and implementation of single-molecule clocks. (a) The equilibrium two-state binding of a conventional probe (as in DNA-PAINT) results in a wide range of bound-state dwell times that follow an exponential distribution whose coefficient of variation (σ/τ) is equal to 1. Complexes with $\tau = 4, 8$, and 12 s (red, blue, and black lines) are predicted to have substantial overlap in their τ distributions. (b) A single-molecule clock exploits a series of irreversible chemical steps that must occur between binding and dissociation of the probe, resulting in a gamma distribution of dwell times whose coefficient of variation is

$1/\sqrt{N}$, where N is the number of irreversible steps). Clock complexes with $N = 40, 80$, and 120 , and $\tau = 4, 8$, and 12 s (red, blue, and black lines) are predicted to have t distributions that only minimally overlap. (c) Implementation of single-molecule clocks using a DNA polymerase (DNAP) with strand displacement activity. A hairpin-loop template (T) binds to an initiator (I) *via* toehold-mediated strand displacement, revealing the primer-binding domain c . Binding of the fluorescent primer (P) yields a localized increase in fluorescence, and is rapidly followed by binding of DNAP. Elongation of the primer by DNAP serves as the rate-limiting process for dissociation, and results in displacement of I from the fluorescent waste complex (W), which diffuses away and results in a loss of fluorescence. As long as elongation is slow relative to DNAP binding, the duration of the localized spike in fluorescence is controlled (or clocked) by the stepwise addition of deoxyribonucleoside triphosphates (dNTPs) to the growing DNA strand. Note that the duplex stem of T contains two mismatched base pairs that are introduced to ensure orthogonal binding between sequences b/b^* and c/c^* (i.e., to prevent binding of P directly to I).

Figure 2.

Demonstration of a polymerase-controlled single-molecule clock. (a) Denaturing polyacrylamide gel showing the DNAP-catalyzed extension of a Cy5-labeled primer (P) in the presence of a 41-nucleotide (nt) loop template, T_{41} . Extension of the Cy5-labeled primer occurs efficiently only in the presence of T_{41} , initiator (I), and DNAP. No detectable leakage occurs in absence of I . Incubation time: 5 min. Gray arrow indicates the direction of migration in the gel. (b) Surface-based assay of single-molecule clock performance using total internal reflection fluorescence (TIRF) microscopy. Each surface-anchored copy of I

undergoes repeated cycles of T_{4I} binding, P binding, extension by DNAP, and displacement of W . Localized fluorescence is only visible between binding of P and the dissociation of W . (c) Representative single-molecule fluorescence trajectory showing four consecutive clock binding cycles in the same location on a coverslip using T_{4I} . (d) Dwell time distribution for T_{4I} in the presence of 2.5 μM dNTPs ($n = 1000$ binding events). The red line represents a nonlinear least-squares fit of a gamma probability distribution to the data, yielding an estimated mean dwell time $\overline{\Delta t}$ of 12.7 ± 0.2 s (95% confidence bounds from fit). (e) Dependence of the mean dwell time on dNTP concentration. The red line represents a nonlinear least-squares fit of a Michaelis-Menten saturation curve to the data, resulting in an estimated K_M of 12.6 ± 2.4 μM and τ_{min} of 2.1 ± 0.3 s (95% confidence bounds from fit).

Figure 3.

Design of polymerase-clocked clocks with tunable duration. (a–d) Design schematics (left) and representative single-molecule fluorescence traces (right) of clocks using four different templates (*T*) of varying loop length. The indicated number of nucleotides (nt) is the total length of each template, including the constant primer- and initiator-binding domains. (e) Dwell time distributions and gamma probability distribution fits for the four clock templates shown in (a–d). $n = 143, 154, 196$, and 155 binding events for the 41 nt, 57 nt, 97 nt, and 153 nt templates, respectively. (f) Dependence of mean dwell time on total template length. A linear regression fit to the data is shown; slope = 0.12 s nt^{-1} , $R^2 = 0.986$. The red dashed line indicates the 14-nt footprint of the primer binding site on the template, which does not contribute to the timing mechanism. All experiments shown in this figure were performed in the presence of a saturating concentration ($100 \mu\text{M}$) of dNTPs.