Published in final edited form as:

Sleep Med Clin. 2018 September ; 13(3): 457–465. doi:10.1016/j.jsmc.2018.04.012.

Impact of Poor Sleep on Physical and Mental Health in Older Women

Katie L. Stone, PhD and Qian Xiao, PhD

California Pacific Medical Center Research Institute, San Francisco, CA 94158, USA. Department of Health and Human Physiology and Department of Epidemiology, University of Iowa, Iowa City, IA 52242, USA

SYNOPSIS

Many aspects of sleep and circadian rhythms change as people age. Older adults usually experience decrease in sleep duration and efficiency, increase in sleep latency and fragmentation, high prevalence of sleep disorders and weakened rest-activity rhythms. Research evidence also suggests that women in particular are more likely to report aging-related sleep problems. In this review, we present epidemiological and clinical evidence on the relationships between sleep deficiency and physical and mental outcomes in older women, explore potential mechanisms underlying such relationships, point out gaps in the literature that warrant future investigations, and consider implications in the clinical and public health settings.

Keywords

Sleep deficiency; circadian rhythms; cardiometabolic disorders; older women; metabolic syndrome; physical health; mental health; falls

INTRODUCTION

Sleep is an important determinant of human health, and healthy sleep is crucial for healthy aging. In a recent joint consensus statement of the American Academy of Sleep Medicine and Sleep Research Society, healthy sleep was defined as “adequate duration, good quality, appropriate timing and regularity, and the absence of sleep disturbances or disorders”.¹ The prevalence of sleep disorders and disturbances increases dramatically with advancing age.² There is growing evidence that sleep disturbances may accelerate the aging process, and contribute to a wide range of chronic diseases. Despite these health consequences, sleep problems frequently are undiagnosed and untreated, particularly in the elderly.³

CORRESPONDING AUTHOR kstone@psg.ucsf.edu Mission Hall, Second Floor, 550 16th Street, San Francisco, CA 94158, USA. AUTHOR CONTACT INFORMATION: kstone@psg.ucsf.edu Mission Hall, Second Floor, 550 16th Street, San Francisco, CA 94158, USA, qian-xiao@uiowa.edu 225 Grand South, E118 Field House, Iowa City, IA 52242, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

DISCLOSURE STATEMENT

The authors declare no conflict of interest

Previous studies have suggested that compared to men, women may be more sensitive to the impact of aging on sleep, and older women are more likely to report sleep problems.^{4,5} As the older population continues to grow in many parts of the world, it is important to understand the health effects of sleep disruption in the context of aging. In this review, we synthesize and present epidemiological and clinical evidence on the relationships between sleep deficiency and various health conditions that are highly prevalent in older women, explore potential mechanisms underlying such relationships, point out gaps in the literature that warrant future investigations, and consider implications for the clinical and public health settings.

SLEEP IN OLDER ADULTS

Many aspects of sleep change as people age. A 2004 meta-analysis by Ohayon and colleagues⁶ synthesized findings from 65 studies with objective measurement of sleep using polysomnography (PSG) or actigraphy. They found that older age was associated with decreases in total sleep time, sleep efficiency and percentage of slow-wave sleep and rapid eye movement (REM) sleep; while sleep latency, percentage of light sleep, and minutes of wake after sleep onset significantly increased with age. Such changes in sleep architecture are consistent with an increase in sleep complaints in the older population. In over 9,000 people aged 65 or older, Foley and colleagues⁷ assessed the frequencies of reporting common sleep disturbances, including trouble falling asleep, waking up, waking too early, and non-restorative sleep. In this study, half of the participants reported at least one complaint as frequently occurring, and up to a third of the population showed symptoms of insomnia.⁷ Similarly high prevalence (15–25%) of insomnia symptoms has also been reported in the Sleep Heart Health Study (SHHS).⁸ Sleep-disordered breathing (SDB), another common sleep disorder, is estimated to affect from 30–60% of older adults, depending on the definition used and specific population.^{9–13} Moreover, aging coincides with altered circadian activity rhythms including decreased amplitude (height of rhythm),¹⁴ fragmentation or loss of rhythms (weakening of rhythmic pattern),^{15,16} and altered timing of peak rhythm activity. Timing changes in older adults frequently result in earlier onset of sleepiness in the evening, and an earlier morning waking time.¹⁶

Interestingly, a growing body of evidence suggests that there are sex differences both in sleep and age-related changes in sleep.⁴ In general, although women tend to have better objectively measured sleep quality, paradoxically, they are more likely than men to report subjective sleep problems, including shorter sleep and poorer sleep quality.¹⁷ A meta-analysis of sex differences in insomnia showed that not only is the risk of insomnia higher in women than in men across all age groups, the difference in insomnia risk between women and men widens with age.¹⁸

SLEEP AND CARDIOMETABOLIC HEALTH

Sleep plays a vital role in numerous physiological processes, including the regulation of metabolic, hormonal, and immune function, all of which are essential for cardiometabolic health. Numerous studies have linked disorders and disturbances of sleep to cardiometabolic

outcomes, including obesity, hypertension, dyslipidemia, diabetes and cardiovascular diseases (CVD).

Sleep Duration—Short sleep duration is associated with obesity in children and younger adults, however such a relationship in older adults remains less clear.^{19–21} The mixed findings in the elderly may be partially due to the high prevalence of chronic conditions in this population, which may both confound and modify the effect of sleep on weight. In a study of over 80,000 healthy men and women aged 51–72 years, those with self-reported short sleep (<7 hours) at baseline were more likely to experience substantial weight gain (> 5kg) and risk of developing obesity over 10 years of follow-up.²² Interestingly, this association may be stronger in older women than in men. Two studies of middle-to-old-aged subjects in Finland and Spain reported an association between short sleep and higher weight gain in women but not in men.^{23,24}

Short sleep duration has also been associated with other cardiometabolic consequences. Two meta-analyses showed that short sleep duration was associated with 23% increase in hypertension risk,²⁵ 48% increase in coronary heart disease and 15% increase in stroke,²⁶ and these associations were stronger in women. Another meta-analysis demonstrated that each one-hour decrease in sleep duration was associated with a significant 9% increase in the risk of type 2 diabetes.²⁷

Several studies have also reported an association between long sleep duration and obesity, diabetes and CVD risk and mortality.^{26,28} In a large observational study of older women, Stone and colleagues²⁹ found that those who reported 10 or more hours of sleep per 24 hours had a 77% increase in risk of cardiovascular-related mortality compared to older women who reported 8–9 hours of sleep. Several lines of evidence suggest that these associations with adverse health outcomes related to long sleep duration may be partially explained by comorbidities.^{30–32}

Studies using objectively-measured sleep duration in the older population are still limited, and their findings are mixed. For example, a cross-sectional relationship between objectively-measured short sleep and higher adiposity was observed in both the Multi-Ethnic Study of Atherosclerosis and Study of Osteoporotic Fractures (SOF).^{33,34} However, thus far prospective analyses in younger adults have found no relationship between actigraphy-measured short sleep duration and weight gain.^{35,36}

Insomnia and Poor Sleep Quality—Several studies reported greater adiposity associated with various measures of sleep quality such as wakefulness, sleep fragmentation, daytime sleepiness, and overall poor sleep quality.^{37–40} Van den Berg and colleagues³⁹ found that greater objectively measured sleep fragmentation was associated with higher BMI and greater risk of obesity in older adults, and the association between short sleep duration and obesity was no longer significant after adjustment for sleep fragmentation. This suggests that, in older adults, the quality of sleep may be more important than the quantity in relation to cardiometabolic outcomes. Poor sleep quality, measured both subjectively and objectively, has also been linked to several markers of sub-clinical CVD,⁴¹ as well as poorer glycemic control measured by increased hemoglobin A_{1c} in patients with type 2 diabetes.⁴² However,

Phillips and colleagues⁴³ found no association of insomnia symptoms with incident hypertension in older women.

Finally, some studies have suggested that objectively measured short sleep duration and insomnia symptoms together increase risk of cardiovascular disease more strongly than either condition alone. For example, Bertisch and colleagues⁴⁴ examined short sleep (objectively measured based on PSG), insomnia or poor sleep quality, and the interaction of these sleep problems for prediction of incident CVD in middle-aged to older adults. Whereas neither condition alone was significantly associated with incident CVD, those with both short sleep and insomnia or poor sleep quality had a significant 30% increase in risk of incident CVD over 11 years of follow-up. These results were similar in both women and men. This finding highlights the importance of considering sleep health as a multi-dimensional exposure in terms of the impact on health in aging.

Snoring and Sleep Disordered Breathing (SDB)—There is a well-established association between SDB and obesity, as well as obesity-related health outcomes.⁴⁵ In addition, several studies also showed that even occasional snoring may be a risk factor for obesity, metabolic syndrome and diabetes in middle-to-old aged women.^{46,47} Among middle-aged women in the Nurses' Health Study, those who reported "occasional" snoring were 40% more likely to develop diabetes during 10 years of follow-up than those who reported never snoring.⁴⁶

Epidemiological studies have also established that SDB increases risk for incident hypertension, heart failure, coronary heart disease, and stroke in men and younger adults.^{48–50} However, effects of SDB on cardiovascular outcomes in older women are uncertain. Early findings from the SHHS showed associations with CVD only in men, while more recent data from the SHHS and Atherosclerotic Risk in Communities (ARIC) cohorts showed that SDB severity predicted higher troponin levels and increased risk for left ventricular hypertrophy, heart failure, and CVD-related mortality among elderly women compared to men.⁵¹ These data, which focused on an older sample for a longer period of time than earlier analyses, underscore the importance of examining this risk in older women.

Weakened Rest-Activity Rhythms—Circadian rhythms are intrinsic physiologic cycles of approximately 24 hours that are critically involved in control of sleep-wake cycles and numerous physiological processes. Circadian and sleep-wake rhythm abnormalities have been observed among those with a wide variety of medical conditions. However, it is not clear whether sleep-wake rhythms directly influence morbidity and mortality in aging, or are biomarkers of advanced physiological age. Indeed, van Hilten and colleagues⁵² studied the relationship of age with nocturnal behavior in 100 healthy older adults and found that in the absence of illness, age itself has only marginal effects on sleep and wake.

Using actigraphy, a previous study of older women in the SOF found that lower amplitude, extreme timing (early or late) of peak activity, and weaker strength in the overall rhythmicity predicted higher cardiovascular mortality.⁵³ Similarly, a more fragmented and less stable rhythm of rest-activity was associated with total mortality in the Rotterdam Study, although the study did not specifically examine cardiovascular deaths.⁵⁴

SLEEP AND DEPRESSION

While the prevalence of major depression in older adults is relatively low at 1–2%, an increasing number of older adults experience clinically significant depressive symptoms,⁵⁵ which are linked to greater risk of functional impairment, disability, and illness.⁵⁶ Furthermore, late-life depression disproportionately impacts older women, and some evidence suggests that this is because older women tend to experience more persistent depressive symptoms over time, and have longer survival compared to men.^{55,57}

Sleep Duration—Cross-sectional studies have reported greater prevalence of depressive symptoms in older adults with both short and long sleep duration.^{30,57,58} While a protective role for sleep deprivation or short sleep in risk of depression has been reported in younger adults, these findings have not been confirmed in older adults.^{59,60} Using data from the Nurses' Health Study, Patel and colleagues³⁰ found that depressive symptoms were strongly correlated with self-reported long sleep duration. Another study reported that an association between self-reported long sleep duration and depressive symptoms was significant in older men, but not older women.⁵⁷

Fewer studies have examined the association of sleep duration and risk of developing depressive symptoms over time. A few longitudinal studies using both objective and subjective assessment of sleep in older adults have reported no significant association of sleep duration (either short or long) and risk of incident depressive symptoms.^{61,62} However, Fernandez-Mendoza and colleagues⁶¹ studied a cohort of adults (mean age 52 years) and found that those with insomnia and objectively measured short sleep duration had the highest risk of developing depressive symptoms over 7.5 years of follow-up. This finding further underscores the importance of considering sleep health more comprehensively rather than focusing on a single domain.

Insomnia and Poor Sleep Quality—Insomnia is common in older adults with depression, but recent evidence suggests that this is a bi-directional association and that insomnia and poor sleep quality may also lead to incident depressive symptoms over time. For example, Maglione and colleagues⁶² studied 952 women aged 70 and older with minimal depressive symptoms at baseline. Sleep was assessed objectively using actigraphy, and subjectively using the Pittsburgh Sleep Quality Index (PSQI). Higher PSQI scores (indicating more sleep disturbance) were associated with greater risk of developing depressive symptoms, with stronger associations for the sleep quality (MOR 1.41; CI 1.13 – 1.77) and sleep latency (MOR 1.21; CI 1.03–1.41) subscales. Objectively, prolonged minutes of wake after sleep onset emerged as a risk factor for incident depressive symptoms, while sleep duration was not a significant predictor. These findings suggest that sleep quality, rather than the absolute quantity of sleep, may be more important for mental health in older women.

The effects of poor sleep on risk of depressive symptoms may not be limited to a single domain of sleep (e.g. sleep duration). Furihata and colleagues⁶³ tested the association of an index of 'sleep health' in relation to risk of incident depressive symptoms in older women. The 0–5 index was created by summing across five dimensions classified as "poor" based on

self-reported sleep, including satisfaction with sleep duration, daytime sleepiness, mid-sleep time, sleep onset latency, and sleep duration. Results showed a strong gradient of increasing risk of developing incident depressive symptoms with increasing number of “poor” sleep dimensions. Despite these intriguing findings, the body of evidence linking sleep to depression has tended to focus on one dimension at a time.

Weakened Rest-Activity Rhythms—Although aging promotes disruption in circadian rhythms, the relationship between depression and circadian rhythm disruption in older adults remains largely unexplored. Maglione and colleagues⁶⁴ examined the relationship between depressive symptoms and circadian activity rhythms in older adults among 3,020 women (mean age 84 years). Greater levels of depressive symptoms were associated with greater desynchronization of circadian activity rhythms, as well as later average time becoming active in the morning, but there was no association with acrophase. Further evidence of an association between circadian activity rhythms and risk of depression was reported in a longitudinal analysis by Smagula and colleagues⁶⁵ Among 2,124 older men with minimal depressive symptoms at baseline, those in the lowest quartile of rhythm robustness were 2.5 times more likely to develop clinically significant depressive symptoms during a 5-year follow-up (OR=2.58; 95% CI 1.11– 5.59). Rhythm timing and amplitude did not significantly predict development of depression. These associations have not been examined in older women.

SLEEP, COGNITIVE DECLINE, AND RISK OF DEMENTIA

Sleep disturbance is common in older adults with dementia, and evidence also suggests that disturbed sleep may contribute to development of cognitive problems and risk of dementia.⁶⁶ While the mechanisms are not completely understood, experimental studies suggest that even a single night of sleep deprivation leads to accumulation of beta-amyloid in the human brain.⁶⁷ Beta-amyloid is a metabolic waste product that may form plaques over time, contributing to Alzheimer Disease (AD). Sleep may therefore play a critical role in the prevention of cognitive decline and risk for AD. Evidence also suggests that SDB may pose a risk for development of cognitive problems in older women.

Sleep Duration and Poor Sleep Quality

Overall, there is little prospective evidence of an association between sleep duration and cognitive decline or risk of incident dementia. Chen and colleagues⁶⁸ studied 7444 community-dwelling older women to test whether self-reported sleep duration predicted incident mild cognitive impairment (MCI) or dementia during follow-up. They found a V-shaped association between sleep duration and risk of MCI/dementia, with both short (< 6 hours/night) and long (> 8 hours/night) sleepers showing 35–36% increase in risk.⁶⁸ However, other studies of both older men and women with actigraphy measures of sleep demonstrated no association of sleep duration and risk of cognitive decline.^{69,70} In contrast, these studies showed that objective measures of sleep fragmentation (for example, sleep efficiency) consistently predicted both cross-sectional cognitive function as well as prospective decline in cognition. Further studies are needed to better define the characteristics of healthy sleep that may prevent cognitive decline in older women, and to

determine if treatment for sleep problems may slow the decline in cognition and development of dementia.

Snoring and Sleep Disordered Breathing (SDB)

SDB results in nocturnal hypoxemia and more fragmented sleep, both of which may have effects on cognitive function. Yaffe and colleagues⁷¹ studied the association of SDB and nocturnal hypoxemia (assessed using overnight in-home polysomnography) with subsequent 5-year risk of developing MCI and dementia among 298 older women (mean age 82.3 years). In this study, older women with SDB (apnea-hypopnea index ≥ 15) had nearly a two-fold increase in risk of developing MCI or dementia over 5 years of follow-up, whereas nocturnal hypoxemia was associated with a 1.71-fold increase in risk of developing MCI or dementia.⁷¹ In another study, Chang and colleagues⁷² used data from a large health insurance database in Taiwan to compare 5-year risk of dementia among adults aged 40 and older with and without sleep apnea diagnosis at baseline. Among 1414 patients studied, presence of sleep apnea was associated with a significant 1.7-fold increase in risk of developing dementia. In this same study, among women aged 70 and older, sleep apnea diagnosis was associated with a 3.2-fold increase in risk of incident dementia.⁷²

There is insufficient evidence for an association between snoring and risk of cognitive decline or onset of dementia.

Weakened Rest-Activity Rhythms

Tranah and colleagues⁷³ examined whether circadian activity rhythms (based on actigraphy) were associated with incident MCI and dementia over 5 years in 1,282 older women. An approximately 50% adjusted higher odds of developing dementia or MCI versus those without any dementia or MCI was observed for those in the lowest quartile of amplitude and rhythm robustness when compared to those in the highest quartile. Timing of activity rhythms also predicted incident MCI/dementia. In particular, older women with delayed acrophase (> 1.5 standard deviations of the population mean) had a significant increase in odds of developing dementia or MCI (OR=1.83, 95% CI, 1.29–2.61) when compared to the mean peak range (mean ± 1.5 standard deviations).

SLEEP AND OTHER AGE-RELATED OUTCOMES AND CONDITIONS

Disturbed sleep in older women, and hypnotics used to treat sleep problems, are independently associated with risk of incident falls.^{74–76} Falls are common in older adults, and frequently lead to injury and increased disability⁷⁷ and mortality.⁷⁸ Further evidence suggests that poor sleep may contribute to decline in physical functioning among older women, independent of falls.⁷⁹ Sleep disturbance is also common in older women with chronic pain, and more recent evidence suggests that this association is bi-directional and that sleep problems may increase the risk of developing chronic pain.⁸⁰ Treatment of insomnia symptoms using cognitive behavioral therapy in older adults with osteoarthritis showed that short-term (2 months) improvements in sleep predicted longer-term improvement in chronic pain outcomes over 9 to 18 months.

IMPLICATIONS FOR CLINICAL PRACTICE

Given the high prevalence of sleep problems in older women, and the debilitating impact on quality of life and physical and mental health, it is imperative for health care professionals to identify, monitor and treat sleep problems in this vulnerable population. Unfortunately, evidence suggests that compared to other lifestyle behaviors, sleep problems have not been screened as frequently in family medicine clinics.⁸¹ Special needs of older women should be considered in determining the best treatment options. In older women with sleep apnea, behavior modification such as avoiding alcohol and certain medications may be particularly important, as these patients may be more sensitive to their effects on upper airway function during sleep.⁸² Hypnotic medications are effective and commonly prescribed by physicians for treatment of insomnia, but have been linked to adverse events in the elderly, including falls and fractures, adverse cognitive and psychomotor events and daytime fatigue.⁸³ Therefore the American Geriatric Society has recommended avoiding hypnotic prescriptions for insomnia in older adults.⁸⁴ Randomized controlled trials have shown that behavioral interventions including cognitive behavioral therapy for insomnia are effective in older adults,⁸⁵ and should be considered as the first-line approach. Moreover, several studies also showed that exercise,^{86,87} social activities,⁸⁸ and cognitive training⁸⁹ may also be beneficial for older adults with insomnia.

REFERENCES

1. Watson NF , Badr MS , Belenky G , et al. Recommended Amount of Sleep for a Healthy Adult: A Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society. *Sleep* 2015;38(6):843–844.26039963
2. Roth T Insomnia: definition, prevalence, etiology, and consequences. *J Clin Sleep Med* 2007;3(5 Suppl):S7–10.17824495
3. Boehlecke BA . Epidemiology and pathogenesis of sleep-disordered breathing. *Curr Opin Pulm Med* 2000;6(6):471–478.11100955
4. Carrier J , Semba K , Deurveilher S , et al. Sex differences in age-related changes in the sleep-wake cycle. *Front Neuroendocrinol* 2017;47:66–85.28757114
5. Reyner LA , Horne JA , Reyner A . Gender- and age-related differences in sleep determined by home-recorded sleep logs and actimetry from 400 adults. *Sleep* 1995;18(2):127–134.7792492
6. Ohayon MM , Carskadon MA , Guilleminault C , Vitiello MV . Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep* 2004;27(7):1255–1273.15586779
7. Foley DJ , Monjan AA , Brown SL , Simonsick EM , Wallace RB , Blazer DG . Sleep complaints among elderly persons: an epidemiologic study of three communities. *Sleep* 1995;18(6):425–432.7481413
8. Unruh ML , Redline S , An MW , et al. Subjective and objective sleep quality and aging in the sleep heart health study. *J Am Geriatr Soc* 2008;56(7):1218–1227.18482295
9. Mehra R , Stone KL , Blackwell T , et al. Prevalence and correlates of sleep-disordered breathing in older men: osteoporotic fractures in men sleep study. *J Am Geriatr Soc* 2007;55(9):1356–1364.17767677
10. Ancoli-Israel S , Kripke DF . Prevalent sleep problems in the aged. *Biofeedback Self Regul* 1991;16(4):349–359.1760457
11. Ancoli-Israel S , Kripke DF , Klauber MR , Mason WJ , Fell R , Kaplan O . Sleep-disordered breathing in community-dwelling elderly. *Sleep* 1991;14(6):486–495.1798880
12. Ancoli-Israel S , Kripke DF , Klauber MR , et al. Natural history of sleep disordered breathing in community dwelling elderly. *Sleep* 1993;16(8 Suppl):S25–29.8178017

13. Ancoli-Israel S , Martin J , Jones DW , et al. Sleep-disordered breathing and periodic limb movements in sleep in older patients with schizophrenia. *Biol Psychiatry* 1999;45(11):1426–1432.10356624
14. Kripke DF , Youngstedt SD , Elliott JA , et al. Circadian phase in adults of contrasting ages. *Chronobiol Int* 2005;22(4):695–709.16147900
15. Buysse DJ , Monk TH , Carrier J , Begley A . Circadian patterns of sleep, sleepiness, and performance in older and younger adults. *Sleep* 2005;28(11):1365–1376.16335481
16. Czeisler CA , Dumont M , Duffy JF , et al. Association of sleep-wake habits in older people with changes in output of circadian pacemaker. *Lancet* 1992;340(8825):933–936.1357348
17. van den Berg JF , Miedema HM , Tulen JH , Hofman A , Neven AK , Tiemeier H . Sex differences in subjective and actigraphic sleep measures: a population-based study of elderly persons. *Sleep* 2009;32(10):1367–1375.19848365
18. Zhang B , Wing YK . Sex differences in insomnia: a meta-analysis. *Sleep* 2006;29(1):85–93.16453985
19. Patel SR , Hu FB . Short sleep duration and weight gain: a systematic review. *Obesity (Silver Spring)* 2008;16(3):643–653.18239586
20. Nielsen LS , Danielsen KV , Sorensen TI . Short sleep duration as a possible cause of obesity: critical analysis of the epidemiological evidence. *Obes Rev* 2011;12(2):78–92.20345429
21. Magee L , Hale L . Longitudinal associations between sleep duration and subsequent weight gain: a systematic review. *Sleep Med Rev* 16(3):231–241.21784678
22. Xiao Q , Arem H , Moore SC , Hollenbeck AR , Matthews CE . A large prospective investigation of sleep duration, weight change, and obesity in the NIH-AARP Diet and Health Study cohort. *Am J Epidemiol* 2013;178(11):1600–1610.24049160
23. Lyytikainen P , Rahkonen O , Lahelma E , Lallukka T . Association of sleep duration with weight and weight gain: a prospective follow-up study. *J Sleep Res* 2011;20(2):298–302.21199039
24. Lopez-Garcia E , Faubel R , Leon-Munoz L , Zuluaga MC , Banegas JR , Rodriguez-Artalejo F . Sleep duration, general and abdominal obesity, and weight change among the older adult population of Spain. *Am J Clin Nutr* 2008;87(2):310–316.18258619
25. Guo X , Zheng L , Wang J , et al. Epidemiological evidence for the link between sleep duration and high blood pressure: a systematic review and meta-analysis. *Sleep Med* 2013;14(4):324–332.23394772
26. Cappuccio FP , Cooper D , D'Elia L , Strazzullo P , Miller MA . Sleep duration predicts cardiovascular outcomes: a systematic review and meta-analysis of prospective studies. *Eur Heart J* 2011;32(12):1484–1492.21300732
27. Shan Z , Ma H , Xie M , et al. Sleep duration and risk of type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care* 2015;38(3):529–537.25715415
28. Tan X , Chapman CD , Cedernaes J , Benedict C . Association between long sleep duration and increased risk of obesity and type 2 diabetes: A review of possible mechanisms. *Sleep Med Rev* 2017.
29. Stone KL , Ewing SK , Ancoli-Israel S , et al. Self-reported sleep and nap habits and risk of mortality in a large cohort of older women. *J Am Geriatr Soc* 2009;57(4):604–611.19220560
30. Patel SR , Malhotra A , Gottlieb DJ , White DP , Hu FB . Correlates of long sleep duration. *Sleep* 2006;29(7):881–889.16895254
31. Krueger PM , Friedman EM . Sleep duration in the United States: a cross-sectional population-based study. *Am J Epidemiol* 2009;169(9):1052–1063.19299406
32. Grandner MA , Drummond SP . Who are the long sleepers? Towards an understanding of the mortality relationship. *Sleep Med Rev* 2007;11(5):341–360.17625932
33. Ogilvie RP , Redline S , Bertoni AG , et al. Actigraphy Measured Sleep Indices and Adiposity: The Multi-Ethnic Study of Atherosclerosis (MESA). *Sleep* 2016;39(9):1701–1708.27306270
34. Patel SR , Blackwell T , Redline S , et al. The association between sleep duration and obesity in older adults. *International journal of obesity* 2008;32(12):1825–1834.18936766

35. Lauderdale DS , Knutson KL , Rathouz PJ , Yan LL , Hulley SB , Liu K . Cross-sectional and longitudinal associations between objectively measured sleep duration and body mass index: the CARDIA Sleep Study. *Am J Epidemiol* 2009;170(7):805–813.19651664
36. Appelhans BM , Janssen I , Cursio JF , et al. Sleep duration and weight change in midlife women: the SWAN sleep study. *Obesity (Silver Spring)* 2013;21(1):77–84.23505171
37. Jennings JR , Muldoon MF , Hall M , Buysse DJ , Manuck SB . Self-reported sleep quality is associated with the metabolic syndrome. *Sleep* 2007;30(2):219–223.17326548
38. Lyytikainen P , Lallukka T , Lahelma E , Rahkonen O . Sleep problems and major weight gain: a follow-up study. *International journal of obesity* 2011;35(1):109–114.20531352
39. van den Berg JF , Knvistingh Neven A , Tulen JH , et al. Actigraphic sleep duration and fragmentation are related to obesity in the elderly: the Rotterdam Study. *International journal of obesity* 2008;32(7):1083–1090.18414418
40. Xiao Q , Gu F , Caporaso N , Matthews CE . Relationship between sleep characteristics and measures of body size and composition in a nationally-representative sample. *BMC Obes* 2016;3:48.27857841
41. Aziz M , Ali SS , Das S , et al. Association of Subjective and Objective Sleep Duration as well as Sleep Quality with Non-Invasive Markers of Sub-Clinical Cardiovascular Disease (CVD): A Systematic Review. *J Atheroscler Thromb* 2017;24(3):208–226.27840384
42. Lee SWH , Ng KY , Chin WK . The impact of sleep amount and sleep quality on glycemic control in type 2 diabetes: A systematic review and meta-analysis. *Sleep Med Rev* 2017;31:91–101.26944909
43. Phillips B , Buzkova P , Enright P , Cardiovascular Health Study Research G. Insomnia did not predict incident hypertension in older adults in the cardiovascular health study. *Sleep* 2009;32(1):65–72.19189780
44. Bertisch SM , Pollock BD , Mittleman MA , et al. Insomnia with Objective Short Sleep Duration and Risk of Incident Cardiovascular Disease and All-Cause Mortality: Sleep Heart Health Study. *Sleep* 2018.
45. Ioachimescu OC , Collop NA . Sleep-disordered breathing. *Neurol Clin* 2012;30(4):1095–1136.23099131
46. Al-Delaimy WK , Manson JE , Willett WC , Stampfer MJ , Hu FB . Snoring as a risk factor for type II diabetes mellitus: a prospective study. *Am J Epidemiol* 2002;155(5):387–393.11867347
47. Shin MH , Kweon SS , Choi BY , et al. Self-reported snoring and metabolic syndrome: the Korean Multi-Rural Communities Cohort Study. *Sleep Breath* 2014;18(2):423–430.24194392
48. Gottlieb DJ , Yenokyan G , Newman AB , et al. Prospective study of obstructive sleep apnea and incident coronary heart disease and heart failure: the sleep heart health study. *Circulation* 2010;122(4):352–360.20625114
49. Shahar E , Whitney CW , Redline S , et al. Sleep-disordered breathing and cardiovascular disease: cross-sectional results of the Sleep Heart Health Study. *Am J Respir Crit Care Med* 2001;163(1):19–25.11208620
50. Stone KL , Blackwell TL , Ancoli-Israel S , et al. Sleep Disordered Breathing and Risk of Stroke in Older Community-Dwelling Men. *Sleep* 2016;39(3):531–540.26943468
51. Querejeta Roca G , Redline S , Punjabi N , et al. Sleep apnea is associated with subclinical myocardial injury in the community. The ARIC-SHHS study. *Am J Respir Crit Care Med* 2013;188(12):1460–1465.24156237
52. van Hilten JJ , Middelkoop HA , Braat EA , et al. Nocturnal activity and immobility across aging (50–98 years) in healthy persons. *J Am Geriatr Soc* 1993;41(8):837–841.8340562
53. Tranah GJ , Blackwell T , Ancoli-Israel S , et al. Circadian activity rhythms and mortality: the study of osteoporotic fractures. *J Am Geriatr Soc* 2010;58(2):282–291.20374404
54. Zuurbier LA , Luik AI , Hofman A , Franco OH , Van Someren EJ , Tiemeier H . Fragmentation and stability of circadian activity rhythms predict mortality: the Rotterdam study. *Am J Epidemiol* 2015;181(1):54–63.25491893
55. Barry LC , Allore HG , Guo Z , Bruce ML , Gill TM . Higher burden of depression among older women: the effect of onset, persistence, and mortality over time. *Arch Gen Psychiatry* 2008;65(2):172–178.18250255

56. Prather AA , Vogelzangs N , Penninx BW . Sleep duration, insomnia, and markers of systemic inflammation: results from the Netherlands Study of Depression and Anxiety (NESDA). *J Psychiatr Res* 2015;60:95–102.25307717
57. Brostrom A , Wahlin A , Alehagen U , Ulander M , Johansson P . Sex-specific associations between self-reported sleep duration, depression, anxiety, fatigue and daytime sleepiness in an older community-dwelling population. *Scand J Caring Sci* 2018;32(1):290–298.28574585
58. Patel SR , Sotres-Alvarez D , Castaneda SF , et al. Social and Health Correlates of Sleep Duration in a US Hispanic Population: Results from the Hispanic Community Health Study/Study of Latinos. *Sleep* 2015;38(10):1515–1522.26085298
59. Kalmbach DA , Arnedt JT , Song PX , Guille C , Sen S . Sleep Disturbance and Short Sleep as Risk Factors for Depression and Perceived Medical Errors in First-Year Residents. *Sleep* 2017;40(3).
60. Roberts RE , Duong HT . The prospective association between sleep deprivation and depression among adolescents. *Sleep* 2014;37(2):239–244.24497652
61. Fernandez-Mendoza J , Shea S , Vgontzas AN , Calhoun SL , Liao D , Bixler EO . Insomnia and incident depression: role of objective sleep duration and natural history. *J Sleep Res* 2015;24(4):390–398.25728794
62. Maglione JE , Ancoli-Israel S , Peters KW , et al. Subjective and objective sleep disturbance and longitudinal risk of depression in a cohort of older women. *Sleep* 2014;37(7):1179–1187.25061246
63. Furihata R , Hall MH , Stone KL , et al. An Aggregate Measure of Sleep Health Is Associated With Prevalent and Incident Clinically Significant Depression Symptoms Among Community-Dwelling Older Women. *Sleep* 2017;40(3).
64. Maglione JE , Ancoli-Israel S , Peters KW , et al. Depressive symptoms and circadian activity rhythm disturbances in community-dwelling older women. *Am J Geriatr Psychiatry* 2014;22(4):349–361.23567424
65. Smagula SF , Ancoli-Israel S , Blackwell T , et al. Circadian rest-activity rhythms predict future increases in depressive symptoms among community-dwelling older men. *Am J Geriatr Psychiatry* 2015;23(5):495–505.25066948
66. Ancoli-Israel S , Vitiello MV . Sleep in dementia. *Am J Geriatr Psychiatry* 2006;14(2):91–94.16473972
67. Shokri-Kojori E , Wang GJ , Wiers CE , et al. beta-Amyloid accumulation in the human brain after one night of sleep deprivation. *Proceedings of the National Academy of Sciences of the United States of America* 2018.
68. Chen JC , Espeland MA , Brunner RL , et al. Sleep duration, cognitive decline, and dementia risk in older women. *Alzheimers Dement* 2016;12(1):21–33.26086180
69. Blackwell T , Yaffe K , Laffan A , et al. Associations of objectively and subjectively measured sleep quality with subsequent cognitive decline in older community-dwelling men: the MrOS sleep study. *Sleep* 2014;37(4):655–663.24899757
70. Blackwell T , Yaffe K , Ancoli-Israel S , et al. Poor sleep is associated with impaired cognitive function in older women: the study of osteoporotic fractures. *J Gerontol A Biol Sci Med Sci* 2006;61(4):405–410.16611709
71. Yaffe K , Laffan AM , Harrison SL , et al. Sleep-disordered breathing, hypoxia, and risk of mild cognitive impairment and dementia in older women. *JAMA* 2011;306(6):613–619.21828324
72. Chang WP , Liu ME , Chang WC , et al. Sleep apnea and the risk of dementia: a population-based 5-year follow-up study in Taiwan. *PloS one* 2013;8(10):e78655.24205289
73. Tranah GJ , Blackwell T , Stone KL , et al. Circadian activity rhythms and risk of incident dementia and mild cognitive impairment in older women. *Ann Neurol* 2011;70(5):722–732.22162057
74. Stone KL , Ancoli-Israel S , Blackwell T , et al. Actigraphy-measured sleep characteristics and risk of falls in older women. *Arch Intern Med* 2008;168(16):1768–1775.18779464
75. Stone KL , Ensrud KE , Ancoli-Israel S . Sleep, insomnia and falls in elderly patients. *Sleep Med* 2008;9 Suppl 1:S18–22.18606568

76. Stone KL , Ewing SK , Lui LY , et al. Self-reported sleep and nap habits and risk of falls and fractures in older women: the study of osteoporotic fractures. *J Am Geriatr Soc* 2006;54(8):1177–1183.16913982
77. Gill TM , Murphy TE , Gahbauer EA , Allore HG . Association of injurious falls with disability outcomes and nursing home admissions in community-living older persons. *Am J Epidemiol* 2013;178(3):418–425.23548756
78. Ensrud KE , Ewing SK , Taylor BC , et al. Frailty and risk of falls, fracture, and mortality in older women: the study of osteoporotic fractures. *J Gerontol A Biol Sci Med Sci* 2007;62(7):744–751.17634322
79. Goldman SE , Stone KL , Ancoli-Israel S , et al. Poor sleep is associated with poorer physical performance and greater functional limitations in older women. *Sleep* 2007;30(10):1317–1324.17969465
80. Finan PH , Goodin BR , Smith MT . The association of sleep and pain: an update and a path forward. *J Pain* 2013;14(12):1539–1552.24290442
81. Sorscher AJ . How is your sleep: a neglected topic for health care screening. *J Am Board Fam Med* 2008;21(2):141–148.18343862
82. Feinsilver SH , Hernandez AB . Sleep in the Elderly: Unanswered Questions. *Clin Geriatr Med* 2017;33(4):579–596.28991652
83. Glass J , Lancot KL , Herrmann N , Sproule BA , Busto UE . Sedative hypnotics in older people with insomnia: meta-analysis of risks and benefits. *BMJ* 2005;331(7526):1169.16284208
84. By the American Geriatrics Society Beers Criteria Update Expert P. American Geriatrics Society 2015 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc* 2015;63(11):2227–2246.26446832
85. Irwin MR , Cole JC , Nicassio PM . Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults and in older adults 55+ years of age. *Health Psychol* 2006;25(1):3–14.16448292
86. Benloucif S , Orbeta L , Ortiz R , et al. Morning or evening activity improves neuropsychological performance and subjective sleep quality in older adults. *Sleep* 2004;27(8):1542–1551.15683146
87. King AC , Oman RF , Brassington GS , Bliwise DL , Haskell WL . Moderate-intensity exercise and self-rated quality of sleep in older adults. A randomized controlled trial. *JAMA* 1997;277(1):32–37.8980207
88. Naylor E , Penev PD , Orbeta L , et al. Daily social and physical activity increases slow-wave sleep and daytime neuropsychological performance in the elderly. *Sleep* 2000;23(1):87–95.10678469
89. Haimov I , Shatil E . Cognitive training improves sleep quality and cognitive function among older adults with insomnia. *PLoS one* 2013;8(4):e61390.23577218

KEY POINTS

- The prevalence of sleep disorders and disturbances increases with age and older women may be more sensitive to the impact of aging on sleep.
- Short sleep duration, poor sleep quality, insomnia, sleep disordered breathing and weakened rest-activity rhythms are associated with adverse cardiometabolic outcomes, including obesity, metabolic disorders and cardiovascular disease.
- Sleep disorders and disturbances, and weakened rest-activity rhythms are associated with adverse mental health outcomes including increased risk of depressive symptoms, dementia, and cognitive decline.
- Disturbances in sleep, and treatments for sleep, contribute to other outcomes such as falls, disability, and chronic pain.
- Clinicians should consider special needs of older women in the diagnosis and treatment of sleep problems.