Risk Factors for Recurrent Shoulder Instability after Arthroscopic Revision Anterior Stabilization

Favian Su, BS, Marcin Kowalczyk, MD, Stephenson Amanze Ikpe, MD, Hannah Lee, MD, PhD, Soheil Sabzevari, MD, Albert Lin, MD

University of Pittsburgh Medical Center, Pittsburgh, PA, USA.

Objectives: Individuals that fail arthroscopic anterior stabilization of the shoulder represent a unique and challenging patient population. To date, there have been few large studies that have investigated failure rates following arthroscopic revision anterior stabilization (ARAS) for failed primary arthroscopic stabilization. This study aims to determine the risk factors for recurrence of shoulder instability following ARAS. We hypothesized that male gender, younger age, participation in contact sports, significant glenoid and/or humeral bone loss, ligamentous laxity, and worker's compensation would increase the risk of revision failure.

Methods: Patients who underwent ARAS after a failed arthroscopic primary Bankart repair and had a minimum of 2-year follow-up were included in this study. Glenoid and humeral bone loss were quantitatively assessed using pre-operative T1-weighted magnetic resonance arthrograms to determine if the lesions were on- or off-track. Failure was defined as a recurrent dislocation or subluxation. Chi-square test and t-test were used to compare demographical and surgical parameters between failure and non-failure groups. The significance level was set to 0.05.

Results: Sixty-five patients [age at revision = 26 years (range, 15 - 57), 44 (68%) male] met the inclusion criteria. The mean follow-up time was 4.7 years (range, 2 - 10.8). Twenty-seven patients (42%) had a failed revision at a mean time of 2.3 years (range, 0.2 - 6.1). Age less than 22 years old, ligamentous laxity, the presence of an off-track lesion, and a concomitant superior labral anterior to posterior were significantly associated with revision failure ($p < 0.05$) (Table 1). No difference was observed in the size of glenoid defect between failure and non-failure groups ($14.1\% \pm 4.4\%$ vs. $13.7\% \pm 3.9\%$, $p = 0.762$). The width and depth of the Hill-Sachs lesions were not significantly different between groups (width: 15.3 ± 5.1 mm vs. 14.2 ± 4.8 mm, $p = 0.432$; depth: 4.2 ± 2.3 mm vs. 3.5 ± 1.8 mm, $p = 0.244$). On multivariate analysis, only the presence of an off-track lesion, age less than 22 years, and ligamentous laxity were independent predictors for recurrent instability (OR = 8.9, $p = 0.022$; OR = 5.4, $p = 0.028$; OR = 7.8, $p = 0.031$, respectively).

Conclusion: The failure rate of arthroscopic revision anterior stabilization was 42% with off-track lesions, age less than 22 years, and ligamentous laxity independent risk factors for recurrent instability. While ARAS may be a viable treatment option in the appropriate setting, our study suggests that considerable thought should be exercised before utilizing this approach given the significant number of patients who suffered recurrent instability at greater than 2-years follow-up. For young patients with off-track lesions and/or evidence of ligamentous laxity on physical exam, strong consideration should be given to either an open Bankart repair, a bony augmentation procedure such as a Bristow-Latarjet procedure, or an arthroscopic revision approach with additional augmentation such as a remplissage.

Table 1. Parameters Tested Against Arthroscopic Revision Anterior Stabilization Failure^a

Parameter	Revision Failure	No Revision Failure	OR	p-value
Age at Revision < 22 y	16/27 (59%)	13/38 (34%)	2.80	0.045
Ligamentous Laxity	10/27 (37%)	5/38 (13%)	3.88	0.024
Off-Track Lesion	10/22 (45%)	4/36 (11%)	6.67	0.003
SLAP Tear	11/27 (41%)	6/38 (16%)	3.67	0.024
Male Gender	17/27 (63%)	27/38 (71%)	0.69	0.492
BMI > 30 kg/m ²	3/24 (13%)	3/27 (11%)	1.14	1.000
Dominant Side Injured	20/27 (74%)	23/38 (61%)	1.86	0.255
Bilateral Anterior Instability	3/27 (11%)	7/38 (18%)	0.55	0.503
Workers' Compensation	3/27 (11%)	9/38 (24%)	0.40	0.331
Athletes	12/27 (44%)	17/38 (45%)	0.99	0.981
Contact Sports	9/12 (75%)	12/17 (71%)	1.25	1.000
Competitive Level	3/12 (25%)	3/17 (18%)	1.56	0.669
Open Primary Bankart Repair	2/23 (9%)	4/35 (11%)	0.74	1.000
Traumatic Repair Failure	7/18 (39%)	16/35 (46%)	0.76	0.635
Time from Failure to Revision < 1 y	21/27 (78%)	31/38 (82%)	0.79	0.706
Total Number of Anchors ≤ 3	6/20 (30%)	10/30 (33%)	0.86	0.804
Number of Anteroinferior Anchors ≤ 3	14/20 (70%)	19/30 (63%)	1.35	0.626
Labral Tear ≥ 120°	13/20 (65%)	20/30 (67%)	0.93	0.903

^aData expressed as count/number of available cases (%). Competitive level is defined as collegiate or higher. Bold denotes significance. SLAP, superior labrum anterior and posterior; BMI, body mass index; OR, odds ratio.