Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>

ORIGINAL RESEARCH

Effects of hammock positioning in behavioral status, vital signs, and pain in preterms: a case series study

Valdecira Rodrigues de Jesus^a, Pricila Mara Novais de Oliveira^b,
Vivian Mara Gonçalves de Oliveira Azevedo^{c,*}

^a Faculdade de Ciências Médicas de Minas Gerais, Hospital Sofia Feldman, Programa de Residência em Neonatologia, Belo Horizonte, MG, Brazil

^b Hospital Universitário da Universidade Federal de Juiz de Fora (HU-UFJF/EBSERH), Pediatria, Juiz de Fora, MG, Brazil

^c Universidade Federal de Uberlândia (UFU), Faculdade de Educação Física e Fisioterapia (FAEFI), Uberlândia, MG, Brazil

Received 5 October 2017; received in revised form 26 February 2018; accepted 2 March 2018

Available online 15 March 2018

KEYWORDS

Newborns;
Prematurity;
Neonatal Intensive
Care Units;
Humanization

Abstract

Background: The hammock positioning within the incubators simulates the intrauterine environment, however, there is little evidence of its benefits and possible risks.

Objectives: The aim of this study was to assess the effects of hammock positioning on behavioral status, vital signs, and pain in very low birth weight preterm newborns.

Methods: This is a *quasi*-experimental/case series study in which premature infants (<1500 g) were positioned in supine for one hour in a hammock. The preterm newborns were assessed 10 min before, during (2, 20, 40, and 60 min), and 10 min after hammock positioning with the Brazelton Neonatal Behavioral Assessment Scale, vital signs and pain by the Neonatal Facial Coding System.

Results: 28 preterm infants between 28 and 36 weeks of gestational age were evaluated. Regarding the behavioral state, the preterm newborns progressively evolved to light or deep sleep during hammock positioning. There was a statistically significant reduction of the heart and respiratory rate from 2 to 60th minute in a hammock, which was maintained after the positioning. The oxygen saturation remained within normal values. No changes in pain scores were observed.

Conclusion: The hammock positioning can be considered a safe method of positioning that can be used to reduce the stress levels in very low birth weight preterm newborns. We did not observe worsening in either pain or vital signs.

© 2018 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

* Corresponding author at: Faculdade de Educação Física e Fisioterapia – FAEFI, Rua Benjamin Constant, 1286, Bairro Aparecida, CEP: 38400-678, Uberlândia, MG, Brazil.

E-mail: viviangazevedo1@gmail.com (V.M. Azevedo).

<https://doi.org/10.1016/j.bjpt.2018.03.002>

1413-3555/© 2018 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

The hammock positioning (Hp) refers to the positioning of newborns in a small rectangle fabric (hammock) fixed by extremities within the incubators. Hp is a simple and low-cost method for the therapeutic positioning of newborns. This technique has been used by physical therapists in several Neonatal Intensive Care Units (NICU), although little is known about Hp.¹

Preterm newborns are exposed to a stressful environment and painful interventions in the NICU.² Low birth weight is also associated with increased morbidity and mortality and motor development delay in preterm infants.³ The adequate therapeutic positioning of the preterm can minimize postural abnormalities and asymmetries related to prematurity and stay in NICU, also favors the development of spontaneous and functional motor abilities of newborns.⁴ Studies indicate that the positioning directly influences the cardiovascular, respiratory, and sleep-wake states.⁴⁻⁶

The Hp intends to promote preterm containment, vestibular stimulus, sensory integration, tonic, and behavioral reorganization, thus reducing the noxious sensory stimuli to which the immature brain is exposed in the NICU.^{4,5} In addition, Hp potentially simulates the intra-uterine environment, provides relaxation, stimulates the harmonization of movements, and reduces the energy expenditure of very low birth weight newborns.⁵ Some studies indicate that the technique promotes the reduction of stress during the period of hospitalization, so it can be considered a strategy to humanize the care in the NICU.⁷⁻⁹

However, some Hp related outcomes such as vital signs,¹⁰ pain and behavioral status¹¹ and motor development are still controversial due to low methodological quality used in most studies.⁶ Some studies also point to the risks of apneas or desaturation during Hp.^{10,11} Recent reviews^{1,12} on this topic highlight the lack of scientific evidence that proves the benefits and possible risks of Hp. Therefore, the objective of this study was to evaluate the effects of Hp on the behavioral states, pain, and vital signs of very low birth weight preterm newborns.

Methods

Design and local of study

It consists of a *quasi*-experimental/case series study carried out in the NICU of the Hospital Sofia Feldman – a philanthropic institution linked to the Unified Health System and located in Belo Horizonte, Minas Gerais, Brazil.

Ethical aspects

The research project was approved by the Ethics and Research Committee of the Hospital Sofia Feldman, Belo Horizonte, MG, Brazil (number 1.684.322). All mothers or legal guardians for the preterm signed the free and informed consent term to authorize their participation in the research.

Subjects

Subjects inclusion criteria were: very low-weight preterms (infants with 1500 g or less who was born before 37 weeks of pregnancy) with more than 24 h of life, respiratory and hemodynamic stability, without continuous sedation, and/or use of amines.

We excluded preterm newborns that presented neurological signs or symptoms, use of continuous diet, gastroesophageal reflux, the presence of malformations, grade 2 or higher of peri-intraventricular hemorrhage, severe heart disease, immediate postoperative period, use of surfactant in less than 72 h, phototherapy or under minimal manipulation.

Data collection

The included preterms were positioned for one hour in the supine position in a hammock for at least 60 min after the last diet. The hammocks were made of cotton fabric 1 cm thick. A fabric was also placed between the cervical and scapular region of the NTBs so there was no hyperflexion or hyperextension of the head, which could impair respiratory function. To fix the hammocks in the incubators, the circular openings were used to pass the ropes and the ropes were tied in the upper part of the incubator as shown in Fig. 1. The behavioral state, vital signs, and pain were evaluated in newborns according to the following instruments.

The pain evaluation was performed with the Neonatal Facing Coding System (NFCS).¹³ This scale is based on the evaluation of the presence of the following facial expression: wrinkled foreheads, compressed palpebral fissures, deep nasolabial groove, half-open lips, mouth vertical or horizontal outstretched, tense tongue, tongue protrusion, and the chin quivering.¹⁴ The presence of each item scores 1 point from the maximum score of 8 points. It is considered the presence of pain when 3 or more facial movements quoted appear during the evaluation.

The sleep/wake cycle was evaluated by the Brazelton Neonatal Behavioral Assessment Scale (NBAS).¹⁵ The scale scores the behavioral state respectively from 1 to 6: deep sleep, light sleep, drowsy, alert, eyes open, and crying.

The heart rate (HR) and peripheral oxygen saturation (SpO₂) was monitored with a multiparametric monitor DX 2022[®] (Dixtal, Brazil) by a sensor positioned on the external side of one foot. The respiratory rate (RR) was assessed by visual inspection of respiratory cycles per minute.

The evaluations were performed 10 min before the positioning in hammock, at the second, twentieth, fortieth, and sixtieth minutes after the positioning of the newborn in hammock, and 10 min after the return of the newborn to your incubator.¹⁶ The newborn when removed from the Hp was placed in the incubator in the same position as the baseline evaluation. Data were collected in the morning and afternoon. The NFCS and NBAS scales were applied at each evaluation, in addition to the vital signs collected.

The study would be discontinued if vital signs of the newborn presented a variation greater than 20% of the baseline or signs of respiratory effort (use of accessory muscles in the neck, intercostal, subcostal, and suprasternal retractions or nasal flaring), or constant crying.

Figure 1 Hammock positioning of newborns within incubators.

Statistical analysis

The sample size was calculated using the G * Power 3.1.9.2 software, considering the comparison of the six measurements in one group using repeated measures ANOVA with alpha level or probability of error of 0.05. The minimum sample for the survey was then calculated in 28 participants.

Data were analyzed by the BioEstat software version 5.3 (Pará, Brazil). The Shapiro–Wilk test analyzed the distribution of study variables. Numerical variables were presented in mean and standard deviation and categorical variables in absolute value (relative). ANOVA for repeated measurements with the Tukey post hoc test was performed to compare the numerical variables with parametric distribution between the evaluations at different times. Friedman test with Dunn's post hoc test was performed to compare numerical variables with no parametric distribution. Cochran's Q and Friedman tests were performed to compare the categorical variables. A significance level of $p < 0.05$ was considered.

Results

Of the eligible population, 28 preterms were included, of which 17 (61%) were female. Only 1 newborn was not included due to lack of consent. The newborns gestational age varied from 24 to 31 weeks. The other characteristics of newborns included in this study are described in Table 1.

Regarding oxygen supplementation, 15 (54%) were breathing ambient air and 4 (14%) using a nasal catheter. Regarding ventilatory support, 9 newborns were in ventilatory support: 6 (21%) were using the nasal continuous

positive airway pressure mode and 3 (11%) noninvasive positive pressure ventilation with two levels of pressure.

Regarding the behavioral assessment (Table 2), a significant difference was observed when comparing the moments before, during, and after the Hp ($p < 0.001$). Most of the preterm newborns gradually evolved into light sleep or deep sleep during the Hp. At 60 min, 7 (25%) of the 28 newborns assessed were in deep sleep and 18 (64%) in light sleep. Results for vital signs are shown in Fig. 2. HR and RR showed a significant reduction from 2° to 60° minute of Hp ($p < 0.001$). However, this reduction in HR and RR values were not clinically relevant. SpO₂ presented non-significant oscillations, remaining between 92% and 100%. Vital signs varied within normal limits.

There was no significant difference in pain scores when compared the moments before, during, and after Hp ($p = 0.42$). Only 1 newborn had a facial expression of pain before Hp, which got better after the 20° minute. In the second minute, 2 newborns had a facial expression of pain and another one had the same expression at 60° minute of positioning.

There were no adverse events or instability of the newborns during Hp that justified their return to the incubator.

Discussion

The hammock positioning of preterm newborns with very low birth weight was safe, induced sleep in preterm infants, maintained vital signs within the normal range and did not cause any pain in our sample. The Hp induced relaxation in preterms, since most of the newborns evolved into a state of sleep, which may favor brain maturity and consequently

Table 1 Characteristics of preterm newborns studied.

Variable (<i>n</i> = 28)	Mean $\pm$ SD	Minimum–maximum
Chronological age (days)	21.8 $\pm$ 16.5	6–67
Gestational age at birth (weeks)	28.1 $\pm$ 1.6	24–31
Post-conceptual age at evaluation (weeks)	31.2 $\pm$ 2.1	28–36
Weight at birth (g)	954.6 $\pm$ 191.4	605–1390
Weight at evaluation (g)	1124.2 $\pm$ 258.9	685–1470

SD, standard deviation.

Table 2 Neonatal behavioral assessment of preterms before, during, and after hammock positioning.

	Before	During				After
		2 min	20 min	40 min	60 min	
Deep sleep	0 (0)	0 (0)	2 (7.1)	4 (14.3)	7 (25.0)	0 (0)
Light sleep	15 (53.6)	2 (7.1)	15 (53.6)	21 (75.0)	18 (64.3)	11 (39.3)
Drowsy	5 (17.9)	9 (32.1)	8 (28.6)	2 (7.1)	1 (3.6)	12 (42.9)
Alert	1 (3.6)	4 (14.3)	1 (3.6)	0 (0)	1 (3.6)	4 (14.3)
Eyes open	7 (25.0)	11 (39.3)	2 (7.1)	1 (3.6)	0 (0)	1 (3.6)
Crying	0 (0)	2 (7.1)	0 (0)	0 (0)	1 (3.6)	0 (0)

Data presented in absolute and relative number: *n* (%). Cochran's *Q* test *p* < 0.001. **Figure 2** Vital signs distributions at different moments of evaluation: (A) HR: heart rate; (B) RR: respiratory rate; (C) SpO₂: peripheral oxygen saturation. There was a significant reduction of HR and RR from 2° to 60° minute.

the neuropsychomotor development.⁵ Moreover, the state of relaxation provided by Hp prevents energy expenditure, facilitates sleep, and may interfere positively in the neuromotor development of preterms.² Non-pharmacological measures that increase preterms sleep time within NICUs should be stimulated in order to regulate the sleep-wake

cycle, reduce stress levels and promote the sensory organization of preterms, which favors weight gain of newborns with very low birth weight.¹⁶

Cavalaria¹⁷ states that removing a preterms from the warmth of its mother and placing it on a flat and sometimes without coatings surface, is to be unaware the extreme

need for support, restraint, and swinging. It is important to implement safe and humanized methods of sensory-motor stimulation within the NICU, which favor the neuropsychomotor development of preterm newborns, especially those who are subject to long periods of hospitalization. The main advantages of the hammock in relation to prone positioning are the greater contention to the newborn, not allowing constant changes as it occurs in the nest; better stimulus to flexor tone, favoring this postural pattern; better self-regulation, guaranteeing autonomic stability; and greater body symmetry stimulating the joining of the hands in the midline.^{11,18} Therefore, Hp could bring both short-term and long-term benefits to the development of preterms. Albuquerque et al.¹⁹ suggest the training of professionals working in the NICU to disseminate the use of hammock positioning.

We observed that Hp did not cause any type of discomfort in preterms. According to Costa et al.,¹¹ Hp contributes to humanization in preterm newborns care. Hp demonstrated superiority compared to positioning in the nest because it promoted a greater reduction of stress levels, posture improvement, and preterms organization, and it may be recommended after a stressful stimulus, such as diaper changes. In our study was verified a reduction of HR and RR throughout the positioning, probably related to the relaxation provided to the preterms evaluated. Our results corroborate the findings of Costa et al.²⁰ that also observed a reduction of HR and RR during Hp. Similar results were obtained by Keller et al.¹⁰ who compared Hp to prone positioning in very low weight preterm newborns. The authors found that the Hp showed better autonomic stability, maintaining the SpO₂ within the normal range, lower RR, and HR and also better neuromuscular maturity compared to prone.¹⁰

Regarding oxygenation during Hp, Bottos et al.¹⁸ found no difference in oxygen levels in the supine position within the incubator and Hp. Another study that compared the positioning of healthy term infants placed in a hammock outside the incubator with the positioning in the crib also did not identify any changes in SpO₂.²¹ Those results corroborate our findings. Still, Zanardo et al.²² evaluated Hp in oxygen-dependent preterms with bronchopulmonary dysplasia and observed a worsening of SpO₂ after Hp. The authors suggest that these group of newborns may present severe hypoxemia during Hp. The preterms who were under ventilatory support in our study maintained the same parameters without adjustment needs. We did not observe any adverse events such as apneas or desaturation during Hp as it has been observed in other studies.^{10,11} We suggest as used in our study the use of a fabric placed between the cervical and scapular region of the preterms in order to reduce cervical hyperflexion, minimizing the risk of apnea and SpO₂ fall.

Despite the absence of complications during the positioning in our study, the indication of Hp needs to be individualized and used in a careful way especially with low birth weight preterms. Newborns may regurgitate during Hp if they are placed in the hammock soon after the diet or if they have major gastroesophageal reflux. Moreover, the fixation of the hammock needs to be done safely so there is no risk of instability or fall of the preterms, thus providing adequate containment.

The absence of a control group may be considered as a limitation of our study. In addition, it would not be possible to blind the investigators to the intervention, but we kept them unaware of the purpose of the study. One possible bias is a subjective assessment of pain and NBAS scale, which evaluates facial expressions and behavioral status of the newborns. The fact of the data collection was not done at the same period of the day in all participants could be another bias. Future randomized controlled trials with a larger sample need to be conducted with the purpose of increasing the external validity of our findings, as well as monitoring newborns for more than one hour to comprehend how vital signs behave after this period.

From our findings, Hp may be indicated as a safe, simple, and low-cost therapeutic option for the humanization of NICUs. Non-pharmacological and low-risk measures that promote the reduction of stress levels and pain in very low birth weight preterm newborns should be stimulated in NICUs.

Conclusion

Hammock positioning seems to induce sleep, maintained the vital signs within the normal range, and did not cause pain in very low birth weight preterm newborns. Hp may be considered as a safe positioning method and a non-pharmacological therapeutic measure for the reduction of stress levels, as well as for the humanization of NICUs. Further randomized controlled trials, with a larger sample and longer follow-up, are needed to evaluate the long-term outcomes of hammock positioning especially concerning neuromotor development.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We are thankful to the multi-professional team of Hospital Sofia Feldman, especially to the physical therapy and nurse team, for the help to conduct this research.

References

1. Gomes NRR, Monteiro RCS. As implicações do uso da "redinha" por bebês prematuros: uma revisão de literatura. *Rev Ciênc Saúde*. 2014;16:94–97.
2. Walter-Nicolet E, Annequin D, Biran V, Mitanchez D, Tourniaire B. Pain management in newborns: from prevention to treatment. *Paediatr Drugs*. 2010;12(6):353–365.
3. Datar A, Jacknowitz A. Birth weight effects on children's mental, motor, and physical development: evidence from twins data. *Matern Child Health J*. 2009;13(6):780–794.
4. Monterosso L, Kristjanson L, Cole J. Neuromotor development and the physiologic effects of positioning in very low birth weight infants. *J Obstet Gynecol Neonatal Nurs*. 2002;31(2):138–146.
5. Yecco GJ. Neurobehavioral development and developmental support of premature infants. *J Perinat Neonatal Nurs*. 1993;7(1):56–65.
6. Bezerra IFD, Torres VB, Lopes JM, Baroni MP, Pereira SA. Assessment of the influence of the hammock on neuromotor

- development in nursing full-term infants. *Rev Bras Crescimento Desenvol Hum*. 2014;24(1):106–111.
7. Lino LH, Coelho PG, Fonseca FLA, Filipini R. Os benefícios da rede de balanço em incubadoras utilizadas em recém-nascidos na UTI neonatal: uma estratégia de humanização. *Rev Enfermagem Revista*. 2015;18(1):88–100.
 8. Lekskulchai R, Cole J. Effect of a developmental program on motor performance in infants born preterm. *Aust J Physiother*. 2001;47(3):169–176.
 9. Guerin IO, Birck MA. A utilização de redes de balanço em incubadoras para recém-nascidos prematuros internados em unidades de terapia intensiva neonatais como um método de humanização: uma revisão de literatura. *Salão do Conhecimento UNIJUL*. Seminário de Iniciação Científica – Ciências da Saúde; 2016.
 10. Keller A, Arbel N, Merlob P, Davidson S. Neurobehavioral and autonomic effects of hammock positioning in infants with very low birth weight. *Pediatr Phys Ther*. 2003;15(1):3–7.
 11. Costa KSF, Beleza LdO, Souza LM, Ribeiro LM. Rede de descanso e ninho: comparação entre efeitos fisiológicos e comportamentais em prematuros. *Rev Gaúcha Enferm*. 2016;37(esp):e62554.
 12. Bortoli AFC, Taglietti M. Efetividade do método hammock nas características pulmonares e maturidade neuromuscular em recém-nascidos internados em unidades de terapias intensivas neonatais: Revisão sistemática da literatura. *FIEP Bull*. 2015;85.
 13. Grunau RE, Oberlander T, Holsti L, Whitfield MF. Bedside application of the Neonatal Facial Coding System in pain assessment of premature neonates. *Pain*. 1998;76(3):277–286.
 14. Melo Gmd, Lélis ALPdA, Moura AFd, Cardoso MVLML, Silva VMd. Escalas de avaliação de dor em recém-nascidos: revisão integrativa. *Rev Paul Pediatr*. 2014;32(4):395–402.
 15. Als H, Tronick E, Lester BM, Brazelton TB. The Brazelton Neonatal Behavioral Assessment Scale (NBAS). *J Abnorm Child Psychol*. 1977;5(3):215–231.
 16. Samra NM, Taweel AE, Cadwell K. Effect of intermittent kangaroo mother care on weight gain of low birth weight neonates with delayed weight gain. *J Perinat Educ*. 2013;22(4):194–200.
 17. Cavalaria SVF. *A terapia ocupacional utilizando redinhas no atendimento de recém-nascidos na uti-neonatal*. Lins – SP: Centro Universitário Católico Salesiano Auxilium; 2009.
 18. Bottos M, Pettenazzo A, Giancola G, et al. The effect of a 'containing' position in a hammock versus the supine position on the cutaneous oxygen level in premature and term babies. *Early Hum Dev*. 1985;11(3–4):265–273.
 19. Albuquerque TM, Albuquerque RC. Estratégias de posicionamento e contenção de recém-nascido pré-termo utilizadas em unidades de terapia intensiva neonatal. *Rev Interinst Bras Ter Ocup*. 2017;1(1):40–51.
 20. Costa DG, Moraes LBA, Nascimento IM. *Estudo comparativo de prematuros posicionados em Hammock (Redinhas) e decúbito ventral*; 2004 <http://www.interfisio.com.br/?artigo&ID=153>, 05.03.17.
 21. Chiu K, Tonkin SL, Gunn AJ, McIntosh CC. Are baby hammocks safe for sleeping babies? A randomised controlled trial. *Acta Paediatr*. 2014;103(7):783–787.
 22. Zanardo V, Trevisanuto D, Dani C, Bottos M, Guglielmi A, Cantarutti F. Oxygen saturation in premature neonates with bronchopulmonary dysplasia in a hammock. *Biol Neonate*. 1995;67(1):54–58.