REVIEW

For reprint orders, please contact: reprints@futuremedicine.com

Dysregulated fatty acid metabolism in hepatocellular carcinoma

Hepatic Oncology

Mingda Wang^{†,1}, Jun Han^{†,1}, Hao Xing^{†,1}, Han Zhang¹, Zhenli Li¹, Lei Liang^{1,2},
Chao Li¹, Shuyang Dai¹, Mengchao Wu¹, Feng Shen^{**1} & Tian Yang^{*1}

Hepatocellular carcinoma (HCC) is one of the most frequent and deadly malignancies worldwide. Studies are urgently needed on its molecular pathogenesis and biological characteristics. Dysregulation of fatty acid (FA) metabolism, in which aberrant activation of oncogenic signaling pathways alters the expression and activity of lipid-metabolizing enzymes, is an emerging hallmark of cancer cells, and it may be involved in HCC development and progression. The current review summarizes what is known about dysregulated FA metabolism in HCC and pathways through which this dysregulation may regulate HCC survival and growth. Our understanding of dysregulated FA metabolism and associated signaling pathways may contribute to the development of novel and efficient antitumor approaches for patients with HCC.

First draft submitted: 12 December 2016; Accepted for publication: 17 February 2017;
Published online: 30 June 2017

Practice points

- Dysregulation of fatty acid (FA) metabolism has increasingly emerged as a hallmark of cancer cells. Aberrant activation of oncogenic signaling pathways also regulates the expression and activity of lipid-metabolizing enzymes, thus reprogramming FA metabolism to promote hepatocellular carcinoma (HCC) development and progression.
- Intracellular FAs are required for biosynthesis of most biological membrane lipids, signaling molecules and post-translational modifications of proteins, and are also used to provide energy to support cancer cells survival and proliferation, when necessary, through β -oxidation process.
- HCC cells can employ appropriate metabolic pathways as different situation demands.
- Intrahepatic cholangiocarcinoma and HCC exhibits differential requirement for *de novo* lipogenesis and distinct response to therapeutic approaches focusing on inhibition of exogenous FA uptake.
- Besides the heightened *de novo* FA biosynthesis, certain cancer cells can also utilize lipolytic pathway to fuel stored fats and acquire free FAs that support tumor growth.
- Nonalcoholic fatty liver disease is characterized by abnormal fat accumulation in hepatic tissues with a number of risk factors regarding metabolic alterations, such as obesity, diabetes and dyslipidemia.

¹Department of Hepatic Surgery, Eastern Hepatobiliary Surgery Hospital, The Second Military Medical University, Shanghai, China

²Department of Hepatic Surgery, The First Affiliated Hospital of Soochow University, Medical College of Soochow University, Suzhou, China

*Author for correspondence: yangtian6666@hotmail.com;

**Author for correspondence: shenfengehbh@sina.com

[†]Co-first authors

KEYWORDS

• carcinogenesis • fatty acid oxidation • fatty acid synthesis • fatty acid uptake • hepatitis C virus • hepatocellular carcinoma • lipolytic pathway • nonalcoholic fatty liver disease

Liver cancer is among the most frequent types of cancer; it ranks as the second leading cause of cancer-related death worldwide, causing nearly 818,000 deaths in 2013 [1,2]. The incidence of liver cancer has increased from 465,000 cases in 1990 to 792,000 cases in 2013, becoming a major public health concern, especially in developing regions [1]. The most common form of liver malignancy is hepatocellular carcinoma (HCC), which accounts for more than 90% of liver cancer cases. HCC is generally a fatal disease, and it is usually diagnosed too late for patients to be eligible for potentially curative therapies such as surgical resection and liver transplantation [3–5]. Noncurative therapies such as transcatheter arterial chemoembolization show limited efficacy in improving overall survival, in part because advanced HCC is quite heterogeneous in clinical manifestations, radiological presentation and biological features [6,7]. While molecular targeted therapies such as sorafenib will undoubtedly improve the treatment of advanced HCC [8], the emergence of chemoresistance renders their clinical benefits temporary and unsatisfactory [8,9]. Developing effective therapeutic strategies for HCCs requires improving our understanding of molecular mechanisms involved in HCC progression.

The rapid, uncontrolled proliferation of malignant cells requires a constant supply of energy and macromolecular building blocks. To meet these demands, cancer cells reprogram their metabolic pathways and show abnormal metabolic alterations [10–13]. Indeed, such metabolic reprogramming has been firmly established as a hallmark of cancer cells [14]. The best understood metabolic alteration in cancer cells, discovered by Otto Warburg in the 1920s and named the ‘Warburg effect’ [15], is enhanced glucose uptake and lactate production through glycolytic pathways, even in conditions of normal oxygen tension.

In addition to changes in glucose metabolism, another defining metabolic alteration in cancer cells is dysregulated fatty acid (FA) metabolism [16,17]. Several reviews have already summarized the fundamental importance of reprogrammed lipid metabolism for cancer cells. Intracellular FAs are important to cancer cells, as well as to normal cells, because they serve as biosynthetic precursors of membrane lipids, signaling molecules and modifying groups that are added to proteins post-translationally (Figure 1). In addition, cancer cells can oxidize FAs to

obtain energy and building blocks. Mounting evidence indicates that aberrant activation of *de novo* FA synthesis contributes to tumor initiation and progression [16,18,19]; this elevated *de novo* synthesis in tumor cells occurs independently of exogenous lipids. Another aspect of dysregulated FA metabolism in various malignancies is dysregulated FA oxidation, also known as β -oxidation. In fact, targeting such oxidation may be an effective antitumor approach [20].

Studies of lipid metabolism in cancer have focused primarily on a few, well-known tumor types such as breast and prostate tumors. Nevertheless, several studies have already established the significant contribution of dysregulated FA metabolism to HCC, and those studies are reviewed here. We also discuss how this dysregulation likely supports HCC progression, and how it may be targeted therapeutically to benefit HCC patients.

De novo FA synthesis in HCC

Compelling evidence identifies enhanced *de novo* FA synthesis as an important feature of various tumor types [16]. This aberrantly upregulated lipogenesis mainly involves significantly increased expression and activity of major lipogenic enzymes, including FASN, ACC, SCD and ACSS [16,21,22] (Figure 2).

FASN is the central enzyme that catalyzes the committed step in *de novo* FA synthesis [16], and it is upregulated in multiple tumor types, where its degree of upregulation correlates closely with tumor aggressiveness [16,23]. The mRNA levels of *FASN*, *ACC*, *ACLY*, and *SCD1* were found to be elevated in a collection of HCC specimens [24]. SREBP1, which transcriptionally activates various enzymes involved in FA biosynthesis [25], is upregulated in HCC tissues, and the level of upregulation correlates with patient outcomes [26]. Aberrant activation of the Akt/mTORC1 inhibits post-transcriptional degradation of *FASN*, *SREBP1* and *SREBP2*, thereby enhancing *de novo* FA synthesis and promoting HCC development [27]. Conversely, genetic depletion of FASN in HCC cell lines *in vitro* strongly arrests growth, promotes apoptosis and decreases FA synthesis. The oncogenic potential of FASN has been confirmed *in vivo* using a mouse model of Akt-driven HCC [28]. In that work, FASN silencing completely impaired HCC carcinogenesis in Akt-overexpressing mice, raising the possibility that blocking FASN may specifically target HCC cells with an activated Akt/mTOR pathway.

Figure 1. Fatty acid metabolism in hepatocellular carcinoma development and progression.

Fatty acids (FAs) can promote various aspects of HCC tumorigenesis. For instance, FAs provide HCC cells with membrane building blocks, signaling molecules and post-translational modifications of proteins. FAs also serve as an energy source to support rapid cell proliferation and survival.

HCC: Hepatocellular carcinoma.

The range of proteins that can trigger *de novo* FA synthesis in HCC is likely to be more extensive. For example, work from our laboratory has shown that forced expression of ACC1 and the resulting *de novo* FA synthesis are required for survival and growth of HCC cells [29]. In fact, ACC1-mediated lipogenesis is associated with cancer aggressiveness and poor prognosis of HCC patients. Dysregulation of the oncogene encoding CD147, a transmembrane glycoprotein frequently expressed in HCCs, can contribute to metabolic reprogramming that helps drive carcinogenesis. CD147 enhances *de novo* FA synthesis, mainly by activating Akt/mTOR signaling, which in turn activates SREBP1c-mediated transcription of lipogenic enzymes such as *FASN* and *ACC* [30]. This is analogous to other examples of oncogene dysregulation that trigger cancer-causing metabolic reprogramming [31,32]. In

HCC-related hepatitis B virus (HBV) infection, the HBV X protein (HBx) induces expression of the liver X receptor and its downstream lipogenic target genes, thereby accelerating lipogenesis and HCC development [33]. HBx-induced *de novo* lipogenesis and HCC development may also involve the Ras family oncogene Rab18 [34]: in this model, HBx drives pathways related to COX-2 and miR-429 to upregulate Rab18 expression, which then increases lipid anabolism [35]. Chronic hepatitis C virus (HCV) infection has become another main cause of HCC [36]. The global population of people infected with HCV has reached an estimated 185 million in 2005, leading to an increasing incidence of HCV-related HCC worldwide [37]. HCV infection is characteristic of disturbed lipid metabolism as HCV induces lipogenesis and steatosis to provide a lipid-rich condition

Figure 2. Overview of fatty acid metabolism in hepatocellular carcinoma cells. This scheme depicts the main regulatory interactions between FA anabolism and catabolism in hepatocellular carcinoma (HCC) cells. See the main text for detailed description of the metabolic pathways shown. Enzymes are in oval boxes with red background. Malonyl-CoA inhibits the activity of CPT, which controls the entry and oxidation of FAs, thus preventing FAO.

CPT: Carnitine palmitoyltransferase; FA: Fatty acid; FAO: Fatty acid oxidation; GLUT: Glucose transporter; HK: Hexokinase; PFK: Phosphofructokinase; PKM: Pyruvate kinase; TCA: Tricarboxylic acid cycle.

to enhance its replication [38]. The expression of HCV core proteins has been proposed for the pathogenesis of hepatic steatosis and HCC formation in transgenic animal models [39]. Further investigations reveal that HCV proteins, including core and nonstructural protein NS4B, upregulates FA synthesis by enhancing the expression and post-translational activation of SREBPs in a proteolytic-dependent pathway [40]. Besides, HCV-induced oxidative stress and the following activation of AKT pathway, or inactivation of phosphatase and tensin homologue (PTEN) could also activate SREBPs expression [40]. Other experimental evidence suggests HCV infection may induce hepatocarcinogenesis by regulating nuclear receptor pathways that facilitate malignant transformation of hepatocytes. Conti *et al.* observes that HCV-stimulated upregulation of SREBP1c and FASN promotes cellular lipid

biosynthesis and transcriptional activates the expression of nuclear receptor member Small heterodimer partner 1 (SHP1), which in turn directly interacts with HCV NS5A viral protein and thereby causes deregulation of several epithelial–mesenchymal transition (EMT) related proteins, thus favoring liver tumorigenesis [41]. These findings highlight the importance of enhanced FA synthesis in HCV-driven HCC carcinogenesis, and imply that interfering with lipogenesis may represent potential therapeutic strategy for HCV-related HCCs.

Potential metabolic biomarkers of HCC have emerged from metabolomic analysis of lipid metabolites and transcriptomic analysis of gene expression patterns in HCCs [26,42–46]. For example, mitochondrial acetate may be one such biomarker [45]: ACSS1 is upregulated in HCC, and this enzyme converts acetate to acetyl-CoA. Since HCC patients can differ substantially in levels of ACSS1 expression, stratifying them based on this expression may prove useful for disease treatment and management. SCD, a key enzyme in FA synthesis, as well as related metabolites may be biomarkers of HCC aggressiveness [42]. It may be that dysregulation of SCD activity causes an imbalance in cellular levels of its substrate, saturated palmitic acid and of its product, monounsaturated palmitic acid [47]. Such an imbalance may contribute to HCC progression [42]. Modulating SCD activity may be clinically useful for treating aggressive HCC.

FA oxidation & HCC tumorigenesis

Like heightened *de novo* FA biosynthesis, FA oxidation also contributes to HCC tumorigenesis, though studies seem to disagree on whether up- or downregulation of such oxidation is involved. FA oxidation shortens FAs by two carbons per reaction cycle, and it generates NADH and FADH₂ for ATP production [20]. While some studies have associated downregulation of FA oxidation with HCC, the expression levels of most FA oxidation-related genes vary greatly among patients [45]. Other work has associated increased catabolism of certain saturated lipids with high α -fetoprotein levels in the serum of HCC patients [43,48]. This discrepancy may reflect, at least in part, the tumor heterogeneity of HCC. Whatever the precise nature of the association between lipid catabolism and HCC, the evidence points to the possibility of targeting lipid catabolic pathways to treat the disease, at least in patients with high α -fetoprotein levels.

Key rate-limiting enzymes in lipid catabolism, such as CPT1 and ACOX1, are involved in cancer progression [49,50], and the same appears to be true in HCC. For example, the transmembrane glycoprotein CD147 activates p38/MARK signaling to downregulate PPAR α , leading in turn to downregulation of both CPT1 and ACOX1 and ultimately inhibiting FA oxidation [30]. In a mouse model of HCC, ACOX1 deficiency is associated with continuous PPAR α activation and subsequent endoplasmic reticulum stress associated with the unfolded protein response; this may contribute to HCC initiation and progression [50]. Therefore, these studies suggest that inhibition of FA oxidation is associated with HCC development.

Pursuing this relationship seems reasonable because it aligns with the fact that solid tumor cells are often short of oxygen and fundamental nutrients because of uncontrolled malignant proliferation and lack of vascularization [51,52]. Under such circumstances, FA oxidation can promote cancer cell survival by maintaining NADPH levels and generating ATP [53–55]. Lu *et al.* characterized the contribution of FAO in maintaining HCC cells survival during energy starvation [56]. They found that FA oxidation is activated in HCC cells to provide energy and substrates via a process involving autophagy activated by C/EBP α , thereby enabling HCC cells to resist energy deprivation. Jeon *et al.* highlighted the regulatory role of AMPK in activating FA oxidation pathways in response to metabolic stress [55]. Our own group has shown that HBx acts via a calcium/CaMKK-dependent pathway to activate AMPK and thereby FA oxidation when nutrients are scarce; etomoxir inhibition of FA oxidation reverses this HBx-mediated resistance to nutrient deficiency [57]. Since chronic HBV infection is the major risk factor for HCC in Asian countries [58], and HBx has been shown to participate in the pathogenesis of HBV-related HCC [59], these data raise the possibility of blocking FA oxidation in order to treat HBV-associated HCC. In addition, the enzyme ACC plays a particularly interesting role in regulating FA oxidation to protect HCC cells against metabolic stress. When nutrients are abundant, ACC forms a complex with the enzyme CPT1 in HCC cells. When nutrients are scarce, however, AMPK phosphorylates ACC, weakening its association with CPT1 [29], which can then dissociate and travel to the mitochondrial membrane where it promotes FA transport

to fuel FA oxidation. This model presents the key factor or enzyme-mediated metabolic switch from lipid anabolism to catabolism in cancer cells under nutritional stress. Another study conducted by Zhang *et al.* has demonstrated that hypoxia-induced HIF-1 expression orchestrates glucose and lipid metabolism to maintain cell survival and fulfill growth demands [60]. Their findings indicate that upregulation of HIF-1 α suppresses FA oxidation by disturbing the expression of medium- and long-chain acyl-CoA dehydrogenases (MCAD and LCAD), two rate-limiting enzymes that catalyze the first step of oxidation in mitochondria. This attenuated FA oxidation process further facilitates HCC progression by reducing ROS generation, promoting Gluts-driven glycolysis and activating cell survival-related pathways. Therefore, it is intriguing to reveal that, although enhanced FA oxidation has no impact on cancer cell proliferation under normoxic conditions, it is not beneficial for cell survival under hypoxic environments. Instead, the cells may employ appropriate metabolic pathways aimed at meeting different situation demands.

Since liver cancer stem cells are thought to give rise to HCC, it would make sense that their FA metabolism should also be dysregulated. Indeed, leukemia-initiating cells and hematopoietic stem cells depend on activation of FA oxidation [61,62]. In liver cancer stem cells, the stem cell marker NANOG activates FA oxidation by regulating the transcription of several mitochondrial genes (*Acadvl*, *Esch1* and *Acads*), helping to meet energy demands and support stem cell-like properties [63]. Consistent with a key role for FA oxidation in HCC, chemically inhibiting FA oxidation renders NANOG-positive cancer stem cells more sensitive to sorafenib. This may be an interesting strategy to eliminate liver cancer stem cells and sensitize HCC to chemotherapy.

The lipolytic pathway in HCC

Although *de novo* FA synthesis has traditionally been regarded as the main source of FAs for most tumors, recent studies show that certain cancer cells can use the lipolytic pathway to generate free FAs from stored lipids in order to support tumor growth [64,65] (Figure 2). In this respect, levels of the lipolytic enzyme MAGL may be important in HCC. MAGL hydrolyzes monoacylglycerols into free FAs and glycerols and it releases FAs from stored lipids, thereby promoting cancer progression [64]. In

HCC, aberrant promoter methylation induces expression of the gene encoding *LATS1*, which enhances dephosphorylation and nuclear transport of YAP, inducing MAGL expression [66]. LPL, widely expressed in adipocytes and muscle tissues, is frequently upregulated in HCC and mouse models of the disease [67]. It participates in the hydrolysis of triglycerides, and it promotes uptake of lipoproteins into cells [68,69]. High LPL expression is closely linked to aggressive tumor features and poor patient outcome in HCC [67]. This raises the possibility of inhibiting LPL to treat HCC, such as by upregulating miR-29, which represses LPL expression [70]. In fact, administration of miR-29 mimic oligonucleotide is of potential value to develop novel therapies for HCC patients [71]. Consistently, overexpressing miR-29 in a mouse model of liver cancer repressed tumorigenesis driven by c-Myc or Akt/Ras [72]. It may be helpful to inhibit both LPL and FASN simultaneously and thereby repress both lipogenic and lipolytic pathways in order to treat HCC [67].

Uptake & transport of exogenous FAs in HCC

A third source of FAs for mammalian cells, in addition to the *de novo* and lipolytic pathways described above, is the external environment (Figure 1). Although most cancer cells exhibit a metabolic shift toward lipogenesis and synthesize nearly all esterified FAs *de novo*, some tumors tend to ‘acquire’ free FAs directly from the external environment. This makes pathways of FA uptake and transport promising therapeutic targets [73–75]. Studies aimed at exploiting lipid metabolism in HCC for therapeutic benefit have focused primarily on inhibiting *de novo* FA synthesis, neglecting possible targets in FA uptake pathways.

External FAs are brought into the cell not by passive diffusion but through a transport mechanism involving specialized enzymes and proteins [76]. FA translocase (FAT, or CD36) is expressed at low levels in normal hepatocytes, but it mediates FA uptake in a variety of malignancies [77,78]. Fatty acid transport proteins FATP2 and FATP5, as well as members of the fatty acid binding-protein family (FABP1, FABP4 and FABP5) are also involved in FA uptake and transport in hepatic tissues [79–81]. Increased expression of CD36 leads to higher FA uptake, which is closely related to induction of the EMT in HCC [79]. The EMT may

also be promoted when the elevated FA levels of HCC patients upregulate inflammation-related oncogenic transcriptional factors (*NF-κB*, *AP-1*, *STAT3* and *HIF-1α*), which activate Wnt and TGF-β signaling pathways [79,82,83]. Given the demand for novel therapeutic targets involved in lipid uptake and transport in HCC, more studies are needed that focus on CD36-mediated FA accumulation.

While inhibiting FASN may make sense for HCC, it does not appear to be suitable for intrahepatic cholangiocarcinoma (ICC), which ranks as the second most prevalent type of primary liver malignancy after HCC and is characterized by highly aggressive features and limited treatment options [84–86]. In different mouse cancer models [87,88], depletion of FASN significantly suppressed HCC development, but did not affect oncogene-driven ICC formation. This likely reflects the fact that ICC features different metabolic requirements from HCCs. ICC cells rely mainly on uptake of exogenous FA rather than *de novo* FA biosynthesis in order to survive and proliferate [89]. ICC cells show higher expression of proteins related to FA uptake and transport, but lower levels of FASN, than do HCC cells. Thus, targeting FASN and associated lipogenesis may not be appropriated for ICC (Figure 3).

Nonalcoholic fatty liver disease & HCC development

Nonalcoholic fatty liver disease (NAFLD) is characterized by an accumulation of fat in the hepatocytes in the absence of obvious alcohol uptake, hereditary disorders or medication history [90]. It comprises a wide spectrum of liver metabolic abnormalities ranging from simple fatty liver to nonalcoholic steatohepatitis (NASH), which can progress into irreversible cirrhosis and ultimately HCC [90]. Epidemiologic data suggest that NAFLD is the most frequent type of chronic liver disease in Western countries, accounting for an estimated prevalence of 20–40% [91,92]. Currently, a close link between NAFLD and clinical features of metabolic syndromes has been widely recognized. Among these syndromes, obesity, Type 2 diabetes mellitus, insulin resistance and dyslipidemia are the most common and well-documented risk factors related to NAFLD [93,94]. According to several population-based studies, obesity and Type 2 diabetes mellitus have increasingly emerged as two major factors responsible for the recent rise in prevalent of HCC worldwide [95,96]. In a

retrospective study including more than 255,702 adults in the USA, it is observed that individuals with obesity and/or diabetes exhibit a 2.47-fold increase in relative risk of HCC, while eliminating obesity/diabetes reduces HCC incidence more than disturbing any other risk factors [97]. Growing evidence also consistently reports a significantly increased risk of HCC occurrence in patients with NAFLD [98,99]. On this condition, the incidence of NAFLD-associated HCC is likely to increase over the next few decades. Of note, several studies have all unveiled much higher risk of HCC carcinogenesis in obese men [100,101]. More recently, a multi-ethnic cohort study including Japanese, white and Latino men further confirms the dramatic differences in HCC incidence modulated by gender disparities [102]. In this scenario, the adipose homeostasis, especially the content of visceral adipose tissue, is involved in the occurrence of metabolic syndromes and its related HCC. It has been proven that visceral adiposity is capable to predict the prognosis of HCC [102]. Previous studies confirm that the accumulation of visceral fat induces insulin resistance and further damage to hepatocytes. The subsequently dysregulated release of free FAs from visceral adipocytes can also lead to the onset of HCC [103]. Considering that men possess approximately 30% more visceral fat than women [104], it is plausible to speculate the strong association between the abnormal lipid metabolism and gender disparities in HCC.

Although great amount of studies have identified the clinical relevance among metabolic syndromes, NAFLD/NASH and HCC development, the exact molecular mechanisms underlying NAFLD/NASH-related HCC carcinogenesis is extremely complicated and remains elusive. The vast majority of current studies have been limited to a handful of well-known metabolic disorders that promote the initiation of HCC, such as lipotoxicity, adipose-derived inflammatory response and insulin resistance [105]. From the perspective of cellular metabolism, mitochondria are regarded as metabolic hub coordinating the cellular anabolic and catabolic processes to maintain energetic homeostasis and meet biosynthesis demand of the cell. It is well documented that nuclear receptor PPARs, the critical modulators of glucose and lipid metabolism in liver, are involved in the pathogenesis of both metabolic abnormalities and malignant neoplasm. Mello *et al.* have

Figure 3. Differences in fatty acid metabolism in liver cancers. HCC and ICC differ greatly in FA metabolic pathways, suggesting the need for fundamentally different therapeutic approaches. Unlike HCC and many other cancers, ICC depends strongly on uptake of exogenous FAs. Therefore, enhanced *de novo* FA synthesis with a concomitant inhibition of FA uptake is not a universal characteristic of liver neoplasm. Stratifying liver cancer patients based on whether FA uptake or *de novo* FA synthesis is the main driver of lipid metabolism in a given patient may be helpful for developing individualized therapies.

FA: Fatty acid; HCC: Hepatocellular carcinoma; ICC: Intrahepatic carcinoma.

summarized the functional roles of PPARs in regulating liver mitochondrial metabolism from NAFLD to HCC [106]: altered PPARs expression mainly induces metabolic dysfunctions in the mitochondria, leading to inhibition of FA oxidation, accumulation of ROS and promotion of *de novo* lipogenesis. This remodeling of cellular metabolism will contribute to the activation of survival/grow pathways and eventually, cell malignant transformation and tumor

promotion. Therefore, unraveling specific metabolic reprogramming accelerating HCC carcinogenesis may shed new light on our current understanding of intricate relationship between NAFLD and HCC development.

Conclusion

Dysregulated FA metabolism in cancer cells have been increasingly recognized in recent years. Generally, cancer cells rely on FAs as cellular building blocks for the generation of membrane structures, the production of signaling molecules, and energy storage. Here, we summarize the characteristics of FA metabolism in liver cancer cells, mainly focusing on the pathways of FA synthesis, oxidation, uptake and transport in HCCs. Furthermore, we also provide a brief review of the relationship between NAFLD and HCC development. Since FA metabolism is complex with various feedback mechanisms and can be regulated by a panel of rate-limiting enzymes, effective therapeutic strategies may be developed based on full understanding of specific metabolic features for a certain cancer type.

Future perspective

HCC, as well as other liver diseases, involves dysregulation of FA metabolism because the liver is the chief metabolizer of fats, carbohydrates and proteins. HCC cells, like other cancer cells, rely heavily on FAs to provide material for cellular membranes, signaling molecules and energy as they proliferate and spread. Many cancer cells meet their demands for energy and building materials by upregulating *de novo* FA synthesis [17], leading researchers to target key lipogenic enzymes, particularly FASN [23], to slow tumor growth. Several chemical inhibitors targeting other key enzymes and pathways in lipid metabolism also show good efficacy against various cancers, but developing inhibitors against HCC lags behind. One reason is that the genotypic and tumor-biological diversity of HCC patients makes it difficult to stratify them based on lipid metabolism. Transcriptomic and metabolomic approaches may help define distinct metabolic subgroups

with similar alterations in FA metabolism, which in turn may lead to the refinement of treatment strategies targeting FA-related pathways. A second reason why the search for FA-related treatments lags for HCC is that FA metabolism is complex, involving a dynamic balance between FA anabolism and catabolism. For instance, acetyl-CoA produced from FA oxidation can be used to synthesize FAs, whereas malonyl-CoA, an intermediate generated during FA synthesis, inhibits FA oxidation (Figure 2). This tight interweaving of positive and negative regulation means that an inhibitor may activate alternative FA metabolic pathways, thereby accelerating HCC progression. Despite these challenges, further work into HCC treatments that exploit FA metabolic pathways is needed. To ensure clinical relevance, these studies should consider pharmacokinetics, selectivity and adverse effects.

The growing literature on FA metabolism in HCC pathogenesis has uncovered novel and potential intervention targets, as well as broadened our understanding of lipid metabolism in the disease. Given the extensive heterogeneity of FA metabolism in HCC cells, therapies may need to target two or more aspects of FA metabolism in order to be effective, or drugs that target FA metabolism may need to be combined with other types of chemotherapy. From this perspective, studies investigating the importance of FA metabolism in HCC may help shift the therapeutic focus from traditional monotherapies to combination approaches.

Financial & competing interests disclosure

This work was supported in part by the National Natural Science Foundation of China (81472284, 81672699), the Shanghai Pujiang Program (16PJD004), the Shanghai Sailing Program (17YF1424900) and the Shanghai Young Talent Program. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

References

Papers of special note have been highlighted as:
• of interest; •• of considerable interest

- 1 Global Burden of Disease Cancer Collaboration, Fitzmaurice C, Dicker D, Pain A *et al.* The Global Burden of Cancer 2013.

- 2 Wang H, Liddell CA, Coates MM *et al.* Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 384(9947), 957–979 (2014).

- 3 Llovet JM, Burroughs A, Bruix J. Hepatocellular carcinoma. *Lancet* 362(9399), 1907–1917 (2003).
- 4 Llovet JM, Bruix J. Molecular targeted therapies in hepatocellular carcinoma. *Hepatology* 48(4), 1312–1327 (2008).

- 5 Carr BI. Hepatocellular carcinoma: current management and future trends. *Gastroenterology* 127(5 Suppl 1), S218–224 (2004).
- 6 Lu LC, Hsu CH, Hsu C, Cheng AL. Tumor heterogeneity in hepatocellular carcinoma: facing the challenges. *Liver Cancer* 5(2), 128–138 (2016).
- 7 Jeng KS, Chang CF, Jeng WJ, Sheen IS, Jeng CJ. Heterogeneity of hepatocellular carcinoma contributes to cancer progression. *Crit. Rev. Oncol. Hematol.* 94(3), 337–347 (2015).
- 8 Llovet JM, Ricci S, Mazzaferro V *et al.* Sorafenib in advanced hepatocellular carcinoma. *N. Engl. J. Med.* 359(4), 378–390 (2008).
- 9 Cheng AL, Kang YK, Chen Z *et al.* Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol.* 10(1), 25–34 (2009).
- 10 Tennant DA, Duran RV, Gottlieb E. Targeting metabolic transformation for cancer therapy. *Nat. Rev. Cancer* 10(4), 267–277 (2010).
- 11 Cairns RA, Harris IS, Mak TW. Regulation of cancer cell metabolism. *Nat. Rev. Cancer* 11(2), 85–95 (2011).
- 12 Ward PS, Thompson CB. Metabolic reprogramming: a cancer hallmark even warburg did not anticipate. *Cancer Cell* 21(3), 297–308 (2012).
- 13 Schulze A, Harris AL. How cancer metabolism is tuned for proliferation and vulnerable to disruption. *Nature* 491(7424), 364–373 (2012).
- 14 Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 144(5), 646–674 (2011).
- The authors revisit, refine and extend the concept of cancer hallmarks, which has provided a useful conceptual framework for understanding the complex biology of cancer.
- 15 Warburg O. On the origin of cancer cells. *Science* 123(3191), 309–314 (1956).
- 16 Menendez JA, Lupu R. Fatty acid synthase and the lipogenic phenotype in cancer pathogenesis. *Nat. Rev. Cancer* 7(10), 763–777 (2007).
- 17 Currie E, Schulze A, Zechner R, Walther TC, Farese RV Jr. Cellular fatty acid metabolism and cancer. *Cell Metab.* 18(2), 153–161 (2013).
- Provides a view of cancer cell metabolism from a lipid perspective, and we summarize evidence that limiting fatty acid availability can control cancer cell proliferation.
- 18 Kuhajda FP, Pizer ES, Li JN, Mani NS, Frehywot GL, Townsend CA. Synthesis and antitumor activity of an inhibitor of fatty acid synthase. *Proc. Natl Acad. Sci. USA* 97(7), 3450–3454 (2000).
- 19 Zhou W, Han WF, Landree LE *et al.* Fatty acid synthase inhibition activates AMP-activated protein kinase in SKOV3 human ovarian cancer cells. *Cancer Res.* 67(7), 2964–2971 (2007).
- 20 Carracedo A, Cantley LC, Pandolfi PP. Cancer metabolism: fatty acid oxidation in the limelight. *Nat. Rev. Cancer* 13(4), 227–232 (2013).
- This Progress article brings to light the important contribution of fatty acid oxidation to cancer cell function.
- 21 Swinnen JV, Brusselmans K, Verhoeven G. Increased lipogenesis in cancer cells: new players, novel targets. *Curr. Opin. Clin. Nutr. Metab. Care* 9(4), 358–365 (2006).
- 22 Mashima T, Seimiya H, Tsuruo T. *De novo* fatty-acid synthesis and related pathways as molecular targets for cancer therapy. *Br. J. Cancer* 100(9), 1369–1372 (2009).
- 23 Flavin R, Peluso S, Nguyen PL, Loda M. Fatty acid synthase as a potential therapeutic target in cancer. *Future Oncol.* 6(4), 551–562 (2010).
- 24 Yahagi N, Shimano H, Hasegawa K *et al.* Co-ordinate activation of lipogenic enzymes in hepatocellular carcinoma. *Eur. J. Cancer* 41(9), 1316–1322 (2005).
- 25 Horton JD. Sterol regulatory element-binding proteins: transcriptional activators of lipid synthesis. *Biochem. Soc. Trans.* 30(Pt 6), 1091–1095 (2002).
- 26 Yamashita T, Honda M, Takatori H *et al.* Activation of lipogenic pathway correlates with cell proliferation and poor prognosis in hepatocellular carcinoma. *J. Hepatol.* 50(1), 100–110 (2009).
- 27 Calvisi DF, Wang C, Ho C *et al.* Increased lipogenesis, induced by AKT-mTORC1-RPS6 signaling, promotes development of human hepatocellular carcinoma. *Gastroenterology* 140(3), 1071–1083 (2011).
- *De novo* lipogenesis has pathogenic and prognostic significance for hepatocellular carcinoma. Inhibitors of lipogenic signaling, including those that inhibit the AKT pathway, might be useful as therapeutics for patients with liver cancer.
- 28 Li L, Pilo GM, Li X *et al.* Inactivation of fatty acid synthase impairs hepatocarcinogenesis driven by AKT in mice and humans. *J. Hepatol.* 64(2), 333–341 (2016).
- 29 Wang MD, Wu H, Fu GB *et al.* Acetyl-coenzyme A carboxylase α promotion of glucose-mediated fatty acid synthesis enhances survival of hepatocellular carcinoma in mice and patients. *Hepatology* 63(4), 1272–1286 (2016).
- Acetyl-CoA carboxylase (ACC) catalyzes the committed step of the fatty acid (FA) synthesis. This study suggests a key role of ACC1 in the integration of glucose and lipid metabolism, and first reports a novel mechanism by which ACC1 can regulate FA oxidation by forming complex with carnitine palmitoyltransferase (CPT) and affecting their mitochondria distribution.
- 30 Li J, Huang Q, Long X *et al.* CD147 reprograms fatty acid metabolism in hepatocellular carcinoma cells through Akt/mTOR/SREBP1c and P38/PPAR α pathways. *J. Hepatol.* 63(6), 1378–1389 (2015).
- 31 Levine AJ, Puzio-Kuter AM. The control of the metabolic switch in cancers by oncogenes and tumor suppressor genes. *Science* 330(6009), 1340–1344 (2010).
- 32 Dang CV. Links between metabolism and cancer. *Genes Dev.* 26(9), 877–890 (2012).
- 33 Na TY, Shin YK, Roh KJ *et al.* Liver X receptor mediates hepatitis B virus X protein-induced lipogenesis in hepatitis B virus-associated hepatocellular carcinoma. *Hepatology* 49(4), 1122–1131 (2009).
- 34 You X, Liu F, Zhang T, Li Y, Ye L, Zhang X. Hepatitis B virus X protein upregulates oncogene Rab18 to result in the dysregulation of lipogenesis and proliferation of hepatoma cells. *Carcinogenesis* 34(7), 1644–1652 (2013).
- 35 Pulido MR, Diaz-Ruiz A, Jimenez-Gomez Y *et al.* Rab18 dynamics in adipocytes in relation to lipogenesis, lipolysis and obesity. *PLoS One* 6(7), e22931 (2011).
- 36 Williams R. Global challenges in liver disease. *Hepatology* 44(3), 521–526 (2006).
- 37 Mohd Hanafiah K, Groeger J, Flaxman AD, Wiersma ST. Global epidemiology of hepatitis C virus infection: new estimates of age-specific antibody to HCV seroprevalence. *Hepatology* 57(4), 1333–1342 (2013).
- 38 Syed GH, Amako Y, Siddiqui A. Hepatitis C virus hijacks host lipid metabolism. *Trends Endocrinol. Metab.* 21(1), 33–40 (2010).
- 39 Moriya K, Fujie H, Shintani Y *et al.* The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice.

- Nat. Med.* 4(9), 1065–1067 (1998).
- 40 Waris G, Felmlee DJ, Negro F, Siddiqui A. Hepatitis C virus induces proteolytic cleavage of sterol regulatory element binding proteins and stimulates their phosphorylation via oxidative stress. *J. Virol.* 81(15), 8122–8130 (2007).
 - 41 Conti B, Porcu C, Viscomi C *et al.* Small heterodimer partner 1 directly interacts with NS5A viral protein and has a key role in HCV related liver cell transformation. *Oncotarget* 7(51), 84575–84586 (2016).
 - 42 Budhu A, Roessler S, Zhao X *et al.* Integrated metabolite and gene expression profiles identify lipid biomarkers associated with progression of hepatocellular carcinoma and patient outcomes. *Gastroenterology* 144(5), 1066–1075; e1061 (2013).
 - 43 Beyoglu D, Imbeaud S, Maurhofer O *et al.* Tissue metabolomics of hepatocellular carcinoma: tumor energy metabolism and the role of transcriptomic classification. *Hepatology* 58(1), 229–238 (2013).
 - 44 Liu Y, Hong Z, Tan G *et al.* NMR and LC/MS-based global metabolomics to identify serum biomarkers differentiating hepatocellular carcinoma from liver cirrhosis. *Int. J. Cancer* 135(3), 658–668 (2014).
 - 45 Bjornson E, Mukhopadhyay B, Asplund A *et al.* Stratification of hepatocellular carcinoma patients based on acetate utilization. *Cell Rep.* 13(9), 2014–2026 (2015).
 - 46 Fages A, Duarte-Salles T, Stepien M *et al.* Metabolomic profiles of hepatocellular carcinoma in a European prospective cohort. *BMC Med.* 13, 242 (2015).
 - 47 Ntambi JM, Miyazaki M. Regulation of stearoyl-CoA desaturases and role in metabolism. *Prog. Lipid Res.* 43(2), 91–104 (2004).
 - 48 Saito S, Ojima H, Ichikawa H, Hirohashi S, Kondo T. Molecular background of alpha-fetoprotein in liver cancer cells as revealed by global RNA expression analysis. *Cancer Sci.* 99(12), 2402–2409 (2008).
 - 49 Rotondo D, Davidson J. Genetics and molecular biology: fatty acid metabolism in cancer cell survival; carnitine palmitoyltransferase-1 as a critical anticancer target. *Curr. Opin. Lipidol.* 22(5), 428–429 (2011).
 - 50 Huang J, Viswakarma N, Yu S *et al.* Progressive endoplasmic reticulum stress contributes to hepatocarcinogenesis in fatty acyl-CoA oxidase 1-deficient mice. *Am. J. Pathol.* 179(2), 703–713 (2011).
 - 51 Hirayama A, Kami K, Sugimoto M *et al.* Quantitative metabolome profiling of colon and stomach cancer microenvironment by capillary electrophoresis time-of-flight mass spectrometry. *Cancer Res.* 69(11), 4918–4925 (2009).
 - 52 Gullino PM, Grantham FH, Courtney AH, Losonczy I. Relationship between oxygen and glucose consumption by transplanted tumors *in vivo*. *Cancer Res.* 27(6), 1041–1052 (1967).
 - 53 Schafer ZT, Grassian AR, Song L *et al.* Antioxidant and oncogene rescue of metabolic defects caused by loss of matrix attachment. *Nature* 461(7260), 109–113 (2009).
 - 54 Zaugg K, Yao Y, Reilly PT *et al.* Carnitine palmitoyltransferase 1C promotes cell survival and tumor growth under conditions of metabolic stress. *Genes Dev.* 25(10), 1041–1051 (2011).
 - 55 Jeon SM, Chandel NS, Hay N. AMPK regulates NADPH homeostasis to promote tumour cell survival during energy stress. *Nature* 485(7400), 661–665 (2012).
 - **AMP-activated protein kinase (AMPK) is an enzyme that plays a role in cellular energy homeostasis. NADPH is the reduced form of NADP⁺, which is a cofactor used in anabolic reactions such as lipid synthesis. This study reveals that AMPK has a key function in NADPH maintenance, which is critical for cancer cell survival under energy stress conditions.**
 - 56 Lu GD, Ang YH, Zhou J *et al.* CCAAT/enhancer binding protein alpha predicts poorer prognosis and prevents energy starvation-induced cell death in hepatocellular carcinoma. *Hepatology* 61(3), 965–978 (2015).
 - 57 Wang MD, Wu H, Huang S *et al.* HBx regulates fatty acid oxidation to promote hepatocellular carcinoma survival during metabolic stress. *Oncotarget* 7(6), 6711–6726 (2016).
 - 58 Brechot C, Pourcel C, Louise A, Rain B, Tiollais P. Presence of integrated hepatitis B virus DNA sequences in cellular DNA of human hepatocellular carcinoma. *Nature* 286(5772), 533–535 (1980).
 - 59 Feitelson MA, Duan LX. Hepatitis B virus X antigen in the pathogenesis of chronic infections and the development of hepatocellular carcinoma. *Am. J. Pathol.* 150(4), 1141–1157 (1997).
 - 60 Huang D, Li T, Li X *et al.* HIF-1-mediated suppression of acyl-CoA dehydrogenases and fatty acid oxidation is critical for cancer progression. *Cell Rep.* 8(6), 1930–1942 (2014).
 - 61 Samudio I, Harmancey R, Fiegl M *et al.* Pharmacologic inhibition of fatty acid oxidation sensitizes human leukemia cells to apoptosis induction. *J. Clin. Invest.* 120(1), 142–156 (2010).
 - 62 Ito K, Carracedo A, Weiss D *et al.* A PML-PPAR-delta pathway for fatty acid oxidation regulates hematopoietic stem cell maintenance. *Nature Med.* 18(9), 1350–1358 (2012).
 - 63 Chen CL, Uthaya Kumar DB, Punj V *et al.* NANOG metabolically reprograms tumor-initiating stem-like cells through tumorigenic changes in oxidative phosphorylation and fatty acid metabolism. *Cell Metab.* 23(1), 206–219 (2016).
 - 64 Nomura DK, Long JZ, Niessen S, Hoover HS, Ng SW, Cravatt BF. Monoacylglycerol lipase regulates a fatty acid network that promotes cancer pathogenesis. *Cell* 140(1), 49–61 (2010).
 - 65 Przybytkowski E, Joly E, Nolan CJ *et al.* Upregulation of cellular triacylglycerol - free fatty acid cycling by oleate is associated with long-term serum-free survival of human breast cancer cells. *Biochem. Cell Biol.* 85(3), 301–310 (2007).
 - 66 Zhang J, Liu Z, Lian Z *et al.* Monoacylglycerol lipase: a novel potential therapeutic target and prognostic indicator for hepatocellular carcinoma. *Sci. Rep.* 6, 35784 (2016).
 - 67 Cao D, Song X, Che L *et al.* Both *de novo* synthesized and exogenous fatty acids support the growth of hepatocellular carcinoma cells. *Liver Int.* 37(1), 80–89 (2016).
 - 68 Rozovski U, Grgurevic S, Bueso-Ramos C *et al.* Aberrant LPL expression, driven by STAT3, mediates free fatty acid metabolism in CLL cells. *Mol. Cancer Res.* 13(5), 944–953 (2015).
 - 69 Rinninger F, Kaiser T, Mann WA, Meyer N, Greten H, Beisiegel U. Lipoprotein lipase mediates an increase in the selective uptake of high density lipoprotein-associated cholesteryl esters by hepatic cells in culture. *J. Lipid Res.* 39(7), 1335–1348 (1998).
 - 70 Mattis AN, Song G, Hitchner K *et al.* A screen in mice uncovers repression of lipoprotein lipase by microRNA-29a as a mechanism for lipid distribution away from the liver. *Hepatology* 61(1), 141–152 (2015).
 - 71 Callegari E, Gramantieri L, Domenicali M, D'Abundo L, Sabbioni S, Negrini M. MicroRNAs in liver cancer: a model for investigating pathogenesis and novel therapeutic approaches. *Cell Death Differ.* 22(1), 46–57 (2015).
 - 72 Tao J, Ji J, Li X *et al.* Distinct anti-oncogenic effect of various microRNAs in different mouse models of liver cancer. *Oncotarget* 6(9), 6977–6988 (2015).
 - 73 Nieman KM, Kenny HA, Penicka CV *et al.* Adipocytes promote ovarian cancer metastasis

- and provide energy for rapid tumor growth. *Nature Med.* 17(11), 1498–1503 (2011).
- 74 Defilippis RA, Chang H, Dumont N *et al.* CD36 repression activates a multicellular stromal program shared by high mammographic density and tumor tissues. *Cancer Discov.* 2(9), 826–839 (2012).
 - 75 Liu Y, Zuckier LS, Ghesani NV. Dominant uptake of fatty acid over glucose by prostate cells: a potential new diagnostic and therapeutic approach. *Anticancer Res.* 30(2), 369–374 (2010).
 - 76 Stremmel W, Pohl L, Ring A, Herrmann T. A new concept of cellular uptake and intracellular trafficking of long-chain fatty acids. *Lipids* 36(9), 981–989 (2001).
 - 77 Ibrahim A, Abumrad NA. Role of CD36 in membrane transport of long-chain fatty acids. *Curr. Opin. Clin. Nutr. Metab. Care* 5(2), 139–145 (2002).
 - 78 Abumrad N, Coburn C, Ibrahim A. Membrane proteins implicated in long-chain fatty acid uptake by mammalian cells: CD36, FATP and FABPm. *Biochim. Biophys. Acta* 1441(1), 4–13 (1999).
 - 79 Nath A, Li I, Roberts LR, Chan C. Elevated free fatty acid uptake via CD36 promotes epithelial-mesenchymal transition in hepatocellular carcinoma. *Sci. Rep.* 5, 14752 (2015).
 - 80 Stahl A, Gimeno RE, Tartaglia LA, Lodish HF. Fatty acid transport proteins: a current view of a growing family. *Trends Endocrinol. Metab.* 12(6), 266–273 (2001).
 - 81 Chmurzynska A. The multigene family of fatty acid-binding proteins (FABPs): function, structure and polymorphism. *J. Appl. Genet.* 47(1), 39–48 (2006).
 - 82 Grivennikov SI, Greten FR, Karin M. Immunity, inflammation, and cancer. *Cell* 140(6), 883–899 (2010).
 - 83 Li Z, Srivastava S, Mittal S, Yang X, Sheng L, Chan C. A Three Stage Integrative Pathway Search (TIPS) framework to identify toxicity relevant genes and pathways. *BMC Bioinformatics* 8, 202 (2007).
 - 84 Razumilava N, Gores GJ. Cholangiocarcinoma. *Lancet* 383(9935), 2168–2179 (2014).
 - 85 Gatto M, Bragazzi MC, Semeraro R *et al.* Cholangiocarcinoma: update and future perspectives. *Dig. Liver Dis.* 42(4), 253–260 (2010).
 - 86 Rizvi S, Borad MJ, Patel T, Gores GJ. Cholangiocarcinoma: molecular pathways and therapeutic opportunities. *Semin. Liver Dis.* 34(4), 456–464 (2014).
 - 87 Ho C, Wang C, Mattu S *et al.* AKT (v-akt murine thymoma viral oncogene homolog 1) and N-Ras (neuroblastoma Ras viral oncogene homolog) coactivation in the mouse liver promotes rapid carcinogenesis by way of mTOR (mammalian target of rapamycin complex 1), FOXM1 (forkhead box M1)/SKP2, and c-Myc pathways. *Hepatology* 55(3), 833–845 (2012).
 - 88 Fan B, Malato Y, Calvisi DF *et al.* Cholangiocarcinomas can originate from hepatocytes in mice. *J. Clin. Invest.* 122(8), 2911–2915 (2012).
 - 89 Li L, Che L, Tharp KM *et al.* Differential requirement for *de novo* lipogenesis in cholangiocarcinoma and hepatocellular carcinoma of mice and humans. *Hepatology* 63(6), 1900–1913 (2016).
 - **Depicts the metabolic requirements of intrahepatic carcinoma cells, and observes that inhibitors of FA uptake, either alone or in association with FASN inhibitors, might be beneficial for the treatment of human intrahepatic carcinoma.**
 - 90 Chalasani N, Younossi Z, Lavine JE *et al.* The diagnosis and management of non-alcoholic fatty liver disease: practice Guideline by the American Association for the Study of Liver Diseases, American College of Gastroenterology, and the American Gastroenterological Association. *Hepatology* 55(6), 2005–2023 (2012).
 - 91 Tiniakos DG, Vos MB, Brunt EM. Nonalcoholic fatty liver disease: pathology and pathogenesis. *Annu. Rev. Pathol.* 5, 145–171 (2010).
 - 92 Chalasani N, Younossi Z, Lavine JE *et al.* The diagnosis and management of non-alcoholic fatty liver disease: practice guideline by the American Gastroenterological Association, American Association for the Study of Liver Diseases, and American College of Gastroenterology. *Gastroenterology* 142(7), 1592–1609 (2012).
 - 93 Hsiao PJ, Kuo KK, Shin SJ *et al.* Significant correlations between severe fatty liver and risk factors for metabolic syndrome. *J. Gastroenterol. Hepatol.* 22(12), 2118–2123 (2007).
 - 94 Cordero P, Campion J, Milagro FI, Martinez JA. Transcriptomic and epigenetic changes in early liver steatosis associated to obesity: effect of dietary methyl donor supplementation. *Mol. Genet. Metab.* 110(3), 388–395 (2013).
 - 95 Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet* 371(9612), 569–578 (2008).
 - 96 Sun B, Karin M. Obesity, inflammation, and liver cancer. *J. Hepatol.* 56(3), 704–713 (2012).
 - 97 Welzel TM, Graubard BI, Quraishi S *et al.* Population-attributable fractions of risk factors for hepatocellular carcinoma in the United States. *Am. J. Gastroenterol.* 108(8), 1314–1321 (2013).
 - 98 Bugianesi E, Leone N, Vanni E *et al.* Expanding the natural history of nonalcoholic steatohepatitis: from cryptogenic cirrhosis to hepatocellular carcinoma. *Gastroenterology* 123(1), 134–140 (2002).
 - 99 White DL, Kanwal F, El-Serag HB. Association between nonalcoholic fatty liver disease and risk for hepatocellular cancer, based on systematic review. *Clin. Gastroenterol. Hepatol.* 10(12), 1342–1359 e1342 (2012).
 - 100 Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N. Engl. J. Med.* 348(17), 1625–1638 (2003).
 - 101 Bhaskaran K, Douglas I, Forbes H, Dos-Santos-Silva I, Leon DA, Smeeth L. Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet* 384(9945), 755–765 (2014).
 - 102 Setiawan VW, Lim U, Lipworth L *et al.* Sex and ethnic differences in the association of obesity with risk of hepatocellular carcinoma. *Clin. Gastroenterol. Hepatol.* 14(2), 309–316 (2016).
 - 103 Ibrahim MM. Subcutaneous and visceral adipose tissue: structural and functional differences. *Obes. Rev.* 11(1), 11–18 (2010).
 - 104 Fujiwara N, Nakagawa H, Kudo Y *et al.* Sarcopenia, intramuscular fat deposition, and visceral adiposity independently predict the outcomes of hepatocellular carcinoma. *J. Hepatol.* 63(1), 131–140 (2015).
 - 105 Streb LA, Vere CC, Rogoveanu I, Streb CT. Nonalcoholic fatty liver disease, metabolic risk factors, and hepatocellular carcinoma: an open question. *World J. Gastroenterol.* 21(14), 4103–4110 (2015).
 - **Summarizes the potential relationships among linking obesity, Type 2 diabetes, systemic inflammation, nonalcoholic fatty liver disease/nonalcoholic steatohepatitis and hepatocellular carcinoma.**
 - 106 Mello T, Materozzi M, Galli A. PPARs and mitochondrial metabolism: from NAFLD to HCC. *PPAR Res.* 2016, 7403230 (2016).