Atypical myopathy in Père David's deer (*Elaphurus davidianus*) associated with ingestion of hypoglycin A

Carolyn Bunert,^{*,†} Sandra Langer,[‡] Dominique M. Votion,[§] François Boemer,[#] Anja Müller,[§]
Kerstin Ternes,^{*} and Annette Liesegang^{†,1}

*Zoo Duisburg AG, Department of Zoo Veterinary Medicine, 47058 Duisburg, Germany; [†]Institute of Animal Nutrition, University of Zurich, 8057 Zurich, Switzerland; [‡]Kölner Zoo, Department of Zoo Veterinary Medicine, 50735 Köln, Germany; [§]Fundamental and Applied Research for Animals & Health (FARAH), Faculty of Veterinary Medicine, University of Liège, 4000-Liège, Belgium; [#]Biochemical Genetics Laboratory, CHU Sart Tilman, University of Liège, 4000-Liège, Belgium; [§]IDEXX Laboratories, 71636 Ludwigsburg, Germany

ABSTRACT: From 2004 until 2016, 21 Père David's deer (*Elaphurus davidianus*) have died for unknown reason at Zoo Duisburg. These deer, also known as milu, have succumbed from a myopathy that occurred seasonally in autumn and in spring. The clinical signs shown by the animals closely resembles those of a disease called equine atypical myopathy (EAM), which is formerly known in horses. The cause for EAM in Europe was found in the ingestion of hypoglycin A, contained in samaras and seedlings of the sycamore maple tree (*Acer pseudoplatanus*). To test the hypothesis that the mortality of milus was caused by ingestion of hypoglycin A, 79 sera from all zoos and wildlife parks that have kept milus in Germany and Austria, including 19 diseased and 60 healthy animals, were used. Selected biochemical values and additionally hypoglycin A, methylenecyclopropyl acetic acid-carnitine (MCPA-carnitine), and acylcarnitines, which have been found in horses suffering from EAM, were determined. The results showed greater values

of serum activities of creatine kinase ($P < 0.001$) and aspartate aminotransferase ($P < 0.001$) in diseased milus comparing to healthy ones confirming a myopathy in affected animals. Moreover, hypoglycin A and MCPA-carnitine were found in the blood of Père David's deer and thus, hypoglycin A intoxication was considered to be a potential cause for the myopathies by ingestion of sycamore maple samaras that were present in the enclosure of the affected animals. Hypoglycin A values were greater in diseased animals ($P < 0.01$) as well as MCPA-carnitine levels ($P < 0.05$). Additionally, affected milus showed greater C5-OH-carnitine ($P < 0.01$) and C6-carnitine ($P < 0.001$) values. Until now hypoglycin A intoxication was only known in the family of *Equidae*, in humans, and in laboratory rats, and it has not been previously described in other zoological families. Comparing to horses, ruminants do have a different digestive tract and it will need further investigation to find out if several factors are involved to trigger an outbreak in ruminants.

Key words: atypical myopathy, *Elaphurus davidianus*, hypoglycin A, MCPA-carnitine, milu, sycamore maple

© The Author(s) 2018. Published by Oxford University Press on behalf of the American Society of Animal Science. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com.

J. Anim. Sci. 2018.96:3537–3547
doi: 10.1093/jas/sky200

INTRODUCTION

The Père David's deer (*Elaphurus davidianus*), also known as milu, is a wild ruminant that is classified as “Extinct in the Wild” on *The IUCN Red List* (Wilson and Mittermeier, 2011). This species,

¹Corresponding author: aliese@nutrivet.uzh.ch
Received March 31, 2018.
Accepted May 11, 2018.

which naturally occurred throughout eastern Asia, became extinct in the 19th century as a result of a flood disaster and turmoil of war during the boxer rebellion (Bannikow et al., 2000). After the extermination, the viability of the species was secured by a captive breeding program. All animals capable of breeding were gathered in England by the Duke of Bedford (Grzimek, 1988). From 1985 to 2012, the reintroduction into China took place, starting with an initial population of five breeding individuals. Today more than 5,000 milus form the world's population (Schürer and Stadler, 2016).

Zoo Duisburg, in Germany, has kept and successfully bred Père David's deer since 1966 with a total number of 77 individuals. From 2004 until 2016, the population has suffered from a disease containing high-grade myopathy. The outbreaks occurred seasonally from September until March and never happened during the summer. To this date, the majority of the milus have died, with only 4 individuals surviving out of 25 affected individuals.

The milus suffered from clinical signs that share similarities with a disease called Equine atypical myopathy (EAM) which is formerly known in horses. EAM can be described as a degeneration of particular muscle groups, clinically characterized by stiffness, weakness, recumbency, myoglobinuria, and a high mortality rate. The disease is seasonal, and the outbreaks mostly occurred in autumn and spring (Votion, 2016). Recent studies showed that EAM in horses in Europe is associated with the ingestion of hypoglycin A contained in samaras and seedlings of the sycamore maple (*Acer pseudo-platanus*) (Votion et al., 2014; Baise et al., 2016).

We hypothesized that the morbidity and mortality of the milus at Zoo Duisburg was caused by ingestion of hypoglycin A.

MATERIALS AND METHODS

Animals and Sampling Sites

In this study, all 11 zoos and wildlife parks in different regions of Germany and Austria that have kept milus were included. Animal husbandry characteristics such as type of enclosure, vegetation, ground coverage, nutrition, and medical history regarding similar symptoms to the diseased animals from Zoo Duisburg were noted in a standardized questionnaire for each holding. The diseased animals included in this study are those presenting signs compatible with a myopathy. The healthy animals are represented by not myopathic animals either from zoos other than Duisburg or milus from

Duisburg that were kept in the same enclosure as the diseased deer.

Pathology

The deceased milus were brought to the Chemical and Veterinary Investigation Office (CVUA) in Krefeld, Germany. Necropsies were performed and muscle samples for histopathology were taken from many different localizations of the body, including thigh, intercostal, masseter, glossal, back, diaphragmatic, shoulder and lumbar muscles.

Blood Samples

Preserved serum samples from a total of 79 animals of different age and sex from five zoos were gathered. Nineteen samples of diseased milus from Zoo Duisburg and 60 samples of clinically healthy animals from other zoos as well as Duisburg were part of the study. The sampling years of the diseased animals were as following: 2004 (three samples), 2005 (two samples), 2007 (one sample), 2010 (two samples), 2012 (one sample), 2013 (two samples), 2014 (four samples), 2015 (two samples), and 2016 (two samples).

Twenty-eight animals under the age of 1 were classified as suckling calves. Twenty-one animals between 1 and 3 yr are representing the subadults by being sexually immature. With over 3 yr of age, 26 milus were considered as adults (Kern, 2008). In four cases, the age could not be traced.

Biochemical Analysis

The sera were frozen at -18°C in Eppendorf vials and 77 samples were sent to a certified laboratory (IDEXX Laboratories, 71636 Ludwigsburg, Germany). The biochemical analysis included Cu, Se, vitamin E, Mg, urea, triglycerides, FFA, and enzyme activities of creatine kinase (CK), alkaline phosphatase (ALP), aspartate aminotransferase (AST), glutamate dehydrogenase (GLDH), and γ -glutamyl-transferase (γ -GT). For analyzing Se and Cu, the inductively coupled plasma mass spectrometry (ICP-MS) method was used. For Se, the isotopes Se77, 78, and 82 and for Cu, the isotopes Cu63 and 65, were analyzed at the same time by ICP-MS (Varian 820 MS). Ultrafast high-performance liquid chromatography (U-HPLC) was taken to determine vitamin E. After protein precipitation, the extracted vitamin E was analyzed on a U-HPLC System (RS-LC 3000, Dionex/Themofisher Scientific, Dreieich, Germany). For

the detection of the vitamin E, a Diodenarry Detector with wavelength 295 nm was used. FFA were measured photometric and the other values (Mg, Urea, CK, ALP, AST, ALT, GLDH, γ -GT) were tested by using a chemical analyzer. FFA as well as Mg, Urea, CK, AP, AST, ALT, GLDH, and γ -GT have been analyzed on a Beckmann AU 5800 (Beckman Coulter, Europark Fichtenhain B13, 47807 Krefeld, Germany).

Twenty-one samples were sent to the Biochemical Genetics Laboratory of the University of Liège, Belgium for hypoglycin A and methylene-cyclopropyl acetic acid-carnitine (MCPA-carnitine) quantification. Hypoglycin A concentration in the defrosted sera was determined by using an aTRAQ kit for AA analysis (Sciex, Framingham, MA; [Boemer et al., 2015](#)). For MCPA-carnitine quantification, the liquid chromatography coupled with mass spectrometry (LC/MS/MS; TQ5500 mass spectrometer; Sciex) was used. Serum acyl-carnitines concentrations (free carnitine, C2-, C3-, C3DC-, C4-, C5-, C5-OH-, C5DC-, C6-, C8-, C8:1-, C10-, C10:1-, C10:2-, C12-, C12:1-, C14-, C14:1-, C16-, C16:1-, C18-, and C18:1-carnitine) were determined by tandem mass spectrometry ([Chace et al., 2003](#)). Briefly, serum proteins were precipitated with a methanol solution containing labeled internal standards. Supernatants were evaporated under nitrogen stream and derivatized with butanolic-HCl. Butylated samples were analyzed with a TQ5500 mass spectrometer (Sciex).

Analyses of Plant Samples

Seedlings (50 g) were collected, at posteriori (spring 2017) from the enclosure where the diseased milus lived and were kept frozen (-18°C) up to analyses. Hypoglycin A was extracted from samaras and seedlings using methanol, and high-performance thin-layer chromatography (HPTLC). Briefly, 5 g of whole maple samaras or seedlings were mechanically ground and mixed with 25 mL

of pure methanol (VWR International, Leuven, Belgium). The mixture was gently agitated for 24 h at room temperature and then centrifuged at 4,500 g for 15 min. Thereafter, 12.5 mL supernatant was removed and evaporated. The residue obtained was dissolved in 3 mL of pure water. Quantification of hypoglycin A was performed by HPTLC using a CAMAG TLC scanner 3 at 490 nm with WinCATS 4.3 software (CAMAG, Muttenz, Switzerland). Additional details about the procedure can be found elsewhere ([Habyarimana et al., 2017](#)).

Statistical Analysis

Data analysis was performed by using the R-program rattle (Rattle 5.0.19). This program's logistic regression function was used to identify those blood measurements which are significantly related to healthy and diseased animals, respectively. Cross-tabulation and Pearsons chi-square test were used to analyze whether sex and age groups were related to disease and healthiness, respectively. XLSTAT (2016.06) was used to calculate the statistics for significant blood measurements including the *t*-test. Test results with a value of $P < 0.05$ were determined to be statistically significant.

RESULTS

A result of the questionnaire was that 25 animals from Zoo Duisburg showed the described symptoms and none of the other zoos or wildlife parks did have similar problems with their milus. The disease appeared in outbreaks in autumn–winter 2004 (11 diseased animals), 2005 (6 diseased animals), 2007 (1 diseased animal), 2010 (2 diseased animals), 2012 (1 diseased animal), 2013 (2 diseased animals), 2014 (3 diseased animals), 2015 (1 diseased animal), and 2016 (1 diseased animal) and never happened during the summer ([Fig. 1](#)). One hind diseased four times over a period of 8 yr and finally died in 2013. The diseased milus from Zoo Duisburg showed

Figure 1. Numbers of diseased milus from year 2004 until 2016 at Zoo Duisburg.

clinical signs with a sudden onset of muscle tremor, apathy, anorexia, salivation, recumbency, labored breathing, and a reduced alert response. Seven individuals additionally showed exophthalmos and two animals emitted dark-colored urine.

Over the years, the myopathic animals from Zoo Duisburg received different supportive therapies including injections of vitamin E and Se, vitamin B complex, butafosfan, menbutone, butylscopolamine and cocktail of vitamins, AA, and trace elements. All of them received fluids either i.v. or s.c. during anesthesia. Different antibiotics and analgesics, including nonsteroidal anti-inflammatory drugs as well as cortisone and diazepam in different combinations were given. Conversions of feeding and supplementation such as reduction of the protein amount and increase of vitamin E, Se, and Cu in the ration have been made.

The enclosure of the milus at Zoo Duisburg, which hosted all of the diseased animals, was surrounded by six maple trees. After the last milu's

death in spring 2017, seedlings of these trees were found on the ground of the enclosure in large numbers (Fig. 2). During botanical inspection, the trees were identified as *A. pseudoplatanus* and digital images of the plants and the enclosure were taken. Analyses of the samples revealed that hypoglycin A was detected in samaras and seedlings of sycamore maple collected within the enclosure at the end of March 2017 was 152.0 ± 4.5 and 202.0 ± 2.9 mg/Kg fresh weight, respectively.

Blood sample analysis showed statistically significant greater values of CK, AST, Se, hypoglycin A, and MCPA-carnitine in diseased animals (Table 1) compared to healthy individuals. The diseased milus exhibited high CK levels in serum and showed an increase in AST levels (Fig. 3). All diseased animals had detectable amount of hypoglycin A and significantly greater values have been computed in diseased milus for hypoglycin A (Fig. 3).

Figure 2. Seedlings of *Acer pseudoplatanus* on the milu enclosure at Zoo Duisburg, March 2017.

Table 1. Significant differences in blood values between diseased and healthy milus (*t*-test)

		Hypoglycin A, $\mu\text{mol/L}$	MCPA-carnitine, ¹ nmol/L	Se, $\mu\text{g/L}$	CK, ² U/L	AST, ³ U/L
Diseased	<i>n</i>	10	10	18	18	18
	Mean	0.248	6.447	146	159,353	7,686
	Median	0.197	2.280	97	116,781	5,949
	Min	0.022	0.000	53	81	43
	Max	0.643	22.600	715	908,707	32,179
	<i>P</i> value	<0.01	<0.05	<0.001	<0.001	<0.001
Healthy	<i>n</i>	11	11	57	59	59
	Mean	0.055	0.000	54	7,964	555
	Median	0.010	0.000	49	189	58
	Min	0.000	0.000	20	1	13
	Max	0.275	0.000	123	211,967	12,534

¹MCPA-carnitine: methylenecyclopropyl acetic acid-carnitine.

²CK = creatine kinase.

³AST = aspartate aminotransferase.

Figure 3. Creatine kinase, aspartate aminotransferase, hypoglycin A, and MCPA-carnitine¹ values in diseased vs. healthy milus. **n* = 59. †*n* = 18. ‡*n* = 11. §*n* = 10. ¹MCPA-carnitine: methylenecyclopropyl acetic acid-carnitine.

Greater values have also been recorded for MCPA-carnitine in diseased animals (Fig. 3). The MCPA-carnitine was only detected in 6 diseased milus out of the 10 tested animals with signs of myopathy. No healthy milus had detectable amount of MCPA-carnitine.

The CK showed significant correlations with AST, MCPA-carnitine, and hypoglycin A. The MCPA-carnitine showed significant correlations with hypoglycin A (Table 2).

The measured concentrations of acylcarnitines in the sera turned out to be higher in diseased milus compared to healthy ones except for two medium chained acylcarnitines (C8:1- and C10:2-carnitine) where the values were equal. Statistically greater values have been found for C5-OH-carnitine and C6-carnitine in diseased vs. healthy milus (Table 3).

There were no statistically significant differences in Cu, vitamin E, Mg, urea, ALP, GLDH, triglycerides, FFA, and γ -GT comparing healthy or diseased animals. Also, there were no statistically significant differences for diseased animals between the two sexes and the three age groups (Tables 4 and 5).

Pathological reports of 15 deceased milus were provided. Forty percent revealed macroscopic lesions consisted of partial pale discoloration of skeletal muscle. Histologically, skeletal muscles of all milus, except for two individuals, demonstrated diffuse low-to-high grade acute-to-subacute hyaline myofiber degeneration (Fig. 4), depending on the localization of the muscle. Severe degenerations were especially found in the intercostal and thigh muscles. In three milus multifocal low-to-high grade acute myocardial fiber, degeneration was additionally found. The kidneys of four individuals exposed

Table 2. Correlation (Pearson) between significant blood values in diseased milus

	AST ¹	MCPA-carnitine ²	CK ³	Hypoglycin A
AST ¹	1	0.46	0.94	0.55
MCPA-carnitine ²	0.46	1	0.88	0.75
CK ³	0.94	0.88	1	0.87
Hypoglycin A	0.55	0.75	0.87	1

¹AST = aspartate aminotransferase.

²MCPA-carnitine = methylenecyclopropyl acetic acid-carnitine.

³CK = creatine kinase.

a tubular epithelial degeneration and the renal pelvis was filled with reddish-brown urine. Five individuals had a collateral degeneration of liver cells. Other pathological findings were unremarkable.

DISCUSSION

The results of this study show the close connection of the symptoms of the milus at Zoo Duisburg to a hypoglycin A intoxication. This shows that our hypothesis that morbidity and mortality of the milus at Zoo Duisburg was caused by ingestion of hypoglycin A was proven due to the following reasons:

Firstly, the milu enclosure was surrounded by sycamore maple trees and every autumn samaras of these trees fell onto the ground of the enclosure in large numbers. Samaras and seedlings of the sycamore maple contain the toxin hypoglycin A which is the etiological factor responsible for EAM in horses. In horses, EAM outbreaks can be divided into two groups. An autumnal outbreak is defined from September up to the end of February mainly caused by the ingestion of samaras. The spring outbreak from mid-March presumably originates from

Table 3. Serum (mean $\pm$ SD) concentrations of free carnitine and acylcarnitines ($\mu\text{mol/L}$) in diseased and healthy milus and in control horses¹ and horses with EAM² (nonsurvivors)¹

	Healthy milus (n = 11)	Diseased milus nonsurvivors (n = 10)	Control horses (n = 44)	Horses with EAM nonsurvivors (n = 40)
Free carnitine				
Free carnitine	79.42 $\pm$ 61.86	121.11 $\pm$ 50.31	26.02 $\pm$ 1.07	87.04 $\pm$ 7.95
Short-chain acylcarnitines (C2 to C5)				
C2-carnitine	13.02 $\pm$ 9.42	20.00 $\pm$ 11.36	8.77 $\pm$ 0.67	55.23 $\pm$ 4.24
C3-carnitine	0.50 $\pm$ 0.44	0.65 $\pm$ 0.40	0.56 $\pm$ 0.03	2.92 $\pm$ 0.27
C3DC-carnitine	0.04 $\pm$ 0.04	0.06 $\pm$ 0.03	0.03 $\pm$ 0.00	0.34 $\pm$ 0.04
C4-carnitine	1.42 $\pm$ 2.70	6.23 $\pm$ 6.44	0.36 $\pm$ 0.02	26.62 $\pm$ 3.67
C5-carnitine	0.68 $\pm$ 1.22	2.06 $\pm$ 1.61	0.21 $\pm$ 0.01	24.71 $\pm$ 3.20
C5-OH-carnitine	0.08 $\pm$ 0.04**	0.11 $\pm$ 0.05**	0.06 $\pm$ 0.0	0.56 $\pm$ 0.07
C5DC-carnitine	0.29 $\pm$ 0.48	0.51 $\pm$ 0.56	0.18 $\pm$ 0.01	2.20 $\pm$ 0.28
Medium-chain acylcarnitines (C6 to C10)				
C6-carnitine	0.06 $\pm$ 0.10***	0.39 $\pm$ 0.45***	0.04 $\pm$ 0.00	5.40 $\pm$ 0.81
C8-carnitine	0.06 $\pm$ 0.09	0.17 $\pm$ 0.15	0.02 $\pm$ 0.00	1.45 $\pm$ 0.18
C8:1-carnitine	0.04 $\pm$ 0.05	0.04 $\pm$ 0.02	0.09 $\pm$ 0.01	1.26 $\pm$ 0.15
C10-carnitine	0.04 $\pm$ 0.07	0.13 $\pm$ 0.12	0.02 $\pm$ 0.00	0.76 $\pm$ 0.09
C10:1-carnitine	0.02 $\pm$ 0.02	0.03 $\pm$ 0.02	0.05 $\pm$ 0.01	0.67 $\pm$ 0.07
C10:2-carnitine	0.01 $\pm$ 0.00	0.01 $\pm$ 0.00	0.03 $\pm$ 0.00	1.31 $\pm$ 0.14
Long-chain acylcarnitines (C12 to C18)				
C12-carnitine	0.16 $\pm$ 0.38	0.34 $\pm$ 0.22	0.04 $\pm$ 0.01	0.28 $\pm$ 0.03
C12:1-carnitine	0.04 $\pm$ 0.09	0.06 $\pm$ 0.04	0.02 $\pm$ 0.00	0.23 $\pm$ 0.03
C14-carnitine	0.22 $\pm$ 0.56	0.63 $\pm$ 0.42	0.03 $\pm$ 0.00	0.21 $\pm$ 0.02
C14:1-carnitine	0.07 $\pm$ 0.15	0.23 $\pm$ 0.15	0.03 $\pm$ 0.00	0.32 $\pm$ 0.04
C16-carnitine	0.80 $\pm$ 2.27	2.59 $\pm$ 1.8	0.14 $\pm$ 0.04	0.68 $\pm$ 0.07
C16:1-carnitine	0.13 $\pm$ 0.36	0.40 $\pm$ 0.28	0.02 $\pm$ 0.00	0.24 $\pm$ 0.03
C18-carnitine	0.33 $\pm$ 0.68	2.81 $\pm$ 1.86	0.07 $\pm$ 0.01	0.28 $\pm$ 0.02
C18:1-carnitine	0.31 $\pm$ 0.85	0.77 $\pm$ 0.44	0.04 $\pm$ 0.01	0.57 $\pm$ 0.06

¹Boemer et al., 2017.²EAM = equine atypical myopathy.** $P < 0.01$; *** $P < 0.001$; superscripts indicate statistically greater values in diseased vs. healthy milus.**Table 4.** Cross-tabulation and chi-square test results show no significance for age group

Age group	Healthy	Diseased
<1 yr	22	9
1 to 3 yr	14	11
>3 yr	19	12
Not applicable	4	0

 χ -squared = 3.7077, df = 3, P value = 0.2948.

mostly ingestion of seedlings (Baise et al., 2016). The outbreaks in the Père David's deer at Zoo Duisburg always occurred from September until early March which suggests that the milus mainly suffered from autumnal outbreaks by ingesting samaras. After the last milu's death in 2017, seedlings of the sycamore maple were seen in large numbers on the former milu enclosure, supporting the hypothesis that the animals may have ingested the samaras in the previous autumn. The concentration of hypoglycin A from autumnal samaras cannot be extrapolated from the one found in spring. At the concentration

Table 5. Cross-tabulation and chi-square test results show no significance for gender

Sex	Healthy	Diseased
Female	35	20
Male	24	12

 χ -squared = 0.087635, df = 1, P value = 0.7672.

found (about 150 mg/Kg fresh weight), almost 170 g of samaras would be sufficient to intoxicate a 500 kg horse as hypothesized based on experiments on laboratory animals (Valberg et al., 2013). The large number of seedlings found within the enclosure (Fig. 2) strengthen the hypothesis of a strong toxic pressure in autumn and thus that hypoglycin A intoxication is a possible explanation for the mortality of the deer.

Secondly, a shared temporality between outbreaks in horses and in milus could be observed. The severe outbreaks in milus happened in autumn–winter 2004 and a year later in 2005. Although there were no official recordings from the “Atypical

Figure 4. Skeletal muscle of a milu, hyaline myofiber degeneration ($\times 10$ magnification) histological specimen: CVUA, Krefeld, Germany.

Myopathy Alert Group” at an European level, there were outbreaks observed in horses in Belgium in both years (Votion et al., 2007). From 2006, where EAM was started to be recorded at an European level, EAM cases were reported yearly (D. Votion, personal communication). So, the outbreaks in milus in 2007, 2010, 2012, 2013, 2014, 2015, and 2016 were temporally accompanied by outbreaks in horses which corroborates our hypothesis.

In addition, the clinical signs shown by horses suffering from EAM and shown by affected milus substantiate a hypoglycin A intoxication. Both horses and milus showed clinical signs like trembling, recumbency, depression, dyspnea, weakness, normothermia, stiffness, and pigmenturia, but in different severities. In EAM horses, 96% showed pigmenturia during the autumnal outbreak in 2013, whereas only two milus out of 25 animals emitted dark-colored urine. Horses additionally showed tachycardia, sweating, heart murmurs, colics, dysphagia, a distended bladder on rectal palpation, and congested mucous membranes that has not occurred in milus (van Galen et al., 2012). It must be mentioned, that all clinical examinations in the milus were performed under anesthesia which might implicate a different depiction of clinical signs. Contrary to horses, the milus showed salivation and anorexia. The reduced alert response which was the first clinical sign in most cases is not comparable with domesticated horses.

Hypoglycin A toxicity is also known in humans suffering from the Jamaican vomiting sickness and in laboratory rats that were exposed to a hypoglycin A diet. In humans, the affected show gastrointestinal distress, hypoglycemia, central nervous system depression, vomiting, hypotonia, and seizures

(Barceloux, 2009), which totally differs from clinical signs in horses. Indicators of hypoglycin A toxicity in rats were abnormal motor movements, arched backs, raised hairs, blackening of fecal matters and a loss in body weight (Blake et al., 2006). Compared to the response of humans and laboratory rats to hypoglycin A, the clinical signs of horses and milus are much more alike. But, even the differences in clinical signs are obviously usual between different kind of species.

Furthermore, the pathological findings in the milus are analogical to the findings in horses. Macroscopic discolorations of certain muscle groups have been found in horses as well as in milus during necropsy. In horses suffering from EAM, a degeneration of particular muscle groups is very distinctive. Preferentially postural, respiratory skeletal muscles, and the myocardium are affected. In the deceased milus, the most severe and frequent lesions were found in the intercostal and thigh muscles which resembles EAM cases. Three individuals showed additionally hyaline myofiber degeneration of the myocardium. The filling of the renal pelvis with reddish-brown urine in four milu cases is matchable with myoglobin casts that could be consistently identified in the kidneys of horses with EAM. In horses, microscopic examination of the liver did not reveal significant changes in contrast to five deer where degenerations of liver cells were observed (Cassart et al. 2007).

The pathological findings in the milus were the first indicators that the animals might suffer from a disease similar to EAM, because no other potential cause of death has been found over the years.

The results of the statistical analysis of the blood samples are equally supportive of this topic.

Creatine kinase is a blood parameter which showed highly increased levels in diseased animals. It is the most muscle-specific enzyme with occurrence in the skeletal and cardiac muscle and is therefore a very sensitive indicator for damage of striated muscles. A high increase of CK may have many reasons such as muscle trauma, exercise myopathy, diet-related deficiency states, or inflammatory processes (myositis) (Moritz, 2013). In wild animals, capture myopathy is a common noninfectious, metabolic disease that is associated with pursuit, capture, restraint, or transportation of animals and goes along with elevations of AST and CK in serum. Clinical signs include ataxia, muscle stiffness, and prostration, and histological findings are characterized by necrotic muscles (West et al., 2014). Thus, the increased activities of CK in combination with AST could be related to

a capture myopathy with necrotizing muscle injury. However, at Zoo Duisburg, the milus developed the symptoms without a trigger like being captured or transported. Moreover, the measured values of the CK are similar to the levels described in horses with EAM which is accompanied by increased serum activity of CK with values from 10,000 U/L to 100,000 U/L (Zuraw et al., 2016).

Diseased milus also showed a significant increase in AST levels. The most abundant causes of AST elevations are hepatopathy, muscle damage (necrosis or inflammation), or hemolysis. In the studied samples, hepatopathy and hemolysis can be excluded by other blood values (Willard and Tvedten, 2006). Glutamate dehydrogenase, ALP, and γ -GT are liver-specific enzymes (Moritz, 2013) and were not significantly greater in the diseased deer. A hemolysis can be clarified by measuring the hematocrit (**hct**) and evaluating the color of serum or plasma (Willard and Tvedten, 2006). Hematocrit was not measured in this study but was always part of measurements taken when examining the diseased milus at Zoo Duisburg. The hct was always within normal reference values for most animals. Most likely the AST increase in combination with the high CK levels is associated with a myopathy. Aspartate aminotransferase is also elevated in horses with EAM (Votion et al., 2007).

Selenium levels were also significantly increased in diseased milus compared to healthy individuals. Selenium is an essential trace element with antioxidant activity and it occurs in the enzyme glutathione peroxidase. Exposure to excessive amounts of Se results in intoxication by severe oxidative tissue damage (Forth et al., 2001). In 2011, Al-Dissi et al. (2011) described Se toxicosis in white-tailed deer (*Odocoileus virginianus*) concomitant with exhibited signs of anorexia and unsteadiness on their feet. In necropsy, myocardial necrosis and mineralization were found. However, it is unlikely that the milus in this present study suffered from a Se intoxication. The high values of Se are much more likely due to the treatment of the diseased deer with vitamin E and Se supplements either oral or via injection. This treatment was chosen, because of the elevated CK and AST serum levels in blood and the degenerative muscle damages that were found during necropsy of other deer. Due to these findings, a deficiency of Se was presumed (McGavin and Zachary, 2009). Thus, an Se intoxication can be excluded.

Moreover, the toxin hypoglycin A has been demonstrated in all diseased milus. Hypoglycin A is a nonproteinogenic AA that is found in the

samaras and seedlings of the sycamore maple (*A. pseudoplatanus*) (Baise et al., 2016). After ingestion, this toxin is metabolized into MCPA-CoA, an inhibitor of acyl-CoA dehydrogenase (Osmundsen and Sherratt, 1975; Ikeda and Tanaka, 1990). This enzyme catalyzes each cycle of fatty acid β -oxidation in the mitochondria. A reduced activity of the dehydrogenase engenders an accumulation of acyl-CoAs in the mitochondria. In the next step, the acyl-CoAs may change into acylcarnitines that leave the mitochondria and circulate in the blood, which leads to a characteristic blood acylcarnitine profile in horses (Boemer et al., 2017). All muscles, including respiratory muscles and myocardium, utilize fatty acids as a primary energy source and the inhibition of fatty acid oxidation results in muscle damage and often leads to death. The toxic metabolite MCPA may form an ester with carnitine to MCPA-carnitine, which has been found in horses suffering from EAM (Votion et al., 2014).

The presence of hypoglycin A in the blood of diseased milus suggests the ingestion of sycamore maple samaras or seedlings.

Compared to horses with EAM (mean $5.47 \pm 1.60 \mu\text{mol/L}$) (Baise et al., 2016) the hypoglycin A value in milus is much lower. In horses, the kinetics of hypoglycin A are unknown at present and above all they are unknown in different species like in deer. In horses, usually AA are reabsorbed in the small intestine, whereas reabsorption in ruminants is taking place in the rumen (von Engelhardt, 2010). If this is also the case for hypoglycin A, this might lead to different concentrations of the toxin in blood despite equal input of samaras or seedlings in horses and ruminants.

Hypoglycin A has also been detected in healthy deer kept at Zoo Duisburg of this study. All *E. davidianus* were kept in the same enclosure the whole time. Certainly, every individual had access to samaras of the surrounding maple trees. In horses, hypoglycin A was also found in the blood of apparently healthy co-grazers which had no clinical signs (Baise et al., 2016). This led to the suggestion that hypoglycin A ingestion alone is not the sole trigger for EAM, but that there may be additional mechanisms that are involved in the pathogenesis of this disease or that there are different toxic levels according to individuals. A possible explanation for individuals that are staying subclinical might be a resistance to the toxic metabolite or blockage of the hypoglycin A conversion into MCPA by intrinsic or extrinsic factors like immunity or antioxidant status (Baise et al., 2016). The demonstration of the presence of hypoglycin A in the blood of all diseased

animals confirms the exposition to the toxin and thus the presumably ingestion of sycamore maple samaras.

In horses, acylcarnitines can be used to confirm a diagnosis of EAM and also some of the acylcarnitines are informative regarding prognosis of survival of the affected horses comparing values between survivors and nonsurvivors (Boemer et al., 2017). The determination of survivors was not possible in this study due to the death of all included milus. The acylcarnitines can be sorted into three groups: short-, medium-, and long-chain acylcarnitines, corresponding to their metabolic pathways. Most of the acylcarnitine mean serum concentrations of diseased milus, including all short- and long-chain acylcarnitines as well as free carnitine, were higher compared to healthy ones. Only the medium-chain acylcarnitine C8:1 and C10:2 values were equal respectively healthy and diseased milus. C5-OH-carnitine and C6-carnitine were statistically significant showing greater values in diseased vs. healthy animals. This leads to the conclusion that acylcarnitines may be used not only in horses, but also in milus to confirm the diagnoses of a myopathy caused by hypoglycin A. Comparing the values of healthy deer with control horses, greater concentrations are noticeable in the milus except in four acylcarnitines (C3, C8:1, C10:1, and C10:2). A possible explanation for these greater values is that the healthy milu group consisted of co-grazers that were likewise exposed to the samaras. Diseased milus showed lower concentrations of short- and medium-chain acylcarnitines, but greater concentrations in free carnitine and in six out of eight long-chain acylcarnitines (C12, C14, C16, C16:1, C18, and C18:1) comparing to nonsurvived horses with EAM.

Finally, the MCPA-carnitine, representing the toxic metabolite of hypoglycin A, was only detected in diseased milus and not in healthy ones.

The measured MCPA-carnitine levels were lower than mean concentrations in diseased horses (20.39 ± 17.24 nmol/L) (Votion et al., 2014). As the metabolism of hypoglycin A in ruminants is uncovered, this might lead to different ranges of MCPA-carnitine in the blood comparing to horses.

Unexpectedly, in 40% of the diseased deer, MCPA-carnitine could not be detected. This matter of fact was also found in horses, where a high concentration of hypoglycin A, but a low concentration of MCPA-carnitine in serum was detected. In this particular case, it was assumed that the early euthanasia of the horse after the beginning of intoxication impede MCPA-carnitine accumulation in

the blood (Bochnia et al., 2015). In the 40% of this study, additionally slightly lower CK values were recorded. Due to these circumstances, it leads to the assumption that there might be several interacting factors followed by the intoxication of the deer. The milus might have suffered from an unresolved disease that only in combination with ingestion of hypoglycin A led to the myopathy. This hypothesis leaves a margin for further studies.

As statistical analysis has pointed out, there is a high correlation between the four values (hypoglycin A, MCPA-carnitine, CK, and AST), especially for MCPA-carnitine and CK ($R = 0.88$). Thus, it can be said that the greater the MCPA value in a deer, the more muscle damage will occur, which will be represented by the high CK value.

CONCLUSION

Hypoglycin A was detected in the blood of all diseased milus thus confirming toxin exposure possibly by ingestion of samaras of *A. pseudoplatanus* at Zoo Duisburg.

Until now EAM was only known in the family of *Equidae* belonging to the order of *Perissodactyla* such as a horse (*Equus ferus caballus*) (Votion et al., 2009). Besides this, hypoglycin A intoxication has only been described in humans suffering from the Jamaican vomiting syndrome (Barceloux, 2009) and in laboratory rats (Blake et al., 2006) and it has not been published for other zoological families or orders. *Elaphurus davidianus* belongs to the family of *Cervidae* within the order of *Artiodactyla* (Gaisler and Zejda, 1997).

As ruminants have a different digestive tract comparing to horses, it will need further investigation to find out if there are several factors involved that trigger an outbreak of this disease in deer and to solidify this hypothesis.

However, there is the need for prevention of this myopathy. As *Acer pseudoplatanus* is ubiquitous in the north of Europe, it is difficult to control the exposure to the toxin (Baise et al., 2016). Prevention is even more complicated in zoos as the susceptible species have not been established at this stage.

ACKNOWLEDGMENTS

Many thanks to Ehrhard Köhn for help in statistics and Linda-Lou Düster for improving this manuscript. The SVWZH is gratefully acknowledged for the financial support. Zoo Wuppertal, Zoo Dresden, Tierpark Berlin, Serengeti-Park Hodenhagen, and the zoo keepers of the milus at Zoo Duisburg are gratefully acknowledged.

LITERATURE CITED

- Al-Dissi, A. N., B. R. Blakley, and M. R. Woodbury. 2011. Selenium toxicosis in a white-tailed deer herd. *Can. Vet. J.* 52:70–73.
- Baise, E., J. A. Habyarimana, H. Amory, F. Boemer, C. Douny, P. Gustin, C. Marcillaud-Pitel, F. Patarin, M. Weber, and D. M. Votion. 2016. Samaras and seedlings of *Acer pseudoplatanus* are potential sources of hypoglycin A intoxication in atypical myopathy without necessarily inducing clinical signs. *Equine Vet. J.* 48:414–417. doi:10.1111/evj.12499
- Bannikow, A. G., B. Grzimek, L. Heck, D. Heinemann, W. G. Heptner, H. G. Klös, E. M. Lang, E. Thenius, and F. Walther. 2000. Grzimeks tierleben. Säugetiere 4. [Animal life encyclopedia. Mammals 4]. Vol. 13. Weltbild Verlag, Augsburg, Germany.
- Barceloux, D. G. 2009. Akee fruit and Jamaican vomiting sickness (*Blighia sapida* König). *Dis. Mon.* 55(6):318–326. doi:10.1016/j.disamonth.2009.03.002
- Blake, O. A., M. R. Bennink, and J. C. Jackson. 2006. Akee (*Blighia sapida*) hypoglycin A toxicity: dose response assessment in laboratory rats. *Food Chem. Toxicol.* 44:207–213. doi:10.1016/j.fct.2005.07.002
- Bochnia, M., J. Ziegler, J. Sander, A. Uhlig, S. Schaefer, S. Vollstedt, M. Glatte, S. Abel, S. Recknagel, G. F. Schusser, et al. 2015. Hypoglycin A content in blood and urine discriminates horses with atypical myopathy from clinically normal horses grazing on the same pasture. *Plos One* 10:e0136785. doi:10.1371/journal.pone.0136785
- Boemer, F., M. Deberg, R. Schoos, E. Baise, H. Amory, G. Gault, J. Carlier, Y. Gaillard, C. Marcillaud-Pitel, and D. Votion. 2015. Quantification of hypoglycin A in serum using aTRAQ® assay. *J. Chromatogr. B. Analyt. Technol. Biomed. Life Sci.* 997:75–80. doi:10.1016/j.jchromb.2015.06.004
- Boemer, F., J. Detilleux, C. Cello, H. Amory, C. Marcillaud-Pitel, E. Richard, G. van Galen, G. van Loon, L. Lefère, and D. M. Votion. 2017. Acylcarnitines profile best predicts survival in horses with atypical myopathy. *Plos One* 12:e0182761. doi:10.1371/journal.pone.0182761
- Cassart, D., E. Baise, Y. Cherel, C. Delguste, N. Antoine, D. Votion, H. Amory, F. Rollin, A. Linden, F. Coignoul, et al. 2007. Morphological alterations in oxidative muscles and mitochondrial structure associated with equine atypical myopathy. *Equine Vet. J.* 39:26–32. doi:10.2746/042516407X157765
- Chace, D. H., R. Pons, C. A. Chiriboga, D. J. McMahon, I. Tein, E. W. Naylor, and D. C. De Vivo. 2003. Neonatal blood carnitine concentrations: normative data by electrospray tandem mass spectrometry. *Pediatr. Res.* 53:823–829. doi:10.1203/01.PDR.0000059220.39578.3D
- von Engelhardt, W. 2010. Physiologie der haustiere. [Physiology of domesticated animals]. 3rd ed. Enke, Stuttgart, Germany. p. 394.
- Forth, W., D. Henschler, W. Rummel, U. Förstermann, and K. Starke. 2001. Allgemeine und spezielle pharmakologie und toxikologie. [General and specific pharmacology and toxicology]. 8th ed. Urban & Fischer, Munich, Germany. p. 1054–1055.
- Gaisler, J., and J. Zejda. 1997. Enzyklopädie der säugetiere. [Encyclopedia of mammals]. Dausien, Hanau, Germany. p. 414.
- van Galen, G., C. Marcillaud Pitel, C. Saegerman, F. Patarin, H. Amory, J. D. Bailly, D. Cassart, V. Gerber, C. Hahn, P. Harris, et al. 2012. European outbreaks of atypical myopathy in grazing equids (2006–2009): spatiotemporal distribution, history and clinical features. *Equine Vet. J.* 44:614–620. doi:10.1111/j.2042-3306.2012.00556.x
- Grzimek, B. 1988. Grzimeks Enzyklopädie. Säugetiere. Band 5 [Grzimeks Encyclopedia. Mammals]. Vol. 5. Kindler, München, Germany. p. 161–164.
- Habyarimana, J. A., E. Baise, C. Douny, M. Weber, F. Boemer, P. De Tullio, T. Franck, C. Marcillaud-Pitel, M. Federich, A. Mouithys-Mickalad, et al. 2017. Development of an HPTLC method for determination of hypoglycin A in aqueous extracts of seedlings and samaras of *Acer* species. *bioRxiv* :148262. doi:10.1101/148262
- Ikeda, Y., and K. Tanaka. 1990. Selective inactivation of various acyl-CoA dehydrogenases by (methylenecyclopropyl)acetyl-CoA. *Biochim. Biophys. Acta.* 1038:216–221. doi:10.1016/0167-4838(90)90208-W
- Kern, C. 2008. Beobachtungen zur Fortpflanzungsbiologie beim Davidshirsch (*Elaphurus davidianus* Milne Edwards, 1866) im Tierpark Berlin [Observations about the biology of reproduction in Père David's deer (*Elaphurus davidianus* Milne Edwards, 1866) at Tierpark Berlin]. In: Tierpark Berlin, editor. Milu. Book 4, Vol. 12. Berlin, Germany. p. 319.
- McGavin, M. D., and J. F. Zachary. 2009. Pathologie der haustiere. [Pathology of domestic animals]. 1st ed. Elsevier, München, Germany. p. 898.
- McKenzie, R. K., F. I. Hill, J. A. Habyarimana, F. Boemer, and D. M. Votion. 2016. Detection of hypoglycin A in the seeds of sycamore (*Acer pseudoplatanus*) and box elder (*A. Negundo*) in New Zealand; the toxin associated with cases of equine atypical myopathy. *N. Z. Vet. J.* 64:182–187. doi:10.1080/00480169.2015.1123123
- Moritz, A. 2013. Klinische labordiagnostik in der tiermedizin. [Clinical laboratory diagnostics in veterinary medicine]. 7th ed. Schattauer, Stuttgart, Germany. p. 254, 320–322.
- Osmundsen, H., and H. S. Sherratt. 1975. A novel mechanism for inhibition of beta-oxidation by methylenecyclopropylacetyl-CoA, a metabolite of hypoglycin. *FEBS Lett.* 55:38–41. doi:10.1016/0014-5793(75)80951-3
- Schürer, U., and A. Stadler. 2016. Pater Davids Hirsch oder Milu (*Elaphurus davidianus* Milne Edwards, 1866), 150 Jahre, 1865 bis 2015 [Père David's Deer or Milu (*Elaphurus davidianus* Milne Edwards, 1866), 150 years, 1865–2015]. In: Kühne, R., editor. Der Zoologische Garten. Issue 6. 2016. [The Zoological Garden. Issue 6. 2016]. Vol. 85. Elsevier, Berlin, Germany. p. 389–396.
- Valberg, S. J., B. T. Sponseller, A. D. Hegemann, J. Earing, J. B. Bender, K. L. Martinson, S. E. Patterson, L. Sweetman. 2013. Seasonal pasture myopathy/atypical myopathy in North America associated with ingestion of hypoglycin A within seeds of the box elder tree. *Equine Vet. J.* 45(4):419–426. doi:10.1111/j.2042-3306.2012.00684.x
- Votion, D. M. 2016. Atypical myopathy: an update. In *Practice* 38:241–246. doi:10.1136/inp.i1859
- Votion, D. M., G. van Galen, L. Sweetman, F. Boemer, P. de Tullio, C. Dopagne, L. Lefère, A. Mouithys-Mickalad, F. Patarin, S. Rouxhet, et al. 2014. Identification of methylenecyclopropyl acetic acid in serum of European horses with atypical myopathy. *Equine Vet. J.* 46:146–149. doi:10.1111/evj.12117
- Votion, D. M., A. Linden, C. Delguste, H. Amory, E. Thiry, P. Engels, G. van Galen, R. Navet, F. Sluse, D. Serteyn, et al. 2009. Atypical myopathy in grazing horses: a first

- exploratory data analysis. *Vet. J.* 180:77–87. doi:10.1016/j.tvjl.2008.01.016
- Votion, D. M., A. Linden, C. Saegerman, P. Engels, M. Erpicum, E. Thiry, C. Delguste, S. Rouxhet, V. Demoulin, R. Navet, et al. 2007. History and clinical features of atypical myopathy in horses in Belgium (2000–2005). *J. Vet. Intern. Med.* 21:1380–1391. doi:10.1111/j.1939-1676.2007.tb01962.x
- West, G., D. Heard, and N. Caulkett. 2014. *Zoo animal and wildlife immobilization and anesthesia*. 2nd ed. Wiley, New York. p. 171.
- Willard, M. D., and H. Tvedten. 2006. *Labordiagnostik in der kleintierpraxis*. [Laboratory diagnostics in small animal medicine]. 1st ed. Elsevier, München, Germany. p. 46, 296–297.
- Wilson, D. E., and R. A. Mittermeier. 2011. *Handbook of the mammals of the world. Hoofed mammals*. Vol. 2. Lynx Edicions, Barcelona, Spain. p. 425.
- Zuraw, A., K. Dietert, S. Kühnel, J. Sander, and R. Klopfleisch. 2016. Equine atypical myopathy caused by hypoglycin A intoxication associated with ingestion of sycamore maple tree seeds. *Equine Vet. J.* 48:418–421. doi:10.1111/evj.12460