Published in final edited form as:

Curr Opin Psychol. 2018 August ; 22: 1–6. doi:10.1016/j.copsyc.2017.07.007.

From Ideation to Action: Recent Advances in Understanding Suicide Capability

Alexis M. May¹ and Sarah E. Victor²

¹University of Utah, Department of Psychology, 380 S 1530 E, Beh S 502, Salt Lake City, UT 84112, USA

²University of Pittsburgh School of Medicine, Department of Psychiatry, 3811 O'Hara Street, Pittsburgh, PA 15213, USA

Abstract

Suicide capability is one of few risk factors associated with suicide attempts among ideators. In the decade since the Interpersonal Psychological Theory of Suicide introduced the concept of acquired capability (i.e., the ability to face the fear and pain associated with death), understanding of the capability to attempt suicide has grown. Acquired (e.g. NSSI), dispositional (e.g., genetic), and practical contributors (e.g., access to firearms) appear to influence suicide capability via mechanisms such as the fear of death, persistence through pain, and familiarity with suicide methods. Self-report methods have shown mixed results, highlighting the importance of developing behavioral measures of suicide capability.

Introduction

The majority of people who think about suicide do not act on their thoughts [1]. However, our understanding of what differentiates risk for suicidal ideation from risk for suicide attempts is limited [2]. The Interpersonal Psychological Theory of suicide (IPTs) [3,4] was the first theory to use an “ideation to action” framework [5], that is, to posit different risk factors for suicide attempts than for suicide ideation. This ran counter to the long held, but rarely articulated, belief that suicide attempts resulted solely from worse ideation. The IPTs suggests that the capability to approach the pain and fear involved in enacting ones’ own death, which opposes the evolutionary goal of survival, must be acquired, primarily via repeated exposure to painful and/or fearsome experiences. Diverse experiences were thought to contribute to “acquiring” capability, including experiences that were endured (e.g., childhood maltreatment), enacted on others (e.g., combat exposure), and, perhaps most perniciously, enacted on oneself (e.g., nonsuicidal self-injury (NSSI)). Importantly, acquired capability was theorized to operate independently of suicidal thoughts; only when suicidal desire interacts with the ability to approach fear of death and pain would a serious suicide attempt occur.

Corresponding author: May, Alexis M. (alexis.m.may@gmail.com).

Conflict of Interest Statement

The authors have no conflicts of interest to report.

In the decade since its publication, new theories expanding on the IPTS have emerged. The Integrated Motivational Volitional Theory [6] hypothesized that impulsivity, intent, access to the lethal means, and social learning were also important in the transition from suicidal thoughts to actions. Klonsky and May [7] recently developed the Three-Step Theory, which identifies three distinct contributors to suicide capability: dispositional, acquired, and practical. In this model, dispositional contributors (e.g., genetics, personality traits) are those that influence baseline levels of suicide capability, whereas acquired contributors, such as those described by Joiner [3], are repeated experiences that wear away at natural barriers to suicide (e.g., fear of death and pain avoidance), thereby increasing suicide capability. Practical contributors are concrete factors that make an attempt easier, such as access to firearms. This review aims to briefly describe the recent literature on suicide capability, including its measurement, mechanisms, contributors, and time course, as well as directions for future research.

Measurement

Until very recently, the only published capability measure was the Acquired Capability for Suicide Scale (ACSS; [8]), developed in conjunction with the IPTS [3]. While the original measure contained 20 items assessing fearlessness and pain tolerance, recent psychometric investigations of the original ACSS and several shortened versions [9,10] support the use of a revised 7 item version specifically assessing fearlessness about death [9]. The Suicide Capacity Scale [7] and the Acquired Capability With Rehearsal for Suicide Scale [11] were recently developed to assess broader definitions of suicide capability.

There has been increasing interest in alternatives to self-report measures of suicide capability. For example, death and suicide-related versions of the Implicit Association Test (IAT; [12]), a behavioral measure of implicit attitudes, have been found to distinguish suicide attempters from ideators [13]. These measures may tap into a subconscious or unconscious comfort with one's own death that is not captured by a questionnaire. Using EEG methodology, a recent study found that a blunted neural response to threatening images was associated with attempts rather than ideation [14]. Another promising tool is the Self-Directed Violence Picture System [15], normed suicide-death specific images that may be used to probe behavioral and cognitive responses to suicide-specific fearsome images. Future work on the measurement of suicide capability will also require use of precise terminology to distinguish between suicide capability itself (i.e., overarching latent construct capturing the ability to act on suicidal thoughts), contributors to suicide capability (i.e., variables influencing capability, for example genetics, NSSI, access to firearms), and the mechanisms by which these contributors may be acting (i.e., reducing fear of death, impacting pain tolerance, or increasing knowledge of lethal means).

Fearlessness about death

Overwhelmingly, recent work has found an absence of evidence that acquired capability or fearlessness about death, as measured by the ACSS or the ACSS-FAD, distinguishes people who attempt suicide from those who ideate. This pattern appears among young adults with psychosis [16], individuals with eating disorders [17], domestic violence offenders [18],

college students [19], and online samples [20,21]. Fearlessness about death also does not distinguish emergency department patients presenting for suicide attempts versus ideation [22]. However, these null results are not universal. Smith et al [23] found that current ideation was only related to past suicide attempt for individuals who reported less fear of pain involved in dying. Further, some recent large studies found higher fearlessness about death among suicide attempters compared to ideators in UK university students [24] and US adults [7]. While self-reported fearlessness about death is not a consistent correlate of suicide capability, further research may clarify whether fearlessness about death relates to suicidal behavior in certain populations or in concert with other aspects of suicide capability.

Pain tolerance

Less research has investigated pain tolerance as a mechanism that increases suicide capability. Generally, self-report measures of pain tolerance have not differentiated individuals with attempts from those with ideation [24], and in at least one study, those with a history of suicide attempts or NSSI actually reported greater pain distress than those with suicide ideation only [25]. Within samples that self-injure, increased pain severity during NSSI was also associated with suicide attempts [26]. These findings run counter to theory, in that they suggest that greater pain severity and distress are associated with attempts as compared to ideation.

Recent innovative work using objective measures of pain in the context of frustrating behavioral tasks has begun to shed light on these paradoxical findings [27,28]. While no relationship was observed between attempt status and pain tolerance or threshold, the relationship between past attempts and current ideation was moderated by task persistence, such that ideation was more strongly related to attempts amongst those with mean or high levels of persistence through a painful and frustrating task [28]. Similarly, the relationship between NSSI and suicide attempts was also strongest among those higher in persistence through painful and distressing tasks [27]. This suggests that, though low distress tolerance may be a risk factor for suicide ideation, it may in fact be protective against suicidal behaviors, as one must be able to persist through physical and emotional distress in order to enact a suicide attempt.

Acquired contributors to suicide capability

Painful and provocative experiences (PPE) are hypothesized to contribute to suicide capability by increasing habituation to pain and decreasing fearlessness about death [4]. Much work has focused on the relationship between these acquired contributors themselves and attempts among those with ideation. PPE exposure relates to suicide attempt history when controlling for suicidal ideation and other relevant covariates in both youth [29] and adults [17,30]. Research in this area has typically focused on two types of PPE: direct exposure to traumatic/threatening events and potentially painful or harmful self-directed behaviors.

Several recent publications have addressed the relationship between threatening or traumatic events and suicide risk. A meta-analytic review of the relationship between combat exposure

and suicidality demonstrated a small, but statistically significant, effect of deployment or combat exposure on suicide-related outcomes [31]. A recent study of U.S. veterans with suicidal ideation compared to those with suicide attempts [32] found no differences in combat exposure; however, veterans with a history of attempted suicide reported greater levels of childhood trauma than veterans who reported passive suicidal ideation, but not compared to veterans with active suicidal ideation. Similarly, undergraduates who reported a suicide attempt with lethal intent reported more childhood physical abuse and neglect than those with only suicidal ideation, but did not differ from participants with a suicide plan; further, childhood sexual abuse did not differentiate participants at any level of suicidality [19]. Jordan et al [33] found that among various traumatic experiences, including being the recipient of violence, only repeatedly committing violent acts against others was associated with suicidal behavior with lethal intent. Similarly, among women seeking shelter from interpersonal violence (IPV), having perpetrated IPV was associated with greater acquired capability, while experiencing interpersonal trauma or violence was not [34]. Together, these findings suggest that, while trauma exposure may be related to suicidality generally, it may not be linked to suicide capability specifically. Furthermore, the act of committing violence against others may contribute more to suicide capability than being the victim of or witnessing violence. Finally, more nuanced measurement of suicidality (e.g., active/passive ideation, plans, intent, lethality) may help in clarifying the relationship between contributors to capability and the suicidal process.

Behaviors that are directly or indirectly self-damaging have also been associated with acquired capability. Two recent studies of adolescents found that health risk behaviors overall were related to suicide capability, either through a significant relationship to attempts when controlling for ideation [35] or by comparing adolescents with ideation only to adolescents with attempts [36]. With respect to substance use, one type of health risk behavior, while use alone may not be linked to suicide capability, certain characteristics of use may be. For example, among undergraduates, alcohol and drug use did not differentiate among participants with suicidal ideation, plans, or attempts [19]. However, in a large, nationally representative survey of participants using potentially injectable substances (i.e., cocaine, heroin, stimulants), choosing a more painful and risky route of administration (i.e., injection) was associated with attempts among those who considered or planned for suicide, even when controlling for depression and substance use disorder symptoms [37]. With respect to disordered eating behavior, a similar pattern has emerged, whereby behaviors that are less directly painful or provocative (such as eating restraint [19] and binge eating [17]) did not distinguish individuals with suicidal thoughts from those with suicide attempts while others (such as self-induced vomiting [17]) did.

Directly self-damaging behaviors have been robustly associated with suicide capability, with research showing that attempted suicide is associated with higher NSSI frequency [19] and use of more NSSI methods [36] compared to ideation only. Further, in undergraduate students, suicidal ideation was only related to suicide attempts among individuals who reported higher levels of NSSI [38]. Among adolescents receiving inpatient psychiatric care, NSSI frequency and number of methods both predicted lifetime suicide attempts in the context of suicidal ideation, though self-reported fearlessness about death, a hypothesized mechanism, was not predictive [39]. Longitudinal research in community adolescent girls

has shown that NSSI in the context of suicidal ideation prospectively predicts higher risk of suicide attempts compared to suicidal ideation alone [40]. NSSI may, therefore, be a particularly important acquired contributor to target when working with suicidal patients.

Practical contributors to suicide capability

The practical contributors to capability make attempting suicide easier, for example, familiarity with and access to suicide methods, or exposure to others' suicidal behavior. Firearms have emerged as a key practical contributor. A psychological autopsy study found that all suicide decedents who died by firearm, owned the firearm, and the majority of decedents who owned a firearm used that as their method [41], indicating a strong relationship between firearm access and suicide death by firearm. Further, epidemiological data suggests that suicide deaths correspond to the repeal or implementation of handgun regulations, and that changes in suicide death rates may be explained by reductions in suicides by firearm [42–44]. Importantly, this work argues against means substitution and emphasizes the importance and specificity of practical contributors to suicide capability overall. Firearm storage also matters, insofar as the relationship between current ideation and the self-reported likelihood of a future attempt is stronger among those who store personal guns loaded and unsecure [45].

Exposure to suicide may also be conceptualized as a practical contributor to suicide capability. Knowing a that family member or friend attempted suicide may not only expose an individual to details about suicide methods, but may also make that action seem more feasible. Among a cluster of proposed contributors to capability, exposure to friends' or family members' suicide attempt or NSSI showed the strongest relationship to attempts among ideators [24]. Exposure to a schoolmate's suicide also prospectively predicts attempts among youth [46]. These findings suggest the importance not only of addressing within-person aspects of capability (such as personality traits or NSSI), but also of addressing public health and means safety programs that could impact practical contributors to capability.

Dispositional contributors to suicide capability

Compared to acquired and practical contributors, less recent work exists on dispositional contributors, though a strong genetic component of suicide capability has been observed [47]. Somewhat more is known about personality traits and capability. Small, but consistent, positive associations have been observed between fearlessness about death and extraversion and openness, as well as negative associations with neuroticism [48,49]. Experimental work has shown that only people higher in sensation-seeking showed a relationship between distress tolerance and pain tolerance, supporting the pathway that dispositional contributors, such as temperament, may increase overall suicide capability by increasing the likelihood of PPEs [50]. Additionally, interoceptive deficits (i.e., difficulty sensing internal physiological sensations) have been associated with suicide attempts among ideators [51].

Time course of suicide capability

Until recently, there has been no longitudinal work on capability. Bryan and colleagues [52] assessed capability among members of the US military. Contrary to expectations, capability did not increase from pre-deployment to post, even when controlling for combat exposure or previous deployments. On the other hand, among undergraduates, NSSI frequency predicted increased capability one year later, while the reverse was not true [53]. This supports the conclusion that NSSI influences capability, rather than higher capability leading to increases in NSSI frequency. Neither adults [54] nor adolescents [55] in outpatient psychotherapy evidenced changes to capability over 1–3 months of treatment. However, one study suggests that capability may be more malleable. Using a newly developed measure, George and colleagues [11] found that, among hospitalized adults with a history of ideation or attempt, mental preparation for suicide and fearlessness about death both dropped significantly over an average of 2 weeks of treatment, while pain tolerance did not change. Differences across studies in population, suicide history, measures, and time course make it difficult to draw firm conclusions, but these intriguing findings suggest that the temporal sequence of capability for suicide is an important area for future study.

Conclusions and future directions

Suicide capability is one of the few factors that distinguishes people who attempt suicide from those who ideate. An explosion of work in recent years has led to a more nuanced understanding of the construct, demonstrating that suicide capability has more contributors than originally conceived, some contributors are more influential than others, capability interacts with other variables in complex ways, and existing self-report measures have limited utility. More work is needed to understand how capability is influenced (e.g., are fearlessness about death and pain tolerance the key mechanisms?), the best measurement approach (e.g., are behavioral tasks, measures of contributors, or measures of the construct itself most useful?), the most pertinent contributors (e.g., among the different contributors to capability, which hold the most power?), and the time course of capability (e.g., what impacts the development of, maintenance of, and changes in capability?). Continuing to refine and improve our understanding of suicide capability is essential to moving forward with the incredibly important and challenging tasks of predicting and preventing suicide attempts and deaths.

References

1. Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. National survey on drug use and health. Vol. 2015. Rockville, MD: 2014.
2. May AM, Klonsky ED. What distinguishes suicide attempters from suicide ideators? A meta-analysis of potential factors. *Clin Psychol Sci Pract*. 2016; 23:5–20.
3. Joiner T. Why people die by suicide. Harvard University Press; 2007.
4. Van Orden KA, Witte TK, Cukrowicz KC, Braithwaite SR, Selby EA, Joiner TE Jr. The interpersonal theory of suicide. *Psychol Rev*. 2010; 117:575–600. [PubMed: 20438238]
5. Klonsky ED, May AM. Differentiating suicide attempters from suicide ideators: a critical frontier for suicidology research. *Suicide Life Threat Behav*. 2014; 44:1–5. [PubMed: 24313594]
6. O'Connor RC. The integrated motivational-volitional model of suicidal behavior. *Crisis*. 2011; 32:295–298. DOI: 10.1027/0227-5910/a000120 [PubMed: 21945841]

- 7*. Klonsky ED, May AM. The three-step theory (3ST): A new theory of suicide rooted in the “ideation-to-action” framework. *Int J Cogn Ther*. 2015; 8(2):114–29. This article presents the most recent ideation-to-action theory of suicide, the 3ST, which expands capability to include acquired, dispositional, and practical contributors. It also presents initial empirical support for the relationship between each of these contributors and attempts among ideators.
8. Van Orden KA, Witte TK, Gordon KH, Bender TW, Joiner TE Jr. Suicidal desire and the capability for suicide: tests of the interpersonal-psychological theory of suicidal behavior among adults. *J Consult Clin Psychol*. 2008; 76:72–83. [PubMed: 18229985]
9. Ribeiro JD, Witte TK, Van Orden KA, Selby EA, Gordon KH, Bender TW, Joiner TE Jr. Fearlessness about death: The psychometric properties and construct validity of the revision to the Acquire Capability for Suicide Scale. *Psychol Assess*. 2014; 26:115. [PubMed: 24274043]
10. Gutierrez PM, Pease J, Matarazzo BB, Monteith LL, Hernandez T, Osman A. Evaluating the psychometric properties of the Interpersonal Needs Questionnaire and the Acquired Capability for Suicide Scale in military veterans. *Psychol Assess*. 2016; 28:1684. [PubMed: 26998894]
- 11*. George SE, Page AC, Hooke GR, Stritzke WG. Multifacet assessment of capability for suicide: Development and prospective validation of the Acquired Capability With Rehearsal for Suicide Scale. *Psychol Assess*. 2016; 28:1452. This paper pushes self-report assessment of capability forward by rigorously developing and testing a new measure that includes an additional contributor to suicide capability, mental preparation. The authors present data from multiple, diverse, longitudinal samples. [PubMed: 27046277]
12. Greenwald AG, McGhee DE, Schwartz JL. Measuring individual differences in implicit cognition: the implicit association test. *J Pers Soc Psychol*. 1998; 74:1464. [PubMed: 9654756]
13. Glenn JJ, Wertz AJ, Slama SJ, Steinman SA, Teachman BA, Nock MK. Suicide and self-injury-related implicit cognition: A large-scale examination and replication. *J Abnorm Psychol*. 2017; 126:199. [PubMed: 27991808]
14. Weinberg A, May AM, Klonsky ED, Kotov R, Hajcak G. Decreased neural response to threat differentiates patients who have attempted suicide from nonattempters with current ideation. *Clin Psychol Sci*. In press.
15. Nazem S, Forster JE, Brenner LA. Initial Validation of the Self-Directed Violence Picture System (SDVPS). *Psychol Assess*. 2017
16. Heelis R, Graham H, Jackson C. A preliminary test of the interpersonal psychological theory of suicidal behavior in young people with a first episode of psychosis. *J Clin Psychol*. 2016; 72:79–87. [PubMed: 26768800]
17. Pisetsky EM, Crow SJ, Peterson CB. An empirical test of the interpersonal theory of suicide in a heterogeneous eating disorder sample. *Int J Eat Disord*. 2017; 50:162–5. [PubMed: 27859487]
18. Wolford-Clevenger C, Brem MJ, Elmquist J, Florimbio AR, Smith PN, Stuart GL. A test of the interpersonal-psychological theory of suicide among arrested domestic violence offenders. *Psychiatry Res*. 2017; 249:195–9. [PubMed: 28119172]
- 19*. Burke TA, Ammerman BA, Knorr AC, Alloy LB, McCloskey MS. Measuring Acquired Capability for Suicide Within an Ideation-to-Action Framework. *Psychol Violence*. 2017. Advance online publication. This study of undergraduates is unique in that it examines differences in fearlessness about death and in specific PPES across multiple levels of suicidality (i.e., ideation, plans, attempts without intent and attempts with intent).
20. Forrest LN, Smith AR. Comparisons of the Interpersonal–Psychological Theory of Suicide Constructs Among Individuals Without Suicidality, Ideators, Planners, and Attempters. *Suicide Life Threat Behav*. 2016:1–12.
21. Khazem LR, Anestis MD. Thinking or doing? An examination of well-established suicide correlates within the ideation-to-action framework. *Psychiatry Res*. 2016; 245:321–6. [PubMed: 27575650]
- 22*. Baertschi M, Costanza A, Richard-Lepouriel H, Pompili M, Sarasin F, Weber K, Canuto A. The application of the interpersonal-psychological theory of suicide to a sample of Swiss patients attending a psychiatric emergency department for a non-lethal suicidal event. *J Affect Disord*. 2017; 210:323–31. This paper recruited participants in the midst of a suicidal crisis, either ideation or a recent attempt, from the Emergency Department. They failed to find the hypothesized relationship between current ideation severity and fearlessness of death predicting

- attempts among ideators. Their results raise important questions about self-report measures of fearlessness, the time course of capability, and impacts of sample selection. [PubMed: 28073040]
23. Smith PN, Stanley IH, Joiner TE Jr, Sachs-Ericsson NJ, Van Orden KA. An aspect of the capability for suicide—Fearlessness of the pain involved in dying—Amplifies the association between suicide ideation and attempts. *Arch Suicide Res.* 2016; 20:650–62. [PubMed: 26984289]
 24. Dhingra K, Boduszek D, O'Connor RC. Differentiating suicide attempters from suicide ideators using the Integrated Motivational–Volitional model of suicidal behaviour. *J Affect Disord.* 2015; 186:211–8. [PubMed: 26247914]
 25. Kirtley OJ, O'Connor RC, O'Carroll RE. Hurting inside and out? Emotional and physical pain in self-harm ideation and enactment. *Int J Cogn Ther.* 2015; 8:156–71.
 26. Ammerman BA, Burke TA, Alloy LB, McCloskey MS. Subjective pain during NSSI as an active agent in suicide risk. *Psychiatry Res.* 2016; 236:80–5. [PubMed: 26747213]
 27. Law KC, Khazem LR, Jin HM, Anestis MD. Non-suicidal self-injury and frequency of suicide attempts: The role of pain persistence. *J Affect Disord.* 2017; 209:254–61. [PubMed: 27936455]
 - 28*. Anestis MD, Capron DW. An investigation of persistence through pain and distress as an amplifier of the relationship between suicidal ideation and suicidal behavior. *J Affect Disord.* 2016; 196:78–82. This paper uses an innovative behavioral task that combines emotional distress and physical pain as a proxy for the type of persistence involved in a suicide attempt, that is, prioritizing the pursuit of death over escaping acute emotional and physical discomfort. Results suggest that ideation was only related to attempts in individuals at mean or high levels of persistence through this task. [PubMed: 26919055]
 29. Grossman AH, Park JY, Russell ST. Transgender youth and suicidal behaviors: applying the interpersonal psychological theory of suicide. *J Gay Lesbian Ment Health.* 2016; 20:329–49. [PubMed: 28344728]
 30. Smith PN, Cukrowicz KC, Poindexter EK, Hobson V, Cohen LM. The acquired capability for suicide: A comparison of suicide attempters, suicide ideators, and non-suicidal controls. *Depress Anxiety.* 2010; 27:871–7. [PubMed: 20821802]
 31. Bryan CJ, Griffith JE, Pace BT, Hinkson K, Bryan AO, Clemans TA, Imel ZE. Combat exposure and risk for suicidal thoughts and behaviors among military personnel and veterans: A systematic review and meta-analysis. *Suicide Life Threat Behav.* 2015; 45:633–49. [PubMed: 29889337]
 32. Hazlett EA, Blair NJ, Fernandez N, Mascitelli K, Perez-Rodriguez MM, New AS, Goetz RR, Goodman M. Startle amplitude during unpleasant pictures is greater in veterans with a history of multiple-suicide attempts and predicts a future suicide attempt. *Psychophys.* 2016; 53:1524–34.
 33. Jordan JT, Samuelson KW. Predicting suicide intent: the roles of experiencing or committing violent acts. *Suicide Life Threat Behav.* 2016; 46:293–300. [PubMed: 26418043]
 34. Wolford-Clevenger C, Smith PN, Kuhlman S, D'Amato D. A preliminary test of the interpersonal-psychological theory of suicide in women seeking shelter from intimate partner violence. *J Interpers Violence.* 2016
 35. Barzilay S, Feldman D, Snir A, Apter A, Carli V, Hoven CW, Wasserman C, Sarchiapone M, Wasserman D. The interpersonal theory of suicide and adolescent suicidal behavior. *J Affect Disord.* 2015; 183:68–74. [PubMed: 26001665]
 - 36*. Stewart JG, Esposito EC, Glenn CR, Gilman SE, Pridgen B, Gold J, Auerbach RP. Adolescent self-injurers: Comparing non-ideators, suicide ideators, and suicide attempters. *J Psychiatric Res.* 2017; 84:105–12. This study features a large sample of high-risk participants - adolescent inpatient self-injurers - and describes a wide range of variables theorized to influence suicide capability.
 - 37*. Cheek SM, Nestor BA, Liu RT. Substance use and suicidality: specificity of substance use by injection to suicide attempts in a nationally representative sample of adults with major depression. *Depress Anxiety.* 2016; 33:541–8. This paper is an excellent example of an analytic design that 1) isolates risk for attempts as opposed to ideation and 2) focuses as precisely as possible on a specific hypothesized contributor to suicide capability. The authors find using drugs by injection (rather than the using the same drugs by a different route of administration) is associated with risk of attempts but not plans or ideation, controlling for a number of important covariates. [PubMed: 26393336]

38. Brackman EH, Morris BW, Andover MS. Predicting risk for suicide: a preliminary examination of non-suicidal self-injury and the acquired capability construct in a college sample. *Arch Suicide Res.* 2016; 20:663–76. [PubMed: 27007698]
39. Matney J, Westers NJ, Horton SE, King JD, Eaddy M, Emslie GJ, Kennard BD, Stewart SM. Frequency and methods of nonsuicidal self-injury in relation to acquired capability for suicide among adolescents. *Arch Suicide Res.* 2017; 21:1–5. [PubMed: 26821811]
40. Scott LN, Pilkonis PA, Hipwell AE, Keenan K, Stepp SD. Non-suicidal self-injury and suicidal ideation as predictors of suicide attempts in adolescent girls: A multi-wave prospective study. *Compr Psychiatry.* 2015; 58:1–10. [PubMed: 25595520]
41. Van Orden KA, Smith PN, Chen T, Conwell Y. A case controlled examination of the interpersonal theory of suicide in the second half of life. *Arch Suicide Res.* 2016; 20:323–35. [PubMed: 26219512]
42. Anestis MD, Anestis JC. Suicide rates and state laws regulating access and exposure to handguns. *Am J Public Health.* 2015; 105:2049–58. [PubMed: 26270305]
43. Anestis MD, Khazem LR, Law KC, Houtsma C, LeTard R, Moberg F, Martin R. The association between state laws regulating handgun ownership and statewide suicide rates. *Am J Public Health.* 2015; 105:2059–67. [PubMed: 25880944]
- 44**. Anestis MD, Selby EA, Butterworth SE. Rising longitudinal trajectories in suicide rates: the role of firearm suicide rates and firearm legislation. *Prev Med.* 2017; 100:159–66. This critically important paper uses epidemiological data to examine the relationship between suicide death rates and firearm legislation in the United States over 16 years. The findings dispute the notion of means substitution (e.g. reductions in firearm suicide deaths were *not* off-set by increases in suicide deaths by other methods) and support the efficacy of legislative and means safety avenues to reduce practical contributors to suicide capability. [PubMed: 28455222]
45. Khazem LR, Houtsma C, Gratz KL, Tull MT, Green BA, Anestis MD. Firearms matter: The moderating role of firearm storage in the association between current suicidal ideation and likelihood of future suicide attempts among United States military personnel. *Mil Psychol.* 2016; 28:25–33.
46. Swanson SA, Colman I. Association between exposure to suicide and suicidality outcomes in youth. *Can Med Assoc J.* 2013; 185:870–7. [PubMed: 23695600]
47. Smith AR, Ribeiro JD, Mikolajewski A, Taylor J, Joiner TE, Iacono WG. An examination of environmental and genetic contributions to the determinants of suicidal behavior among male twins. *Psychiatry Res.* 2012; 197:60–5. [PubMed: 22417928]
48. Cramer RJ, Moore CE, Bryson CN. A test of the trait-interpersonal model of suicide proneness in emerging adults. *Pers Individ Dif.* 2016; 102:252–9.
49. Christensen H, Batterham PJ, Mackinnon AJ, Donker T, Soubelet A. Predictors of the risk factors for suicide identified by the interpersonal-psychological theory of suicidal behaviour. *Psychiatry Res.* 2014; 219:290–7. [PubMed: 24947914]
50. Bender TW, Anestis MD, Anestis JC, Gordon KH, Joiner TE. Affective and behavioral paths toward the acquired capacity for suicide. *J Soc Clin Psychol.* 2012; 31:81–100.
51. Forrest LN, Smith AR, White RD, Joiner TE. (Dis) connected: An examination of interoception in individuals with suicidality. *J Abnorm Psychol.* 2015; 124:754–63. [PubMed: 26147323]
- 52**. Bryan CJ, Sinclair S, Heron EA. Do military personnel “acquire” the capability for suicide from combat? A test of the interpersonal-psychological theory of suicide. *Clin Psychol Sci.* 2016; 4:376–85. This study of 168 military personnel was the first to test whether capability for suicide is acquired as a result of exposure to military training and combat. Participants completed a 5-item version of the ACSS 6 times over predeployment training, deployment to Iraq, and 1 year of follow up. Contrary to expectations, capability was a stable construct and was not related to combat exposure. Both traditional statistical approaches and dynamical systems approaches produced the same conclusions.
- 53**. Willoughby T, Heffer T, Hamza CA. The link between nonsuicidal self-injury and acquired capability for suicide: A longitudinal study. *J Abnorm Psychol.* 2015; 124:1110–15. This study of 782 undergraduates represents the first longitudinal test of the hypothesis that NSSI is linked to reductions in fear of death and pain sensitivity. Using a 5-item version of the ACSS, the authors

found that greater frequency of NSSI was indeed related to greater capability 1 year later and that this relationship was unidirectional. [PubMed: 26372006]

54. Wachtel S, Vocks S, Edel MA, Nyhuis P, Willutzki U, Teismann T. Validation and psychometric properties of the German Capability for Suicide Questionnaire. *Compr Psychiatry*. 2014; 55:1292–302. [PubMed: 24721192]
55. King JD, Horton SE, Hughes JL, Eaddy M, Kennard BD, Emslie GJ, Stewart SM. The Interpersonal–Psychological Theory of Suicide in Adolescents: A Preliminary Report of Changes Following Treatment. *Suicide Life Threat Behav*. 2017

Highlights

- The majority of people who think about suicide never act on their thoughts, however little is known about risk factors for attempts specifically. Suicide capability is one of few factors repeatedly linked to suicide attempts among ideators.
- In the decade since the publication of the Interpersonal Theory, research on capability for suicide has grown rapidly. The construct of capability has expanded beyond acquired contributors, such as painful and provocative events, to include dispositional and practical contributors as well.
- Of the many painful and provocative events that may contribute to suicide capability enacting violence against oneself (i.e., nonsuicidal self injury) and against others (i.e., perpetrating interpersonal violence) appear to be particularly important.
- Practical contributors to capability present a promising avenue for intervention. For example, recent work supports the efficacy of legislative and means safety approaches to reduce the impact of firearms, a practical contributor to suicide capability.
- Existing self-report measures of capability have produced inconsistent results, thus behavioral measurement approaches and innovative self-report measures are needed. Additionally longitudinal designs will be particularly useful in understanding the development and maintenance of suicide capability.