Original Article

Correlation of mitosis obtained by using 1% crystal violet stain with Ki67LI in histological grades of oral squamous cell carcinoma

Priyanka Kadoo^{a,e,*}, Rishikesh Dandekar^b, Meena Kulkarni^b, Aarti Mahajan^b, Ramniwas Kumawat^c, Neeta Parate^d^a Department of Oral Pathology and Microbiology, MGV's KBH Dental College and Hospital, Nashik, Maharashtra, India^b Oral Pathology and Microbiology, MGV's KBH Dental College and Hospital, Nashik, Maharashtra, India^c CDER, All India Institute of Medical Sciences, AIIMS, Delhi, India^d Ministry of Health, Bahrain^e Swargiya Dadasaheb Kalmegh Smruti Dental College and Hospital, Nagpur, Maharashtra, India

ARTICLE INFO

Article history:

Received 27 August 2016

Received in revised form 17 January 2017

Accepted 28 September 2017

Available online 12 October 2017

Keywords:

Mitosis

Ki67

Crystal violet stain

Oral squamous cell carcinoma

ABSTRACT

Aim: The purpose of this study was to use a simple, cost-effective technique for studying mitosis in various grades of oral squamous cell carcinoma (OSCC) using 1% Crystal-violet stain and to correlate mitotic frequency index (MFI) obtained by it with Ki67 labeling index (Ki67LI) so as to validate its usefulness as a selective stain for evaluating proliferation.

Materials and Methods: The invasive front grading score (IFG Score) was recorded in 40 patients of OSCC. Mitotic figures were assessed in hematoxylin and eosin (H and E) stained section as well as in 1% crystal-violet stained section using MFI (MFI = Mitosis/total number of cell counted X100). Comparison between MFI obtained by 1% crystal violet stain and H and E stain was done. Ki67LI was assessed using Ki67 immunohistochemical (IHC) marker. Correlation between Ki67LI and MFI obtained by using 1% crystal-violet stain was performed.

Results: There was statistically significant increase in MFI obtained by using 1% crystal-violet stain compared to routine H and E stain. Statistically significant positive correlation was observed between Ki67LI and mitosis in well and moderately differentiated OSCC. Positive correlation was also observed in poorly differentiated OSCC, but it was not statistically significant. Both mean MFI and mean Ki67LI significantly increased from grade I to grade II to grade III OSCC.

Conclusion: 1% crystal violet stain provides a definite advantage over the H and E stained sections. Thus crystal violet stain is easy and cost effective to evaluate proliferation when compared with expensive proliferative IHC marker.

© 2018 Craniofacial Research Foundation. Published by Elsevier B.V. All rights reserved.

1. Introduction

Oral Squamous Cell Carcinoma (OSCC) is the commonest malignant tumor of the oral cavity, accounting to about 96% of all oral malignancies in both males and females. It is one of the leading causes of death in developing countries.^{1,2} In order to improve the prognosis to some extent, it is important to govern and regulate the aggressiveness and prognostic factors that will aid in appropriate treatment plan. Although many histopathologic characteristics of OSCC have been known as prognostic factors, accurate and unambiguous factors have not yet been clearly identified. Oral

carcinogenesis is a multifactorial process involving numerous genetic events which is the root cause for the progression from a normal cell to a cancer cell, referred to as multi-step carcinogenesis. This event may be concomitant with deregulated control of cell proliferation and apoptosis.³ Thus proliferation is one of the most significant features exemplifying the malignant phenotype and aggressive biological feature.

The assessment of cell proliferation activity in tumors is an important tool for diagnosis, clinical behavior, and therapy. Numerous techniques have been established to identify and measure proliferating cells. The pioneer method was based on counting of mitotic figures using a light microscope. Other techniques include tritium labeled thymidine, the use of another pyrimidine analogue, the bromodeoxyuridine.⁴ Newer prognosticators like flow cytometry, autoradiography, DNA ploidy measurements are now at the leading edge.⁵ These techniques are

* Corresponding author at: Plot no. 21, Kukde layout, Wanjarinagar, Nagpur, Maharashtra, 440027, India.

E-mail address: priyanka82kadoo@gmail.com (P. Kadoo).

challenging, expensive, time consuming and often requiring sophisticated equipment, restraining their routine use by pathologists.⁴ Immunohistochemistry is another advanced technique in use. Ki67, is a reliable immunohistochemical (IHC) marker which shows a marked variation within different tumor grades, indicating that it has a crucial role in the aggressive behavior of malignant tumors and thus can be used for assessment of proliferation.^{6,7} Though it is practical compared to other methods of proliferation, it is quite expensive when compared to counting of mitotic figures. Thus cost and time factor makes it less feasible for many laboratories.

Counting of mitotic figures is the oldest, economical way of assessing proliferation. Increased number and abnormal mitosis indicate genetic damage. Thus identification and quantitation of mitotic cells forms an indivisible part of histological grading systems which is further valuable in the assessment of prognosis. However strict and precise morphologic criteria should be applied for recognition of mitotic figures.⁵ They can be identified and counted under conventional light microscope. However routine Hematoxylin and Eosin (H and E) stained section may pose a problem in distinguishing mitotic cells from apoptotic cells. This may further weaken the reliability of histological grading because of inappropriate use of morphologic criteria. Therefore, to obtain more reliable and reproducible count, appropriate histochemical techniques needs to be standardized which may include combination or modification of stains. Stains to visualize atypical and typical mitotic figures include H and E, Crystal violet, Giemsa stain and fluorescent microscopy.^{8,9} These stains have been used in tissues apart from oral tissue.^{10,11} Crystal violet is a basic dye having high affinity for the highly acidic chromatin of mitotic cells which are stained magenta and stand out clearly against a light blue background of resting cells.⁹ Literature has revealed very few studies which used these special stains for studying mitosis in oral dysplasia and oral carcinoma.^{9,12,13} However, these studies neither compared their results with established proliferative marker like Ki67 nor they compared mitosis in different grades of OSCC.

The aim of this study was to validate the usefulness of Crystal violet as a special stain for evaluating proliferation. This was done by correlating mitotic frequency index (MFI) obtained by it with that obtained by H and E stain as well as with Ki67 labeling index (Ki67LI), an established proliferative IHC marker, in various grades of oral squamous cell carcinoma.

2. Materials and methods

The study was carried out at the Department of Oral and Maxillofacial Pathology and Microbiology, MGV's KBH Dental College, Nashik, Maharashtra, India. The study protocol was approved by the Institutional Ethics Committee. 40 patients with lesions histologically diagnosed as oral squamous cell carcinoma were included in the study. After clinical examination and prior informed consent of the patient, biopsy was performed as a part of the treatment protocol and in order to obtain histopathological diagnosis of the lesion. Grading of the subjects was done according to Bryne (1991)¹⁴ using invasive front grading score (IFG Score). Accordingly patients were grouped as grade I OSCC (IFG score <5), grade II OSCC (IFG score 5–12) and grade III OSCC (IFG score >12). Table 1 summarizes clinical records of the patients studied. Three histological sections of each case were made. First two sections were stained with H and E stain and with 1% Crystal violet stain respectively. Mitotic figures were evaluated in both the sections.

Third section was stained with antibody against Ki67 nuclear antigen, and Ki67LI was assessed. Using binocular compound microscope, both H and E stained and Crystal violet stained sections were studied at 40 X. To assess inter-observer variation in interpretation, counting of mitotic figures was done by two

Table 1

Clinical And Pathological Data Of The Patients Studied.

			n	%
Age	Mean	48.25 years	40	
	Range	20 to 72 years		
Sex	Males	26	40	
	Females	14		
T	Mean	2.02 cm		
	Range	1 to 3		
N	Positive		31	77.5
	Negative		9	22.5
M	Positive		0	0
	Negative		40	100
Stage	I		4	10
	II		6	15
	III		20	50
	IV		10	25

T – tumor size.

N – node.

M – Metastasis.

examiner and no statistically difference was observed between the readings.

The areas of the tissue with most invasive and cellular part were selected whereas areas demonstrating tissue folds, necrosis, inflammation, and calcifications were excluded. Following of precise protocols are of great help in determining the accuracy of mitotic counting.¹⁵ Therefore they were recognized according to the criteria as given by Vandiest et al (1998).¹⁶ Mitosis was studied using MFI which was expressed as the percentage of mitotic nuclei.

MFI = Number of mitotic figure × 100 / Total number of cells observed.

Counting the 1000 tumor cells was simplified by placing 10 × 10 squared grid. Only clear nuclei within the grid squares and/or hit by the right or bottom line of the squares were counted in order to avoid double count.

For preparing crystal violet stain, solution A was prepared by mixing 2 grams of crystal violet (HiMedia Laboratories Pvt. Ltd. Crystal violet powder) and 20 ml of 95% ethyl alcohol. Solution B was prepared by mixing 0.8 gram of ammonium oxalate and 80 ml of distilled water. Both the solutions, A and B were mixed and kept for 24 hours. The resulting mixture was filtered and stored in an amber colored bottle. 1N HCl was prepared by adding 8.7 ml of 35%–37% HCl in 90 ml of distilled water. The sections were deparaffinized in hot air oven which was followed by two changes of xylene. Then the sections were hydrated through descending grades of absolute alcohol (90%, 80%, 70%, 60%, and 50%) for 5 minutes each. The sections were further washed well in distilled water for about 10 minutes and were hydrolyzed in 1 N HCl at 60 °C for 10 min (not an essential step). Thereafter the sections were washed well in distilled water for about 10 minutes and stained with 1% crystal violet solution for 30 min. Finally the sections were again washed in distilled water and were differentiated in 30% ethanol containing 1 to 2 drops of 35%–37% HCL, cleared in xylene and mounted in DPX. Known hyperplastic lymphnode was used as positive control whereas negative controls were not used.

For immunohistochemistry, the sections were deparaffinized with xylene and hydrated in graded alcohol. Antigen retrieval for Ki67 was performed by heating tissue sections in microwave oven in citrate buffer (pH 6.0). In order to inhibit internal peroxidase activity, the sections were incubated with 3% H2O2 in methanol for 30 minutes. Further they were incubated with 10% serum to prevent non-specific reactions and lastly with Ki67 antibody (Prediluted, Clone MIB-1; Product code: N1633, Dako, Denmark) for 90 min. After cleaning the sections in PBS, the HRP Labelled Polymer Antimouse (Dako EnVision + System HRP Labelled

Polymer Antimouse, Product Code: K4000, Dako North America Inc.), was used as secondary antibody. Thereafter, solution of 3, 3'Diaminobenzidine (DAB) in provided buffer was used for color development. Finally, the sections were counterstained in Mayer's hematoxylin. Known hyperplastic lymph node showing Ki67 was used as positive control and omission of the primary antibody was used as negative control.

IHC stained sections were evaluated using the light microscope where intranuclear brown color was considered as positive for Ki67 antigen. All stained nuclei were scored as positive regardless of intensity of staining whereas cells that lacked a clear nucleus were omitted. The suitable counting protocol was followed.¹⁷ Minimum 1000 tumor cells were counted at X40 magnification. Ki67LI was assessed by expressing it as a percentage of total number of tumor cells counted per section.

$Ki-67\ LI = \text{Number of IHC Positive Cells (Ki-67)} \times 100 / \text{Total number of cells observed.}$

3. Results

Finally all data was tabulated and the following statistical analysis was done.

1. MFI obtained by using 1% Crystal violet stain and H and E stain were compared using Maan Whitney U test
2. Correlation between Ki67LI and MFI obtained by using 1% Crystal violet stain was performed using Pearson correlation test.
3. Ki67LI as well as MFI obtained by using 1% Crystal violet stain was further correlated with histological grades using Pearson correlation test.

Out of 40 patients, 17 were diagnosed as grade I, 16 as grade II and 7 as grade III OSCC. Statistically significant (P- value: 0.008) increased MFI was found in Crystal violet stained sections compared to Hand E counterparts (Table 2, Graph 1).

Significant positive correlation was observed between Ki67LI and MFI obtained by using 1% Crystal violet stain in well and moderately differentiated OSCC. Positive correlation was also observed in poorly differentiated OSCC, but it was not statistically significant (Table 3).

The mean MFI significantly increased from grade I to grade II to grade III OSCC (Table 4). There was a significant positive correlation (p value- 0.006) between IFG score and MFI within three groups (Table 5, Graph 1).

Similarly the mean Ki67LI significantly increased from grade I to grade II to grade III OSCC (Table 6). Also, there was a significant positive correlation (p-value: 0.040) between IFG score and Ki67LI within three groups (Table 7, Graph 2).

4. Discussion

Cell proliferation is one of the most important biological mechanisms in oncogenesis.⁶ In fact it is a vital biological process, having potential significance as an indicator of treatment response and relapse. As reported by various studies, abnormal cell proliferation appears to be a precursor and may be a prognosticator

of tumorigenesis.⁷ Although many other advanced ways of assessing proliferation have become available, popularity of counting of mitotic figures is increasing because of the ease with which it can be identified without any special equipment except microscope and a well stained section.⁵ Since H and E stain may pose a problem in identification of mitotic figure, the present study compared mitotic count obtained by H and E with that of Crystal violet stain. Our study revealed an overall significantly increased mitosis in Crystal violet-stained sections (Table 2, Graph 1). It was easy to locate mitotic figures in Crystal violet stained sections even at low magnification. This suggests that Crystal violet stain provides a crisp staining simplifying the identification of mitotic figures compared to H and E stained counterparts (Fig. 1). Not only different phases of mitosis (Fig. 2) were easily and quickly appreciated in sections stained with Crystal violet stain compared to H and E stained section, but also it provided easy detection of mitotic abnormalities which occur naturally in tumors (Fig. 3). The mitotic abnormalities found in our study were further categorized according to Koller PC¹⁸ into following 3 types:

1. Structural alteration in chromosomes: stickiness of chromosomes (Fig. 3a)
2. Numerical changes in chromosome complement: abnormal distribution of chromosomes (Fig. 3b)
3. Complete or partial suppression of spindles: unipolar and tripolar (Fig. 3c)

Mitotic figures having combination of above three were also found in our study (Fig. 3d). This increased mitotic count with 1% Crystal violet stain was in accordance with previous studies.^{9,12,13} However they didn't compare their results with an established proliferative marker.

Ki67, the most extensively applied cell proliferation markers, is considered more reliable which not only reflects the activity of cell proliferation, but also quantitatively analyzes the growth fraction. Its expression is intimately associated with the differentiated degrees of the tumor. Thus it is of great importance in the tumor grading, and determining the proliferative degrees and prognosis.¹⁹ So we correlated MFI obtained by 1% crystal violet stain with Ki67LI (Table 3). Ki67 was expressed in all the 40 tumors examined (Fig. 4). Significant positive correlation was observed in well and moderately differentiated OSCC. These results were in accordance with previous studies conducted in tissues apart from oral cavity which showed a significant correlation between mitotic indices and Ki67 LI.^{20,21,22} Positive correlation was also observed in poorly differentiated OSCC, but it was not statistically significant. This discrepancy in poorly differentiated cases might be due to less sample size (n=7). The present study thus revealed that when mitotic figures were carefully counted and expressed as a percentage of tumor cells (MFI), the measure of mitotic count correlated intensely with Ki67 antigen labeling, which also was counted and expressed in the same fashion (Ki67LI).

We further correlated MFI as well as Ki67LI with tumor grade and found that the mean MFI obtained by using 1% crystal violet stain significantly increased from grade I to grade II to grade OSCC (Table 4, Table 5 and Graph 1). This was in accordance with the results of the earlier studies²³ which reported that strong correlation exists between total malignancy grading scores including mitosis based on several pathologic parameters and the prognosis in OSCC. Thus it can be concluded that mitotic activity is an important factor in grading system.

Further our study also revealed that there was a significant positive correlation between histological grade and Ki67LI within three groups. (Table 6, Table 7 and Graph 2) These results were in accordance with previous studies which reported that the expression of Ki67 increased progressively according to the grades

Table 2
Comparison of MFI in Crystal Violet and H and E.

MFI (Crystal Violet)	MFI (H and E)	U	p-value	Remark
1.24 ± 0.40	0.99 ± 0.38	2.64	0.008	Significant

*MFI – Mitotic frequency index.

Graph 1. Comparison of MFI in Crystal Violet stain and H and E stain.

*MFI – Mitotic frequency index.

Table 3

Correlation Of Ki67 LI With MFI using Crystal Violet stain in OSCC.

Grades of OSCC/IFG score	Ki67 LI	MFI (%) Crystal violet stain	r	p-value	Remark
Grade I OSCC (IFG score <5)	36 ± 10.0	1.02 ± 0.32	0.86	0.000	Significant
Grade II OSCC (IFG score 5-12)	45.6 ± 9.79	1.40 ± 0.36	0.8	0.000	Significant
Grade III OSCC (IFG score >12)	45.8 ± 15.7	1.42 ± 0.40	0.69	0.085	Nonsignificant

*MFI – Mitotic frequency index.

* OSCC – Oral Squamous cell Carcinoma.

* Ki67LI – Ki67 Labeling index.

*IFG score – invasive front grading score.

Table 4

Correlation of IFG score with MFI using crystal violet stain in different grades of OSCC:.

Grades of OSCC/IFG score	IFG grading	MFI (%)	r	p-value	Remark
Grade I OSCC (IFG score <5)	6.64 ± 1.16	1.12 ± 0.33	0.41	0.09	Significant
Grade II OSCC (IFG score 5-12)	10.31 ± 1.13	1.40 ± 0.36	0.95	0	Significant
Grade III OSCC (IFG score >12)	16 ± 2.16	1.45 ± 0.44	0.98	0	Significant

* OSCC – Oral Squamous cell Carcinoma.

*MFI – Mitotic frequency index.

*IFG score – invasive front grading score.

Table 5

Comparison of MFI using crystal violet stain in different grades of OSCC.

Grades of OSCC/IFG score	N	Mean	Std. Deviation	F	p-value	Remark
Grade I OSCC (IFG score <5)	17	1.02	0.32	5.79	0.006	Significant
Grade II OSCC (IFG score 5-12)	16	1.4	0.36			
Grade III OSCC (IFG score >12)	7	1.45	0.44			

* OSCC – Oral Squamous cell Carcinoma.

*MFI – Mitotic frequency index.

*IFG score – invasive front grading score.

Table 6

Correlation Of IFG score with Ki67LI in OSCC.

Grades of OSCC/IFG score	IFG Grading	Ki67 LI	γ	R ²	p-value	Remark
Grade I OSCC (IFG score <5)	6.64 ± 1.16	36 ± 10.94	0.78	0.61	0.000	Significant
Grade II OSCC (IFG score 5-12)	10.31 ± 1.13	45.65 ± 9.79	0.73	0.54	0.001	Significant
Grade III OSCC (IFG score >12)	16.28 ± 1.97	45.82 ± 15.73	0.78	0.61	0.038	Significant

* OSCC – Oral Squamous cell Carcinoma.

* Ki67LI – Ki67 Labeling index.

*IFG score – invasive front grading score.

Table 7

Comparison of Ki67 LI in different grades of OSCC.

Grades of OSCC/IFG score	N	Mean	Std. Deviation	F	p-value	Remark
Grade I OSCC (IFG score <5)	17	36	10.9	3.51	0.04	Significant
Grade II OSCC (IFG score 5-12)	16	45.6	9.79			
Grade III OSCC (IFG score >12)	7	45.8	15.7			

* OSCC – Oral Squamous cell Carcinoma.

* Ki67LI – Ki67 Labeling index.

*IFG score – invasive front grading score.

Graph 2. Comparison of Ki67 LI in different grades of OSCC.

*OSCC – Oral Squamous cell Carcinoma.

*Ki67LI – Ki67 Labeling index.

Fig. 1. H&E Staining (40X), (100X); Crystal Violet Staining (40X) (100X).

Fig. 2. Phases of Mitosis (H&E stain (100X) Crystal violet stain (100X)).

- a. PROMETAPHASE.
- b. METAPHASE.
- c. EARLY ANAPHASE.
- d. LATE ANAPHASE.
- e. TELOPHASE.

Fig. 3. Abnormal Mitosis – Crystal Violet stain (100X).

- a. Structural Alteration In Chromosomes: Stickiness Of Chromosomes.
- b. Numerical Changes In Chromosome Complement: Abnormal Distribution Of Chromosomes.
- c. Complete Or Partial Suppression Of Spindles: Unipolar And Tripolar Or Polypoid Spindles.
- d. Combination Of Sticky Chromosomes And Abnormal Distribution Of Chromosomes And Complete Or Partial Suppression Of Spindles.

of OSCC²⁴ and thus have a positive association with the histologic grading.²⁵

5. Conclusion

Our study has shown a significant positive correlation between Ki67LI and MFI. Since MFI is technically more convenient and

economically less expensive than Ki67LI, it remains a simple, reliable and apparently the most convenient indicator of proliferation which is one of the vital processes of carcinogenesis. Compared to H and E stained section we easily found more mitotic figures using 1% Crystal violet stain. Crystal violet also helps in the distinction between a pyknotic nucleus, an apoptotic cell and a mitotic cell compared to routinely used H and E stained section.

a. Well Differentiated Squamous Cell Carcinoma

b. Moderately Differentiated Squamous Cell Carcinoma.

c. Poorly Differentiated Squamous Cell Carcinoma

Fig. 4. Ki67 Labelled Cells (40X).

- a. Well Differentiated Squamous Cell Carcinoma.
b. Moderately Differentiated Squamous Cell Carcinoma.
c. Poorly Differentiated Squamous Cell Carcinoma.

Thus it can be concluded that the use of 1% Crystal violet as a selective stain for mitotic figures is validated. Hence, we recommend its routine use for assessing proliferation.

Future perspectives

The quantification of mitotic figures has taken a backseat over the decades especially in oral squamous cell carcinoma patients. Nonetheless, multicentric studies with large sample sizes are required to positively correlate mitosis with TNM staging, prognosis and survival rate in oral cancer patients.

Conflict of interest

We do not have any conflict of interest.

Acknowledgments

We wish to thank to the members of the NAMCO cancer institute, Nashik for their help in this work.

References

- Warnakulasuriya S. Review-Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol.* 2009;45:309–316.
- Siddiqui MS, Chandra R, Aziz A, Suman S. Epidemiology and histopathological spectrum of head and neck cancers in Bihar, a state of Eastern India. *Asian Pacific J Cancer Prev.* 2012;13:3949–3953.
- Fukuda M, Kusama K, Sakashita H. Review article molecular insights into the proliferation and progression mechanisms of the oral cancer: strategies for the effective and personalized therapy. *Jpn Dent Sci Rev.* 2012;48(1):23–41.
- De Moraes M, Monteiro Maia CAD, de Almeida Freitas R, Galvão HC. Cell proliferation markers in oral squamous cell carcinoma. *J Mol Biomark Diagn.* 2012;10.4172/2155-9929.S2-006 S2:006.
- Kapoor K, Puri A, Prakash A, Jazib Sharma G. Mitotic counting and its significance in histopathological grading of OSCC & oral epithelial dysplasia. *Head talk.* 2013;5(6):35–37.
- Tadbir AA, Pardis S, Ashkavandi ZJ, et al. Expression of Ki67 and CD105 as proliferation and angiogenesis markers in salivary gland tumors. *Asian Pacific J Cancer Prev.* 2012;13:5155–5159.
- Birajdar SS, Radhika MB, Paremala K, Sudhakara M, Saumya M, Gadivan M. Expression of Ki-67 in normal oral epithelium, leukoplakic oral epithelium and oral squamous cell carcinoma. *JOMFP.* 2014;18(2):169–176.
- Bavle RM. Mitosis at a glance. *J Oral Maxillofac Pathol.* 2014;18(Suppl. 1):S2–S5.
- Ankle M, Kale A, Charantin S. Comparison of staining of mitotic figures by haematoxylin and eosin and crystal violet stains, in oral epithelial dysplasia and squamous cell carcinoma. *Indian J Dent Res.* 2007;18(3):101–105.
- Gander DW, Meyer J. A simple two dye basic stain facilitating recognition of mitosis of plastic embedded tissue sections. *Stain Technol.* 1988;63:75–81.
- Fraser FJ. Selective stain for mitotic figures, particularly in the developing brain. *Stain Technol.* 1982;57(4):219–224.
- Jadhav KB, Ahmed Mujib BR, Gupta N. Crystal violet stain as a selective stain for the assessment of mitotic figures in oral epithelial dysplasia and oral squamous cell carcinoma? *Indian J Pathol Microbiol.* 2012;55(3):283–287.
- Palaskar SJ, Patil S, Narang B, et al. Efficacy of various stains to study mitotic figures in oral epithelial dysplasia – a pilot study? *J Dent Allied Sci.* 2013;2(2):46–48.
- Bryne M. Prognostic value of various molecular and cellular features in oral squamous cell carcinomas: a review. *J Oral Pathol Med.* 1991;20:413–420.
- Yadav KS, Gonuguntla S, Reddy Ealla KK, et al. Assessment of interobserver variability in mitotic figure counting in different histological grades of oral squamous cell carcinoma. *J Contemp Dent Pract.* 2012;13(3):339–344.
- Van Diest PJ, Brugal G, Baak JPA. Proliferation markers in tumours: interpretation and clinical value. *J Clin Pathol.* 1998;51:716–724.
- Kurokawa H, Zhang M, Matsumoto S, et al. The relationship of histological grade at the deep invasive front and the expression of Ki67 antigen and p53 protein in oral squamous cell carcinoma. *J Oral Pathol Med.* 2005;34:602–607.
- Koller PC. Abnormal mitosis in tumours. *Br J Cancer.* 1947;1(1):38–47.
- Yu JQ, Zhou Q, Zheng YF, Bao Y. Expression of vimentin and Ki-67 proteins in cervical squamous cell carcinoma and their relationships with clinicopathological features. *Asian Pac J Cancer Prev.* 2015;16(10):4271–4275.
- Yip CH, Bhoopathy N, Daniel JM, Mohamed AK, Abdullah MM, Ng YS, Yap BK, Pathmanathan R. Roles of Ki67 in breast cancer – important for management. *Asian Pac J Cancer Prev.* 2016;17(3):1077–1082.
- Kamoi S, Ohaki Y, Amamo Y, et al. Pre-treatment mitotic index versus computer-quantitated Ki-67 nuclear antigen labeling index as predictors of response to neoadjuvant chemotherapy in uterine cervical carcinoma. *J Nippon Med Sch.* 2003;70(3):219–226.
- Goyal S, Singh UR, Sharma S, Kaur N. Correlation of mitotic indices, AgNor count, Ki-67 and Bcl-2 with grade and stage in papillary urothelial bladder cancer. *Urol J.* 2012;11(1):1238–1247.
- Bryne M, Koppan HS, Lilling R, Kjaerhei A. Malignancy grading of the deep invasive margins of oral squamous cell carcinomas has high prognostic value. *J Pathol.* 1992;166(4):375–381.
- Dwivedi N, Chandra S, Kashyap B, Raj V, Agrawal A. Suprabasal expression of Ki67 as a marker for the severity of epithelial dysplasia and oral squamous cell carcinoma. *Contemp. Clin Dent.* 2013;4(1):7–12.
- Premalatha BR, Uma K. Analysis of Ki-67 antigen in human oral squamous cell carcinoma – an immunohistochemical study. *J Int Oral Health.* 2010;2(1):9–16.